

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre <u>West Siang, Basar – 791 101</u>

(<u>Changlang</u>)



Dated: 19th April, 2013

<u>No: -4/2013-14/Tue</u>

Weather during the period

- <u>Rainfall</u>: -
 - Changlang NA
 - Miao– NA

Weather Forecast

- <u>Rainfall</u> Clear/partly cloudy weather for 2/3 days are forecasted.
- <u>Temperature</u> The temperature is likely to be in the range of 17-30 °C.

• <u>Relative Humidity</u> - The RH is likely to be in the range of 65-88 %.

Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
	U		VEGETABLE CROPS
Cucurbits	Seedling/ Vegetative	Damping off	 Avoid water stagnation around the pit of the plant, Light interculture operations is also advised, Provide proper support for healthy trailing.
French Bean	Vegetative/ Flowering		 Avoid water stagnation in the crop field, Light interculture operations is required,
Ladies Finger	Vegetative		Avoid water stagnation in the crop field,Light interculture operations,
		1	FIELD CROPS
Maize	Vegetative		• Avoid water stagnation in the crop field,
Soybean	Vegetative		• Avoid water stagnation in the crop field,
Black gram	Vegetative		 In present weather conditions reports of collar rot/wilting/damping off may occur. Spraying of Carbandazim 2gms per litre of water or Endofil M45 @ 3 gms per litre of water advised, Treat the seeds before sowing. Avoid water stagnation in the crop field,
x (.)			FRUIT CROPS
Lemon/other citrus fruits	Planting Fruiting		 Planting of new sapling may be done, Apply first dose of NPK fertilizers. Spray micronutrients. Apply any contact insecticide to ward off sucking insect pests,

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department

Contact: - <u>IAAS.BASAR@gmail.com</u>, Phone Number: - 03795-226247/37, -96 (Fax)



ICAR Research Complex for NEH Region, Arunachal Pradesh Centre <u>West Siang, Basar – 791 101</u>

(Dibang Valley)



Dated: 19th April, 2013

<u>No: -4/2013-14/Tue</u>

Weather during the period

- <u>Rainfall</u>: -
 - <u>Roing</u> NA
 - <u>Anini</u> 99.0 mm

Weather Forecast

- <u>Rainfall</u> Cloudy weather with light rains in few places are forecasted.
- <u>Temperature</u> The temperature is likely to be in the range of 12-26 °C.
- <u>Relative Humidity</u> The RH is likely to be in the range of 60-90 %.

Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
			VEGETABLE CROPS
Cucurbits	Sowing/ Nursery/ Vegetative	Damping off	 Avoid water stagnation around the pit of the plant, Light interculture operations is also advised,
French Bean	Vegetative/ Flowering		 Avoid water stagnation in the crop field, Light interculture operations is required,
Ladies Finger	Vegetative		 Avoid water stagnation in the crop field, Light interculture operations, FIELD CROPS
Jhum Rice	Sowing/ Four leaf stage	Yellow stem borer	 Make structures to avoid runoff in the slope Install yellow stem borer pheromone traps to trap the adults
Maize	Vegetative		• Avoid water stagnation in the crop field,
Soybean	Vegetative		• Avoid water stagnation in the crop field,
Black gram	Vegetative		 In present weather conditions reports of collar rot/wilting/damping off may occur. Spraying of Carbandazim 2gms per litre of water or Endofil M45 @ 3 gms per litre of water advised, Treat the seeds before sowing. Avoid water stagnation in the crop field,
Green gram	Vegetative	Damping off Wilting Collar Rot	 In present weather conditions reports of collar rot/wilting/damping off may occur. Spraying of Carbandazim 2gms per litre of water or Endofil M45 @ 3 gms per litre of water advised, Treat the seeds before sowing. Avoid water stagnation in the crop field, FRUIT CROPS
Khasi Mandarin	Fruiting		 Planting of new sapling may be done, Apply first dose of NPK fertilizers. Spray micronutrients. Apply any contact insecticide to ward off sucking insect pests,
Pineapple /Strawberry	Planting		• Treat the plantlets with DithaneM45 @ 2 grams per litre of water for half an hour before planting,
Apple	Fruiting		Mulching is advised
Kiwi	Flowering		 Mulching is advised, Fertilizer application,
Banana	Planting		• Planting of new saplings may be done in present weather,
	I	1	LIVESTOCKS

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department Contact: - <u>IAAS.BASAR@gmail.com</u>, Phone Number: - 03795-226247/37, -96 (Fax)



ICAR Research Complex for NEH Region, Arunachal Pradesh Centre West Siang, Basar – 791 101



(East Kameng)

<u>No:-4/2013-14/Tue</u>

Weather during the period

Dated: 19th April, 2013

<u>Rainfall</u>: -

• Seppa – 0.0 mm

Weather Forecast

- <u>Rainfall</u> Cloudy/partly cloudy weather with light rain in few places is forecasted for next 2/3 days.
 - <u>Temperature</u> The temperature is likely to be in the range of 17-29°C.
- <u>Relative Humidity</u> The RH is likely to be in the range of 79-91 %.

Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
			VEGETABLE CROPS
Cucurbits	Seedling/ Vegetative	Damping off	 Avoid water stagnation around the pit of the plant, Light interculture operations is also advised, Provide proper support for healthy trailing.
French Bean	Vegetative/ Flowering		 Avoid water stagnation in the crop field, Light interculture operations is required,
Ladies Finger	Vegetative		Avoid water stagnation in the crop field,Light interculture operations,
		T	FIELD CROPS
Jhum Rice	Sowing/ Four leaf stage	Yellow stem borer	 Make structures to avoid runoff in the slope Install yellow stem borer pheromone traps to trap the adults,
Maize	Vegetative		 Avoid water stagnation in the crop field,
Soybean	Vegetative		• Avoid water stagnation in the crop field,
Black gram	Vegetative		 In present weather conditions reports of collar rot/wilting/damping off may occur. Spraying of Carbandazim 2gms per litre of water or Endofil M45 @ 3 gms per litre of water advised, Treat the seeds before sowing. Avoid water stagnation in the crop field,
Green gram	Vegetative	Damping off Wilting Collar Rot	 In present weather conditions reports of collar rot/wilting/damping off may occur. Spraying of Carbandazim 2gms per litre of water or Endofil M45 @ 3 gms per litre of water advised, Treat the seeds before sowing. Avoid water stagnation in the crop field, FRUIT CROPS
	Dlanting		
Khasi Mandarin	Planting Fruiting		 Planting of new sapling may be done, Apply first dose of NPK fertilizers. Spray micronutrients. Apply any contact insecticide at recommended dose to ward off sucking insect pests,
Banana	Planting		• Planting of new saplings may be done in present weather,
			LIVESTOCKS

Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795226247/37

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department Contact: - <u>IAAS.BASAR@gmail.com</u>, Phone Number: - 03795-226247/37, -96 (Fax)

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre <u>West Siang, Basar – 791 101</u>

(East Siang)

<u>No:-4/2013-14/Tue</u>

Dated: 19th April, 2013

Weather during the period

- Maximum temperature: 29.4°C
- Minimum Temperature: 17.9°C
- <u>Rainfall</u>:
 - <u>Pasighat</u> 30.6 mm
- <u>Relative humidity (%)</u>:69-98

Weather Forecast

• <u>Rainfall</u> - Cloudy weather are forecasted for next three/four days. Light rains may occur at few places.

• <u>Temperature</u> - The temperature is likely to be in the range of 18-29 °C.

• <u>Relative Humidity</u> - The RH is likely to be in the range of 67-83 %.

Weather Based Agro-meteorological Advisories Bulletin

Сгор	Stage	Pest/Disease	Precautionary Practices to be followed
		V	EGETABLE CROPS
Cucurbits	Seedling/	Damping off	• Avoid water stagnation around the pit of the plant,
	Vegetative		• Light interculture operations is also advised,
			• Provide proper support for healthy trailing.
French Bean	Vegetative/		 Avoid water stagnation in the crop field,
	Flowering		Light interculture operations is required,
Ladies Finger	Vegetative		 Avoid water stagnation in the crop field,
			Light interculture operations,
			FIELD CROPS
Jhum Rice	Sowing/	Yellow stem	 Make structures to avoid runoff in the slope
	Four leaf	borer	• Install yellow stem borer pheromone traps to trap the
	stage		adults,
Maize	Vegetative		 Avoid water stagnation in the crop field,
Soybean	Vegetative		• Avoid water stagnation in the crop field,
Black gram	Vegetative		 In present weather conditions reports of collar rot/wilting/damping off may occur. Spraying of Carbandazim 2gms per litre of water or Endofil M45 @ 3 gms per litre of water advised, Treat the seeds before sowing. Avoid water stagnation in the crop field,
Green gram	Vegetative	Damping off Wilting Collar Rot	 In present weather conditions reports of collar rot/wilting/damping off may occur. Spraying of Carbandazim 2gms per litre of water or Endofil M45 @ 3 gms per litre of water advised, Treat the seeds before sowing. Avoid water stagnation in the crop field,
			FRUIT CROPS
Khasi Mandarin	Planting Fruiting		 Planting of new sapling may be done, Apply first dose of NPK fertilizers. Spray micronutrients. Apply any contact insecticide at recommended dose to ward off sucking insect pests,
Pineapple /Strawberry	Planting		• Treat the plantlets with DithaneM45 @ 2 grams per litre of water for half an hour before planting,
Banana	Planting		• Planting of new saplings may be done in present weathe
(LIVESTOCKS

or call at 03795226247/37

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department Contact: - <u>IAAS.BASAR@gmail.com</u>, Phone Number: - 03795-226247/37, -96 (Fax)







ICAR Research Complex for NEH Region, Arunachal Pradesh Centre <u>West Siang, Basar – 791 101</u>

(<u>Lohit</u>)



<u>No:-4/2013-14/Tue</u>

Weather during the period

Dated: 19th April, 2013

- <u>Rainfall</u>: -
 - Tezu –85.2 mm
 - Namsai –NA

Weather Forecast

- <u>Rainfall</u> Clear/partly cloudy weather is forecasted for next three/four days.
- <u>Temperature</u> The temperature is likely to be in the range of 17-30 °C.
- <u>Relative Humidity</u> The RH is likely to be in the range of 61-74 %.

Weather Based Agro-meteorological Advisories Bulletin

Light in	terculture op	erations is adv	ised,
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
			VEGETABLE CROPS
Cucurbits	Seedling/ Vegetative	Damping off	 Avoid water stagnation around the pit of the plant, Light interculture operations is also advised, Provide proper support for healthy trailing.
French Bean	Vegetative/ Flowering		Avoid water stagnation in the crop field,Light interculture operations is required,
Ladies Finger	Vegetative		 Avoid water stagnation in the crop field, Light interculture operations,
Maize	Vegetative		 FIELD CROPS Avoid water stagnation in the crop field,
Soybean	Vegetative		• Avoid water stagnation in the crop field,
Black gram	Vegetative		 In present weather conditions reports of collar rot/wilting/damping off may occur. Spraying of Carbandazim 2gms per litre of water or Endofil M45 @ 3 gms per litre of water advised, Treat the seeds before sowing. Avoid water stagnation in the crop field,
			FRUIT CROPS
Lemon/other citrus fruits	Planting Fruiting		 Planting of new sapling may be done, Apply first dose of NPK fertilizers. Spray micronutrients. Apply any contact insecticide to ward off sucking insect pests,
	ientists (ICAR 95226247/37	,	Staff or KVK staff for measures against adverse conditions or

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department Contact: - <u>IAAS.BASAR@gmail.com</u>, Phone Number: - 03795-226247/37, -96 (Fax).



ICAR Research Complex for NEH Region, Arunachal Pradesh Centre <u>West Siang, Basar – 791 101</u>

(Lower Subansiri)

<u>No:-4/2013-14/Tue</u>

Dated: 19th April, 2013

Weather during the period

- <u>Rainfall</u>: -
 - 1. Ziro –NA

Weather Forecast

- <u>Rainfall</u> Cloudy weather with rains is forecasted for next three/four days.
- <u>Temperature</u> The temperature is likely to be in the range of 12-22 °C.
- <u>Relative Humidity</u> The RH is likely to be in the range of 81-94 %.

Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
			VEGETABLE CROPS
Cucurbits	Seedling/ Vegetative	Damping off	 Avoid water stagnation around the pit of the plant, Light interculture operations is also advised, Provide proper support for healthy trailing.
French Bean	Vegetative/ Flowering		 Avoid water stagnation in the crop field, Light interculture operations is required, FIELD CROPS
Jhum Rice	Sowing/ Four leaf stage	Yellow stem borer	 Make structures to avoid runoff in the slope Install yellow stem borer pheromone traps to trap the adults,
Maize	Vegetative		• Avoid water stagnation in the crop field,
Soybean	Vegetative		Avoid water stagnation in the crop field,
		1	FRUIT CROPS
Khasi Mandarin	Planting Fruiting		 Planting of new sapling may be done, Apply first dose of NPK fertilizers. Spray micronutrients. Apply any contact insecticide at recommended dose to ward off sucking insect pests,
Apple	Fruiting		Mulching is advised
Kiwi	Flowering		 Mulching is advised, Fertilizer application,
Banana	Planting		• Planting of new saplings may be done in present weather,
		1	LIVESTOCKS

Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795226247/37

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department Contact: - <u>IAAS.BASAR@gmail.com</u>, Phone Number: - 03795-226247/37,-96 (Fax)





ICAR Research Complex for NEH Region, Arunachal Pradesh Centre <u>West Siang, Basar – 791 101</u>

(<u>Papum Pare</u>)



No:-4/2013-14/Tue

Weather during the period

Dated: 19th April, 2013

- Maximum temperature: 28.6°C
- Minimum Temperature: 18.5°C
- Relative Humidity (%):66-96
- <u>Rainfall</u>: -
 - 1. Itanagar NA
 - 2. Naharlagun 50.9 mm

Weather Forecast

- <u>Rainfall</u> Cloudy weather is forecasted for next three/four days. Light rain may occur at few places.
- <u>Temperature</u> The temperature is likely to be in the range of 18-30 °C.
- <u>Relative Humidity</u> The RH is likely to be in the range of 76-86%.

Weather Based Agro-meteorological Advisories Bulletin

Сгор	Stage	Pest/Disease	Precautionary Practices to be followed
			VEGETABLE CROPS
Cucurbits	Seedling/	Damping off	• Avoid water stagnation around the pit of the plant,
	Vegetative		• Light interculture operations is also advised,
			Provide proper support for healthy trailing.
French Bean	Vegetative/		 Avoid water stagnation in the crop field,
	Flowering		Light interculture operations is required,
Ladies Finger	Vegetative		• Avoid water stagnation in the crop field,
			Light interculture operations,
r		1	FIELD CROPS
Jhum Rice	Sowing/	Yellow stem	 Make structures to avoid runoff in the slope
	Four leaf stage	borer	Install yellow stem borer pheromone traps to trap the adults,
Maize	Vegetative		 Avoid water stagnation in the crop field,
Soybean	Vegetative		Avoid water stagnation in the crop field,
Black gram	Vegetative		 In present weather conditions reports of collar rot/wilting/damping off may occur. Spraying of Carbendazim 2gms per litre of water or Endofil M45 @ 3 gms per litre of water advised, Treat the seeds before sowing. Avoid water stagnation in the crop field,
Green gram	Vegetative	Damping off Wilting Collar Rot	 In present weather conditions reports of collar rot/wilting/damping off may occur. Spraying of Carbendazim 2gms per litre of water or Endofil M45 @ 3 gms per litre of water advised, Treat the seeds before sowing. Avoid water stagnation in the crop field, FRUIT CROPS
	Dlanting		
Khasi Mandarin	Planting Fruiting		 Planting of new sapling may be done, Apply first dose of NPK fertilizers. Spray micronutrients. Apply any contact insecticide at recommended dose to ward off sucking insect pests,
Pineapple /Strawberry	Planting		• Treat the plantlets with DithaneM45 @ 2 grams per litre of water for half an hour before planting,
Kiwi	Flowering		Mulching is advised,Fertilizer application,
Banana	Planting		Planting of new saplings may be done in present weather,
		I	LIVESTOCKS

Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795226247/37

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department Contact: - <u>IAAS.BASAR@gmail.com</u>, Phone Number: - 03795-226247/37, -96 (Fax)



ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

<u>West Siang, Basar – 791 101</u>



(Tawang)

<u>No: -4/2013-14/Tue</u>

Weather during the period

Dated: 19th April, 2013

<u>Rainfall</u>: 1. Tawang – 90.0 mm

Weather Forecast

- <u>Rainfall</u> Cloudy weather is forecasted for next 3/4 days.
- <u>Temperature</u> The temperature is likely to be in the range of 4-15 °C.
- <u>Relative Humidity</u> The RH is likely to be in the range of 81-92 %.

Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
Apple	Fruiting		• Avoid water stagnation in the basin
			• Apply manures and fertilizers in the crop basin,
Kiwi	Flowering		<u>Fertilizer application</u>
			Basin cleaning
			Basin cleaning LIVESTOCKS

Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795222247 (KVK, West Siang)

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department Contact: - <u>IAAS.BASAR@gmail.com</u>, Phone Number: - 03795-226247/37, -96 (Fax)



ICAR Research Complex for NEH Region, Arunachal Pradesh Centre West Siang, Basar - 791 101

(Tirap)



No: -4/2013-14/Tue

Weather during the period

Dated: 19th April, 2013

- Rainfall: -
 - <u>Khonsa</u>–NA
 - <u>Deomali</u> 75.2 mm

Weather Forecast

- <u>Rainfall</u> Clear /partial cloudy weather is forecasted. •
- <u>Temperature</u> The temperature is likely to be in the range of 16-25 °C. •
- <u>Relative Humidity</u> The RH is likely to be in the range of 66-79 %. •

Weather Based Agro-meteorological Advisories Bulletin

• Light in	terculture or	perations is adv	ised,
Сгор	Stage	Pest/Disease	Precautionary Practices to be followed
			VEGETABLE CROPS
Cucurbits	Seedling/ Vegetative	Damping off	 Avoid water stagnation around the pit of the plant, Light interculture operations is also advised, Provide proper support for healthy trailing.
French Bean	Vegetative/ Flowering		 Avoid water stagnation in the crop field, Light interculture operations is required,
Ladies Finger	Vegetative		 Avoid water stagnation in the crop field, Light interculture operations,
			FIELD CROPS
Maize	Vegetative		Avoid water stagnation in the crop field,
Soybean	Vegetative		Avoid water stagnation in the crop field,
Black gram	Vegetative		 In present weather conditions reports of collar rot/wilting/damping off may occur. Spraying of Carbandazim 2gms per litre of water or Endofil M45 @ 3 gms per litre of water advised, Treat the seeds before sowing. Avoid water stagnation in the crop field,
			FRUIT CROPS
Lemon/other citrus fruits	Planting Fruiting		 Planting of new sapling may be done, Apply first dose of NPK fertilizers. Spray micronutrients. Apply any contact insecticide to ward off sucking insect pests,

call at 03795226247/37

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)



ICAR Research Complex for NEH Region, Arunachal Pradesh Centre





(Upper Siang)

<u>No: -4/2013-14/Tue</u>

Weather during the period

Dated: 19th April, 2013

• <u>Rainfall</u>: -

• Tuting – 31.0 mm

Weather Forecast

- <u>Rainfall</u> Partially cloudy/cloudy weather with light rain is forecasted.
- <u>Temperature</u> The temperature is likely to be in the range of 12-22 °C.
- <u>Relative Humidity</u> The RH is likely to be in the range of 72-92 %.

Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
			VEGETABLE CROPS
Cucurbits	Seedling/ Vegetative	Damping off	 Avoid water stagnation around the pit of the plant, Light interculture operations is also advised, Provide proper support for healthy trailing.
French Bean	Vegetative/ Flowering		 Avoid water stagnation in the crop field, Light interculture operations is required,
Ladies Finger	Vegetative		Avoid water stagnation in the crop field,Light interculture operations,
			FIELD CROPS
Jhum Rice	Sowing/ Four leaf stage	Yellow stem borer	 Make structures to avoid runoff in the slope Install yellow stem borer pheromone traps to trap the adults,
Maize	Vegetative		• Avoid water stagnation in the crop field,
Soybean	Vegetative		Avoid water stagnation in the crop field,
Black gram	Vegetative		 In present weather conditions reports of collar rot/wilting/damping off may occur. Spraying of Carbandazim 2gms per litre of water or Endofil M45 @ 3 gms per litre of water advised, Treat the seeds before sowing. Avoid water stagnation in the crop field,
Green gram	Vegetative	Damping off Wilting Collar Rot	 In present weather conditions reports of collar rot/wilting/damping off may occur. Spraying of Carbandazim 2gms per litre of water or Endofil M45 @ 3 gms per litre of water advised, Treat the seeds before sowing. Avoid water stagnation in the crop field,
T		T	FRUIT CROPS
Khasi Mandarin	Planting Fruiting		 Planting of new sapling may be done, Apply first dose of NPK fertilizers. Spray micronutrients. Apply any contact insecticide at recommended dose to ward off sucking insect pests,
Pineapple /Strawberry	Planting		• Treat the plantlets with DithaneM45 @ 2 grams per litre of water for half an hour before planting,
Kiwi	Flowering		Mulching is advised,Fertilizer application,
Banana	Planting		• Planting of new saplings may be done in present weather,
1			LIVESTOCKS

Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795226247/37

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department Contact: - <u>IAAS.BASAR@gmail.com</u>, Phone Number: - 03795-226247/37, -96 (Fax)



Integrated Agro-Meteorological Advisory Services ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

<u>West Siang, Basar – 791 101</u>



(<u>Upper Subansiri</u>)

<u>No: -4/2013-14/Tue</u>

Dated: 19th April, 2013

Weather during the period

- <u>Rainfall</u>: -
 - 1. Daporijo- NA

Weather Forecast

- <u>Rainfall</u> Cloudy weather with light/moderate rain is forecasted for next three/four days.
- <u>Temperature</u> The temperature is likely to be in the range of 14-24°C.
- <u>Relative Humidity</u> The RH is likely to be in the range of 81-91 %.

Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
			VEGETABLE CROPS
Cucurbits	Seedling/ Vegetative	Damping off	 Avoid water stagnation around the pit of the plant, Light interculture operations is also advised, Provide proper support for healthy trailing.
French Bean	Vegetative/ Flowering		 Avoid water stagnation in the crop field, Light interculture operations is required,
Ladies Finger	Vegetative		Avoid water stagnation in the crop field,Light interculture operations,
			FIELD CROPS
Jhum Rice	Sowing/ Four leaf stage	Yellow stem borer	 Make structures to avoid runoff in the slope Install yellow stem borer pheromone traps to trap the adults,
Maize	Vegetative		Treat the seeds before sowing.Avoid water stagnation in the crop field,
Soybean	Vegetative		Treat the seeds before sowing.Avoid water stagnation in the crop field,
Black gram	Vegetative		 In present weather conditions reports of collar rot/wilting/damping off may occur. Spraying of Carbandazim 2gms per litre of water or Endofil M45 @ 3 gms per litre of water advised, Treat the seeds before sowing. Avoid water stagnation in the crop field,
Green gram	Vegetative	Damping off Wilting Collar Rot	 In present weather conditions reports of collar rot/wilting/damping off may occur. Spraying of Carbandazim 2gms per litre of water or Endofil M45 @ 3 gms per litre of water advised, Treat the seeds before sowing. Avoid water stagnation in the crop field,
		I	FRUIT CROPS
Khasi Mandarin	Planting Fruiting		 Planting of new sapling may be done, Apply first dose of NPK fertilizers. Spray micronutrients. Apply any contact insecticide at recommended dose to ward off sucking insect pests,
Pineapple /Strawberry	Planting		• Treat the plantlets with DithaneM45 @ 2 grams per litre of water for half an hour before planting,
Banana	Planting		• Planting of new saplings may be done in present weather,
			LIVESTOCKS

Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795226247/37

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department Contact: - <u>IAAS.BASAR@gmail.com</u>, Phone Number: - 03795-226247/37, -96 (Fax)



ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar - 791 101

(West Kameng)



<u>No: -4/2013-14/Tue</u>

Weather during the period

Dated: 19th April, 2013

• <u>Rainfall</u>: - Bomdila – 2.0 mm Bhalokpung –NA

Weather Forecast

- <u>Rainfall</u>-Cloudy weather with light rains in few places is forecasted for next three/four days.
- <u>Temperature</u> The temperature is likely to be in the range of 17-29°C.
- <u>Relative Humidity</u> The RH is likely to be in the range of 79-91 %.

Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
Apple	Fruiting		• Avoid water stagnation in the basin
			• Apply manures and fertilizers in the crop basin,
Kiwi	Flowering		<u>Fertilizer application</u>
			Basin cleaning
			LIVESTOCKS

Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795222247 (KVK, West Siang)

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department Contact: - <u>IAAS.BASAR@gmail.com</u>, Phone Number: - 03795-226247/37, -96 (Fax)



ICAR Research Complex for NEH Region, Arunachal Pradesh Centre <u>West Siang, Basar – 791 101</u>

(West Siang)

<u>No: -4/2013-14/Tue</u>

Weather during the period

Dated: 19th April, 2013

- During the period from 13th -16th April, weather was cloudy and rains were recorded.
- Maximum temperature: 22.0°C
- Minimum Temperature:15.0 °C
- Rainfall: 26.0 mm
- Evapotranspiration: Not Available
- Relative Humidity (%):- 82-99%
- Soil moisture (m²/ m²):-0.40
- Prominent Wind Direction: South West
- Average Wind Speed : 0.1kmph
- Average Sunshine hours : Not Available

(Source-Agromet Observatory, Research Farm, Gori, ICAR, AP Centre, Basar)

Weather Forecast

- <u>Rainfall</u> Partly cloudy/cloudy weather for next two/three days with rains in few places are forecasted.
- <u>Temperature</u> The temperature is likely to be in the range of 16-26 °C.
- <u>Relative Humidity</u> The RH is likely to be in the range of 70-87 %.

Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
	U		VEGETABLE CROPS
Cucurbits	Seedling/	Damping off	• Avoid water stagnation around the pit of the plant,
	Vegetative		• Light interculture operations is also advised,
			• Provide proper support for healthy trailing.
French Bean	Vegetative/		• Avoid water stagnation in the crop field,
	Flowering		Light interculture operations is required,
Ladies Finger	Vegetative		• Avoid water stagnation in the crop field,
			Light interculture operations,
			FIELD CROPS
Jhum Rice	Sowing/	Yellow stem	 Make structures to avoid runoff in the slope
	Four leaf	borer	• Install yellow stem borer pheromone traps to trap the adults,
	stage		
Maize	Vegetative		• Avoid water stagnation in the crop field,
Soybean	Vegetative		• Avoid water stagnation in the crop field,
Black gram	Vegetative		In present weather conditions reports of collar
			rot/wilting/damping off may occur. Spraying of
			Carbandazim 2gms per litre of water or Endofil M45 @ 3
			gms per litre of water advised,
			Treat the seeds before sowing.
			Avoid water stagnation in the crop field,
Green gram	Vegetative	Damping off	In present weather conditions reports of collar
		Wilting	rot/wilting/damping off may occur. Spraying of
		Collar Rot	Carbandazim 2gms per litre of water or Endofil M45 @ 3
			gms per litre of water advised,
			 Treat the seeds before sowing. Avaid water stagnation in the gran field
			Avoid water stagnation in the crop field, FRUIT CROPS
Khasi	Planting		 Planting of new sapling may be done,
Mandarin	Fruiting		 Apply first dose of NPK fertilizers. Spray micronutrients.
			 Apply any contact insecticide at recommended dose to ward
			off sucking insect pests,
Pineapple	Planting		 Treat the plantlets with DithaneM45 @ 2 grams per litre of
/Strawberry	0		water for half an hour before planting,

Fruiting	Mulching is advised
Flowering	Mulching is advised,
	Fertilizer application,
Planting	• Planting of new saplings may be done in present weather,
	LIVESTOCKS
	Flowering

Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795226247/37

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department Contact: - <u>IAAS.BASAR@gmail.com</u>, Phone Number: - 03795-226247/37, -96 (Fax)