



Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar – 791 101



(East Siang)

No:-60/2013-14/Fri

Dated: 29th November, 2013

Weather during the period

- Maximum temperature: NA
- Minimum Temperature: NA
- Rainfall:
Pasighat – NA
- Relative humidity (%): NA

Weather Forecast

- Rainfall – Clear/sunny/dry weather fcsted for next 3-4 days with shallow morning fog.
- Temperature – Temperature will vary between 15-28° C
- Relative Humidity - The RH is likely to be in the range of 38-91 %.

Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
VEGETABLE CROPS			
Cauliflower/ Cabbage/ Broccoli/ Knolkhol	Vegetative/ Maturing/	Leaf eating/ Cabbage caterpillars	<ul style="list-style-type: none">• Light irrigation is essential,• In present weather conditions severe attack of leaf eating caterpillars has been reported. Spray <i>Chlorpyrifos/Imidacloprid</i> in recommended dose in the next one or two days when weather is clear. Dose- Imidacloprid 17SL @ 1-2 ml per litre depending upon the population,
Radish	Vegetative Harvesting	Leaf eating caterpillars	<ul style="list-style-type: none">• Light interculture operations may be done,• Spray Chlorpyrifos/Imidacloprid in recommended dose.
Carrot	Vegetative		<ul style="list-style-type: none">• Light irrigation is essential,• Light interculture operations should be done,
Spinach	Vegetative Harvesting		<ul style="list-style-type: none">• Sowing of certified/locally available seeds in well prepared nursery may be done,
FIELD CROPS			
WRC rice – Main crop (Mipun)	Harvesting/ Threshing		<ul style="list-style-type: none">• Drying of grains/seed to appropriate level of moisture,• Seeds/grains should be stored in dry place and if available use some <i>neem</i> leaves,
Potato	Vegetative		<ul style="list-style-type: none">• Light irrigation followed by earthing-up,• Monitor the crop against attack of Cutworm,
Turmeric and ginger	Harvesting		<ul style="list-style-type: none">• Proper cleaning of the rhizomes before storing,
Mustard	Flowering		<ul style="list-style-type: none">• Need based irrigation for those in flowering stage• Thinning may be done to maintain proper plant count,
Winter Maize	Vegetative	Cut worm	<ul style="list-style-type: none">• Monitor for attack of cut worm in the germinated fields,
Pea	Vegetative		<ul style="list-style-type: none">• Manual collection of larvae during the evening hours may be done, (If damage level is not serious). Use of Carbaryl dust in severely damaged fields may be done,
FRUIT CROPS			
Khasi Mandarin	Maturing Harvesting	Trunk borer/ Fruit fly	<ul style="list-style-type: none">• In present weather conditions regularly collect and dispose off fallen fruits,• Farmers are advised to use <i>Jaggary</i>/Malathion traps or pheromone traps to monitor the pest population and collect and destroy the adult insects. Installation of pheromone traps may also be done,
Banana	All stages	Pseudostem weevil	<ul style="list-style-type: none">• Removal of heavily infested plants and deep burial in the pits,• Cleaning around the crop,
Honey bee			<ul style="list-style-type: none">• Cover hives with insulating materials/reduce hive entrance/provide adequate ventilation/keep in sunny place
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages		<ul style="list-style-type: none">• In egg laying birds feed, add calcium,

			<ul style="list-style-type: none"> Schedule vaccination should be followed,
Pigs	All stages		<ul style="list-style-type: none"> Care pregnant animals,
Cattle	All stages		<ul style="list-style-type: none"> Good time to handle parasites, consult vet for deworming/vaccinations Provide paddy straws for cushioning to the newly born calves, Lighting to maintain warmth in the shade, As green feeds supply will decrease, provide quality feeds,
Rabbits	All stages	Snuffles	<ul style="list-style-type: none"> Keep strict vigil Consult experts,
Mithun	All stages	FMD Black quarter	<ul style="list-style-type: none"> Vaccination should be done post monsoon, In young ones, BQ may be observed, vaccinate the animals,
Dog	All stages	Rabies,	<ul style="list-style-type: none"> Vaccinate against rabies, Deworming,
<ul style="list-style-type: none"> Contact Scientists (ICAR) or State Agril Staff or KVK staff for any queries/ measures against adverse conditions or call at 03795226247/37 			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
 Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

Agro Meteorological Field Unit
 ICAR, BASAR



Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar – 791 101



(Lohit)

No:-60/2013-14/Fri

Dated: 29th November, 2013

Weather during the period

- Rainfall: -
 - Tezu – NA
 - Namsai – NA

Weather Forecast

- Rainfall: Clear/Sunny weather fcsted for next 3-4 days.
- Temperature –The temperature is likely to be in the range of 10-28°C.
- Relative Humidity - The RH is likely to be in the range of 38-92 %.

Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
VEGETABLE CROPS			
Cauliflower/ Cabbage/ Broccoli/ Knolkhol	Vegetative/ Maturing/	Leaf eating/ Cabbage caterpillars	<ul style="list-style-type: none">• Light irrigation is essential,• In present weather conditions severe attack of leaf eating caterpillars has been reported. Spray Chlorpyrifos/Imidacloprid in recommended dose in the next one or two days when weather is clear. Dose- Imidacloprid 17SL @ 1-2 ml per litre depending upon the population,
FIELD CROPS			
Cauliflower/ Cabbage/ Broccoli/ Knolkhol	Vegetative/ Maturing/	Leaf eating/ Cabbage caterpillars	<ul style="list-style-type: none">• Light irrigation is essential,• In present weather conditions severe attack of leaf eating caterpillars has been reported. Spray Chlorpyrifos/Imidacloprid in recommended dose in the next one or two days when weather is clear. Dose- Imidacloprid 17SL @ 1-2 ml per litre depending upon the population,
Potato	Vegetative		<ul style="list-style-type: none">• Light irrigation followed by earthing-up,• Monitor the crop against attack of Cutworm,
Mustard	Flowering		<ul style="list-style-type: none">• Need based irrigation for those in flowering stage• Thinning may be done to maintain proper plant count,
FRUIT CROPS			
Khasi Mandarin	Maturing Harvesting	Trunk borer/ Fruit fly	<ul style="list-style-type: none">• In present weather conditions regularly collect and dispose off fallen fruits,• Farmers are advised to use <i>Jaggary</i>/Malathion traps or pheromone traps to monitor the pest population and collect and destroy the adult insects. Installation of pheromone traps may also be done,
Honey bee			<ul style="list-style-type: none">• Cover hives with insulating materials/reduce hive entrance/provide adequate ventilation/keep in sunny place
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages		<ul style="list-style-type: none">• In egg laying birds feed, add calcium,• Schedule vaccination should be followed,
Pigs	All stages		<ul style="list-style-type: none">• Care pregnant animals,
Cattle	All stages		<ul style="list-style-type: none">• Good time to handle parasites, consult vet for deworming/vaccinations• Provide paddy straws for cushioning to the newly born calves,• Lighting to maintain warmth in the shade,• As green feeds supply will decrease, provide quality feeds,
<ul style="list-style-type: none">• Contact Scientists (ICAR) or State Agril Staff or KVK staff for any queries/measures against adverse conditions or call at 03795226247/37			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit
ICAR, BASAR



Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar – 791 101



(Lower Subansiri)

No:-60/2013-14/Fri

Dated: 29th November, 2013

Weather during the period

- Rainfall: -
1. Ziro – NA

Weather Forecast

• <u>Rainfall</u> – Clear/sunny/dry weather fcted for next 3-4 days with shallow morning fog. Parially cloudy weather may persist on some days.
• <u>Temperature</u> - The temperature is likely to be in the range of 2-12°C.
• <u>Relative Humidity</u> - The RH is likely to be in the range of 55-96 %.

Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
VEGETABLE CROPS			
Cauliflower/ Cabbage/ Broccoli/ Knolkhol	Vegetative/ Maturing/	Leaf eating/ Cabbage caterpillars	<ul style="list-style-type: none">• Light irrigation is essential,• In present weather conditions severe attack of leaf eating caterpillars has been reported. Spray <i>Chlorpyrifos/Imidacloprid</i> in recommended dose in the next one or two days when weather is clear. Dose- Imidacloprid 17SL @ 1-2 ml per litre depending upon the population,
Radish	Vegetative Harvesting	Leaf eating caterpillars	<ul style="list-style-type: none">• Light interculture operations may be done,• Spray Chlorpyrifos/Imidacloprid in recommended dose.
FIELD CROPS			
WRC rice – Main crop (Mipun)	Harvesting/ Threshing		<ul style="list-style-type: none">• Drying of grains/seed to appropriate level of moisture,• Seeds/grains should be stored in dry place and if available use some <i>neem</i> leaves,
Potato	Vegetative		<ul style="list-style-type: none">• Light irrigation followed by earthing-up,• Monitor the crop against attack of Cutworm,
Turmeric and ginger	Harvesting		<ul style="list-style-type: none">• Proper cleaning of the rhizomes before storing,
Mustard	Flowering		<ul style="list-style-type: none">• Need based irrigation for those in flowering stage• Thinning may be done to maintain proper plant count,
Winter Maize	Vegetative	Cut worm	<ul style="list-style-type: none">• Monitor for attack of cut worm in the germinated fields,• Manual collection of larvae during the evening hours may be done, (If damage level is not serious). Use of Carbaryl dust in severely damaged fields may be done,
Pea	Vegetative		
FRUIT CROPS			
Khasi Mandarin	Maturing Harvesting	Trunk borer/ Fruit fly	<ul style="list-style-type: none">• In present weather conditions regularly collect and dispose off fallen fruits,• Farmers are advised to use <i>Jaggary</i>/Malathion traps or pheromone traps to monitor the pest population and collect and destroy the adult insects. Installation of pheromone traps may also be done,
Banana	All stages	Pseudostem weevil	<ul style="list-style-type: none">• Removal of heavily infested plants and deep burial in the pits,• Cleaning around the crop,
Honey bee			<ul style="list-style-type: none">• Cover hives with insulating materials/reduce hive entrance/provide adequate ventilation/keep in sunny place
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages		<ul style="list-style-type: none">• In egg laying birds feed, add calcium,• Schedule vaccination should be followed,
Pigs	All stages		<ul style="list-style-type: none">• Care pregnant animals,
Cattle	All stages		<ul style="list-style-type: none">• Good time to handle parasites, consult vet for deworming/vaccinations• Provide paddy straws for cushioning to the newly born calves.

			<ul style="list-style-type: none">• Lighting to maintain warmth in the shade,• As green feeds supply will decrease, provide quality feeds,
Rabbits	All stages	Snuffles	<ul style="list-style-type: none">• Keep strict vigil• Consult experts,
Mithun	All stages	FMD Black quarter	<ul style="list-style-type: none">• Vaccination should be done post monsoon,• In young ones, BQ may be observed, vaccinate the animals,
Dog	All stages	Rabies,	<ul style="list-style-type: none">• Vaccinate against rabies,• Deworming,
<ul style="list-style-type: none">• Contact Scientists (ICAR) or State Agril Staff or KVK staff for any queries/measures against adverse conditions or call at 03795226247/37			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37,-96 (Fax)

**Agrometeorological Field Unit
ICAR, BASAR**



Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar – 791 101



(Papum Pare)

No:-60/2013-14/Fri

Dated: 29th November, 2013

Weather during the period

- Maximum temperature: NA
- Minimum Temperature: NA
- Relative Humidity (%):NA
- Rainfall: -
 1. Itanagar -NA
 2. Naharlagun -NA

Weather Forecast

- Rainfall Clear/sunny/dry weather fcted for next 3-4 days with shallow morning fog.Parially cloudy weather may persist on some days.
- Temperature - The temperature is likely to be in the range of 11-28°C.
- Relative Humidity - The RH is likely to be in the range of 47-95 %.

Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
VEGETABLE CROPS			
Cauliflower/ Cabbage/ Brocolli/ Knolkhol	Vegetative/ Maturing/	Leaf eating/ Cabbage caterpillars	<ul style="list-style-type: none">• Light irrigation is essential,• In present weather conditions severe attack of leaf eating caterpillars has been reported. Spray Chlorpyriphos/Imidacloprid in recommended dose in the next one or two days when weather is clear. Dose- Imidacloprid 17SL @ 1-2 ml per litre depending upon the population,
Radish	Vegetative Harvesting	Leaf eating caterpillars	<ul style="list-style-type: none">• Light interculture operations may be done,• Spray Chlorpyriphos/Imidacloprid in recommended dose.
Carrot	Vegetative		<ul style="list-style-type: none">• Light irrigation is essential,• Light interculture operations should be done,
Spinach	Vegetative Harvesting		<ul style="list-style-type: none">• Sowing of certified/locally available seeds in well prepared nursery may be done,
FIELD CROPS			
WRC rice – Main crop (Mipun)	Harvesting/ Threshing		<ul style="list-style-type: none">• Drying of grains/seed to appropriate level of moisture,• Seeds/grains should be stored in dry place and if available use some <i>neem</i> leaves,
Potato	Vegetative		<ul style="list-style-type: none">• Light irrigation followed by earthing-up,• Monitor the crop against attack of Cutworm,
Turmeric and ginger	Harvesting		<ul style="list-style-type: none">• Proper cleaning of the rhizomes before storing,
Mustard	Flowering		<ul style="list-style-type: none">• Need based irrigation for those in flowering stage• Thinning may be done to maintain proper plant count,
Winter Maize	Vegetative	Cut worm	<ul style="list-style-type: none">• Monitor for attack of cut worm in the germinated fields,• Manual collection of larvae during the evening hours may be done, (If damage level is not serious). Use of Carbaryl dust in severely damaged fields may be done,
Pea	Vegetative		
FRUIT CROPS			
Khasi Mandarin	Maturing Harvesting	Trunk borer/ Fruit fly	<ul style="list-style-type: none">• In present weather conditions regularly collect and dispose off fallen fruits,• Farmers are advised to use <i>Jaggary</i>/Malathion traps or pheromone traps to monitor the pest population and collect and destroy the adult insects. Installation of pheromone traps may also be done,
Banana	All stages	Pseudostem weevil	<ul style="list-style-type: none">• Removal of heavily infested plants and deep burial in the pits,• Cleaning around the crop,
Honey bee			<ul style="list-style-type: none">• Cover hives with insulating materials/reduce hive entrance/provide adequate ventilation/keep in sunny place
LIVESTOCKS			

Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages		<ul style="list-style-type: none"> In egg laying birds feed, add calcium, Schedule vaccination should be followed,
Pigs	All stages		<ul style="list-style-type: none"> Care pregnant animals,
Cattle	All stages		<ul style="list-style-type: none"> Good time to handle parasites, consult vet for deworming/vaccinations Provide paddy straws for cushioning to the newly born calves, Lighting to maintain warmth in the shade, As green feeds supply will decrease, provide quality feeds,
Rabbits	All stages	Snuffles	<ul style="list-style-type: none"> Keep strict vigil Consult experts,
Mithun	All stages	FMD Black quarter	<ul style="list-style-type: none"> Vaccination should be done post monsoon, In young ones, BQ may be observed, vaccinate the animals,
Dog	All stages	Rabies,	<ul style="list-style-type: none"> Vaccinate against rabies, Deworming,
<ul style="list-style-type: none"> Contact Scientists (ICAR) or State Agril Staff or KVK staff for any queries/measures against adverse conditions or call at 03795226247/37 			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit
ICAR, BASAR



Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar – 791 101



(Tawang)

No:-60/2013-14/Fri

Dated: 29th November, 2013

Weather during the period

- Rainfall: -
1. Tawang –NA

Weather Forecast

- Rainfall- – Clear/partially cloudy weather fctsted for next 3-4 days with fog.
- Temperature - The temperature is likely to be in the range of 1-12 °C.
- Relative Humidity - The RH is likely to be in the range of 52-93 %.

Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FRUITS			
Apple	Lean period		<ul style="list-style-type: none">• <i>Pruning of the trees</i>,• Timely and periodic inspection of trees for its health,• Urgent/immediate treatment of affected parts of trees,
VEGETABLE CROPS			
Cauliflower / Cabbage/ Radish	Harvesting	Cutworm Diamond back moth	<ul style="list-style-type: none">• Monitor crop against attack of cutworm in the field,• In present weather conditions, monitor the crop against attack of DBM caterpillars,• If damage level is high, spraying of <i>Bt</i> formulation may be done,
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages		<ul style="list-style-type: none">• Schedule vaccination should be followed,
Cattle	All stages		<ul style="list-style-type: none">• Provide paddy straws for cushioning to the newly born calves,• Lighting to maintain warmth in the shade,• As green feeds supply will decrease, provide quality feeds,
Goat/Sheep	All stages	PPR	<ul style="list-style-type: none">• Vaccinate against the PPR,• Provide mineral mixture,
Pigs	All stages		<ul style="list-style-type: none">• Provide clean water free from ice/rapid or laboured breathing needs immediate attention & treatment• Care pregnant animals,

Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795222247 (KVK, West Siang)

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department

Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit
ICAR, BASAR



Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar – 791 101



(Tirap)

No:-60/2013-14/Fri

Dated: 29th November, 2013

Weather during the period

- Rainfall: -
 - Khonsa –NA
 - Deomali –NA

Weather Forecast

• <u>Rainfall</u> : Clear/Sunny weather fcted for next 3-4 days.
• <u>Temperature</u> - The temperature is likely to be in the range of 8-24°C.
• <u>Relative Humidity</u> - The RH is likely to be in the range of 39-99 %.

Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
VEGETABLE CROPS			
Cauliflower/ Cabbage/ Broccoli/ Knolkhol	Vegetative/ Maturing/	Leaf eating/ Cabbage caterpillars	<ul style="list-style-type: none"> • Light irrigation is essential, • In present weather conditions severe attack of leaf eating caterpillars has been reported. Spray <i>Chlorpyrifos/Imidacloprid</i> in recommended dose in the next one or two days when weather is clear. Dose- Imidacloprid 17SL @ 1-2 ml per litre depending upon the population,
FIELD CROPS			
Cauliflower/ Cabbage/ Broccoli/ Knolkhol	Vegetative/ Maturing/	Leaf eating/ Cabbage caterpillars	<ul style="list-style-type: none"> • Light irrigation is essential, • In present weather conditions severe attack of leaf eating caterpillars has been reported. Spray <i>Chlorpyrifos/Imidacloprid</i> in recommended dose in the next one or two days when weather is clear. Dose- Imidacloprid 17SL @ 1-2 ml per litre depending upon the population,
Potato	Vegetative		<ul style="list-style-type: none"> • Light irrigation followed by earthing-up, • Monitor the crop against attack of Cutworm,
Mustard	Flowering		<ul style="list-style-type: none"> • Need based irrigation for those in flowering stage • Thinning may be done to maintain proper plant count,
FRUIT CROPS			
Khasi Mandarin	Maturing Harvesting	Trunk borer/ Fruit fly	<ul style="list-style-type: none"> • In present weather conditions regularly collect and dispose off fallen fruits, • Farmers are advised to use <i>Jaggary</i>/Malathion traps or pheromone traps to monitor the pest population and collect and destroy the adult insects. Installation of pheromone traps may also be done,
Honey bee			<ul style="list-style-type: none"> • Cover hives with insulating materials/reduce hive entrance/provide adequate ventilation/keep in sunny place
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages		<ul style="list-style-type: none"> • In egg laying birds feed, add calcium, • Schedule vaccination should be followed,
Pigs	All stages		<ul style="list-style-type: none"> • Care pregnant animals,
Cattle	All stages		<ul style="list-style-type: none"> • Good time to handle parasites, consult vet for deworming/vaccinations • Provide paddy straws for cushioning to the newly born calves, • Lighting to maintain warmth in the shade, • As green feeds supply will decrease, provide quality feeds,
<ul style="list-style-type: none"> • Contact Scientists (ICAR) or State Agril Staff or KVK staff for any queries/measures against adverse conditions or call at 03795226247/37 			



Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar – 791 101



(Upper Siang)

No:-60/2013-14/Fri

Dated: 29th November, 2013

Weather during the period

- Rainfall: -
N Tuting – NA

Weather Forecast

• <u>Rainfall</u> – Clear/sunny/dry weather fcsted for next 3-4 days with shallow morning fog.Parially cloudy weather may persist on some days
• <u>Temperature</u> - The temperature is likely to be in the range of 5-20°C.
• <u>Relative Humidity</u> - The RH is likely to be in the range of 29-94 %.

Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
VEGETABLE CROPS			
Cauliflower/ Cabbage/ Brocolli/ Knolkhol	Vegetative/ Maturing/	Leaf eating/ Cabbage caterpillars	<ul style="list-style-type: none">• Light irrigation is essential,• In present weather conditions severe attack of leaf eating caterpillars has been reported. Spray <i>Chlorpyrifos/Imidacloprid</i> in recommended dose in the next one or two days when weather is clear. Dose- Imidacloprid 17SL @ 1-2 ml per litre depending upon the population,
Radish	Vegetative Harvesting	Leaf eating caterpillars	<ul style="list-style-type: none">• Light interculture operations may be done,• Spray Chlorpyrifos/Imidacloprid in recommended dose.
FIELD CROPS			
WRC rice – Main crop (Mipun)	Harvesting/ Threshing		<ul style="list-style-type: none">• Drying of grains/seed to appropriate level of moisture,• Seeds/grains should be stored in dry place and if available use some <i>neem</i> leaves,
Potato	Vegetative		<ul style="list-style-type: none">• Light irrigation followed by earthing-up,• Monitor the crop against attack of Cutworm,
Turmeric and ginger	Harvesting		<ul style="list-style-type: none">• Proper cleaning of the rhizomes before storing,
Mustard	Flowering		<ul style="list-style-type: none">• Need based irrigation for those in flowering stage• Thinning may be done to maintain proper plant count,
Winter Maize	Vegetative	Cut worm	<ul style="list-style-type: none">• Monitor for attack of cut worm in the germinated fields,• Manual collection of larvae during the evening hours may be done, (If damage level is not serious). Use of Carbaryl dust in severely damaged fields may be done,
Pea	Vegetative		
FRUIT CROPS			
Khasi Mandarin	Maturing Harvesting	Trunk borer/ Fruit fly	<ul style="list-style-type: none">• In present weather conditions regularly collect and dispose off fallen fruits,• Farmers are advised to use <i>Jaggary</i>/Malathion traps or pheromone traps to monitor the pest population and collect and destroy the adult insects. Installation of pheromone traps may also be done,
Banana	All stages	Pseudostem weevil	<ul style="list-style-type: none">• Removal of heavily infested plants and deep burial in the pits,• Cleaning around the crop,
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages		<ul style="list-style-type: none">• In egg laying birds feed, add calcium,• Schedule vaccination should be followed,
Pigs	All stages		<ul style="list-style-type: none">• Care pregnant animals,
Cattle	All stages		<ul style="list-style-type: none">• Good time to handle parasites, consult vet for deworming/vaccinations• Provide paddy straws for cushioning to the newly born calves,• Lighting to maintain warmth in the shade,• As green feeds supply will decrease, provide quality feeds,

Rabbits	All stages	Snuffles	<ul style="list-style-type: none">• Keep strict vigil• Consult experts,
Mithun	All stages	FMD Black quarter	<ul style="list-style-type: none">• Vaccination should be done post monsoon,• In young ones, BQ may be observed, vaccinate the animals,
Dog	All stages	Rabies,	<ul style="list-style-type: none">• Vaccinate against rabies,• Deworming,
<ul style="list-style-type: none">• Contact Scientists (ICAR) or State Agril Staff or KVK staff for any queries/measures against adverse conditions or call at 03795226247/37			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

Agro Meteorological Field Unit
ICAR, BASAR



Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar – 791 101



(Upper Subansiri)

No:-60/2013-14/Fri

Dated: 29th November, 2013

Weather during the period

- Rainfall: -
 - Daporijo- NA

Weather Forecast

- Rainfall** – Clear/sunny/dry weather fcted for next 3-4 days with shallow morning fog.Parially cloudy weather may persist on some days.
- Temperature** - The temperature is likely to be in the range of 2-12°C.
- Relative Humidity** - The RH is likely to be in the range of 48-94 %.

Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
VEGETABLE CROPS			
Cauliflower/ Cabbage/ Broccoli/ Knolkhol	Vegetative/ Maturing/	Leaf eating/ Cabbage caterpillars	<ul style="list-style-type: none">• Light irrigation is essential,• In present weather conditions severe attack of leaf eating caterpillars has been reported. Spray <i>Chlorpyrifos/Imidacloprid</i> in recommended dose in the next one or two days when weather is clear. Dose- Imidacloprid 17SL @ 1-2 ml per litre depending upon the population,
Radish	Vegetative Harvesting	Leaf eating caterpillars	<ul style="list-style-type: none">• Light interculture operations may be done,• Spray Chlorpyrifos/Imidacloprid in recommended dose.
FIELD CROPS			
WRC rice – Main crop (<i>Mipun</i>)	Harvesting/ Threshing		<ul style="list-style-type: none">• Drying of grains/seed to appropriate level of moisture,• Seeds/grains should be stored in dry place and if available use some <i>neem</i> leaves,
Potato	Vegetative		<ul style="list-style-type: none">• Light irrigation followed by earthing-up,• Monitor the crop against attack of Cutworm,
Turmeric and ginger	Harvesting		<ul style="list-style-type: none">• Proper cleaning of the rhizomes before storing,
Mustard	Flowering		<ul style="list-style-type: none">• Need based irrigation for those in flowering stage• Thinning may be done to maintain proper plant count,
Winter Maize	Vegetative	Cut worm	<ul style="list-style-type: none">• Monitor for attack of cut worm in the germinated fields,• Manual collection of larvae during the evening hours may be done, (If damage level is not serious). Use of Carbaryl dust in severely damaged fields may be done,
Pea	Vegetative		
FRUIT CROPS			
Khasi Mandarin	Fruit development	Trunk borer/ Fruit fly	<ul style="list-style-type: none">• In present weather conditions regularly collect and dispose off fallen fruits,• Farmers are advised to use <i>Jaggary</i>/Malathion traps or pheromone traps to monitor the pest population and collect and destroy the adult insects. Installation of pheromone traps may also be done,
Banana	All stages	Pseudostem weevil	<ul style="list-style-type: none">• Removal of heavily infested plants and deep burial in the pits,• Cleaning around the crop,
Honey bee			<ul style="list-style-type: none">• Cover hives with insulating materials/reduce hive entrance/provide adequate ventilation/keep in sunny place
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages		<ul style="list-style-type: none">• In egg laying birds feed, add calcium,• Schedule vaccination should be followed,
Pigs	All stages		<ul style="list-style-type: none">• Care pregnant animals,
Cattle	All stages		<ul style="list-style-type: none">• Good time to handle parasites, consult vet for deworming/vaccinations• Provide paddy straws for cushioning to the newly born calves,

			<ul style="list-style-type: none"> • Lighting to maintain warmth in the shade, • As green feeds supply will decrease, provide quality feeds,
Rabbits	All stages	Snuffles	<ul style="list-style-type: none"> • Keep strict vigil • Consult experts,
Mithun	All stages	FMD Black quarter	<ul style="list-style-type: none"> • Vaccination should be done post monsoon, • In young ones, BQ may be observed, vaccinate the animals,
Dog	All stages	Rabies,	<ul style="list-style-type: none"> • Vaccinate against rabies, • Deworming,
<ul style="list-style-type: none"> • Contact Scientists (ICAR) or State Agril Staff or KVK staff for any queries/measures against adverse conditions or call at 03795226247/37 			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

Agro Meteorological Field Unit
 ICAR, BASAR



Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar – 791 101



(West Kameng)

No:-60/2013-14/Fri

Dated: 29th November, 2013

Weather during the period

- Rainfall: - Bomdila –NA
Bhalokpung – NA

Weather Forecast

• <u>Rainfall</u> - Clear/sunny/dry weather fested for next 3-4 days with fog during morning hours. Partially cloudy weather may persist on some days.
• <u>Temperature</u> - The temperature is likely to be in the range of 5-14°C.
• <u>Relative Humidity</u> - The RH is likely to be in the range of 53-96 %.

Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FRUITS			
Apple	Lean period		<ul style="list-style-type: none">• <i>Pruning of the trees</i>,• Timely and periodic inspection of trees for its health,• Urgent/immediate treatment of affected parts of trees,
VEGETABLE CROPS			
Cauliflower/ Cabbage/ Radish	Harvesting	Cutworm Diamond back moth	<ul style="list-style-type: none">• Monitor crop against attack of cutworm in the field,• In present weather conditions, monitor the crop against attack of DBM caterpillars,• If damage level is high, spraying of <i>Bt</i> formulation may be done,
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages		<ul style="list-style-type: none">• Schedule vaccination should be followed,
Cattle	All stages		<ul style="list-style-type: none">• Provide paddy straws for cushioning to the newly born calves,• Lighting to maintain warmth in the shade,• As green feeds supply will decrease, provide quality feeds,
Goat/Sheep	All stages	PPR	<ul style="list-style-type: none">• Vaccinate against the PPR,• Provide mineral mixture,
Pigs	All stages		<ul style="list-style-type: none">• Provide clean water free from ice/rapid or laboured breathing needs immediate attention & treatment• Care pregnant animals,

Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795222247 (KVK, West Siang)

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department

Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit
ICAR, BASAR



Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar – 791 101



(West Siang)

No:-60/2013-14/Fri

Dated: 29th November, 2013

Weather during the period

- During the period weather was clear and dry during the day time and morning shallow fog.
 - Maximum temperature: **28.5°C**
 - Minimum Temperature: **9.0°C**
 - Rainfall: **0.0 mm**
 - Evapotranspiration: - -
 - Relative Humidity (%): **-44-100%**
 - Soil moisture (m^2/m^2): **0.25**
 - Prominent Wind Direction: **North West (Morning hours) & North West (Afternoon)**
 - Average Wind Speed: **0.3 metres per second**
 - Average Sunshine hours : **5.8 hours**
- (Source-Agromet Observatory, Research Farm, Gori, ICAR RC for NEH Region, AP Centre, Basar)

Weather Forecast

- **Rainfall:** Clear/sunny/dry weather fcsted for next 3-4 days with shallow morning fog. Parially cloudy weather may persist on some days.
- **Temperature** – Temperature will vary between 9-24°C.
- **Relative Humidity** - The RH is likely to be in the range of 41-94 %.

Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
VEGETABLE CROPS			
Cauliflower/ Cabbage/ Brocolli/ Knolkhol	Vegetative/ Maturing/	Leaf eating/ Cabbage caterpillars	<ul style="list-style-type: none">• Light irrigation is essential,• In present weather conditions severe attack of leaf eating caterpillars has been reported. Spray Chlorpyriphos/Imidacloprid in recommended dose in the next one or two days when weather is clear. Dose-Imidacloprid 17SL @ 1-2 ml per litre depending upon the population,
Radish	Vegetative Harvesting	Leaf eating caterpillars	<ul style="list-style-type: none">• Light interculture operations may be done,• Spray Chlorpyriphos/Imidacloprid in recommended dose.
Carrot	Vegetative		<ul style="list-style-type: none">• Light irrigation is essential,• Light interculture operations should be done,
Spinach	Vegetative Harvesting		<ul style="list-style-type: none">• Sowing of certified/locally available seeds in well prepared nursery may be done,
FIELD CROPS			
WRC rice – Main crop (Mipun)	Harvesting/ Threshing		<ul style="list-style-type: none">• Drying of grains/seed to appropriate level of moisture,• Seeds/grains should be stored in dry place and if available use some <i>neem</i> leaves,
Potato	Vegetative		<ul style="list-style-type: none">• Light irrigation followed by earthing-up,• Monitor the crop against attack of Cutworm,
Turmeric and ginger	Harvesting		<ul style="list-style-type: none">• Proper cleaning of the rhizomes before storing,
Mustard	Flowering		<ul style="list-style-type: none">• Need based irrigation for those in flowering stage• Thinning may be done to maintain proper plant count,
Honey bee			<ul style="list-style-type: none">• Cover hives with insulating materials/reduce hive entrance/provide adequate ventilation/keep in sunny place,
FRUIT CROPS			
Khasi Mandarin	Maturing Harvesting	Trunk borer/ Fruit fly	<ul style="list-style-type: none">• In present weather conditions regularly collect and dispose off fallen fruits,• Farmers are advised to use <i>Jaggary</i>/Malathion traps or pheromone traps to monitor the pest population and collect and destroy the adult insects. Installation of pheromone traps may also be done,
Banana	All stages	Pseudostem weevil	<ul style="list-style-type: none">• Removal of heavily infested plants and deep burial in the pits,• Cleaning around the crop,

LIVESTOCKS and VETERINARY ASPECT			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages		<ul style="list-style-type: none"> In egg laying birds feed, add calcium, Schedule vaccination should be followed,
Pigs	All stages		<ul style="list-style-type: none"> Care pregnant animals,
Cattle	All stages		<ul style="list-style-type: none"> Good time to handle parasites, consult vet for deworming/vaccinations Provide paddy straws for cushioning to the newly born calves, Lighting to maintain warmth in the shade, As green feeds supply will decrease, provide quality feeds,
Rabbits	All stages	Snuffles	<ul style="list-style-type: none"> Keep strict vigil Consult experts,
Mithun	All stages	FMD Black quarter	<ul style="list-style-type: none"> Vaccination should be done post monsoon, In young ones, BQ may be observed, vaccinate the animals,
Dog	All stages	Rabies,	<ul style="list-style-type: none"> Vaccinate against rabies, Deworming,
Goat/Sheep	All stages	PPR	<ul style="list-style-type: none"> Vaccinate against the PPR, Provide mineral mixture,
Fish	All stages		<ul style="list-style-type: none"> Maintain adequate water level in the pond, Pond sides should be properly cleaned, Harvest of mature fish,
<ul style="list-style-type: none"> Contact Scientists (ICAR) or State Agril Staff or KVK staff for any queries/measures against adverse conditions or call at 03795226247/37 			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

Agro Meteorological Field Unit
 ICAR, BASAR



Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar – 791 101



(Changlang)

No:-60/2013-14/Fri

Dated: 29th November, 2013

Weather during the period

- **Rainfall:** -
 - Changlang –NA
 - Miao – NA

Weather Forecast

- **Rainfall** – Clear/sunny/dry weather fcsted for next 3-4 days with shallow morning fog.Partially cloudy weather may persist on some days.
- **Temperature** - The temperature is likely to be in the range of 10-23 °C.
- **Relative Humidity** - The RH is likely to be in the range of 37-92 %.

Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
VEGETABLE CROPS			
Cauliflower/ Cabbage/ Brocolli/ Knolkhol	Vegetative/ Maturing/	Leaf eating/ Cabbage caterpillars	<ul style="list-style-type: none">• Light irrigation is essential,• In present weather conditions severe attack of leaf eating caterpillars has been reported. Spray Chlorpyrifos/Imidacloprid in recommended dose in the next one or two days when weather is clear. Dose- Imidacloprid 17SL @ 1-2 ml per litre depending upon the population,
FIELD CROPS			
Cauliflower/ Cabbage/ Brocolli/ Knolkhol	Vegetative/ Maturing/	Leaf eating/ Cabbage caterpillars	<ul style="list-style-type: none">• Light irrigation is essential,• In present weather conditions severe attack of leaf eating caterpillars has been reported. Spray Chlorpyrifos/Imidacloprid in recommended dose in the next one or two days when weather is clear. Dose- Imidacloprid 17SL @ 1-2 ml per litre depending upon the population,
Potato	Vegetative		<ul style="list-style-type: none">• Light irrigation followed by earthing-up,• Monitor the crop against attack of Cutworm,
Mustard	Flowering		<ul style="list-style-type: none">• Need based irrigation for those in flowering stage• Thinning may be done to maintain proper plant count,
FRUIT CROPS			
Khasi Mandarin	Maturing Harvesting	Trunk borer/ Fruit fly	<ul style="list-style-type: none">• In present weather conditions regularly collect and dispose off fallen fruits,• Farmers are advised to use <i>Jaggary</i>/Malathion traps or pheromone traps to monitor the pest population and collect and destroy the adult insects. Installation of pheromone traps may also be done,
Honey bee			<ul style="list-style-type: none">• Cover hives with insulating materials/reduce hive entrance/provide adequate ventilation/keep in sunny place
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages		<ul style="list-style-type: none">• In egg laying birds feed, add calcium,• Schedule vaccination should be followed,
Pigs	All stages		<ul style="list-style-type: none">• Care pregnant animals,
Cattle	All stages		<ul style="list-style-type: none">• Good time to handle parasites, consult vet for deworming/vaccinations• Provide paddy straws for cushioning to the newly born calves,• Lighting to maintain warmth in the shade,• As green feeds supply will decrease, provide quality feeds,
<ul style="list-style-type: none">• Contact Scientists (ICAR) or State Agril Staff or KVK staff for any queries/measures against adverse conditions or call at 03795226247/37			



Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar – 791 101



(Dibang Valley)

No:-60/2013-14/Fri

Dated: 29th November, 2013

Weather during the period

- **Rainfall:** -
 - Roing – NA
 - Anini – NA

Weather Forecast

- **Rainfall:** Clear/sunny/dry weather fsted for next 3-4 days with shallow morning fog.
- **Temperature** – Temperature will vary between 4-18°C.
- **Relative Humidity** - The RH is likely to be in the range of 34-84 %.

Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
VEGETABLE CROPS			
Cauliflower/ Cabbage/ Broccoli/ Knolkhol	Vegetative/ Maturing/	Leaf eating/ Cabbage caterpillars	<ul style="list-style-type: none"> • Light irrigation is essential, • In present weather conditions severe attack of leaf eating caterpillars has been reported. Spray Chlorpyrifos/Imidacloprid in recommended dose in the next one or two days when weather is clear. Dose-Imidacloprid 17SL @ 1-2 ml per litre depending upon the population,
Radish	Vegetative Harvesting	Leaf eating caterpillars	<ul style="list-style-type: none"> • Light interculture operations may be done, • Spray Chlorpyrifos/Imidacloprid in recommended dose.
Carrot	Vegetative		<ul style="list-style-type: none"> • Light irrigation is essential, • Light interculture operations should be done,
Spinach	Vegetative Harvesting		<ul style="list-style-type: none"> • Sowing of certified/locally available seeds in well prepared nursery may be done,
FIELD CROPS			
WRC rice – Main crop (Mipun)	Harvesting/ Threshing		<ul style="list-style-type: none"> • Drying of grains/seed to appropriate level of moisture, • Seeds/grains should be stored in dry place and if available use some <i>neem</i> leaves,
Potato	Vegetative		<ul style="list-style-type: none"> • Light irrigation followed by earthing-up, • Monitor the crop against attack of Cutworm,
Turmeric and ginger	Harvesting		<ul style="list-style-type: none"> • Proper cleaning of the rhizomes before storing,
Mustard	Flowering		<ul style="list-style-type: none"> • Need based irrigation for those in flowering stage • Thinning may be done to maintain proper plant count,
Honey bee			<ul style="list-style-type: none"> • Cover hives with insulating materials/reduce hive entrance/provide adequate ventilation/keep in sunny place,
FRUIT CROPS			
Khasi Mandarin	Maturing Harvesting	Trunk borer/ Fruit fly	<ul style="list-style-type: none"> • In present weather conditions regularly collect and dispose off fallen fruits, • Farmers are advised to use <i>Jaggary</i>/Malathion traps or pheromone traps to monitor the pest population and collect and destroy the adult insects. Installation of pheromone traps may also be done,
Banana	All stages	Pseudostem weevil	<ul style="list-style-type: none"> • Removal of heavily infested plants and deep burial in the pits, • Cleaning around the crop,
LIVESTOCKS and VETERINARY ASPECT			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages		<ul style="list-style-type: none"> • In egg laying birds feed, add calcium, • Schedule vaccination should be followed,
Pigs	All stages		<ul style="list-style-type: none"> • Care pregnant animals,
Cattle	All stages		<ul style="list-style-type: none"> • Good time to handle parasites, consult vet for deworming/vaccinations • Provide paddy straws for cushioning to the newly born calves, • Lighting to maintain warmth in the shade,

			<ul style="list-style-type: none">As green feeds supply will decrease, provide quality feeds,
Mithun	All stages	FMD Black quarter	<ul style="list-style-type: none">Vaccination should be done post monsoon,In young ones, BQ may be observed, vaccinate the animals,
Goat/Sheep	All stages	PPR	<ul style="list-style-type: none">Vaccinate against the PPR,Provide mineral mixture,
Fish	All stages		<ul style="list-style-type: none">Maintain adequate water level in the pond,Pond sides should be properly cleaned,Harvest of mature fish,
<ul style="list-style-type: none">Contact Scientists (ICAR) or State Agril Staff or KVK staff for any queries/measures against adverse conditions or call at 03795226247/37			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

Agro Meteorological Field Unit
ICAR, BASAR



Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar – 791 101



(East Kameng)

No:-60/2013-14/Fri

Dated: 29th November, 2013

Weather during the period

Rainfall: -

- Seppa -NA

Weather Forecast

- **Rainfall** : Clear/sunny/dry weather fested for next 3-4 days with shallow morning fog. Parially cloudy weather may persist on some days
- **Temperature** - The temperature is likely to be in the range of 4-13 °C.
- **Relative Humidity** - The RH is likely to be in the range of 53-97 %.

Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
VEGETABLE CROPS			
Cauliflower/ Cabbage/ Broccoli/ Knolkhol	Vegetative/ Maturing/	Leaf eating/ Cabbage caterpillars	<ul style="list-style-type: none">• Light irrigation is essential,• In present weather conditions severe attack of leaf eating caterpillars has been reported. Spray <i>Chlorpyrifos/Imidacloprid</i> in recommended dose in the next one or two days when weather is clear. Dose- Imidacloprid 17SL @ 1-2 ml per litre depending upon the population,
Radish	Vegetative Harvesting	Leaf eating caterpillars	<ul style="list-style-type: none">• Light interculture operations may be done,• Spray Chlorpyrifos/Imidacloprid in recommended dose.
Carrot	Vegetative		<ul style="list-style-type: none">• Light irrigation is essential,• Light interculture operations should be done,
Spinach	Vegetative Harvesting		<ul style="list-style-type: none">• Sowing of certified/locally available seeds in well prepared nursery may be done,
FIELD CROPS			
WRC rice – Main crop (<i>Mipun</i>)	Harvesting/ Threshing		<ul style="list-style-type: none">• Drying of grains/seed to appropriate level of moisture,• Seeds/grains should be stored in dry place and if available use some <i>neem</i> leaves,
Potato	Vegetative		<ul style="list-style-type: none">• Light irrigation followed by earthing-up,• Monitor the crop against attack of Cutworm,
Turmeric and ginger	Harvesting		<ul style="list-style-type: none">• Proper cleaning of the rhizomes before storing,
Mustard	Flowering		<ul style="list-style-type: none">• Need based irrigation for those in flowering stage• Thinning may be done to maintain proper plant count,
Winter Maize	Vegetative	Cut worm	<ul style="list-style-type: none">• Monitor for attack of cut worm in the germinated fields,• Manual collection of larvae during the evening hours may be done, (If damage level is not serious). Use of Carbaryl dust in severely damaged fields may be done,
Pea	Vegetative		
FRUIT CROPS			
Khasi Mandarin	Maturing Harvesting	Trunk borer/ Fruit fly	<ul style="list-style-type: none">• In present weather conditions regularly collect and dispose off fallen fruits,• Farmers are advised to use <i>Jaggary</i>/Malathion traps or pheromone traps to monitor the pest population and collect and destroy the adult insects. Installation of pheromone traps may also be done,
Banana	All stages	Pseudostem weevil	<ul style="list-style-type: none">• Removal of heavily infested plants and deep burial in the pits,• Cleaning around the crop,
Honey bee			<ul style="list-style-type: none">• Cover hives with insulating materials/reduce hive entrance/provide adequate ventilation/keep in sunny place
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages		<ul style="list-style-type: none">• In egg laying birds feed, add calcium,• Schedule vaccination should be followed.

Pigs	All stages		<ul style="list-style-type: none"> Care pregnant animals,
Cattle	All stages		<ul style="list-style-type: none"> Good time to handle parasites, consult vet for deworming/vaccinations Provide paddy straws for cushioning to the newly born calves, Lighting to maintain warmth in the shade, As green feeds supply will decrease, provide quality feeds,
Rabbits	All stages	Snuffles	<ul style="list-style-type: none"> Keep strict vigil Consult experts,
Mithun	All stages	FMD Black quarter	<ul style="list-style-type: none"> Vaccination should be done post monsoon, In young ones, BQ may be observed, vaccinate the animals,
Dog	All stages	Rabies,	<ul style="list-style-type: none"> Vaccinate against rabies, Deworming,
<ul style="list-style-type: none"> Contact Scientists (ICAR) or State Agril Staff or KVK staff for any queries/measures against adverse conditions or call at 03795226247/37 			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit
ICAR, BASAR