



# Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar - 791 101



## (Changlang)

No: -73/2012-13/Tue

Dated: 8<sup>th</sup> January, 2013

### Weather during the last four days

- Rainfall: -
  - Changlang -NA
  - Miao- 0.2 mm

### Weather Forecast

• <u>Rainfall</u> - Mainly clear weather is forecasted with possibility of cloudy development on some day.
• <u>Temperature</u> - The temperature is likely to be in the range of 8-20°C.
• <u>Relative Humidity</u> - The RH is likely to be in the range of 62-83%.

### Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>VEGETABLE CROPS</b>			
Cauliflower /Cabbage/Knolkhol/Broccoli/	Vegetative/ Maturity		<ul style="list-style-type: none"> <li>• Mulching with locally available weeds, paddy straws to conserve the soil moisture and to prevent the affect of low temperature</li> </ul>
<b>FIELD CROPS</b>			
Potato	Vegetative	Late Blight	<ul style="list-style-type: none"> <li>• <b>Avoid water logging.</b></li> <li>• <b>Monitor the crop for symptoms of blight.</b></li> <li>• Foliar spraying with RidomilMZ 72 WP @ 2.5 grams per litre of water</li> <li>• Earthing up may be done,</li> </ul>
Pea	Vegetative	Aphid	<ul style="list-style-type: none"> <li>• Spray Dimethoate @ 2 ml per litre of water</li> </ul>
Rapeseed/ Mustard	Flowering	Aphid Painted bug	<ul style="list-style-type: none"> <li>• Collect and dispose off the heavily infested twigs,</li> <li>• For painted bug dusting with wood ash is advised,</li> <li>• Mulching to conserve the moisture</li> </ul>
<b>FRUIT CROPS</b>			
Khasi Mandarin	Harvest Post harvest	Lichens/ mosses	<ul style="list-style-type: none"> <li>• In present weather conditions remove lichens/mosses from the affected trees using gunny bags smoothly,</li> <li>• Prune the diseased and dried branches,</li> <li>• Collect and bury fallen fruits,</li> </ul>
<b>LIVESTOCKS</b>			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages	Cold  Coccidiosis	<ul style="list-style-type: none"> <li>• Day old chick-Maintain warmth by lighting,</li> <li>• Provide heat by lighting 100W filament bulb @ 1 bulb per 20 newly born chicks,</li> <li>• In present weather conditions frequently turn the litters/avoid hardening of litters,</li> <li>• Provide limited ventilation,</li> <li>• Add multivitamin feeds (More of Vit C)</li> <li>• Add Amprolium to the ration,</li> </ul>
Pigs	New born	Cold	<ul style="list-style-type: none"> <li>• Warming of pig sheds using gunny bags, straw, enclosure,</li> </ul>
Cattle	All stages	FMD  Respiratory	<ul style="list-style-type: none"> <li>• Vaccination - of healthy animals</li> <li>• Treatment - sick animals,</li> <li>• Intra muscular injection of Oxi-tetracycline @ 20ml/400 kg body weight animal (<b>Avoid injecting to Pregnant animals</b>)</li> </ul>
<ul style="list-style-type: none"> <li>• <b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795222247 (KVK, West Siang)</b></li> </ul>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department

Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

Integrated Agromet Advisory Services,  
Basar, Arunachal Pradesh.



# Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar - 791 101



(Lower Dibang Valley)

No: -73/2012-13/Tue

Dated: 8<sup>th</sup> January, 2013

## Weather during the last four days

- Rainfall: -
  - Roing - NA

## Weather during the last four days

• <u>Rainfall</u> – Mainly clear weather is forecasted with possibility of cloudy development on some day with light rain.
• <u>Temperature</u> - The temperature is likely to be in the range of 5-17°C.
• <u>Relative Humidity</u> - The RH is likely to be in the range of 57-88%.

## Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>VEGETABLE CROPS</b>			
Cauliflower /Cabbage/Knolkhol/Broccoli/	Vegetative/ Maturity		<ul style="list-style-type: none"> <li>• Mulching with locally available weeds, paddy straws to conserve the soil moisture and to prevent the affect of low temperature</li> </ul>
<b>FIELD CROPS</b>			
Potato	Vegetative	Late Blight	<ul style="list-style-type: none"> <li>• <b>Avoid water logging.</b></li> <li>• <b>Monitor the crop for symptoms of blight.</b></li> <li>• Foliar spraying with RidomilMZ 72 WP @ 2.5 grams per litre of water</li> <li>• Earthing up may be done,</li> </ul>
Pea	Vegetative	Aphid	<ul style="list-style-type: none"> <li>• Spray Dimethoate @ 2 ml per litre of water</li> </ul>
Rapeseed/ Mustard	Flowering	Aphid Painted bug	<ul style="list-style-type: none"> <li>• Collect and dispose off the heavily infested twigs,</li> <li>• For painted bug dusting with wood ash is advised,</li> <li>• Mulching to conserve the moisture</li> </ul>
<b>FRUIT CROPS</b>			
Khasi Mandarin	Harvest Post harvest	Lichens/ mosses	<ul style="list-style-type: none"> <li>• In present weather conditions remove lichens/mosses from the affected trees using gunny bags smoothly,</li> <li>• Prune the diseased and dried branches,</li> <li>• Collect and bury fallen fruits,</li> </ul>
<b>LIVESTOCKS</b>			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages	Cold  Coccidiosis	<ul style="list-style-type: none"> <li>• Day old chick-Maintain warmth by lighting,</li> <li>• Provide heat by lighting 100W filament bulb @ 1 bulb per 20 newly born chicks,</li> <li>• In present weather conditions frequently turn the litters/avoid hardening of litters,</li> <li>• Provide limited ventilation,</li> <li>• Add multivitamin feeds (More of Vit C)</li> <li>• Add Amprolium to the ration,</li> </ul>
Pigs	New born	Cold	<ul style="list-style-type: none"> <li>• Warming of pig sheds using gunny bags, straw, enclosure,</li> </ul>
Cattle	All stages	FMD  Respiratory	<ul style="list-style-type: none"> <li>• Vaccination – of healthy animals</li> <li>• Treatment – sick animals,</li> <li>• Intra muscular injection of Oxi-tetracycline @ 20ml/400 kg body weight animal <b>(Avoid injecting to Pregnant animals)</b></li> </ul>
<ul style="list-style-type: none"> <li>• <b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795222247 (KVK, West Siang)</b></li> </ul>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department

Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

Integrated Agromet Advisory Services,  
Basar, Arunachal Pradesh.



# Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar - 791 101



(East Kameng)

No: -73/2012-13/Tue

Dated: 8<sup>th</sup> January, 2013

## Weather during the last four days

Rainfall: -

- Seppa - 0.0 mm

## Weather Forecast

• <b>Rainfall</b> - Mainly clear weather is forecasted with possibility of cloudy development on some day
• <b>Temperature</b> - The temperature is likely to be in the range of 8-20°C
• <b>Relative Humidity</b> - The RH is likely to be in the range of 63-80%.

## Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>VEGETABLE CROPS</b>			
Cauliflower /Cabbage/Knol khol/Broccoli/	Vegetative/ Maturity		<ul style="list-style-type: none"> <li>• Mulching with locally available weeds, paddy straws to conserve the soil moisture and to prevent the affect of low temperature</li> </ul>
Ginger/ Turmeric	Maturity		<ul style="list-style-type: none"> <li>• Cure should be taken at harvesting so that the rhizome is not damaged,</li> </ul>
Leafy rapeseed mustard	Vegetative	Flea Beetle	<ul style="list-style-type: none"> <li>• Apply wood ash in the morning hours</li> </ul>
<b>FIELD CROPS</b>			
Potato	Vegetative	Late Blight	<ul style="list-style-type: none"> <li>• <b>Avoid water logging.</b></li> <li>• <b>Monitor the crop for symptoms of blight.</b></li> <li>• Foliar spraying with RidomilMZ 72 WP @ 2.5 grams per litre of water</li> <li>• Earthing up may be done,</li> </ul>
Pea	Vegetative	Aphid	<ul style="list-style-type: none"> <li>• Spray Dimethoate @ 2 ml per litre of water</li> </ul>
Rapeseed/ Mustard	Flowering	Aphid Painted bug	<ul style="list-style-type: none"> <li>• Collect and dispose off the heavily infested twigs,</li> <li>• For painted bug dusting with wood ash is advised,</li> <li>• Mulching to conserve the moisture</li> </ul>
<b>FRUIT CROPS</b>			
Khasi Mandarin	Harvest Post harvest	Lichens/ mosses	<ul style="list-style-type: none"> <li>• In present weather conditions remove lichens/mosses from the affected trees using gunny bags smoothly,</li> <li>• Prune the diseased and dried branches,</li> <li>• Collect and bury fallen fruits,</li> </ul>
<b>LIVESTOCKS</b>			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages	Cold Coccidiosis	<ul style="list-style-type: none"> <li>• Day old chick-Maintain warmth by lighting,</li> <li>• Provide heat by lighting 100W filament bulb @ 1 bulb per 20 newly born chicks,</li> <li>• In present weather conditions frequently turn the litters/avoid hardening of litters,</li> <li>• Provide limited ventilation,</li> <li>• Add multivitamin feeds (More of Vit C)</li> <li>• Add Amprolium to the ration,</li> </ul>
Pigs	New born	Cold	<ul style="list-style-type: none"> <li>• Warming of pig sheds using gunny bags, straw, enclosure,</li> </ul>
Cattle	All stages	FMD Respiratory	<ul style="list-style-type: none"> <li>• Vaccination - of healthy animals</li> <li>• Treatment - sick animals,</li> <li>• Intra muscular injection of Oxi-tetracycline @ 20ml/400 kg body weight animal (<b>Avoid injecting to Pregnant animals</b>)</li> </ul>
Goat/Sheep	All stages	Goat pox	<ul style="list-style-type: none"> <li>• Vaccinate with PPX,</li> <li>• Provide standard feeds,</li> </ul>
<ul style="list-style-type: none"> <li>• <b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795222247 (KVK, West Siang)</b></li> </ul>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

Integrated Agromet Advisory Services



# Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar - 791 101



(East-Siang)

No: -73/2012-13/Tue

Dated: 8<sup>th</sup> January, 2013

## Weather during the last four days

- Maximum temperature: 21.8 °C
- Minimum Temperature: 12.9 °C
- Rainfall:  
    Passihat -0.2 mm
- Relative humidity (%):64-89

## Weather Forecast

- Rainfall- Mainly clear weather is forecasted with possibility of cloudy development on some day.
- Temperature - The temperature is likely to be in the range of 8-19°C.
- Relative Humidity - The RH is likely to be in the range of 62-82%.

## Weather Based Agro-meteorological Advisories Bulletin

## Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>VEGETABLE CROPS</b>			
Cauliflower /Cabbage/Knol khol/Broccoli/	Vegetative/ Maturity		<ul style="list-style-type: none"> <li>• Mulching with locally available weeds, paddy straws to conserve the soil moisture and to prevent the affect of low temperature</li> </ul>
Ginger/ Turmeric	Maturity		<ul style="list-style-type: none"> <li>• Cure should be taken at harvesting so that the rhizome is not damaged,</li> </ul>
Leafy rapeseed mustard	Vegetative	Flea Beetle	<ul style="list-style-type: none"> <li>• Apply wood ash in the morning hours</li> </ul>
<b>FIELD CROPS</b>			
Potato	Vegetative	Late Blight	<ul style="list-style-type: none"> <li>• <b><u>Avoid water logging.</u></b></li> <li>• <b><u>Monitor the crop for symptoms of blight.</u></b></li> <li>• Foliar spraying with RidomilMZ 72 WP @ 2.5 grams per litre of water</li> <li>• Earthing up may be done,</li> </ul>
Pea	Vegetative	Aphid	<ul style="list-style-type: none"> <li>• Spray Dimethoate @ 2 ml per litre of water</li> </ul>
Rapeseed/ Mustard	Flowering	Aphid Painted bug	<ul style="list-style-type: none"> <li>• Collect and dispose off the heavily infested twigs,</li> <li>• For painted bug dusting with wood ash is advised,</li> <li>• Mulching to conserve the moisture</li> </ul>
<b>FRUIT CROPS</b>			
Khasi Mandarin	Harvest Post harvest	Lichens/ mosses	<ul style="list-style-type: none"> <li>• In present weather conditions remove lichens/mosses from the affected trees using gunny bags smoothly,</li> <li>• Prune the diseased and dried branches,</li> <li>• Collect and bury fallen fruits,</li> </ul>
<b>LIVESTOCKS</b>			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages	Cold  Coccidiosis	<ul style="list-style-type: none"> <li>• Day old chick-Maintain warmth by lighting,</li> <li>• Provide heat by lighting 100W filament bulb @ 1 bulb per 20 newly born chicks,</li> <li>• In present weather conditions frequently turn the litters/avoid hardening of litters,</li> <li>• Provide limited ventilation,</li> <li>• Add multivitamin feeds (More of Vit C)</li> <li>• Add Amprolium to the ration,</li> </ul>
Pigs	New born	Cold	<ul style="list-style-type: none"> <li>• Warming of pig sheds using gunny bags, straw, enclosure,</li> </ul>
Cattle	All stages	FMD  Respiratory	<ul style="list-style-type: none"> <li>• Vaccination - of healthy animals</li> <li>• Treatment - sick animals,</li> <li>• Intra muscular injection of Oxi-tetracycline @ 20ml/400 kg body weight animal <b>(Avoid injecting to Pregnant animals)</b></li> </ul>
Goat/Sheep	All stages	Goat pox	<ul style="list-style-type: none"> <li>• Vaccinate with PPX,</li> </ul>

			• Provide standard feeds,
--	--	--	---------------------------

- |   |  |  |  |
|---|--|--|--|
| • <b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795222247 (KVK, West Siang)</b> |  |  |  |
|---|--|--|--|

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

**Integrated Agromet Advisory Services,  
Basar, Arunachal Pradesh.**



# Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre  
West Siang, Basar – 791 101



(Lohit)

No: -73/2012-13/Tue

Dated: 8<sup>th</sup> January, 2013

## Weather during the last four days

- Rainfall: -
  - Tezu -27.2 mm
  - Namsai -0.0 mm

## Weather Forecast

- |  |
|--|
| • <u>Rainfall</u> - Weather will be mainly clear with possibility of partly cloudy development on some days. |
| • <u>Temperature</u> - The temperature is likely to be in the range of 8-20°C.                               |
| • <u>Relative Humidity</u> - The RH is likely to be in the range of 62-84%.                                  |

## Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>VEGETABLE CROPS</b>			
Cauliflower /Cabbage/Knol khol/Broccoli/	Vegetative/ Maturity		<ul style="list-style-type: none"> <li>• Mulching with locally available weeds, paddy straws to conserve the soil moisture and to prevent the affect of low temperature</li> </ul>
<b>FIELD CROPS</b>			
Potato	Vegetative	Late Blight	<ul style="list-style-type: none"> <li>• <b>Avoid water logging.</b></li> <li>• <b>Monitor the crop for symptoms of blight.</b></li> <li>• Foliar spraying with RidomilMZ 72 WP @ 2.5 grams per litre of water</li> <li>• Earthing up may be done,</li> </ul>
Pea	Vegetative	Aphid	<ul style="list-style-type: none"> <li>• Spray Dimethoate @ 2 ml per litre of water</li> </ul>
Rapeseed/ Mustard	Flowering	Aphid Painted bug	<ul style="list-style-type: none"> <li>• Collect and dispose off the heavily infested twigs,</li> <li>• For painted bug dusting with wood ash is advised,</li> <li>• Mulching to conserve the moisture</li> </ul>
<b>FRUIT CROPS</b>			
Khasi Mandarin	Harvest Post harvest	Lichens/ mosses	<ul style="list-style-type: none"> <li>• In present weather conditions remove lichens/mosses from the affected trees using gunny bags smoothly,</li> <li>• Prune the diseased and dried branches,</li> <li>• Collect and bury fallen fruits,</li> </ul>
<b>LIVESTOCKS</b>			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages	Cold  Coccidiosis	<ul style="list-style-type: none"> <li>• Day old chick-Maintain warmth by lighting,</li> <li>• Provide heat by lighting 100W filament bulb @ 1 bulb per 20 newly born chicks,</li> <li>• In present weather conditions frequently turn the litters/avoid hardening of litters,</li> <li>• Provide limited ventilation,</li> <li>• Add multivitamin feeds (More of Vit C)</li> <li>• Add Amprolium to the ration,</li> </ul>
Pigs	New born	Cold	<ul style="list-style-type: none"> <li>• Warming of pig sheds using gunny bags, straw, enclosure,</li> </ul>
Cattle	All stages	FMD  Respiratory	<ul style="list-style-type: none"> <li>• Vaccination – of healthy animals</li> <li>• Treatment – sick animals,</li> <li>• Intra muscular injection of Oxi-tetracycline @ 20ml/400 kg body weight animal (<b>Avoid injecting to Pregnant animals</b>)</li> </ul>
<ul style="list-style-type: none"> <li>• <b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795222247 (KVK, West Siang)</b></li> </ul>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department

Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

**Integrated Agromet Advisory Services, Basar, Arunachal Pradesh.**



# Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar - 791 101



## (Lower Subansiri)

No: -73/2012-13/Tue

Dated: 8<sup>th</sup> January, 2013

### Weather during the last four days

- Rainfall: -  
1. Ziro - NA

### Weather Forecast

- Rainfall- Partial Cloudiness with possibility of light rains for the next ½ days and thereafter weather will be clear.
- Temperature - The temperature is likely to be in the range of 3-15°C.
- Relative Humidity - The RH is likely to be in the range of 58-82%.

### Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>VEGETABLE CROPS</b>			
Cauliflower /Cabbage/Knol khol/Broccoli/	Vegetative/ Maturity		<ul style="list-style-type: none"> <li>• Mulching with locally available weeds, paddy straws to conserve the soil moisture and to prevent the affect of low temperature</li> </ul>
Leafy rapeseed mustard	Vegetative	Flea Beetle	<ul style="list-style-type: none"> <li>• Apply wood ash in the morning hours</li> </ul>
<b>FIELD CROPS</b>			
Rapeseed/ Mustard	Flowering	Aphid Painted bug	<ul style="list-style-type: none"> <li>• Collect and dispose off the heavily infested twigs,</li> <li>• For painted bug dusting with wood ash is advised,</li> <li>• Mulching to conserve the moisture</li> </ul>
<b>FRUIT CROPS</b>			
Khasi Mandarin	Harvest Post harvest	Lichens/ mosses	<ul style="list-style-type: none"> <li>• In present weather conditions remove lichens/mosses from the affected trees using gunny bags smoothly,</li> <li>• Prune the diseased and dried branches,</li> <li>• Collect and bury fallen fruits,</li> </ul>
<b>LIVESTOCKS</b>			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages	Cold  Coccidiosis	<ul style="list-style-type: none"> <li>• Day old chick-Maintain warmth by lighting,</li> <li>• Provide heat by lighting 100W filament bulb @ 1 bulb per 20 newly born chicks,</li> <li>• In present weather conditions frequently turn the litters/avoid hardening of litters,</li> <li>• Provide limited ventilation,</li> <li>• Add multivitamin feeds (More of Vit C)</li> <li>• Add Amprolium to the ration,</li> </ul>
Pigs	New born	Cold	<ul style="list-style-type: none"> <li>• Warming of pig sheds using gunny bags, straw, enclosure,</li> </ul>
Cattle	All stages	FMD  Respiratory	<ul style="list-style-type: none"> <li>• Vaccination - of healthy animals</li> <li>• Treatment - sick animals,</li> <li>• Intra muscular injection of Oxi-tetracycline @ 20ml/400 kg body weight animal <b>(Avoid injecting to Pregnant animals)</b></li> </ul>
<ul style="list-style-type: none"> <li>• <b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 0379522247 (KVK, West Siang)</b></li> </ul>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department.

Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-222247/37, -96 (Fax)

**Integrated Agromet Advisory Services,  
Basar, Arunachal Pradesh.**



# Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre  
West Siang, Basar - 791 101



## (Papum Pare)

No: -73/2012-13/Tue

Dated: 8<sup>th</sup> January, 2013

### Weather during the last four days

- Maximum temperature: 23.2°C
- Minimum Temperature: 9.1°C
- Relative Humidity (%):82-93
- Rainfall: -
  1. Itanagar - 1.4 mm
  2. Naharlagun - 2.0 mm

### Weather Forecast

• <u>Rainfall</u> - Clear/Partially cloudy weather is forecasted.
• <u>Temperature</u> - The temperature is likely to be in the range of 9-22°C
• <u>Relative Humidity</u> - The RH is likely to be in the range of 60-75%.

### Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>VEGETABLE CROPS</b>			
Cauliflower /Cabbage/Knol khol/Broccoli/	Vegetative/ Maturity		<ul style="list-style-type: none"> <li>• Mulching with locally available weeds, paddy straws to conserve the soil moisture and to prevent the affect of low temperature</li> </ul>
Ginger/ Turmeric	Maturity		<ul style="list-style-type: none"> <li>• Cure should be taken at harvesting so that the rhizome is not damaged,</li> </ul>
Leafy rapeseed mustard	Vegetative	Flea Beetle	<ul style="list-style-type: none"> <li>• Apply wood ash in the morning hours</li> </ul>
<b>FIELD CROPS</b>			
Potato	Vegetative	Late Blight	<ul style="list-style-type: none"> <li>• <b>Avoid water logging.</b></li> <li>• <b>Monitor the crop for symptoms of blight.</b></li> <li>• Foliar spraying with RidomilMZ 72 WP @ 2.5 grams per litre of water</li> <li>• Earthing up may be done,</li> </ul>
Pea	Vegetative	Aphid	<ul style="list-style-type: none"> <li>• Spray Dimethoate @ 2 ml per litre of water</li> </ul>
Rapeseed/ Mustard	Flowering	Aphid Painted bug	<ul style="list-style-type: none"> <li>• Collect and dispose off the heavily infested twigs,</li> <li>• For painted bug dusting with wood ash is advised,</li> <li>• Mulching to conserve the moisture</li> </ul>
<b>FRUIT CROPS</b>			
Khasi Mandarin	Harvest Post harvest	Lichens/ mosses	<ul style="list-style-type: none"> <li>• In present weather conditions remove lichens/mosses from the affected trees using gunny bags smoothly,</li> <li>• Prune the diseased and dried branches,</li> <li>• Collect and bury fallen fruits,</li> </ul>
Kiwi	Post Harvest		<ul style="list-style-type: none"> <li>• Training and pruning is advised,</li> </ul>
<b>LIVESTOCKS</b>			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages	Cold Coccidiosis	<ul style="list-style-type: none"> <li>• Day old chick-Maintain warmth by lighting,</li> <li>• Provide heat by lighting 100W filament bulb @ 1 bulb per 20 newly born chicks,</li> <li>• In present weather conditions frequently turn the litters/avoid hardening of litters,</li> <li>• Provide limited ventilation,</li> <li>• Add multivitamin feeds (More of Vit C)</li> <li>• Add Amprolium to the ration,</li> </ul>
Pigs	New born	Cold	<ul style="list-style-type: none"> <li>• Warming of pig sheds using gunny bags, straw, enclosure,</li> </ul>
Cattle	All stages	FMD Respiratory	<ul style="list-style-type: none"> <li>• Vaccination - of healthy animals</li> <li>• Treatment - sick animals,</li> <li>• Intra muscular injection of Oxi-tetracycline @ 20ml/400 kg body weight animal (<b>Avoid injecting to Pregnant animals</b>)</li> </ul>
Goat/Sheep	All stages	Goat pox	<ul style="list-style-type: none"> <li>• Vaccinate with PPX,</li> <li>• Provide standard feeds,</li> </ul>



- **Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 0379522247 (KVK, West Siang)**

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

**Integrated Agromet Advisory Services,  
Basar, Arunachal Pradesh.**



# Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar - 791 101



(Tawang)

No: -73/2012-13/Tue

Dated: 8<sup>th</sup> January, 2013

## Weather during the last four days

- Rainfall: -  
1. Tawang -NA

## Weather Forecast

• <b>Rainfall</b> - Weather will be cloudy/partly cloudy with possibility of light showers/snow for next ½ days and thereafter clear weather will be there.
• <b>Temperature</b> - The temperature is likely to be in the range of -1-9°C.
• <b>Relative Humidity</b> - The RH is likely to be in the range of 58-92%.

## Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
Apple	Post harvest	Trunk Borer	<ul style="list-style-type: none"> <li>• Apply Bordeaux mixture to the tree trunk,</li> <li>• Burn the cut branches, don't heap in and around the crop field,</li> </ul>
		Training	<ul style="list-style-type: none"> <li>• Train the plant and remove dead branches and stem cankers and apply Bordeaux paste on cut portions,</li> </ul>
Kiwi	Post Harvest		<ul style="list-style-type: none"> <li>• Training and pruning is advised,</li> </ul>

### LIVESTOCKS

Livestock	Stage	Pest/Disease	Precautionary Practices to be followed
Poultry	All stages	Cold	<ul style="list-style-type: none"> <li>• Day old chick-Maintain warmth by lighting,</li> <li>• Provide heat by lighting 100W filament bulb @ 1 bulb per 20 newly born chicks,</li> <li>• In present weather conditions frequently turn the litters/avoid hardening of litters,</li> <li>• Provide limited ventilation,</li> <li>• Add multivitamin feeds (More of Vit C)</li> <li>• Add Amprolium to the ration,</li> </ul>
		Coccidiosis	
Pigs	New born	Cold	<ul style="list-style-type: none"> <li>• Warming of pig sheds using gunny bags, straw, enclosure,</li> </ul>
Cattle	All stages	FMD	<ul style="list-style-type: none"> <li>• Vaccination - of healthy animals</li> <li>• Treatment - sick animals,</li> <li>• Intra muscular injection of Oxi-tetracycline @ 20ml/400 kg body weight animal <b>(Avoid injecting to Pregnant animals)</b></li> </ul>
		Respiratory	

Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795222247 (KVK, West Siang)

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department

Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

Integrated Agromet Advisory Services  
Basar, Arunachal Pradesh



# Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre  
West Siang, Basar – 791 101



(Tirap)

No: -73/2012-13/Tue

Dated: 8<sup>th</sup> January, 2013

## Weather during the last four days

- Rainfall: -
  - Khonsa -NA
  - Deomali -1.6 mm

## Weather Forecast

• <u>Rainfall</u> : Weather will be mainly clear with possibility of partly cloudy on some days.
• <u>Temperature</u> - The temperature is likely to be in the range of 7-17°C.
• <u>Relative Humidity</u> - The RH is likely to be in the range of 60-76%.

## Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>VEGETABLE CROPS</b>			
Cauliflower /Cabbage/Knolkhol/Broccoli/	Vegetative/ Maturity		<ul style="list-style-type: none"> <li>• Mulching with locally available weeds, paddy straws to conserve the soil moisture and to prevent the affect of low temperature</li> </ul>
<b>FIELD CROPS</b>			
Potato	Vegetative	Late Blight	<ul style="list-style-type: none"> <li>• <b>Avoid water logging.</b></li> <li>• <b>Monitor the crop for symptoms of blight.</b></li> <li>• Foliar spraying with RidomilMZ 72 WP @ 2.5 grams per litre of water</li> <li>• Earthing up may be done,</li> </ul>
Pea	Vegetative	Aphid	<ul style="list-style-type: none"> <li>• Spray Dimethoate @ 2 ml per litre of water</li> </ul>
Rapeseed/ Mustard	Flowering	Aphid Painted bug	<ul style="list-style-type: none"> <li>• Collect and dispose off the heavily infested twigs,</li> <li>• For painted bug dusting with wood ash is advised,</li> <li>• Mulching to conserve the moisture</li> </ul>
<b>FRUIT CROPS</b>			
Khasi Mandarin	Harvest Post harvest	Lichens/ mosses	<ul style="list-style-type: none"> <li>• In present weather conditions remove lichens/mosses from the affected trees using gunny bags smoothly,</li> <li>• Prune the diseased and dried branches,</li> <li>• Collect and bury fallen fruits,</li> </ul>
<b>LIVESTOCKS</b>			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages	Cold  Coccidiosis	<ul style="list-style-type: none"> <li>• Day old chick-Maintain warmth by lighting,</li> <li>• Provide heat by lighting 100W filament bulb @ 1 bulb per 20 newly born chicks,</li> <li>• In present weather conditions frequently turn the litters/avoid hardening of litters,</li> <li>• Provide limited ventilation,</li> <li>• Add multivitamin feeds (More of Vit C)</li> <li>• Add Amprolium to the ration,</li> </ul>
Pigs	New born	Cold	<ul style="list-style-type: none"> <li>• Warming of pig sheds using gunny bags, straw, enclosure,</li> </ul>
Cattle	All stages	FMD  Respiratory	<ul style="list-style-type: none"> <li>• Vaccination - of healthy animals</li> <li>• Treatment - sick animals,</li> <li>• Intra muscular injection of Oxi-tetracycline @ 20ml/400 kg body weight animal (<b>Avoid injecting to Pregnant animals</b>)</li> </ul>
<ul style="list-style-type: none"> <li>• <b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 0379522247 (KVK, West Siang)</b></li> </ul>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department

Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

Integrated Agromet Advisory Services, Basar, Arunachal Pradesh.



# Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar - 791 101



## (Upper-Siang)

No: -73/2012-13/Tue

Dated: 8<sup>th</sup> January, 2013

### Weather during the last four days

- Rainfall: -
  - Tuting – 3.6 mm

### Weather Forecast

- Rainfall – Cloudy weather with possibility of light rain is forecasted for the next 1/2 days and thereafter weather will be clear.
- Temperature - The temperature is likely to be in the range of 2-15°C.
- Relative Humidity - The RH is likely to be in the range of 64-88%.

### Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>VEGETABLE CROPS</b>			
Cauliflower /Cabbage/Knolkhol/Broccoli/	Vegetative/ Maturity		<ul style="list-style-type: none"> <li>• Mulching with locally available weeds, paddy straws to conserve the soil moisture and to prevent the affect of low temperature</li> </ul>
Ginger/ Turmeric	Maturity		<ul style="list-style-type: none"> <li>• Cure should be taken at harvesting so that the rhizome is not damaged,</li> </ul>
Leafy rapeseed mustard	Vegetative	Flea Beetle	<ul style="list-style-type: none"> <li>• Apply wood ash in the morning hours</li> </ul>
<b>FIELD CROPS</b>			
Potato	Vegetative	Late Blight	<ul style="list-style-type: none"> <li>• <b>Avoid water logging.</b></li> <li>• <b>Monitor the crop for symptoms of blight.</b></li> <li>• Foliar spraying with RidomilMZ 72 WP @ 2.5 grams per litre of water</li> <li>• Earthing up may be done,</li> </ul>
<b>FRUIT CROPS</b>			
Khasi Mandarin	Harvest Post harvest	Lichens/ mosses	<ul style="list-style-type: none"> <li>• In present weather conditions remove lichens/mosses from the affected trees using gunny bags smoothly,</li> <li>• Prune the diseased and dried branches,</li> <li>• Collect and bury fallen fruits,</li> </ul>
<b>LIVESTOCKS</b>			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages	Cold Coccidiosis	<ul style="list-style-type: none"> <li>• Day old chick-Maintain warmth by lighting,</li> <li>• Provide heat by lighting 100W filament bulb @ 1 bulb per 20 newly born chicks,</li> <li>• In present weather conditions frequently turn the litters/avoid hardening of litters,</li> <li>• Provide limited ventilation,</li> <li>• Add multivitamin feeds (More of Vit C)</li> <li>• Add Amprolium to the ration,</li> </ul>
Pigs	New born	Cold	<ul style="list-style-type: none"> <li>• Warming of pig sheds using gunny bags, straw, enclosure,</li> </ul>
Cattle	All stages	FMD Respiratory	<ul style="list-style-type: none"> <li>• Vaccination – of healthy animals</li> <li>• Treatment – sick animals,</li> <li>• Intra muscular injection of Oxi-tetracycline @ 20ml/400 kg body weight animal (<b>Avoid injecting to Pregnant animals</b>)</li> </ul>
Goat/Sheep	All stages	Goat pox	<ul style="list-style-type: none"> <li>• Vaccinate with PPX,</li> <li>• Provide standard feeds,</li> </ul>
<ul style="list-style-type: none"> <li>• <b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795222247 (KVK, West Siang)</b></li> </ul>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
 Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)



# Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar - 791 101



## (Upper-Subansiri)

No: -73/2012-13/Tue

Dated: 8<sup>th</sup> January, 2013

### Weather during the last four days

- Rainfall: -  
1. Daporijo-NA

### Weather Forecast

• <u>Rainfall</u> - Cloudy/partially cloudy weather is forecasted for the next 2/3 days.
• <u>Temperature</u> - The temperature is likely to be in the range of 4-17°C.
• <u>Relative Humidity</u> - The RH is likely to be in the range of 64-84%.

### Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>VEGETABLE CROPS</b>			
Cauliflower /Cabbage/Knolkhol/Broccoli/	Vegetative/ Maturity		<ul style="list-style-type: none"> <li>• Mulching with locally available weeds, paddy straws to conserve the soil moisture and to prevent the affect of low temperature</li> </ul>
Ginger/ Turmeric	Maturity		<ul style="list-style-type: none"> <li>• Cure should be taken at harvesting so that the rhizome is not damaged,</li> </ul>
Leafy rapeseed mustard	Vegetative	Flea Beetle	<ul style="list-style-type: none"> <li>• Apply wood ash in the morning hours</li> </ul>
<b>FIELD CROPS</b>			
Potato	Vegetative	Late Blight	<ul style="list-style-type: none"> <li>• <b>Avoid water logging.</b></li> <li>• <b>Monitor the crop for symptoms of blight.</b></li> <li>• Foliar spraying with RidomilMZ 72 WP @ 2.5 grams per litre of water</li> <li>• Earthing up may be done,</li> </ul>
Pea	Vegetative	Aphid	<ul style="list-style-type: none"> <li>• Spray Dimethoate @ 2 ml per litre of water</li> </ul>
Rapeseed/ Mustard	Flowering	Aphid Painted bug	<ul style="list-style-type: none"> <li>• Collect and dispose off the heavily infested twigs,</li> <li>• For painted bug dusting with wood ash is advised,</li> <li>• Mulching to conserve the moisture</li> </ul>
<b>FRUIT CROPS</b>			
Khasi Mandarin	Harvest Post harvest	Lichens/ mosses	<ul style="list-style-type: none"> <li>• In present weather conditions remove lichens/mosses from the affected trees using gunny bags smoothly,</li> <li>• Prune the diseased and dried branches,</li> <li>• Collect and bury fallen fruits,</li> </ul>
<b>LIVESTOCKS</b>			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages	Cold  Coccidiosis	<ul style="list-style-type: none"> <li>• Day old chick-Maintain warmth by lighting,</li> <li>• Provide heat by lighting 100W filament bulb @ 1 bulb per 20 newly born chicks,</li> <li>• In present weather conditions frequently turn the litters/avoid hardening of litters,</li> <li>• Provide limited ventilation,</li> <li>• Add multivitamin feeds (More of Vit C)</li> <li>• Add Amprolium to the ration,</li> </ul>
Pigs	New born	Cold	<ul style="list-style-type: none"> <li>• Warming of pig sheds using gunny bags, straw, enclosure,</li> </ul>
Cattle	All stages	FMD  Respiratory	<ul style="list-style-type: none"> <li>• Vaccination - of healthy animals</li> <li>• Treatment - sick animals,</li> <li>• Intra muscular injection of Oxi-tetracycline @ 20ml/400 kg body weight animal (<b>Avoid injecting to Pregnant animals</b>)</li> </ul>
Goat/Sheep	All stages	Goat pox	<ul style="list-style-type: none"> <li>• Vaccinate with PPX,</li> <li>• Provide standard feeds,</li> </ul>
<ul style="list-style-type: none"> <li>• <b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 0379522247 (KVK, West Siang)</b></li> </ul>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department

Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)





# Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar - 791 101



## (West Kameng)

No: -73/2012-13/Tue

Dated: 8<sup>th</sup> January, 2013

### Weather during the last three days

- Rainfall: - Bomdila - 1.0 mm  
Bhalokpong - 0.0 mm

### Weather Forecast

- |   |
|---|
| • <u>Rainfall</u> - Clear/Partially cloudy weather is forecasted with possibility of light rains in some areas, |
| • <u>Temperature</u> - The temperature is likely to be in the range of 7-20°C.                                  |
| • <u>Relative Humidity</u> - The RH is likely to be in the range of 69-84%.                                     |

### Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
Apple	Post harvest	Trunk Borer	<ul style="list-style-type: none"> <li>• Apply Bordeaux mixture to the tree trunk,</li> <li>• Burn the cut branches, don't heap in and around the crop field,</li> </ul>
		Training	<ul style="list-style-type: none"> <li>• Train the plant and remove dead branches and stem cankers and apply Bordeaux paste on cut portions,</li> </ul>
Kiwi	Post Harvest		<ul style="list-style-type: none"> <li>• Training and pruning is advised,</li> </ul>
<b>LIVESTOCKS</b>			
Livestock	Stage	Pest/Disease	Precautionary Practices to be followed
Poultry	All stages	Cold	<ul style="list-style-type: none"> <li>• Day old chick-Maintain warmth by lighting,</li> <li>• Provide heat by lighting 100W filament bulb @ 1 bulb per 20 newly born chicks,</li> <li>• In present weather conditions frequently turn the litters/avoid hardening of litters,</li> <li>• Provide limited ventilation,</li> <li>• Add multivitamin feeds (More of Vit C)</li> <li>• Add Amprolium to the ration,</li> </ul>
		Coccidiosis	
Pigs	New born	Cold	<ul style="list-style-type: none"> <li>• Warming of pig sheds using gunny bags, straw, enclosure,</li> </ul>
Cattle	All stages	FMD	<ul style="list-style-type: none"> <li>• Vaccination - of healthy animals</li> <li>• Treatment - sick animals,</li> <li>• Intra muscular injection of Oxi-tetracycline @ 20ml/400 kg body weight animal <b>(Avoid injecting to Pregnant animals)</b></li> </ul>
		Respiratory	
<b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795222247 (KVK, West Siang)</b>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department'

Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-222247/37, -96 (Fax)

Integrated Agromet Advisory Services  
Basar, Arunachal Pradesh



# Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar - 791 101



## (West Siang)

No: -73/2012-13/Tue

Dated: 8<sup>th</sup> January, 2013

### Weather during the period

- During the last two days weather was cloudy and rain was recorded.
- Maximum temperature:14.5°C
- Minimum Temperature:7.5°C
- Rainfall: **15.0 mm**
- Evapotranspiration: NA
- Relative Humidity (%):-69-99
- Soil moisture (m<sup>2</sup>/ m<sup>2</sup>):-0.28
- Prominent Wind Direction: Morning-South East & Evening-North West
- Average Wind Speed : 0.2 kmph

(Source-Agromet Observatory, Research Farm, Gori, ICAR, AP Centre, Basar)

### Weather Forecast

- Rainfall:-Clear/Partially cloudy weather is forecasted.
- Temperature - The temperature is likely to be in the range of 7-18°C.
- Relative Humidity - The RH is likely to be in the range of 63-82%.

### Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>VEGETABLE CROPS</b>			
Cauliflower /Cabbage/Knol khol/Broccoli/	Vegetative/ Maturity		<ul style="list-style-type: none"> <li>• Mulching with locally available weeds, paddy straws to conserve the soil moisture and to prevent the affect of low temperature</li> </ul>
Ginger/ Turmeric	Maturity		<ul style="list-style-type: none"> <li>• Cure should be taken at harvesting so that the rhizome is not damaged,</li> </ul>
Leafy rapeseed mustard	Vegetative	Flea Beetle	<ul style="list-style-type: none"> <li>• Apply wood ash in the morning hours</li> </ul>
<b>FIELD CROPS</b>			
Potato	Vegetative	Late Blight	<ul style="list-style-type: none"> <li>• <b>Avoid water logging.</b></li> <li>• <b>Monitor the crop for symptoms of blight.</b></li> <li>• Foliar spraying with RidomilMZ 72 WP @ 2.5 grams per litre of water</li> <li>• Earthing up may be done,</li> </ul>
Pea	Vegetative	Aphid	<ul style="list-style-type: none"> <li>• Spray Dimethoate @ 2 ml per litre of water</li> </ul>
Rapeseed/ Mustard	Flowering	Aphid Painted bug	<ul style="list-style-type: none"> <li>• Collect and dispose off the heavily infested twigs,</li> <li>• For painted bug dusting with wood ash is advised,</li> <li>• Mulching to conserve the moisture</li> </ul>
<b>FRUIT CROPS</b>			
Khasi Mandarin	Harvest Post harvest	Lichens/ mosses	<ul style="list-style-type: none"> <li>• In present weather conditions remove lichens/mosses from the affected trees using gunny bags smoothly,</li> <li>• Prune the diseased and dried branches,</li> <li>• Collect and bury fallen fruits,</li> </ul>
<b>LIVESTOCKS</b>			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages	Cold  Coccidiosis	<ul style="list-style-type: none"> <li>• Day old chick-Maintain warmth by lighting,</li> <li>• Provide heat by lighting 100W filament bulb @ 1 bulb per 20 newly born chicks,</li> <li>• In present weather conditions frequently turn the litters/avoid hardening of litters,</li> <li>• Provide limited ventilation,</li> <li>• Add multivitamin feeds (More of Vit C)</li> <li>• Add Amprolium to the ration,</li> </ul>
Pigs	New born	Cold	<ul style="list-style-type: none"> <li>• Warming of pig sheds using gunny bags, straw, enclosure,</li> </ul>
Cattle	All stages	FMD  Respiratory	<ul style="list-style-type: none"> <li>• Vaccination - of healthy animals</li> <li>• Treatment - sick animals,</li> <li>• Intra muscular injection of Oxi-tetracycline @ 20ml/400 kg body weight animal <b>(Avoid injecting to Pregnant animals)</b></li> </ul>



Rabbits	All stages	Snuffles	<ul style="list-style-type: none"> <li>Use broad spectrum antibiotic treatment like Gentamycin, anrofloxacin under proper supervision of veterinary officials</li> </ul>
Goat/Sheep	All stages	Goat pox	<ul style="list-style-type: none"> <li>Vaccinate with PPX,</li> <li>Provide standard feeds,</li> </ul>
<ul style="list-style-type: none"> <li><b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795222247 (KVK, West Siang)</b></li> </ul>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

**Integrated Agromet Advisory Services  
Basar, Arunachal Pradesh**