



# Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar – 791 101



(Lohit)

No: 55/2014-15/Fri

Dated: 10<sup>th</sup> October, 2014

## Agro Advisory Bulletin for the period 11<sup>th</sup> – 15<sup>th</sup> October 2014

### Weather summery of the previous week

Max Temp: 22.0<sup>0</sup>C, Min Temp: 9.0-10.0<sup>0</sup>C, Wind Speed: 7-8Kmph, Wind Direction: 68-71Degree, Rainfall: 0.0mm.

### Weather forecast valid up to 15<sup>th</sup> October, 2014

The possibility of Rainfall is 21.0mm for the next five days. Max Temp: 22.0<sup>0</sup>C, Min Temp: 9.0-10.0<sup>0</sup>C, Cloud Cover: sky will be partly cloudy i.e. 2-5 Okta, Max RH: 87-98%, Min RH: 46-74%, Wind Speed: 3-8Kmph, Wind Direction: 65-169Degree.

*Weekly cumulative rainfall : 21.0mm*

## Weather Based Agro-meteorological Advisories Bulletin

- **Obtain seeds from reputed/certified/reliable source,**

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>FIELD CROPS</b>			
Kharif Rice (Early transplanted)	Harvesting		<ul style="list-style-type: none"><li>• Threshing followed by proper drying for safe storage,</li></ul>
Kharif Rice (Normal transplanted)	Tillering/ Booting	Blast/Gundhi bug	<ul style="list-style-type: none"><li>• Need based alternate foliar spray of Hexaconazole 5 EC(2ml/l) &amp; Carbendazim 12WP+Mancozeb 63WP(2g/l) at 15days interval beginning from first appearance of the disease,</li><li>• Dust with Malathion 5D @ 20kg/ha. Repeat weekly,</li><li>• Properly maintain the water level in the field,</li></ul>
<b>FRUIT CROPS</b>			
Lemon/other citrus fruits	Fruit Bearing	Fruit fly/Trunk borer	<ul style="list-style-type: none"><li>• Monitor fruit fly population with Methyl eugenol traps,</li><li>• Spray Malathion (2ml/l) in the early stage to avoid fruit fly infestation,</li><li>• Drenching of base soil with Dichlorvos 76 EC @ 4ml/l of water. Repeat weekly till the tree regain its greenery,</li></ul>
Pineapple	Fruiting/ Harvesting		<ul style="list-style-type: none"><li>• Clean the vicinity of plants by weeding and loosen up the soil.</li><li>• Remove sly and use it as a new planting material.</li><li>• Sucker dipping treatment with Bavistin @ 3g + Chlorpyriphos 20EC@ 2ml/l of water to protect against soil borne disease and insect.</li></ul>
<b>LIVESTOCKS</b>			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages		<ul style="list-style-type: none"><li>• Avoid dampness, litter should be periodically changed.</li><li>• Provide clean and fresh water daily,</li><li>• Vaccination of birds as per the recommended age,</li><li>• Artificial heating through lighting of electric bulb during night time to protect the poultry birds (day old to 25 days old) from low temperature.</li></ul>
Pigs	New born	Cold	<ul style="list-style-type: none"><li>• Pig shed/stay should keep dry and avoid dampness during rainy days,</li><li>• Avoid stagnant water near the shed,</li><li>• Vaccinate against swine fever,</li><li>• Avoid of eating human excreta,</li><li>• Provide ad libitum water,</li></ul>
Cattle	All stages	FMD	<ul style="list-style-type: none"><li>• Vaccination of animal with FMD,</li><li>• Treatment – sick animals,</li></ul>
Fish	Stocking	-	<ul style="list-style-type: none"><li>• Before stocking, make free the weeds and predators in the pond</li></ul>
<ul style="list-style-type: none"><li>• <b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795226247/37</b></li></ul>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax).

**Agrometeorological Field Unit  
ICAR, BASAR**



# Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar – 791 101



**(Lower-Dibang- Valley)**

**No: 55/2014-15/Fri**

**Dated: 10<sup>th</sup> October, 2014**

## **Agro Advisory Bulletin for the period 11<sup>th</sup> – 15<sup>th</sup> October 2014**

### **Weather summary of the previous week**

Max Temp: 20.0-21.0°C, Min Temp: 5.0-6.0°C, Wind Speed: 6-9Kmph, Wind Direction: 53-96Degree, Rainfall: 0.0mm.

### **Weather forecast valid up to 15<sup>th</sup> October, 2014**

The possibility of Rainfall is 0.0mm for the next five days. Max Temp: 20.0-21.0°C, Min Temp: 5.0-6.0°C, Cloud Cover: sky will be partly cloudy i.e. 1-5 Okta, Max RH: 76-94%, Min RH: 38-59%, Wind Speed: 4-8Kmph, Wind Direction: 66-106Degree.

***Weekly cumulative rainfall : 0.0mm***

## **Weather Based Agro-meteorological Advisories Bulletin**

- **Obtain seeds from reputed/certified/reliable source,**

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>FIELD CROPS</b>			
Kharif Rice (Early transplanted)	Harvesting		<ul style="list-style-type: none"><li>• Threshing followed by proper drying for safe storage,</li></ul>
Kharif Rice (Normal transplanted)	Tillering/ Booting	Blast/Gundhi bug	<ul style="list-style-type: none"><li>• Need based alternate foliar spray of Hexaconazole 5 EC(2ml/l) &amp; Carbendazim 12WP+Mancozeb 63WP(2g/l) at 15days interval beginning from first appearance of the disease,</li><li>• Dust with Malathion 5D @ 20kg/ha. Repeat weekly,</li><li>• Properly maintain the water level in the field,</li></ul>
<b>FRUIT CROPS</b>			
Lemon/other citrus fruits	Fruit Bearing	Fruit fly/Trunk borer	<ul style="list-style-type: none"><li>• Monitor fruit fly population with Methyl eugenol traps,</li><li>• Spray Malathion (2ml/l) in the early stage to avoid fruit fly infestation,</li><li>• Drenching of base soil with Dichlorvos 76 EC @ 4ml/l of water. Repeat weekly till the tree regain its greenery,</li></ul>
Pineapple	Fruiting/ Harvesting		<ul style="list-style-type: none"><li>• Clean the vicinity of plants by weeding and loosen up the soil.</li><li>• Remove sly and use it as a new planting material.</li><li>• Sucker dipping treatment with Bavistin @ 3g + Chlorpyriphos 20EC@ 2ml/lt of water to protect against soil borne disease and insect.</li></ul>
Apple	Fruit Development		<ul style="list-style-type: none"><li>• Spray Boric Acid 100g / 100lt water and Urea 500 g / 100 lt water at green bud or pink bud stage.</li><li>• Protection of Scab/ Fungal infection/core rot- spray Mencozeb (600g/ 200 lt water) or Diefenconazole (300 ml/ 200lt water) at pink bub stage. Spray Hexaconazole (100 ml/ 200 lt water) or carbendazime (100 gm/ 200 lt water) after flower shedding or just start of fruit development.</li><li>• Spray Bavistin (0.5g/l) to control scab</li><li>• Spray Boron (1g/l)</li><li>• Spray 50ml Megister/Megistic in 200 litres of water in case of mite attack is noticed in apple orchards,</li></ul>
<b>LIVESTOCKS</b>			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages		<ul style="list-style-type: none"><li>• Avoid dampness, litter should be periodically changed,</li><li>• Provide clean and fresh water daily,</li><li>• Vaccination of birds as per the recommended age,</li><li>• Artificial heating through lighting of electric bulb during night time to protect the poultry birds (day old to 25 days old) from low temperature,</li></ul>
Pigs	New born	Cold	<ul style="list-style-type: none"><li>• Pig shed/stay should keep dry and avoid dampness during rainy days,</li><li>• Avoid stagnant water near the shed,</li><li>• Vaccinate against swine fever,</li><li>• Avoid of eating human excreta,</li><li>• Provide ad libitum water,</li></ul>

Cattle	All stages	FMD	<ul style="list-style-type: none"><li>• Vaccination of animal with FMD,</li><li>• Treatment – sick animals,</li></ul>
Goat/Sheep	All stages	Goat pox	<ul style="list-style-type: none"><li>• Vaccinate with PPR,</li><li>• Provide standard feeds,</li><li>• Vaccination &amp; deworming at regular interval,</li></ul>
Fish	Stocking	-	<ul style="list-style-type: none"><li>• Before stocking, make free the weeds and predators in the pond</li></ul>
<ul style="list-style-type: none"><li>• <b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795226247/37</b></li></ul>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit**  
**ICAR, BASAR**



# Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar – 791 101



**(Lower Subansiri)**

**No: 55/2014-15/Fri**

**Dated: 10<sup>th</sup> October, 2014**

## **Agro Advisory Bulletin for the period 11<sup>th</sup> – 15<sup>th</sup> October 2014**

### **Weather summery of the previous week**

Max Temp: 16.0-17.0<sup>0</sup>C, Min Temp: 7.0-8.0<sup>0</sup>C, Wind Speed: 4-5Kmph, Wind Direction: 151-281Degree, Rainfall: 4.0mm.

### **Weather forecast valid up to 15<sup>th</sup> October, 2014**

The possibility of Rainfall is 42.0mm for the next five days. Max Temp: 16.0-17.0<sup>0</sup>C, Min Temp: 7.0-8.0<sup>0</sup>C, Cloud Cover: sky will be partly cloudy i.e. 3-7 Okta, Max RH: 88-97%, Min RH: 53-80%, Wind Speed: 2-4Kmph, Wind Direction: 185-261Degree.

***Weekly cumulative rainfall : 42.0mm***

## **Weather Based Agro-meteorological Advisories Bulletin**

- Obtain seeds from reputed/certified/reliable source,**

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>FIELD CROPS</b>			
Kharif Rice (Early transplanted)	Harvesting		<ul style="list-style-type: none"><li>Threshing followed by proper drying for safe storage,</li></ul>
Kharif Rice (Normal transplanted)	Tillering/ Booting	Blast/Gundhi bug	<ul style="list-style-type: none"><li>Need based alternate foliar spray of Hexaconazole 5 EC(2ml/l) &amp; Carbendazim 12WP+Mancozeb 63WP(2g/l) at 15days interval beginning from first appearance of the disease,</li><li>Dust with Malathion 5D @ 20kg/ha. Repeat weekly,</li><li>Properly maintain the water level in the field,</li></ul>
<b>FRUIT CROPS</b>			
Lemon/other citrus fruits	Fruit Bearing	Fruit fly/Trunk borer	<ul style="list-style-type: none"><li>Monitor fruit fly population with Methyl eugenol traps,</li><li>Spray Malathion (2ml/l) in the early stage to avoid fruit fly infestation,</li><li>Drenching of base soil with Dichlorvos 76 EC @ 4ml/l of water. Repeat weekly till the tree regain its greenery,</li></ul>
Pineapple	Fruiting/ Harvesting		<ul style="list-style-type: none"><li>Clean the vicinity of plants by weeding and loosen up the soil.</li><li>Remove sly and use it as a new planting material.</li><li>Sucker dipping treatment with Bavistin @ 3g + Chlorpyriphos 20EC@ 2ml/l of water to protect against soil borne disease and insect.</li></ul>
Apple	Fruit Development		<ul style="list-style-type: none"><li>Spray Boric Acid 100g / 100lt water and Urea 500 g / 100 lt water at green bud or pink bud stage,</li><li>Protection of Scab/ Fungal infection/core rot- spray Mencozeb (600g/ 200 lt water) or Diefenconazole (300 ml/ 200lt water) at pink bub stage. Spray Hexaconazole (100 ml/ 200 lt water) or carbendazime (100 gm/ 200 lt water) after flower shedding or just start of fruit development.</li><li>Spray Bavistin (0.5g/l) to control scab</li><li>Spray Boron (1g/l)</li><li>Spray 50ml Megister/Megistic in 200 litres of water in case of mite attack is noticed in apple orchards,</li></ul>
<b>LIVESTOCKS</b>			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages		<ul style="list-style-type: none"><li>Avoid dampness, litter should be periodically changed,</li><li>Provide clean and fresh water daily,</li><li>Vaccination of birds as per the recommended age,</li><li>Artificial heating through lighting of electric bulb during night time to protect the poultry birds (day old to 25 days old) from low temperature.</li></ul>
Pigs	New born	Cold	<ul style="list-style-type: none"><li>Pig shed/stay should keep dry and avoid dampness during rainy days,</li><li>Avoid stagnant water near the shed,</li><li>Vaccinate against swine fever,</li><li>Avoid of eating human excreta,</li><li>Provide ad libitum water,</li></ul>

Cattle	All stages	FMD	<ul style="list-style-type: none"><li>• Vaccination of animal with FMD,</li><li>• Treatment – sick animals,</li></ul>
Goat/Sheep	All stages	Goat pox	<ul style="list-style-type: none"><li>• Vaccinate with PPR,</li><li>• Provide standard feeds,</li><li>• Vaccination &amp; deworming at regular interval,</li></ul>
Fish	Stocking	-	<ul style="list-style-type: none"><li>• Before stocking, make free the weeds and predators in the pond</li></ul>
<ul style="list-style-type: none"><li>• <b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795226247/37</b></li></ul>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37,-96 (Fax)

**Agrometeorological Field Unit**  
**ICAR, BASAR**



# Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar – 791 101



**(Papum Pare)**

**No: 55/2014-15/Fri**

**Dated: 10<sup>th</sup> October, 2014**

## **Agro Advisory Bulletin for the period 11<sup>th</sup> – 15<sup>th</sup> October 2014**

### **Weather summery of the previous week**

Max Temp: 29.0-30.0<sup>0</sup>C, Min Temp: 19.0-20.0<sup>0</sup>C, Wind Speed: 5-6Kmph, Wind Direction: 68-277Degree, Rainfall: 3.0mm.

### **Weather forecast valid up to 15<sup>th</sup> October, 2014**

The possibility of Rainfall is 40.0mm for the next five days. Max Temp: 29.0-30.0<sup>0</sup>C, Min Temp: 19.0-20.0<sup>0</sup>C, Cloud Cover: sky will be partly cloudy i.e. 2-6 Okta, Max RH: 80-92%, Min RH: 45-71%, Wind Speed: 2-6Kmph, Wind Direction: 193-280Degree.

***Weekly cumulative rainfall : 40.0mm***

### **Weather Based Agro-meteorological Advisories Bulletin**

- Obtain seeds from reputed/certified/reliable source,**

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>FIELD CROPS</b>			
Kharif Rice (Early transplanted)	Harvesting		<ul style="list-style-type: none"><li>Threshing followed by proper drying for safe storage,</li></ul>
Kharif Rice (Normal transplanted)	Tillering/ Booting	Blast/Gundhi bug	<ul style="list-style-type: none"><li>Need based alternate foliar spray of Hexaconazole 5 EC(2ml/l) &amp; Carbendazim 12WP+Mancozeb 63WP(2g/l) at 15days interval beginning from first appearance of the disease,</li><li>Dust with Malathion 5D @ 20kg/ha. Repeat weekly,</li><li>Properly maintain the water level in the field,</li></ul>
<b>FRUIT CROPS</b>			
Lemon/other citrus fruits	Fruit Bearing	Fruit fly/Trunk borer	<ul style="list-style-type: none"><li>Monitor fruit fly population with Methyl eugenol traps,</li><li>Spray Malathion (2ml/l) in the early stage to avoid fruit fly infestation,</li><li>Drenching of base soil with Dichlorvos 76 EC @ 4ml/l of water. Repeat weekly till the tree regain its greenery,</li></ul>
Pineapple	Fruiting/ Harvesting		<ul style="list-style-type: none"><li>Clean the vicinity of plants by weeding and loosen up the soil.</li><li>Remove sly and use it as a new planting material.</li><li>Sucker dipping treatment with Bavistin @ 3g + Chlorpyriphos 20EC@ 2ml/lt of water to protect against soil borne disease and insect.</li></ul>
<b>LIVESTOCKS</b>			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages		<ul style="list-style-type: none"><li>Day old chick-Maintain warmth by lighting,</li><li>Provide heat by lighting 100W filament bulb @ 1 bulb per 20 newly born chicks,</li><li>Vaccination of birds as per the recommended age,</li><li>Prevent from Coccidiosis give Anti coucidid drops,</li><li>During day time provide proper ventilation,</li><li>Add multivitamin feeds (More of Vit C)</li><li>Add Amprolium to the ration,</li><li>Regular check the litter materials to prevent from caking,</li></ul>
Pigs	New born/ All stages	Cold	<ul style="list-style-type: none"><li>Vaccinate against swine fever,</li><li>Avoid stagnant water near the shed,</li><li>Avoid of eating human excreta,</li><li>Provide ad libitum water,</li><li>Deworming of piglets after 2 months,</li></ul>
Cattle	All stages	FMD	<ul style="list-style-type: none"><li>Vaccination of animal with FMD,</li><li>Treatment – sick animals,</li></ul>
Fish	Stocking	-	<ul style="list-style-type: none"><li>Before stocking, make free the weeds and predators in the pond</li></ul>
Dog	All stages		<ul style="list-style-type: none"><li>Vaccinate against rabies ,</li></ul>
Sheep/Goat	All stages		<ul style="list-style-type: none"><li>Vaccination &amp; deworming at regular interval.</li><li>Vaccinate with PPR,</li></ul>
<ul style="list-style-type: none"><li><b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795226247/37</b></li></ul>			

**Agrometeorological Field Unit**  
**ICAR, BASAR**





# Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar – 791 101



**(Tawang)**

**No: 55/2014-15/Fri**

**Dated: 10<sup>th</sup> October, 2014**

## **Agro Advisory Bulletin for the period 11<sup>th</sup> – 15<sup>th</sup> October 2014**

### **Weather summary of the previous week**

Max Temp: 16.0<sup>0</sup>C, Min Temp: 4.0-6.0<sup>0</sup>C,  
Wind Speed: 5-8Kmph, Wind Direction: 99-  
261Degree, Rainfall: 4.0mm.

### **Weather forecast valid up to 15<sup>th</sup> October, 2014**

The possibility of Rainfall is 25.0mm for the next five days. Max Temp: 16.0<sup>0</sup>C, Min Temp: 4.0-6.0<sup>0</sup>C, Cloud Cover: sky will be partly cloudy i.e. 2-7 Okta, Max RH: 84-92%, Min RH: 46-80%, Wind Speed: 4-8Kmph, Wind Direction: 98-259Degree.

***Weekly cumulative rainfall : 25.0mm***

## **Weather Based Agro-meteorological Advisories Bulletin**

<b>Crop</b>	<b>Stage</b>	<b>Pest/Disease</b>	<b>Precautionary Practices to be followed</b>
Apple	Fruit Development		<ul style="list-style-type: none"><li>• Protection of Scab/ Fungal infection/core rot- spray Mencozeb (600g/ 200 lt water) or Diefenconazole (300 ml/ 200lt water) at pink bub stage. Spray Hexaconazole (100 ml/ 200 lt water) or carbendazime (100 gm/ 200 lt water) after flower shedding or just start of fruit development.</li><li>• Spray Bavistin (0.5g/l) to control scab</li><li>• Spray Boron (1g/l)</li><li>• Spray 50ml Megister/Megistic in 200 litres of water in case of mite attack is noticed in apple orchards,</li></ul>
<b>LIVESTOCKS</b>			
<b>Livestock</b>	<b>Stage</b>	<b>Pest/Disease</b>	<b>Precautionary Practices to be followed</b>
Poultry	All stages		<ul style="list-style-type: none"><li>• Day old chick-Maintain warmth by lighting,</li><li>• Vaccination of birds as per the recommended age,</li><li>• Provide heat by lighting 100W filament bulb @ 1 bulb per 20 newly born chicks,</li><li>• Prevent from Coccidiosis give anti coucidid drops,</li><li>• Provide limited ventilation,</li><li>• Add multivitamin feeds (More of Vit C)</li><li>• Add Amprolium to the ration,</li><li>Regular check the litter materials to prevent from caking,</li></ul>
Pigs	New born	Cold	<ul style="list-style-type: none"><li>• Warming of pig sheds using gunny bags, straw, enclosure,</li><li>• Vaccinate against swine fever,</li><li>• Avoid stagnant water near the shed,</li><li>• Avoid of eating human excreta,</li><li>• Provide ad libitum water,</li><li>• Deworming of piglets after 2 months,</li></ul>
Cattle	All stages	FMD	<ul style="list-style-type: none"><li>• Vaccination of animal with FMD,</li><li>• Treatment – sick animals,</li></ul>
Fish	Stocking	-	<ul style="list-style-type: none"><li>• Before stocking, make free the weeds and predators in the pond</li></ul>
Dog	All stages		<ul style="list-style-type: none"><li>• Vaccinate against rabies ,</li></ul>
Sheep/Goat	All stages		<ul style="list-style-type: none"><li>• Vaccination &amp; deworming at regular interval.</li><li>• Vaccinate with PPR,</li></ul>

**Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795226247 (KVK, West Siang)**

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department

Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit  
ICAR, BASAR**





# Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar – 791 101



**(Tirap)**

**No: 55/2014-15/Fri**

**Dated: 10<sup>th</sup> October, 2014**

## **Agro Advisory Bulletin for the period 11<sup>th</sup> – 15<sup>th</sup> October 2014**

### **Weather summery of the previous week**

### **Weather forecast valid up to 15<sup>th</sup> October, 2014**

Max Temp: 30.0-31.0<sup>0</sup>C, Min Temp: 14.0-16.0<sup>0</sup>C, Wind Speed: 2-4Kmph, Wind Direction: 136-145Degree, Rainfall: 0.0mm.

The possibility of Rainfall is 0.0mm for the next five days. Max Temp: 30.0-31.0<sup>0</sup>C, Min Temp: 14.0-16.0<sup>0</sup>C, Cloud Cover: sky will be partly clear i.e. 1-5 Okta, Max RH: 92-98%, Min RH: 44-74%, Wind Speed: 0-4Kmph, Wind Direction: 115-267Degree.

***Weekly cumulative rainfall : 0.0mm***

## **Weather Based Agro-meteorological Advisories Bulletin**

- **Obtain seeds from reputed/certified/reliable source,**

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>FIELD CROPS</b>			
Kharif Rice (Early transplanted)	Harvesting		<ul style="list-style-type: none"><li>• Threshing followed by proper drying for safe storage,</li></ul>
Kharif Rice (Normal transplanted)	Tillering/ Booting	Blast/Gundhi bug	<ul style="list-style-type: none"><li>• Need based alternate foliar spray of Hexaconazole 5 EC(2ml/l) &amp; Carbendazim 12WP+Mancozeb 63WP(2g/l) at 15days interval beginning from first appearance of the disease,</li><li>• Dust with Malathion 5D @ 20kg/ha. Repeat weekly,</li><li>• Properly maintain the water level in the field,</li></ul>
<b>FRUIT CROPS</b>			
Lemon/other citrus fruits	Fruit Bearing	Fruit fly/Trunk borer	<ul style="list-style-type: none"><li>• Monitor fruit fly population with Methyl eugenol traps,</li><li>• Spray Malathion (2ml/l) in the early stage to avoid fruit fly infestation,</li><li>• Drenching of base soil with Dichlorvos 76 EC @ 4ml/l of water. Repeat weekly till the tree regain its greenery,</li></ul>
Pineapple	Fruiting/ Harvesting		<ul style="list-style-type: none"><li>• Clean the vicinity of plants by weeding and loosen up the soil.</li><li>• Remove sly and use it as a new planting material.</li><li>• Sucker dipping treatment with Bavistin @ 3g + Chlorpyriphos 20EC@ 2ml/l of water to protect against soil borne disease and insect.</li></ul>
<b>LIVESTOCKS</b>			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages		<ul style="list-style-type: none"><li>• Avoid dampness, litter should be periodically changed.</li><li>• Provide clean and fresh water daily,</li><li>• Vaccination of birds as per the recommended age,</li><li>• Prevent from Coccidiosis give anti coucidid drops,</li><li>• Artificial heating through lighting of electric bulb during night time to protect the poultry birds (day old to 25 days old) from low temperature.</li></ul>
Pigs	New born	Cold	<ul style="list-style-type: none"><li>• Pig shed/stay should keep dry and avoid dampness during rainy days,</li><li>• Avoid stagnant water near the shed,</li><li>• Vaccinate against swine fever,</li><li>• Avoid of eating human excreta,</li><li>• Provide ad libitum water,</li></ul>
Cattle	All stages	FMD	<ul style="list-style-type: none"><li>• Vaccination of animal with FMD,</li><li>• Treatment – sick animals,</li></ul>
Fish	Stocking	-	<ul style="list-style-type: none"><li>• Before stocking, make free the weeds and predators in the pond</li></ul>
<ul style="list-style-type: none"><li>• <b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795226247/37</b></li></ul>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department

Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit  
ICAR, BASAR**



# Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar – 791 101



(Upper Siang)

No: 55/2014-15/Fri

Dated: 10<sup>th</sup> October, 2014

## Agro Advisory Bulletin for the period 11<sup>th</sup> – 15<sup>th</sup> October 2014

### Weather summary of the previous week

Max Temp: 19.0-20.0<sup>0</sup>C, Min Temp: 9.0-10.0<sup>0</sup>C, Wind Speed: 4-6Kmph, Wind Direction: 71-130Degree, Rainfall: 0.0mm.

### Weather forecast valid up to 15<sup>th</sup> October, 2014

The possibility of Rainfall is 12.0mm for the next five days. Max Temp: 19.0-20.0<sup>0</sup>C, Min Temp: 9.0-10.0<sup>0</sup>C, Cloud Cover: sky will be partly cloudy i.e. 1-6 Okta, Max RH: 81-91%, Min RH: 39-63%, Wind Speed: 2-5Kmph, Wind Direction: 117-165Degree.

**Weekly cumulative rainfall : 12.0mm**

## Weather Based Agro-meteorological Advisories Bulletin

- **Obtain seeds from reputed/certified/reliable source,**

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>FIELD CROPS</b>			
Kharif Rice (Early transplanted)	Harvesting		<ul style="list-style-type: none"><li>• Threshing followed by proper drying for safe storage,</li></ul>
Kharif Rice (Normal transplanted)	Tillering/Booting	Blast/Gundhi bug	<ul style="list-style-type: none"><li>• Need based alternate foliar spray of Hexaconazole 5 EC(2ml/l) &amp; Carbendazim 12WP+Mancozeb 63WP(2g/l) at 15days interval beginning from first appearance of the disease,</li><li>• Dust with Malathion 5D @ 20kg/ha. Repeat weekly,</li><li>• Properly maintain the water level in the field,</li></ul>
<b>FRUIT CROPS</b>			
Lemon/other citrus fruits	Fruit Bearing	Fruit fly/Trunk borer	<ul style="list-style-type: none"><li>• Monitor fruit fly population with Methyl eugenol traps,</li><li>• Spray Malathion (2ml/l) in the early stage to avoid fruit fly infestation,</li><li>• Drenching of base soil with Dichlorvos 76 EC @ 4ml/l of water. Repeat weekly till the tree regain its greenery,</li></ul>
Pineapple	Fruiting/Harvesting		<ul style="list-style-type: none"><li>• Clean the vicinity of plants by weeding and loosen up the soil.</li><li>• Remove sly and use it as a new planting material.</li><li>• Sucker dipping treatment with Bavistin @ 3g + Chlorpyrifos 20EC@ 2ml/l of water to protect against soil borne disease and insect.</li></ul>
<b>LIVESTOCKS</b>			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages		<ul style="list-style-type: none"><li>• Avoid dampness, litter should be periodically changed.</li><li>• Provide clean and fresh water daily,</li><li>• Vaccination of birds as per the recommended age,</li><li>• Prevent from Coccidiosis give anti coucidid drops,</li><li>• Artificial heating through lighting of electric bulb during night time to protect the poultry birds (day old to 25 days old) from low temperature.</li></ul>
Pigs	New born	Cold	<ul style="list-style-type: none"><li>• Pig shed/stay should keep dry and avoid dampness during rainy days,</li><li>• Avoid stagnant water near the shed,</li><li>• Vaccinate against swine fever,</li><li>• Avoid of eating human excreta,</li><li>• Provide ad libitum water,</li></ul>
Cattle	All stages	FMD	<ul style="list-style-type: none"><li>• Vaccination of animal with FMD,</li><li>• Treatment – sick animals,</li></ul>
Dog	All stages		<ul style="list-style-type: none"><li>• Vaccinate against rabies,</li></ul>
Rabbits	All stages	Snuffles	<ul style="list-style-type: none"><li>• Use broad spectrum antibiotic treatment like Gentamycin, anrofloxacin under proper supervision of veterinary officials</li></ul>

- **Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795226247/37**

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department

Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit  
ICAR, BASAR



# Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar – 791 101



**(Upper Subansiri)**

**No: 55/2014-15/Fri**

**Dated: 10<sup>th</sup> October, 2014**

## **Agro Advisory Bulletin for the period 11<sup>th</sup> – 15<sup>th</sup> October 2014**

### **Weather summery of the previous week**

Max Temp: 15.0-16.0<sup>0</sup>C, Min Temp: 5.0-6.0<sup>0</sup>C, Wind Speed: 3-6Kmph, Wind Direction: 229-253Degree, Rainfall: 6.0mm.

### **Weather forecast valid up to 15<sup>th</sup> October, 2014**

The possibility of Rainfall is 32.0mm for the next five days. Max Temp: 15.0-16.0<sup>0</sup>C, Min Temp: 5.0-6.0<sup>0</sup>C, Cloud Cover: sky will be partly cloudy i.e. 2-7 Okta, Max RH: 87-95%, Min RH: 48-80%, Wind Speed: 2-5Kmph, Wind Direction: 200-250Degree.

***Weekly cumulative rainfall : 32.0mm***

## **Weather Based Agro-meteorological Advisories Bulletin**

- Obtain seeds from reputed/certified/reliable source,**

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>FIELD CROPS</b>			
Kharif Rice (Early transplanted)	Harvesting		<ul style="list-style-type: none"><li>Threshing followed by proper drying for safe storage,</li></ul>
Kharif Rice (Normal transplanted)	Tillering/ Booting	Blast/Gundhi bug	<ul style="list-style-type: none"><li>Need based alternate foliar spray of Hexaconazole 5 EC(2ml/l) &amp; Carbendazim 12WP+Mancozeb 63WP(2g/l) at 15days interval beginning from first appearance of the disease,</li><li>Dust with Malathion 5D @ 20kg/ha. Repeat weekly,</li><li>Properly maintain the water level in the field,</li></ul>
<b>FRUIT CROPS</b>			
Lemon/other citrus fruits	Fruit Bearing	Fruit fly/Trunk borer	<ul style="list-style-type: none"><li>Monitor fruit fly population with Methyl eugenol traps,</li><li>Spray Malathion (2ml/l) in the early stage to avoid fruit fly infestation,</li><li>Drenching of base soil with Dichlorvos 76 EC @ 4ml/l of water. Repeat weekly till the tree regain its greenery,</li></ul>
Pineapple	Fruiting/ Harvesting		<ul style="list-style-type: none"><li>Clean the vicinity of plants by weeding and loosen up the soil.</li><li>Remove sly and use it as a new planting material.</li><li>Sucker dipping treatment with Bavistin @ 3g + Chlorpyriphos 20EC@ 2ml/lt of water to protect against soil borne disease and insect.</li></ul>
Apple	Fruit Development		<ul style="list-style-type: none"><li>Spray Boric Acid 100g / 100lt water and Urea 500 g / 100 lt water at green bud or pink bud stage,</li><li>Protection of Scab/ Fungal infection/core rot- spray Mencozeb (600g/ 200 lt water) or Diefenconazole (300 ml/ 200lt water) at pink bub stage. Spray Hexaconazole (100 ml/ 200 lt water) or carbendazime (100 gm/ 200 lt water) after flower shedding or just start of fruit development.</li><li>Spray Bavistin (0.5g/l) to control scab,</li><li>Spray Boron (1g/l),</li><li>Spray 50ml Megister/Megistic in 200 litres of water in case of mite attack is noticed in apple orchards,</li></ul>
<b>LIVESTOCKS</b>			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages		<ul style="list-style-type: none"><li>Avoid dampness, litter should be periodically changed.</li><li>Provide clean and fresh water daily,</li><li>Vaccination of birds as per the recommended age,</li><li>Prevent from Coccidiosis give anti coucidid drops,</li><li>Artificial heating through lighting of electric bulb during night time to protect the poultry birds (day old to 25 days old) from low temperature.</li></ul>
Pigs	New born	Cold	<ul style="list-style-type: none"><li>Pig shed/stay should keep dry and avoid dampness during rainy days,</li><li>Avoid stagnant water near the shed,</li><li>Vaccinate against swine fever,</li><li>Avoid of eating human excreta,</li></ul>

			<ul style="list-style-type: none"> <li>• Provide ad libitum water,</li> </ul>
Cattle	All stages	FMD	<ul style="list-style-type: none"> <li>• Vaccination of animal with FMD,</li> <li>• Treatment – sick animals,</li> </ul>
Goat/Sheep	All stages	Goat pox	<ul style="list-style-type: none"> <li>• Vaccinate with PPR,</li> <li>• Provide standard feeds,</li> <li>• Vaccination &amp; deworming at regular interval.</li> </ul>
<ul style="list-style-type: none"> <li>• <b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795226247/37</b></li> </ul>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit**  
**ICAR, BASAR**



Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar – 791 101



(West Kameng)

No: 55/2014-15/Fri

Dated: 10<sup>th</sup> October, 2014

Agro Advisory Bulletin for the period 11<sup>th</sup> – 15<sup>th</sup> October 2014

Weather summery of the previous week

Weather forecast valid up to 15<sup>th</sup> October, 2014

Max Temp: 17.0-18.0 <sup>0</sup> C, Min Temp: 7.0-8.0 <sup>0</sup> C, Wind Speed: 6-7Kmph, Wind Direction: 46-209Degree, Rainfall: 11.0mm.	The possibility of Rainfall is 45.0mm for the next five days. Max Temp: 17.0-18.0 <sup>0</sup> C, Min Temp: 7.0-8.0 <sup>0</sup> C, Cloud Cover: sky will be partly cloudy i.e. 2-7Okta, Max RH: 87-96%, Min RH: 50-78%, Wind Speed: 2-8Kmph, Wind Direction: 48-212Degree.
--	---

Weekly cumulative rainfall : 45.0mm

Weather Based Agro-meteorological Advisories Bulletin

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
Apple	Fruit Development		<ul style="list-style-type: none"><li>Protection of Scab/ Fungal infection/core rot- spray Mencozeb (600g/ 200 lt water) or Diefenconazole (300 ml/ 200lt water) at pink bub stage. Spray Hexaconazole (100 ml/ 200 lt water) or carbendazime (100 gm/ 200 lt water) after flower shedding or just start of fruit development.</li><li>Spray Bavistin (0.5g/l) to control scab</li><li>Spray Boron (1g/l)</li><li>Spray 50ml Megister/Megistic in 200 litres of water in case of mite attack is noticed in apple orchards,</li></ul>
LIVESTOCKS			
Livestock	Stage	Pest/Disease	Precautionary Practices to be followed
Poultry	All stages		<ul style="list-style-type: none"><li>Day old chick-Maintain warmth by lighting,</li><li>Vaccination of birds as per the recommended age,</li><li>Provide heat by lighting 100W filament bulb @ 1 bulb per 20 newly born chicks,</li><li>Prevent from Coccidiosis give anti coucidid drops,</li><li>Provide limited ventilation,</li><li>Add multivitamin feeds (More of Vit C)</li></ul>
Pigs	New born	Cold	<ul style="list-style-type: none"><li>Vaccinate against swine fever,</li><li>Avoid stagnant water near the shed,</li><li>Avoid of eating human excreta,</li><li>Provide ad libitum water,</li><li>Deworming of piglets after 2 months,</li></ul>
Cattle	All stages	FMD	<ul style="list-style-type: none"><li>Vaccination of animal with FMD,</li><li>Treatment – sick animals,</li></ul>
Fish	Stocking	-	<ul style="list-style-type: none"><li>Before stocking, make free the weeds and predators in the pond</li></ul>
Dog	All stages		<ul style="list-style-type: none"><li>Vaccinate against rabies,</li></ul>
Sheep/Goat	All stages		<ul style="list-style-type: none"><li>Vaccination &amp; deworming at regular interval.</li><li>Vaccinate with PPR,</li></ul>

Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795226247 (KVK, West Siang)

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit  
ICAR, BASAR





# Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar – 791 101



**(West Siang)**

No: 55/2014-15/Fri

Dated: 10<sup>th</sup> October, 2014

## Agro Advisory Bulletin for the period 11<sup>th</sup> – 15<sup>th</sup> October 2014

### Weather summery of the previous week

Rainfall: 4.0mm, Max and min temp varied between 19.0-20.0°C and 10.0°C respectively. The sky was partly clear *i.e.* 1-3 Okta; Wind speed: 4-6kmph, Wind direction: 130-267degree, Max RH: 87-91% and Min RH: 44-47%.

### Weather forecast valid up to 15<sup>th</sup> October, 2014

The possibility of Rainfall is 25.0mm for next five days. Max and min temp will be varied between 19.0-20.0°C and 10.0°C respectively. The sky will be partly cloudy *i.e.* 1-7 Okta; Relative humidity will be varied between 80-92% and 45-70% in the morning and afternoon respectively. The wind will be 2-5kmph. The wind will blow from variable direction at 141-263degree.

*Weekly cumulative rainfall: 25.0mm*

### Weather Based Agro-meteorological Advisories Bulletin

- **Obtain seeds from reputed/certified/reliable source,**

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>FIELD CROPS</b>			
Kharif Rice (Early transplanted)	Harvesting		<ul style="list-style-type: none"><li>• Threshing followed by proper drying for safe storage,</li></ul>
Kharif Rice (Normal transplanted)	Tillering/ Booting	Blast/Gundhi bug	<ul style="list-style-type: none"><li>• Need based alternate foliar spray of Hexaconazole 5 EC(2ml/l) &amp; Carbendazim 12WP+Mancozeb 63WP(2g/l) at 15days interval beginning from first appearance of the disease,</li><li>• Dust with Malathion 5D @ 20kg/ha. Repeat weekly,</li><li>• Properly maintain the water level in the field,</li></ul>
<b>FRUIT CROPS</b>			
Lemon/other citrus fruits	Fruit Bearing	Fruit fly/Trunk borer	<ul style="list-style-type: none"><li>• Monitor fruit fly population with Methyl eugenol traps,</li><li>• Spray Malathion (2ml/l) in the early stage to avoid fruit fly infestation,</li><li>• Drenching of base soil with Dichlorvos 76 EC @ 4ml/l of water. Repeat weekly till the tree regain its greenery,</li></ul>
Pineapple	Fruiting/ Harvesting		<ul style="list-style-type: none"><li>• Clean the vicinity of plants by weeding and loosen up the soil.</li><li>• Remove sly and use it as a new planting material.</li><li>• Sucker dipping treatment with Bavistin @ 3g + Chlorpyrifos 20EC@ 2ml/l of water to protect against soil borne disease and insect.</li></ul>
Apple	Fruit Development		<ul style="list-style-type: none"><li>• Spray Boric Acid 100g / 100lt water and Urea 500 g / 100 lt water at green bud or pink bud stage,</li><li>• Protection of Scab/ Fungal infection/core rot- spray Mencozeb (600g/ 200 lt water) or Diefenconazole (300 ml/ 200lt water) at pink bub stage. Spray Hexaconazole (100 ml/ 200 lt water) or carbendazime (100 gm/ 200 lt water) after flower shedding or just start of fruit development.</li><li>• Spray Bavistin (0.5g/l) to control scab,</li><li>• Spray Boron (1g/l)</li><li>• Spray 50ml Megister/Megistic in 200 litres of water in case of mite attack is noticed in apple orchards,</li></ul>
<b>LIVESTOCKS</b>			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages		<ul style="list-style-type: none"><li>• Avoid dampness, litter should be periodically changed,</li><li>• Provide clean and fresh water daily,</li><li>• Vaccination of birds as per the recommended age,</li><li>• Prevent from Coccidiosis give anti coucidid drops,</li><li>• Artificial heating through lighting of electric bulb during night time to protect the poultry birds (day old to 25 days</li></ul>



			old) from low temperature.
Pigs	New born	Cold	<ul style="list-style-type: none"> <li>Pig shed/stay should keep dry and avoid dampness during rainy days,</li> <li>Avoid stagnant water near the shed,</li> <li>Vaccinate against swine fever,</li> <li>Avoid of eating human excreta,</li> <li>Provide ad libitum water,</li> </ul>
Cattle	All stages	FMD	<ul style="list-style-type: none"> <li>Vaccination of animal with FMD,</li> <li>Treatment – sick animals,</li> </ul>
Rabbits	All stages	Snuffles	<ul style="list-style-type: none"> <li>Use broad spectrum antibiotic treatment like Gentamycin, anrofloxacin under proper supervision of veterinary officials</li> </ul>
Goat/Sheep	All stages	Goat pox	<ul style="list-style-type: none"> <li>Vaccinate with PPR,</li> <li>Provide standard feeds,</li> <li>Vaccination &amp; deworming at regular interval.</li> </ul>
Fish	Stocking	-	<ul style="list-style-type: none"> <li>Before stocking, make free the weeds and predators in the pond</li> </ul>
Dog	All stages		<ul style="list-style-type: none"> <li>Vaccinate against rabies,</li> </ul>
<ul style="list-style-type: none"> <li><b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795226247/37</b></li> </ul>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

Agro Meteorological Field Unit  
 ICAR, BASAR



# Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar – 791 101



(Anjaw)

No: 55/2014-15/Fri

Dated: 10<sup>th</sup> October, 2014

## Agro Advisory Bulletin for the period 11<sup>th</sup> – 15<sup>th</sup> October 2014

### Weather summary of the previous week

Max Temp: 15.0-16.0<sup>0</sup>C, Min Temp: 7.0-8.0<sup>0</sup>C, Wind Speed: 6-9Kmph, Wind Direction: 83-99Degree, Rainfall: 0.0mm.

### Weather forecast valid up to 15<sup>th</sup> October, 2014

The possibility of Rainfall is 3.0mm for the next five days. Max Temp: 15.0-16.0<sup>0</sup>C, Min Temp: 7.0-8.0<sup>0</sup>C, Cloud Cover: sky will be partly cloudy i.e. 2-5Okta, Max RH: 87-97%, Min RH: 44-72%, Wind Speed: 3-8Kmph, Wind Direction: 81-227Degree.

*Weekly cumulative rainfall : 3.0mm*

## Weather Based Agro-meteorological Advisories Bulletin

• Obtain seeds from reputed/certified/reliable source,			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>FIELD CROPS</b>			
Kharif Rice (Early transplanted)	Harvesting		<ul style="list-style-type: none"><li>Threshing followed by proper drying for safe storage,</li></ul>
Kharif Rice (Normal transplanted)	Tillering/ Booting	Blast/Gundhi bug	<ul style="list-style-type: none"><li>Need based alternate foliar spray of Hexaconazole 5 EC(2ml/l) &amp; Carbendazim 12WP+Mancozeb 63WP(2g/l) at 15days interval beginning from first appearance of the disease,</li><li>Dust with Malathion 5D @ 20kg/ha. Repeat weekly,</li><li>Properly maintain the water level in the field,</li></ul>
<b>FRUIT CROPS</b>			
Lemon/other citrus fruits	Fruit Bearing	Fruit fly/Trunk borer	<ul style="list-style-type: none"><li>Monitor fruit fly population with Methyl eugenol traps,</li><li>Spray Malathion (2ml/l) in the early stage to avoid fruit fly infestation,</li><li>Drenching of base soil with Dichlorvos 76 EC @ 4ml/l of water. Repeat weekly till the tree regain its greenery,</li></ul>
Pineapple	Fruiting/ Harvesting		<ul style="list-style-type: none"><li>Clean the vicinity of plants by weeding and loosen up the soil.</li><li>Remove sly and use it as a new planting material.</li><li>Sucker dipping treatment with Bavistin @ 3g + Chlorpyriphos 20EC@ 2ml/l of water to protect against soil borne disease and insect.</li></ul>
<b>LIVESTOCKS</b>			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages		<ul style="list-style-type: none"><li>Vaccination of birds as per the recommended age,</li><li>Avoid dampness, litter should be periodically changed.</li><li>Provide clean and fresh water daily.</li><li>Artificial heating through lighting of electric bulb during night time to protect the poultry birds (day old to 25 days old) from low temperature.</li></ul>
Pigs	New born	Cold	<ul style="list-style-type: none"><li>Pig shed/stay should keep dry and avoid dampness during rainy days,</li><li>Vaccinate against swine fever,</li><li>Avoid of eating human excreta,</li><li>Provide ad libitum water,</li><li>Avoid stagnant water near the shed,</li></ul>
Cattle	All stages	FMD	<ul style="list-style-type: none"><li>Vaccination of animal with FMD,</li><li>Treatment – sick animals,</li><li>Vaccination &amp; deworming at regular interval,</li></ul>
Fish	Stocking	-	<ul style="list-style-type: none"><li>Before stocking, make free the weeds and predators in the pond</li></ul>
• Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795226247/37			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department

Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit  
ICAR, BASAR





# Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar – 791 101



**(Changlang)**

**No: 55/2014-15/Fri**

**Dated: 10<sup>th</sup> October, 2014**

## **Agro Advisory Bulletin for the period 11<sup>th</sup> – 15<sup>th</sup> October 2014**

### **Weather summary of the previous week**

Max Temp: 27.0-28.0<sup>o</sup>C, Min Temp: 12.0<sup>o</sup>C,  
Wind Speed: 6-7Kmph, Wind Direction: 90-  
97Degree, Rainfall: 0.0mm.

### **Weather forecast valid up to 15<sup>th</sup> October, 2014**

The possibility of Rainfall is 7.0mm for the next five days. Max Temp: 27.0-28.0<sup>o</sup>C, Min Temp: 12.0<sup>o</sup>C, Cloud Cover: sky will be partly cloudy i.e. 2-5 Okta, Max RH: 86-98%, Min RH: 43-75%, Wind Speed: 3-7Kmph, Wind Direction: 85-133Degree.

***Weekly cumulative rainfall : 7.0mm***

## **Weather Based Agro-meteorological Advisories Bulletin**

- Obtain seeds from reputed/certified/reliable source,**

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>FIELD CROPS</b>			
Kharif Rice (Early transplanted)	Harvesting		<ul style="list-style-type: none"><li>Threshing followed by proper drying for safe storage,</li></ul>
Kharif Rice (Normal transplanted)	Tillering/ Booting	Blast/Gundhi bug	<ul style="list-style-type: none"><li>Need based alternate foliar spray of Hexaconazole 5 EC(2ml/l) &amp; Carbendazim 12WP+Mancozeb 63WP(2g/l) at 15days interval beginning from first appearance of the disease,</li><li>Dust with Malathion 5D @ 20kg/ha. Repeat weekly,</li><li>Properly maintain the water level in the field,</li></ul>
<b>FRUIT CROPS</b>			
Lemon/other citrus fruits	Fruit Bearing	Fruit fly/Trunk borer	<ul style="list-style-type: none"><li>Monitor fruit fly population with Methyl eugenol traps,</li><li>Spray Malathion (2ml/l) in the early stage to avoid fruit fly infestation,</li><li>Drenching of base soil with Dichlorvos 76 EC @ 4ml/l of water. Repeat weekly till the tree regain its greenery,</li></ul>
Pineapple	Fruiting/ Harvesting		<ul style="list-style-type: none"><li>Clean the vicinity of plants by weeding and loosen up the soil.</li><li>Remove sly and use it as a new planting material.</li><li>Sucker dipping treatment with Bavistin @ 3g + Chlorpyrifos 20EC@ 2ml/lt of water to protect against soil borne disease and insect.</li></ul>
<b>LIVESTOCKS</b>			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages		<ul style="list-style-type: none"><li>Vaccination of birds as per the recommended age,</li><li>Prevent from Coccidiosis give anti coucidid drops,</li><li>Provide heat by lighting 100W filament bulb @ 1 bulb per 20 newly born chicks,</li><li>During day time provide proper ventilation,</li><li>Add multivitamin feeds (More of Vit C)</li><li>Add Amprolium to the ration,</li><li>Regular check the litter materials to prevent from caking,</li></ul>
Pigs	New born/ All stages	Cold	<ul style="list-style-type: none"><li>Vaccinate against swine fever,</li><li>Avoid of eating human excreta,</li><li>Provide ad libitum water,</li><li>Avoid stagnant water near the shed,</li></ul>
Cattle	All stages	FMD	<ul style="list-style-type: none"><li>Vaccination of animal with FMD,</li><li>Treatment – sick animals,</li></ul>
Fish	Stocking	-	<ul style="list-style-type: none"><li>Before stocking, make free the weeds and predators in the pond</li></ul>
Dog	All stages		<ul style="list-style-type: none"><li>Vaccinate against rabies,</li></ul>
Sheep/Goat	All stages		<ul style="list-style-type: none"><li>Vaccination &amp; deworming at regular interval.</li><li>Vaccinate with PPR,</li></ul>
<ul style="list-style-type: none"><li><b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795226247/37</b></li></ul>			

Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone  
Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit  
ICAR, BASAR**



# Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar – 791 101



**(Upper-Dibang- Valley)**

**No: 55/2014-15/Fri**

**Dated: 10<sup>th</sup> October, 2014**

## **Agro Advisory Bulletin for the period 11<sup>th</sup> – 15<sup>th</sup> October 2014**

### **Weather summery of the previous week**

Max Temp: 16.0-17.0<sup>0</sup>C, Min Temp: 8.0-9.0<sup>0</sup>C, Wind Speed: 3-7Kmph, Wind Direction: 83-101Degree, Rainfall: 0.0mm.

### **Weather forecast valid up to 15<sup>th</sup> October, 2014**

The possibility of Rainfall is 3.0mm for the next five days. Max Temp: 16.0-17.0<sup>0</sup>C, Min Temp: 8.0-9.0<sup>0</sup>C, Cloud Cover: sky will be partly cloudy i.e. 1-6 Okta, Max RH: 85-94%, Min RH: 38-75%, Wind Speed: 5-7Kmph, Wind Direction: 98-139Degree.

***Weekly cumulative rainfall : 3.0mm***

## **Weather Based Agro-meteorological Advisories Bulletin**

- **Obtain seeds from reputed/certified/reliable source,**

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>FIELD CROPS</b>			
Kharif Rice (Early transplanted)	Harvesting		<ul style="list-style-type: none"><li>• Threshing followed by proper drying for safe storage,</li></ul>
Kharif Rice (Normal transplanted)	Tillering/ Booting	Blast/Gundhi bug	<ul style="list-style-type: none"><li>• Need based alternate foliar spray of Hexaconazole 5 EC(2ml/l) &amp; Carbendazim 12WP+Mancozeb 63WP(2g/l) at 15days interval beginning from first appearance of the disease,</li><li>• Dust with Malathion 5D @ 20kg/ha. Repeat weekly,</li><li>• Properly maintain the water level in the field,</li></ul>
<b>FRUIT CROPS</b>			
Lemon/other citrus fruits	Fruit Bearing	Fruit fly/Trunk borer	<ul style="list-style-type: none"><li>• Monitor fruit fly population with Methyl eugenol traps,</li><li>• Spray Malathion (2ml/l) in the early stage to avoid fruit fly infestation,</li><li>• Drenching of base soil with Dichlorvos 76 EC @ 4ml/l of water. Repeat weekly till the tree regain its greenery,</li></ul>
Apple	Fruit Development		<ul style="list-style-type: none"><li>• Spray Boric Acid 100g / 100lt water and Urea 500 g / 100 lt water at green bud or pink bud stage.</li><li>• Protection of Scab/ Fungal infection/core rot- spray Mencozeb (600g/ 200 lt water) or Diefenconazole (300 ml/ 200lt water) at pink bub stage. Spray Hexaconazole (100 ml/ 200 lt water) or carbendazime (100 gm/ 200 lt water) after flower shedding or just start of fruit development.</li><li>• Spray Bavistin (0.5g/l) to control scab</li><li>• Spray Boron (1g/l)</li><li>• Spray 50ml Megister/Megistic in 200 litres of water in case of mite attack is noticed in apple orchards,</li></ul>
<b>LIVESTOCKS</b>			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages		<ul style="list-style-type: none"><li>• Avoid dampness, litter should be periodically changed.</li><li>• Provide clean and fresh water daily.</li><li>• Artificial heating through lighting of electric bulb during night time to protect the poultry birds (day old to 25 days old) from low temperature.</li><li>• Vaccination of birds as per the recommended age,</li></ul>
Pigs	New born	Cold	<ul style="list-style-type: none"><li>• Pig shed/stay should keep dry and avoid dampness during rainy days,</li><li>• Vaccinate against swine fever,</li><li>• Avoid of eating human excreta,</li><li>• Provide ad libitum water,</li><li>• Avoid stagnant water near the shed,</li></ul>
Cattle	All stages	FMD	<ul style="list-style-type: none"><li>• Vaccination of animal with FMD,</li><li>• Treatment – sick animals,</li></ul>
Goat/Sheep	All stages	Goat pox	<ul style="list-style-type: none"><li>• Vaccinate with PPR,</li><li>• Provide standard feeds,</li></ul>
Fish	Stocking	-	<ul style="list-style-type: none"><li>• Before stocking, make free the weeds and predators in the pond</li></ul>



- **Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795226247/37**

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit  
ICAR, BASAR**



# Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar – 791 101



**(East Kameng)**

No: 55/2014-15/Fri

Dated: 10<sup>th</sup> October, 2014

## Agro Advisory Bulletin for the period 11<sup>th</sup> – 15<sup>th</sup> October 2014

### Weather summery of the previous week

Max Temp: 16.0-17.0<sup>0</sup>C, Min Temp: 8.0-9.0<sup>0</sup>C, Wind Speed: 5-6Kmph, Wind Direction: 60-169Degree, Rainfall: 11.0mm.

### Weather forecast valid up to 15<sup>th</sup> October, 2014

The possibility of Rainfall is 55.0mm for the next five days. Max Temp: 16.0-17.0<sup>0</sup>C, Min Temp: 8.0-9.0<sup>0</sup>C, Cloud Cover: sky will be partly cloudy i.e. 2-6 Okta, Max RH: 85-97%, Min RH: 50-75%, Wind Speed: 2-6Kmph, Wind Direction: 85-154Degree.

**Weekly cumulative rainfall : 55.0mm**

## Weather Based Agro-meteorological Advisories Bulletin

- **Obtain seeds from reputed/certified/reliable source,**

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>FIELD CROPS</b>			
Kharif Rice (Early transplanted)	Harvesting		<ul style="list-style-type: none"><li>• Threshing followed by proper drying for safe storage,</li></ul>
Kharif Rice (Normal transplanted)	Tillering/ Booting	Blast/Gundhi bug	<ul style="list-style-type: none"><li>• Need based alternate foliar spray of Hexaconazole 5 EC(2ml/l) &amp; Carbendazim 12WP+Mancozeb 63WP(2g/l) at 15days interval beginning from first appearance of the disease,</li><li>• Dust with Malathion 5D @ 20kg/ha. Repeat weekly,</li><li>• Properly maintain the water level in the field,</li></ul>
<b>FRUIT CROPS</b>			
Lemon/other citrus fruits	Fruit Bearing	Fruit fly/Trunk borer	<ul style="list-style-type: none"><li>• Monitor fruit fly population with Methyl eugenol traps,</li><li>• Spray Malathion (2ml/l) in the early stage to avoid fruit fly infestation,</li><li>• Drenching of base soil with Dichlorvos 76 EC @ 4ml/l of water. Repeat weekly till the tree regain its greenery,</li></ul>
Pineapple	Fruiting/ Harvesting		<ul style="list-style-type: none"><li>• Clean the vicinity of plants by weeding and loosen up the soil.</li><li>• Remove sly and use it as a new planting material.</li><li>• Sucker dipping treatment with Bavistin @ 3g + Chlorpyriphos 20EC@ 2ml/l of water to protect against soil borne disease and insect.</li></ul>
<b>LIVESTOCKS</b>			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages		<ul style="list-style-type: none"><li>• Avoid dampness, litter should be periodically changed.</li><li>• Provide clean and fresh water daily.</li><li>• Vaccination of birds as per the recommended age,</li><li>• Artificial heating through lighting of electric bulb during night time to protect the poultry birds (day old to 25 days old) from low temperature.</li></ul>
Pigs	New born	Cold	<ul style="list-style-type: none"><li>• Pig shed/stay should keep dry and avoid dampness during rainy days,</li><li>• Avoid stagnant water near the shed,</li><li>• Vaccinate against swine fever,</li><li>• Avoid of eating human excreta,</li><li>• Provide ad libitum water,</li></ul>
Cattle	All stages	FMD	<ul style="list-style-type: none"><li>• Vaccination of animal with FMD,</li><li>• Treatment – sick animals,</li></ul>
Fish	Stocking	-	<ul style="list-style-type: none"><li>• Before stocking, make free the weeds and predators in the pond</li></ul>

- **Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795226247/37**

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit  
ICAR, BASAR**





# Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar – 791 101



**(East Siang)**

**No: 55/2014-15/Fri**

**Dated: 10<sup>th</sup> October, 2014**

## **Agro Advisory Bulletin for the period 11<sup>th</sup> – 15<sup>th</sup> October 2014**

### **Weather summery of the previous week**

Max Temp: 32.0-33.0<sup>0</sup>C, Min Temp: 22.0-23.0<sup>0</sup>C, Wind Speed: 6Kmph, Wind Direction: 14-286Degree, Rainfall: 0.0mm.

### **Weather forecast valid up to 15<sup>th</sup> October, 2014**

The possibility of Rainfall is 4.0mm for the next five days. Max Temp: 32.0-33.0<sup>0</sup>C, Min Temp: 22.0-23.0<sup>0</sup>C, Cloud Cover: sky will be mainly cloudy i.e. 1-7 Okta, Max RH: 70-91%, Min RH: 41-55%, Wind Speed: 4-7Kmph, Wind Direction: 26-286Degree.

***Weekly cumulative rainfall : 4.0mm***

## **Weather Based Agro-meteorological Advisories Bulletin**

- Obtain seeds from reputed/certified/reliable source,**

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>FIELD CROPS</b>			
Kharif Rice (Early transplanted)	Harvesting		<ul style="list-style-type: none"><li>Threshing followed by proper drying for safe storage,</li></ul>
Kharif Rice (Normal transplanted)	Tillering/ Booting	Blast/Gundhi bug	<ul style="list-style-type: none"><li>Need based alternate foliar spray of Hexaconazole 5 EC(2ml/l) &amp; Carbendazim 12WP+Mancozeb 63WP(2g/l) at 15days interval beginning from first appearance of the disease,</li><li>Dust with Malathion 5D @ 20kg/ha. Repeat weekly,</li><li>Properly maintain the water level in the field,</li></ul>
<b>FRUIT CROPS</b>			
Lemon/other citrus fruits	Fruit Bearing	Fruit fly/Trunk borer	<ul style="list-style-type: none"><li>Monitor fruit fly population with Methyl eugenol traps,</li><li>Spray Malathion (2ml/l) in the early stage to avoid fruit fly infestation,</li><li>Drenching of base soil with Dichlorvos 76 EC @ 4ml/l of water. Repeat weekly till the tree regain its greenery,</li></ul>
Pineapple	Fruiting/ Harvesting		<ul style="list-style-type: none"><li>Clean the vicinity of plants by weeding and loosen up the soil.</li><li>Remove sly and use it as a new planting material.</li><li>Sucker dipping treatment with Bavistin @ 3g + Chlorpyriphos 20EC@ 2ml/l of water to protect against soil borne disease and insect.</li></ul>
<b>LIVESTOCKS</b>			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages		<ul style="list-style-type: none"><li>Avoid dampness, litter should be periodically changed.</li><li>Provide clean and fresh water daily,</li><li>Vaccination of birds as per the recommended age,</li><li>Artificial heating through lighting of electric bulb during night time to protect the poultry birds (day old to 25 days old) from low temperature.</li></ul>
Pigs	New born	Cold	<ul style="list-style-type: none"><li>Pig shed/stay should keep dry and avoid dampness during rainy days,</li><li>Vaccinate against swine fever,</li><li>Avoid of eating human excreta,</li><li>Provide ad libitum water,</li><li>Avoid stagnant water near the shed,</li></ul>
Cattle	All stages	FMD	<ul style="list-style-type: none"><li>Vaccination of animal with FMD,</li><li>Treatment – sick animals,</li></ul>
Goat/Sheep	All stages	Goat pox	<ul style="list-style-type: none"><li>Vaccinate with PPR,</li><li>Provide standard feeds,</li><li>Vaccination &amp; deworming at regular interval,</li></ul>
Fish	Stocking	-	<ul style="list-style-type: none"><li>Before stocking, make free the weeds and predators in the pond</li></ul>
<b>• Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795226247/37</b>			





# Integrated Agro-Meteorological Advisory Services

ICAR Research Complex for NEH Region, Arunachal Pradesh Centre

West Siang, Basar – 791 101



**(Kurung-Kumey)**

**No: 55/2014-15/Fri**

**Dated: 10<sup>th</sup> October, 2014**

## **Agro Advisory Bulletin for the period 11<sup>th</sup> – 15<sup>th</sup> October 2014**

### **Weather summery of the previous week**

Max Temp: 15.0<sup>0</sup>C, Min Temp: 6.0-7.0<sup>0</sup>C,  
Wind Speed: 3-6Kmph, Wind Direction: 150-  
279Degree, Rainfall: 0.0mm.

### **Weather forecast valid up to 15<sup>th</sup> October, 2014**

The possibility of Rainfall is 17.0mm for the next five days. Max Temp: 15.0<sup>0</sup>C, Min Temp: 6.0-7.0<sup>0</sup>C, Cloud Cover: sky will be mainly cloudy i.e. 3-7 Okta, Max RH: 90-96%, Min RH: 56-79%, Wind Speed: 3-4Kmph, Wind Direction: 170-249Degree.

***Weekly cumulative rainfall : 17.0mm***

## **Weather Based Agro-meteorological Advisories Bulletin**

- Obtain seeds from reputed/certified/reliable source,**

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>FIELD CROPS</b>			
Kharif Rice (Early transplanted)	Harvesting		<ul style="list-style-type: none"><li>Threshing followed by proper drying for safe storage,</li></ul>
Kharif Rice (Normal transplanted)	Tillering/ Booting	Blast/Gundhi bug	<ul style="list-style-type: none"><li>Need based alternate foliar spray of Hexaconazole 5 EC(2ml/l) &amp; Carbendazim 12WP+Mancozeb 63WP(2g/l) at 15days interval beginning from first appearance of the disease,</li><li>Dust with Malathion 5D @ 20kg/ha. Repeat weekly,</li><li>Properly maintain the water level in the field,</li></ul>
<b>FRUIT CROPS</b>			
Lemon/other citrus fruits	Fruit Bearing	Fruit fly/Trunk borer	<ul style="list-style-type: none"><li>Monitor fruit fly population with Methyl eugenol traps,</li><li>Spray Malathion (2ml/l) in the early stage to avoid fruit fly infestation,</li><li>Drenching of base soil with Dichlorvos 76 EC @ 4ml/l of water. Repeat weekly till the tree regain its greenery,</li></ul>
Pineapple	Fruiting/ Harvesting		<ul style="list-style-type: none"><li>Clean the vicinity of plants by weeding and loosen up the soil.</li><li>Remove sly and use it as a new planting material.</li><li>Sucker dipping treatment with Bavistin @ 3g + Chlorpyriphos 20EC@ 2ml/l of water to protect against soil borne disease and insect.</li></ul>
<b>LIVESTOCKS</b>			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages		<ul style="list-style-type: none"><li>Avoid dampness, litter should be periodically changed.</li><li>Provide clean and fresh water daily,</li><li>Vaccination of birds as per the recommended age,</li><li>Artificial heating through lighting of electric bulb during night time to protect the poultry birds (day old to 25 days old) from low temperature.</li></ul>
Pigs	New born	Cold	<ul style="list-style-type: none"><li>Pig shed/stay should keep dry and avoid dampness during rainy days,</li><li>Vaccinate against swine fever,</li><li>Avoid of eating human excreta,</li><li>Provide ad libitum water,</li><li>Avoid stagnant water near the shed,</li></ul>
Cattle	All stages	FMD	<ul style="list-style-type: none"><li>Vaccination of animal with FMD,</li><li>Treatment – sick animals,</li></ul>
Goat/Sheep	All stages	Goat pox	<ul style="list-style-type: none"><li>Vaccinate with PPR,</li><li>Provide standard feeds,</li><li>Vaccination &amp; deworming at regular interval,</li></ul>
Fish	Stocking	-	<ul style="list-style-type: none"><li>Before stocking, make free the weeds and predators in the pond</li></ul>
<ul style="list-style-type: none"><li><b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795226247/37</b></li></ul>			



