

# Agro-met Advisory Bulletin

(Period: 5<sup>th</sup> to 9<sup>th</sup> February, 2016)



## GRAMIN KRISHI MAUSAM SEWA (Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region  
Arunachal Pradesh Centre, Basar-791101

### District: Lower Dibang Valley

No: 11/2015-16/Fri



Dated: 5<sup>th</sup> February, 2016



Weather summery of the previous week	Weather forecast valid up to 9 <sup>th</sup> February, 2016
Rainfall = 0.0 mm.	Partly cloudy sky with possibility of light rain in few isolated places. Max Temp=23°C and Min Temp=10°C, RH=23-70%, Wind Speed=5-6 kmph, Wind direction: E

### Weather Based Agro-meteorological Advisories Bulletin

• Obtain seeds from reputed/certified/reliable source			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
VEGETABLES			
Tomato	Vegetative		<ul style="list-style-type: none"><li>Cover the crop with polyethene/cloths to protect from frost</li><li>Place 2-3 inch crop residue mulch around the base of the plant</li></ul>
Pea/Cauliflower/Cabbage	Vegetative	Leaf eating insects 	<ul style="list-style-type: none"><li>On clear weather, spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in cabbage/cauliflower/pea &amp; other vegetables is advocated</li></ul>

Potato	Harvesting	Ants 	<ul style="list-style-type: none"><li>• Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants.</li></ul>
FRUIT CROPS			
Citrus	Maturity/ Harvesting		<ul style="list-style-type: none"><li>• Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plant</li><li>• Go for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is over</li><li>• Spray Imagalil 34WP @ 1g/lit against blue &amp; green mauld in Khasi mandarin when the harvesting is over</li><li>• Remove all water sprouts from the base of the plant</li></ul>
Banana			<ul style="list-style-type: none"><li>• Timely/regular removal of extra suckers, dry leaves &amp; pseudo-stem</li></ul>
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Pigs	All stages		<ul style="list-style-type: none"><li>• Provide proper bedding materials for newly born piglets to protect them from cold</li><li>• Vaccination as per the schedule with proper consultation with vet.</li><li>• Deworm at regular interval</li><li>• During scarcity of fodder provide hay/silage/feedblocks.</li></ul>
Poultry	All stages		<ul style="list-style-type: none"><li>• Provide sufficient bedding materials to keep them warm</li><li>• Provide adequate clean drinking water+electrolyte</li><li>• Vaccination as per the schedule with proper consultation with vet.</li><li>• Provide anti-coccidial drugs</li><li>• Remove wet litter/provide artificial heating and light</li></ul>
Fishery			<ul style="list-style-type: none"><li>• Use soil-water probiotics in fish pond for making disease free environment</li></ul>
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit  
ICAR, BASAR

# Agro-met Advisory Bulletin

(Period: 5<sup>th</sup> to 9<sup>th</sup> February, 2016)



## GRAMIN KRISHI MAUSAM SEWA (Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region  
Arunachal Pradesh Centre, Basar-791101

### District: Lower Subansiri



No: 11/2015-16/Fri



Dated: 5<sup>th</sup> February, 2016



Weather summary of the previous week	Weather forecast valid up to 9 <sup>th</sup> February, 2016
Rainfall = 0.0 mm.	Moderately cloudy sky with possibility of light rain. Max Temp=17°C and Min Temp=10°C, RH=35-89%, Wind Speed=2-3 kmph, Wind direction: WSW

### Weather Based Agro-meteorological Advisories Bulletin

• Obtain seeds from reputed/certified/reliable source			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
VEGETABLES			
Tomato	Vegetative		<ul style="list-style-type: none"><li>Cover the crop with polyethene/cloths to protect from frost</li><li>Place 2-3 inch crop residue mulch around the base of the plant</li></ul>
Pea/Cauliflower/Cabbage	Vegetative	Leaf eating insects 	<ul style="list-style-type: none"><li>On clear weather, spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in cabbage/cauliflower/pea &amp; other vegetables is advocated</li></ul>

Potato	Harvesting	Ants 	<ul style="list-style-type: none"> <li>• Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants.</li> </ul>
<b>FRUIT CROPS</b>			
Citrus	Maturity/ Harvesting		<ul style="list-style-type: none"> <li>• Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plant</li> <li>• Go for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is over</li> <li>• Spray Imagalil 34WP @ 1g/lit against blue &amp; green mould in Khasi mandarin when the harvesting is over</li> <li>• Remove all water sprouts from the base of the plant</li> </ul>
Banana			<ul style="list-style-type: none"> <li>• Timely/regular removal of extra suckers, dry leaves &amp; pseudo-stem</li> </ul>
Apple/Kiwi			<ul style="list-style-type: none"> <li>• Training &amp; Pruning on priority to remove diseased and crisscross branches to maintain health and shape of plants</li> </ul>
<b>LIVESTOCKS</b>			
<b>Livestock</b>	<b>Stage</b>	<b>Disease</b>	<b>Precautionary Practices to be followed</b>
Pigs	All stages		<ul style="list-style-type: none"> <li>• Provide proper bedding materials for newly born piglets to protect them from cold</li> <li>• Vaccination as per the schedule with proper consultation with vet.</li> <li>• Deworm at regular interval</li> <li>• During scarcity of fodder provide hay/silage/feedblocks.</li> </ul>
Poultry	All stages		<ul style="list-style-type: none"> <li>• Provide sufficient bedding materials to keep them warm</li> <li>• Provide adequate clean drinking water+electrolyte</li> <li>• Vaccination as per the schedule with proper consultation with vet.</li> <li>• Provide anti-coccidial drugs</li> <li>• Remove wet litter/provide artificial heating and light</li> </ul>
Fishery			<ul style="list-style-type: none"> <li>• Use soil-water probiotics in fish pond for making disease free environment</li> </ul>
<b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247</b>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
 Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit  
 ICAR. BASAR

# Agro-met Advisory Bulletin

(Period: 5<sup>th</sup> to 9<sup>th</sup> February, 2016)



## GRAMIN KRISHI MAUSAM SEWA (Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region  
Arunachal Pradesh Centre, Basar-791101

### District: Papum Pare



No: 11/2015-16/Fri

Dated: 5<sup>th</sup> February, 2016





Weather summary of the previous week	Weather forecast valid up to 9 <sup>th</sup> February, 2016
Rainfall = 0.0 mm. Max Temp:26.1°C and Min. Temp:10.5°C, RH: 60-79%	Partly cloudy sky with possibility of very light rain in some places. Max Temp=28°C and Min Temp=13°C, RH=34-82%, Wind Speed=3-4 kmph, Wind direction: E

### Weather Based Agro-meteorological Advisories Bulletin

• Obtain seeds from reputed/certified/reliable source			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
VEGETABLES			
Tomato	Vegetative		<ul style="list-style-type: none"><li>Cover the crop with polyethene/cloths to protect from frost</li><li>Place 2-3 inch crop residue mulch around the base of the plant</li></ul>
Pea/Cauliflower/Cabbage	Vegetative	Leaf eating insects 	<ul style="list-style-type: none"><li>On clear weather, spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in cabbage/cauliflower/pea &amp; other vegetables is advocated</li></ul>



Potato	Harvesting	Ants 	<ul style="list-style-type: none"> <li>• Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants.</li> </ul>
<b>FRUIT CROPS</b>			
Citrus	Maturity/ Harvesting		<ul style="list-style-type: none"> <li>• Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plant</li> <li>• Go for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is over</li> <li>• On dry weather, spray Imagalil 34WP @ 1g/lit against blue &amp; green mauld in Khasi mandarin when the harvesting is over</li> <li>• Remove all water sprouts from the base of the plant</li> </ul>
Banana			<ul style="list-style-type: none"> <li>• Timely/regular removal of extra suckers, dry leaves &amp; pseudo-stem</li> </ul>
<b>LIVESTOCKS</b>			
<b>Livestock</b>	<b>Stage</b>	<b>Disease</b>	<b>Precautionary Practices to be followed</b>
Pigs	All stages		<ul style="list-style-type: none"> <li>• Provide proper bedding materials for newly born piglets to protect them from cold</li> <li>• Vaccination as per the schedule with proper consultation with vet.</li> <li>• Deworm at regular interval</li> <li>• During scarcity of fodder provide hay/silage/feedblocks.</li> </ul>
Poultry	All stages		<ul style="list-style-type: none"> <li>• Provide sufficient bedding materials to keep them warm</li> <li>• Provide adequate clean drinking water+electrolyte</li> <li>• Vaccination as per the schedule with proper consultation with vet.</li> <li>• Provide anti-coccidial drugs</li> <li>• Remove wet litter/provide artificial heating and light</li> </ul>
Fishery			<ul style="list-style-type: none"> <li>• Use soil-water probiotics in fish pond for making disease free environment</li> </ul>
<b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247</b>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
 Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit  
 ICAR, BASAR

# Agro-met Advisory Bulletin

(Period: 5<sup>th</sup> to 9<sup>th</sup> February, 2016)



## GRAMIN KRISHI MAUSAM SEWA (Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region  
Arunachal Pradesh Centre, Basar-791101




### District: Tawang


No: 11/2015-16/Tue Dated: 5<sup>th</sup> February, 2016



Weather summery of the previous week	Weather forecast valid up to 9 <sup>th</sup> February, 2016
Rainfall = 2.0 mm.	Partly cloudy sky with possibility of very light rain. Max Temp=14°C and Min Temp=3°C, RH=37-87%, Wind Speed=4-9 kmph, Wind direction: E

### Weather Based Agro-meteorological Advisories Bulletin

• Obtain seeds from reputed/certified/reliable source			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
VEGETABLES			
Tomato	Vegetative		<ul style="list-style-type: none"><li>Cover the crop with polyethene/cloths to protect from frost</li><li>Place 2-3 inch crop residue mulch around the base of the plant</li><li>Need based irrigation</li></ul>
Pea/Cauliflower/Cabbage	Vegetative	Leaf eating insects 	<ul style="list-style-type: none"><li>On clear weather, spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in cabbage/cauliflower/pea &amp; other vegetables is advocated</li></ul>
Potato	Harvesting	Ants 	<ul style="list-style-type: none"><li>Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants.</li></ul>

FRUIT CROPS			
Citrus	Maturity/ Harvesting		<ul style="list-style-type: none"> <li>• Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plant</li> <li>• Go for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is over</li> <li>• Spray Imagalil 34WP @ 1g/lit against blue &amp; green mauld in Khasi mandarin when the harvesting is over</li> <li>• Remove all water sprouts from the base of the plant</li> </ul>
Banana			<ul style="list-style-type: none"> <li>• Timely/regular removal of extra suckers, dry leaves &amp; pseudo-stem</li> </ul>
Apple/Kiwi			<ul style="list-style-type: none"> <li>• Traning &amp; Pruning on priority to remove diseased and crisscross branches to maintain health and shape of plants</li> </ul>
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Pigs	All stages		<ul style="list-style-type: none"> <li>• Provide proper bedding materials for newly born piglets to protect them from cold</li> <li>• Vaccination as per the schedule with proper consultation with vet.</li> <li>• Deworm at regular interval</li> <li>• During scarcity of fodder provide hay/silage/feedblocks.</li> </ul>
Poultry	All stages		<ul style="list-style-type: none"> <li>• Provide sufficient bedding materials to keep them warm</li> <li>• Provide adequate clean drinking water+electrolyte</li> <li>• Vaccination as per the schedule with proper consultation with vet.</li> <li>• Provide anti-coccidial drugs</li> <li>• Remove wet litter/provide artificial heating and light</li> </ul>
Fishery			<ul style="list-style-type: none"> <li>• Use soil-water probiotics in fish pond for making disease free environment</li> </ul>
<b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247</b>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
 Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit  
 ICAR, BASAR



# Agro-met Advisory Bulletin

(Period: 5<sup>th</sup> to 9<sup>th</sup> February, 2016)



## GRAMIN KRISHI MAUSAM SEWA (Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region  
Arunachal Pradesh Centre, Basar-791101

**District: Tirap**

No: 11/2015-16/Fri



Dated: 5<sup>th</sup> February, 2016



Weather summery of the previous week	Weather forecast valid up to 9 <sup>th</sup> February, 2016
Rainfall = 0.6 mm.	Partly cloudy sky with possibility of no rain. Max Temp=25°C and Min Temp=11°C, RH=39-100%, Wind Speed=2 kmph, Wind direction: S

### Weather Based Agro-meteorological Advisories Bulletin

• Obtain seeds from reputed/certified/reliable source			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
VEGETABLES			
Tomato	Vegetative		<ul style="list-style-type: none"><li>Cover the crop with polyethene/cloths to protect from frost</li><li>Place 2-3 inch crop residue mulch around the base of the plant</li></ul>
Pea/Cauliflower/Cabbage	Vegetative	Leaf eating insects 	<ul style="list-style-type: none"><li>On clear weather, spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in cabbage/cauliflower,/pea &amp; other vegetables is advocated</li></ul>

Potato	Harvesting	Ants 	<ul style="list-style-type: none"> <li>• Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants.</li> </ul>
<b>FRUIT CROPS</b>			
Citrus	Maturity/ Harvesting		<ul style="list-style-type: none"> <li>• Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plant</li> <li>• Go for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is over</li> <li>• Spray Imagalil 34WP @ 1g/lit against blue &amp; green mauld in Khasi mandarin when the harvesting is over</li> <li>• Remove all water sprouts from the base of the plant</li> </ul>
Banana			<ul style="list-style-type: none"> <li>• Timely/regular removal of extra suckers, dry leaves &amp; pseudo-stem</li> </ul>
<b>LIVESTOCKS</b>			
<b>Livestock</b>	<b>Stage</b>	<b>Disease</b>	<b>Precautionary Practices to be followed</b>
Pigs	All stages		<ul style="list-style-type: none"> <li>• Provide proper bedding materials for newly born piglets to protect them from cold</li> <li>• Vaccination as per the schedule with proper consultation with vet.</li> <li>• Deworm at regular interval</li> <li>• During scarcity of fodder provide hay/silage/feedblocks.</li> </ul>
Poultry	All stages		<ul style="list-style-type: none"> <li>• Provide sufficient bedding materials to keep them warm</li> <li>• Provide adequate clean drinking water+electrolyte</li> <li>• Vaccination as per the schedule with proper consultation with vet.</li> <li>• Provide anti-coccidial drugs</li> <li>• Remove wet litter/provide artificial heating and light</li> </ul>
Fishery			<ul style="list-style-type: none"> <li>• Use soil-water probiotics in fish pond for making disease free environment</li> </ul>
<b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247</b>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
 Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit  
 ICAR, BASAR

# Agro-met Advisory Bulletin

(Period: 5<sup>th</sup> to 9<sup>th</sup> February, 2016)



## GRAMIN KRISHI MAUSAM SEWA (Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region  
Arunachal Pradesh Centre, Basar-791101

### District: Upper Siang



No: 11/2015-16/Fri




Dated: 5<sup>th</sup> January, 2016



Weather summery of the previous week	Weather forecast valid up to 9 <sup>th</sup> February, 2016
Rainfall = 0.0 mm.	Moderate cloud cover with possibility of light rain in some places. Max Temp=17°C and Min Temp=9°C, RH=32-75%, Wind Speed=5-6 kmph, Wind direction: SSE

### Weather Based Agro-meteorological Advisories Bulletin

• Obtain seeds from reputed/certified/reliable source,			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
VEGETABLES			
Tomato	Vegetative		<ul style="list-style-type: none"><li>Cover the crop with polyethene/cloths to protect from frost</li><li>Place 2-3 inch crop residue mulch around the base of the plant</li></ul>
Pea/Cauliflower/Cabbage	Vegetative	Leaf eating insects 	<ul style="list-style-type: none"><li>Avoid spray operations</li><li>On clear weather, spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in cabbage/cauliflower,pea &amp; other vegetables is advocated, use stickers if spray is must</li></ul>

Potato	Harvesting	Ants 	<ul style="list-style-type: none"> <li>• Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants.</li> </ul>
<b>FRUIT CROPS</b>			
Citrus	Maturity/ Harvesting		<ul style="list-style-type: none"> <li>• Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plant</li> <li>• Go for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is over on clear day</li> <li>• On dry weather, spray Imagalil 34WP @ 1g/lit against blue &amp; green mauld in Khasi mandarin when the harvesting is over</li> <li>• Remove all water sprouts from the base of the plant on clear day using stickers</li> </ul>
Banana			<ul style="list-style-type: none"> <li>• Timely/regular removal of extra suckers, dry leaves &amp; pseudo-stem</li> </ul>
Apple /Kiwi			<ul style="list-style-type: none"> <li>• Traning &amp; Pruning on priority to remove diseased and crisscross branches to maintain health and shape of plants</li> </ul>
<b>LIVESTOCKS</b>			
<b>Livestock</b>	<b>Stage</b>	<b>Disease</b>	<b>Precautionary Practices to be followed</b>
Pigs	All stages		<ul style="list-style-type: none"> <li>• Provide proper bedding materials for newly born piglets to protect them from cold</li> <li>• Vaccination as per the schedule with proper consultation with vet.</li> <li>• Deworm at regular interval</li> <li>• During scarcity of fodder provide hay/silage/feedblocks.</li> </ul>
Poultry	All stages		<ul style="list-style-type: none"> <li>• Provide sufficient bedding materials to keep them warm</li> <li>• Provide adequate clean drinking water+electrolyte</li> <li>• Vaccination as per the schedule with proper consultation with vet.</li> <li>• Provide anti-coccidial drugs</li> <li>• Remove wet litter/provide artificial heating and light</li> </ul>
Fishery			<ul style="list-style-type: none"> <li>• Use soil-water probiotics in fish pond for making disease free environment</li> </ul>
<b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247</b>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit  
ICAR, BASAR

# Agro-met Advisory Bulletin

(Period: 5<sup>th</sup> to 9<sup>th</sup> February, 2016)



## GRAMIN KRISHI MAUSAM SEWA (Integrated Agro-Meteorological Advisory Services)




ICAR Research Complex for NEH Region  
Arunachal Pradesh Centre, Basar-791101

### District: Upper Subansiri

No: 11/2015-16/Fri

Dated: 5<sup>th</sup> February, 2016





Weather summery of the previous week	Weather forecast valid up to 9 <sup>th</sup> February, 2016
Rainfall = 0.0 mm.	Partly cloudy sky with possibility of light rain. Max Temp=20°C and Min Temp=4°C, RH=42-90%, Wind Speed=2-5 kmph, Wind direction: WSW

### Weather Based Agro-meteorological Advisories Bulletin

• Obtain seeds from reputed/certified/reliable source			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
VEGETABLES			
Tomato	Vegetative		<ul style="list-style-type: none"><li>Cover the crop with polyethene/cloths to protect from frost</li><li>Place 2-3 inch crop residue mulch around the base of the plant</li><li>Need based irrigation</li></ul>
Pea/Cauliflower/Cabbage	Vegetative	<p>Leaf eating insects</p>	<ul style="list-style-type: none"><li>On clear weather, spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in cabbage/cauliflower/pea &amp; other vegetables is advocated</li></ul>



Potato	Harvesting	<div>Ants</div> 	<ul style="list-style-type: none"> <li>• Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants.</li> </ul>
<b>FRUIT CROPS</b>			
Citrus	Maturity/ Harvesting		<ul style="list-style-type: none"> <li>• Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plant</li> <li>• Go for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is over</li> <li>• Spray Imagalil 34WP @ 1g/lit against blue &amp; green mauld in Khasi mandarin when the harvesting is over</li> <li>• Remove all water sprouts from the base of the plant</li> </ul>
Banana			<ul style="list-style-type: none"> <li>• Timely/regular removal of extra suckers, dry leaves &amp; pseudo-stem</li> </ul>
<b>LIVESTOCKS</b>			
<b>Livestock</b>	<b>Stage</b>	<b>Disease</b>	<b>Precautionary Practices to be followed</b>
Pigs	All stages		<ul style="list-style-type: none"> <li>• Provide proper bedding materials for newly born piglets to protect them from cold</li> <li>• Vaccination as per the schedule with proper consultation with vet.</li> <li>• Deworm at regular interval</li> <li>• During scarcity of fodder provide hay/silage/feedblocks.</li> </ul>
Poultry	All stages		<ul style="list-style-type: none"> <li>• Provide sufficient bedding materials to keep them warm</li> <li>• Provide adequate clean drinking water+electrolyte</li> <li>• Vaccination as per the schedule with proper consultation with vet.</li> <li>• Provide anti-coccidial drugs</li> <li>• Remove wet litter/provide artificial heating and light</li> </ul>
Fishery			<ul style="list-style-type: none"> <li>• Use soil-water probiotics in fish pond for making disease free environment</li> </ul>
<b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247</b>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
 Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit  
 ICAR, BASAR

# Agro-met Advisory Bulletin

(Period: 5<sup>th</sup> to 9<sup>th</sup> February, 2016)



## GRAMIN KRISHI MAUSAM SEWA (Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region  
Arunachal Pradesh Centre, Basar-791101



### District: West Kameng



No: 11/2015-16/Fri Dated: 5<sup>th</sup> February, 2016



Weather summery of the previous week	Weather forecast valid up to 9 <sup>th</sup> February, 2016
Rainfall = 0.0 mm.	Partly cloudy sky with possibility of no rain. Max Temp=8°C and Min Temp=1°C, RH=21-79%, Wind Speed=5-8 kmph, Wind direction: E

### Weather Based Agro-meteorological Advisories Bulletin

• Obtain seeds from reputed/certified/reliable source			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
VEGETABLES			
Pea/Cauliflower/Cabbage	Vegetative	Leaf eating insects 	• On clear weather, spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in cabbage/cauliflower/pea & other vegetables is advocated
Potato	Harvesting	Ants 	• Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants.

FRUIT CROPS			
Citrus	Maturity/ Harvesting		<ul style="list-style-type: none"> <li>• Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plant</li> <li>• Go for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is over</li> <li>• Spray Imagalil 34WP @ 1g/lit against blue &amp; green mauld in Khasi mandarin when the harvesting is over</li> <li>• Remove all water sprouts from the base of the plant</li> </ul>
Banana			<ul style="list-style-type: none"> <li>• Timely/regular removal of extra suckers, dry leaves &amp; pseudo-stem</li> </ul>
Apple/Kiwi			<ul style="list-style-type: none"> <li>• Traning &amp; Pruning on priority to remove diseased and crisscross branches to maintain health and shape of plants</li> </ul>
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Pigs	All stages		<ul style="list-style-type: none"> <li>• Provide proper bedding materials for newly born piglets to protect them from cold</li> <li>• Vaccination as per the schedule with proper consultation with vet.</li> <li>• Deworm at regular interval</li> <li>• During scarcity of fodder provide hay/silage/feedblocks.</li> </ul>
Poultry	All stages		<ul style="list-style-type: none"> <li>• Provide sufficient bedding materials to keep them warm</li> <li>• Provide adequate clean drinking water+electrolyte</li> <li>• Vaccination as per the schedule with proper consultation with vet.</li> <li>• Provide anti-coccidial drugs</li> <li>• Remove wet litter/provide artificial heating and light</li> </ul>
Fishery			<ul style="list-style-type: none"> <li>• Use soil-water probiotics in fish pond for making disease free environment</li> </ul>
<b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247</b>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit  
ICAR. BASAR

# Agro-met Advisory Bulletin

(Period: 5<sup>th</sup> to 9<sup>th</sup> February, 2016)




## GRAMIN KRISHI MAUSAM SEWA (Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region  
Arunachal Pradesh Centre, Basar-791101

### District: West Siang




No: 11/2015-16/Fri Dated: 5<sup>th</sup> February, 2016



Weather summery of the previous week	Weather forecast valid up to 9 <sup>th</sup> February, 2016
Rainfall = 0.0 mm. T. Max: 19.9°C, T. Min: 7.5 °C, RH: 58-98 %, Wind Speed: 2.4 kmph, Wind Direction: NE	Partly cloudy sky with possibility of rain in few places. Max Temp=18°C and Min Temp=7°C, RH=42-85%, Wind Speed=3-5 kmph, Wind direction: W

### Weather Based Agro-meteorological Advisories Bulletin

• Obtain seeds from reputed/certified/reliable source			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
VEGETABLES			
Tomato	Vegetative		<ul style="list-style-type: none"><li>Cover the crop with polyethene/cloths to protect from frost</li><li>Place 2-3 inch crop residue mulch around the base of the plant</li></ul>
Pea/Cauliflow er/Cabbage	Vegetative	Leaf eating insects 	<ul style="list-style-type: none"><li>On clear weather, spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in cabbage/cauliflower,/pea &amp; other vegetables is advocated</li></ul>

Potato	Harvesting	Ants 	<ul style="list-style-type: none"> <li>• Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants.</li> </ul>
Mustard			<ul style="list-style-type: none"> <li>• Provide provision for drainage in the crop in WRC field to protect in case of heavy rainfall</li> </ul>
<b>FRUIT CROPS</b>			
Citrus	Maturity/ Harvesting		<ul style="list-style-type: none"> <li>• Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plant</li> <li>• Go for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is over</li> <li>• Spray Imagalil 34WP @ 1g/lit against blue &amp; green mauld in Khasi mandarin when the harvesting is over</li> <li>• Remove all water sprouts from the base of the plant</li> </ul>
Banana			<ul style="list-style-type: none"> <li>• Timely/regular removal of extra suckers, dry leaves &amp; pseudo-stem</li> </ul>
Apple/Kiwi			<ul style="list-style-type: none"> <li>• Traning &amp; Pruning on priority to remove diseased and crisscross branches to maintain health and shape of plants</li> </ul>
<b>LIVESTOCKS</b>			
<b>Livestock</b>	<b>Stage</b>	<b>Disease</b>	<b>Precautionary Practices to be followed</b>
Pigs	All stages		<ul style="list-style-type: none"> <li>• Provide proper bedding materials for newly born piglets to protect them from cold</li> <li>• Vaccination as per the schedule with proper consultation with vet.</li> <li>• Deworm at regular interval</li> <li>• During scarcity of fodder provide hay/silage/feedblocks.</li> </ul>
Poultry	All stages		<ul style="list-style-type: none"> <li>• Provide sufficient bedding materials to keep them warm</li> <li>• Provide adequate clean drinking water+electrolyte</li> <li>• Vaccination as per the schedule with proper consultation with vet.</li> <li>• Provide anti-coccidial drugs</li> <li>• Remove wet litter/provide artificial heating and light</li> </ul>
Fishery			<ul style="list-style-type: none"> <li>• Use soil-water probiotics in fish pond for making disease free environment</li> </ul>
<b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247</b>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit

ICAR. BASAR



# Agro-met Advisory Bulletin

(Period: 5<sup>th</sup> to 9<sup>th</sup> February, 2016)



## GRAMIN KRISHI MAUSAM SEWA (Integrated Agro-Meteorological Advisory Services)




ICAR Research Complex for NEH Region  
Arunachal Pradesh Centre, Basar-791101

### District: Anjaw



No: 11/2015-16/Fri




Dated: 5<sup>th</sup> February, 2016



Weather summery of the previous week	Weather forecast valid up to 9 <sup>th</sup> February, 2016
Rainfall = 0.0 mm.	Partially cloudy sky with possibility of very light rain in few isolated places. Max Temp=16°C and Min Temp=6°C, RH=20-73%, Wind Speed=6-8 kmph, Wind direction: E

### Weather Based Agro-meteorological Advisories Bulletin

• Obtain seeds from reputed/certified/reliable source,			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
VEGETABLES			
Tomato	Vegetative		<ul style="list-style-type: none"><li>Cover the crop with polyethene/cloths to protect from frost</li><li>Place 2-3 inch crop residue mulch around the base of the plant</li></ul>
Pea/Cauliflower/Cabbage	Vegetative	Leaf eating insects 	<ul style="list-style-type: none"><li>On clear weather, spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in cabbage/cauliflower,/pea &amp; other vegetables is advocated</li></ul>

Potato	Harvesting	<div>Ants</div> 	<ul style="list-style-type: none"> <li>• Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants.</li> </ul>
FRUIT CROPS			
Citrus	Maturity/ Harvesting		<ul style="list-style-type: none"> <li>• Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plant</li> <li>• Go for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is over</li> <li>• Spray Imagalil 34WP @ 1g/lit against blue &amp; green mauld in Khasi mandarin when the harvesting is over</li> <li>• Remove all water sprouts from the base of the plant</li> </ul>
Banana			<ul style="list-style-type: none"> <li>• Timely/regular removal of extra suckers, dry leaves &amp; pseudo-stem</li> </ul>
Apple/Kiwi			<ul style="list-style-type: none"> <li>• Traning &amp; Pruning on priority to remove diseased and crisscross branches to maintain health and shape of plants</li> </ul>
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Pigs	All stages		<ul style="list-style-type: none"> <li>• Provide proper bedding materials for newly born piglets to protect them from cold</li> <li>• Vaccination as per the schedule with proper consultation with vet.</li> <li>• Deworm at regular interval</li> <li>• During scarcity of fodder provide hay/silage/feedblocks.</li> </ul>
Poultry	All stages		<ul style="list-style-type: none"> <li>• Provide sufficient bedding materials to keep them warm</li> <li>• Provide adequate clean drinking water+electrolyte</li> <li>• Vaccination as per the schedule with proper consultation with vet.</li> <li>• Provide anti-coccidial drugs</li> <li>• Remove wet litter/provide artificial heating and light</li> </ul>
Fishery			<ul style="list-style-type: none"> <li>• Use soil-water probiotics in fish pond for making disease free environment</li> </ul>
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
 Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit  
 ICAR, BASAR

# Agro-met Advisory Bulletin

(Period: 5<sup>th</sup> to 9<sup>th</sup> February, 2016)



## GRAMIN KRISHI MAUSAM SEWA (Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region  
Arunachal Pradesh Centre, Basar-791101



### District: Changlang


No: 11/2015-16/Fri Dated: 5<sup>th</sup> February, 2016



Weather summary of the previous week	Weather forecast valid up to 9 <sup>th</sup> February, 2016
Rainfall = 0.2 mm.	Partly cloudy sky with possibility of light rain in few places. Max Temp=25°C and Min Temp=10°C, RH=25-80%, Wind Speed=2-4 kmph, Wind direction: E

### Weather Based Agro-meteorological Advisories Bulletin

• Obtain seeds from reputed/certified/reliable source,			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
VEGETABLES			
Tomato	Vegetative		• Cover the crop with polyethene/cloths to protect from frost
Pea/Cauliflower/Cabbage	Vegetative	Leaf eating insects 	• On clear weather, spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in cabbage/cauliflower,pea & other vegetables is advocated
Potato	Harvesting	Ants 	• Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants.

FRUIT CROPS			
Citrus	Maturity/ Harvesting		<ul style="list-style-type: none"> <li>• Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plant</li> <li>• Go for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is over</li> <li>• On dry weather, spray Imagalil 34WP @ 1g/lit against blue &amp; green mauld in Khasi mandarin when the harvesting is over</li> <li>• Remove all water sprouts from the base of the plant</li> </ul>
Banana			<ul style="list-style-type: none"> <li>• Timely/regular removal of extra suckers, dry leaves &amp; pseudo-stem</li> </ul>
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Pigs	All stages		<ul style="list-style-type: none"> <li>• Provide proper bedding materials for newly born piglets to protect them from cold</li> <li>• Vaccination as per the schedule with proper consultation with vet.</li> <li>• Deworm at regular interval</li> <li>• During scarcity of fodder provide hay/silage/feedblocks.</li> </ul>
Poultry	All stages		<ul style="list-style-type: none"> <li>• Provide sufficient bedding materials to keep them warm</li> <li>• Provide adequate clean drinking water+electrolyte</li> <li>• Vaccination as per the schedule with proper consultation with vet.</li> <li>• Provide anti-coccidial drugs</li> <li>• Remove wet litter/provide artificial heating and light</li> </ul>
Fishery			<ul style="list-style-type: none"> <li>• Use soil-water probiotics in fish pond for making disease free environment</li> </ul>
<b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247</b>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit  
ICAR, BASAR

# Agro-met Advisory Bulletin

(Period: 5<sup>th</sup> to 9<sup>th</sup> February, 2016)



## GRAMIN KRISHI MAUSAM SEWA (Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region  
Arunachal Pradesh Centre, Basar-791101

### District: Dibang Valley


No: 11/2015-16/Fri

Dated: 5<sup>th</sup> February, 2016





Weather summery of the previous week	Weather forecast valid up to 9 <sup>th</sup> February, 2016
Rainfall = 0.0 mm.	Partly cloudy sky with possibility of vey light rain. Max Temp=17°C and Min Temp=8°C, RH=24-80%, Wind Speed=2-8 kmph, Wind direction: E

### Weather Based Agro-meteorological Advisories Bulletin

• Obtain seeds from reputed/certified/reliable source			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
VEGETABLES			
Tomato	Vegetative		<ul style="list-style-type: none"><li>Cover the crop with polyethene/cloths to protect from frost</li><li>Place 2-3 inch crop residue mulch around the base of the plant</li></ul>
Pea/Cauliflow er/Cabbage	Vegetative	Leaf eating insects 	<ul style="list-style-type: none"><li>On clear weather, spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in cabbage/cauliflower/pea &amp; other vegetables is advocated using stickers.</li></ul>



Potato	Harvesting	Ants 	<ul style="list-style-type: none"> <li>• Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants.</li> </ul>
<b>FRUIT CROPS</b>			
Citrus	Maturity/ Harvesting		<ul style="list-style-type: none"> <li>• Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plant</li> <li>• Go for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is over</li> <li>• Remove all water sprouts from the base of the plant</li> </ul>
Banana			<ul style="list-style-type: none"> <li>• Timely/regular removal of extra suckers, dry leaves &amp; pseudo-stem</li> </ul>
<b>LIVESTOCKS</b>			
<b>Livestock</b>	<b>Stage</b>	<b>Disease</b>	<b>Precautionary Practices to be followed</b>
Pigs	All stages		<ul style="list-style-type: none"> <li>• Provide proper bedding materials for newly born piglets to protect them from cold</li> <li>• Vaccination as per the schedule with proper consultation with vet.</li> <li>• Deworm at regular interval</li> <li>• During scarcity of fodder provide hay/silage/feedblocks.</li> </ul>
Poultry	All stages		<ul style="list-style-type: none"> <li>• Provide sufficient bedding materials to keep them warm</li> <li>• Provide adequate clean drinking water+electrolyte</li> <li>• Vaccination as per the schedule with proper consultation with vet.</li> <li>• Provide anti-coccidial drugs</li> <li>• Remove wet litter/provide artificial heating and light</li> </ul>
Fishery			<ul style="list-style-type: none"> <li>• Use soil-water probiotics in fish pond for making disease free environment</li> </ul>
<b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247</b>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit  
ICAR. BASAR

# Agro-met Advisory Bulletin

(Period: 5<sup>th</sup> to 9<sup>th</sup> February, 2016)



## GRAMIN KRISHI MAUSAM SEWA

(Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region  
Arunachal Pradesh Centre, Basar-791101



### District: East Kameng



No: 11/2015-16/Fri Dated: 5<sup>th</sup> February, 2016



Weather summary of the previous week	Weather forecast valid up to 9 <sup>th</sup> February, 2016
Rainfall = 0.0 mm.	Partly cloudy sky with possibility of light rain in some places. Max Temp=19°C and Min Temp=9°C, RH=33-85%, Wind Speed=4-6 kmph, Wind direction: SE

### Weather Based Agro-meteorological Advisories Bulletin

• Obtain seeds from reputed/certified/reliable source,			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
VEGETABLES			
Tomato	Vegetative		<ul style="list-style-type: none"><li>Cover the crop with polyethene/cloths to protect from frost</li><li>Place 2-3 inch crop residue mulch around the base of the plant</li></ul>
Pea/Cauliflower/Cabbage	Vegetative	Leaf eating insects 	<ul style="list-style-type: none"><li>On clear weather, spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in cabbage/cauliflower/pea &amp; other vegetables is advocated</li></ul>

Potato	Harvesting	Ants 	<ul style="list-style-type: none"><li>• Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants.</li></ul>
FRUIT CROPS			
Citrus	Maturity/ Harvesting		<ul style="list-style-type: none"><li>• Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plant</li><li>• Go for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is over</li><li>• Spray Imagalil 34WP @ 1g/lit against blue &amp; green mauld in Khasi mandarin when the harvesting is over</li><li>• Remove all water sprouts from the base of the plant</li></ul>
Banana			<ul style="list-style-type: none"><li>• Timely/regular removal of extra suckers, dry leaves &amp; pseudo-stem</li></ul>
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Pigs	All stages		<ul style="list-style-type: none"><li>• Provide proper bedding materials for newly born piglets to protect them from cold</li><li>• Vaccination as per the schedule with proper consultation with vet.</li><li>• Deworm at regular interval</li><li>• During scarcity of fodder provide hay/silage/feedblocks.</li></ul>
Poultry	All stages		<ul style="list-style-type: none"><li>• Provide sufficient bedding materials to keep them warm</li><li>• Provide adequate clean drinking water+electrolyte</li><li>• Vaccination as per the schedule with proper consultation with vet.</li><li>• Provide anti-coccidial drugs</li><li>• Remove wet litter/provide artificial heating and light</li></ul>
Fishery			<ul style="list-style-type: none"><li>• Use soil-water probiotics in fish pond for making disease free environment</li></ul>
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit  
ICAR. BASAR

# Agro-met Advisory Bulletin

(Period: 5<sup>th</sup> to 9<sup>th</sup> February, 2016)



## GRAMIN KRISHI MAUSAM SEWA (Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region  
Arunachal Pradesh Centre, Basar-791101

### District: East Siang


No: 11/2015-16/Fri



Dated: 5<sup>th</sup> February, 2016



Weather summary of the previous week	Weather forecast valid up to 9 <sup>th</sup> February, 2016
Rainfall = 0.0 mm; RH: 60-79%, T. Max: 26.1°C and T. Min: 10.5°C	Moderately cloudy sky with possibility of light rain in some places. Max Temp=26°C and Min Temp=14°C, RH=33-81%, Wind Speed=3-6 kmph, Wind direction: WNW

### Weather Based Agro-meteorological Advisories Bulletin

• Obtain seeds from reputed/certified/reliable source,			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
VEGETABLES			
Tomato	Vegetative		<ul style="list-style-type: none"><li>Cover the crop with polyethene/cloths to protect from frost</li><li>Place 2-3 inch crop residue mulch around the base of the plant</li></ul>
Pea/Cauliflower/Cabbage	Vegetative	Leaf eating insects 	<ul style="list-style-type: none"><li>On clear weather, spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in cabbage/cauliflower/pea &amp; other vegetables is advocated</li></ul>

Potato	Harvesting	Ants 	<ul style="list-style-type: none"> <li>• Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants.</li> </ul>
<b>FRUIT CROPS</b>			
Citrus	Maturity/ Harvesting		<ul style="list-style-type: none"> <li>• Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plant</li> <li>• Go for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is over</li> <li>• Spray Imagalil 34WP @ 1g/lit against blue &amp; green mauld in Khasi mandarin when the harvesting is over</li> <li>• Remove all water sprouts from the base of the plant</li> </ul>
Banana			<ul style="list-style-type: none"> <li>• Timely/regular removal of extra suckers, dry leaves &amp; pseudo-stem</li> </ul>
<b>LIVESTOCKS</b>			
<b>Livestock</b>	<b>Stage</b>	<b>Disease</b>	<b>Precautionary Practices to be followed</b>
Pigs	All stages		<ul style="list-style-type: none"> <li>• Provide proper bedding materials for newly born piglets to protect them from cold</li> <li>• Vaccination as per the schedule with proper consultation with vet.</li> <li>• Deworm at regular interval</li> <li>• During scarcity of fodder provide hay/silage/feedblocks.</li> </ul>
Poultry	All stages		<ul style="list-style-type: none"> <li>• Provide sufficient bedding materials to keep them warm</li> <li>• Provide adequate clean drinking water+electrolyte</li> <li>• Vaccination as per the schedule with proper consultation with vet.</li> <li>• Provide anti-coccidial drugs</li> <li>• Remove wet litter/provide artificial heating and light</li> </ul>
Fishery			<ul style="list-style-type: none"> <li>• Use soil-water probiotics in fish pond for making disease free environment</li> </ul>
<b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247</b>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
 Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit  
 ICAR, BASAR



# Agro-met Advisory Bulletin

(Period: 5<sup>th</sup> to 9<sup>th</sup> February, 2016)



## GRAMIN KRISHI MAUSAM SEWA

(Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region  
Arunachal Pradesh Centre, Basar-791101



**District: Kurung Kumey**



**No: 11/2015-16/Fri** **Dated: 5<sup>th</sup> February, 2016**



Weather summery of the previous week	Weather forecast valid up to 9 <sup>th</sup> February, 2016
Rainfall = 0.0 mm.	Moderate cloud cover with possibility of light rain in few places. Max Temp=19°C and Min Temp=10°C, RH=28-90%, Wind Speed=2-4 kmph, Wind direction: W

### Weather Based Agro-meteorological Advisories Bulletin

<ul style="list-style-type: none"><li>Obtain seeds from reputed/certified/reliable source,</li></ul>			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
VEGETABES			
Tomato	Vegetative		<ul style="list-style-type: none"><li>Cover the crop with polyethene/cloths to protect from frost</li></ul>
Pea/Cauliflower/Cabbage	Vegetative	Leaf eating insects 	<ul style="list-style-type: none"><li>On clear weather, spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in cabbage/cauliflower,/pea &amp; other vegetables is advocated</li></ul>
Potato	Harvesting	Ants 	<ul style="list-style-type: none"><li>Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants.</li></ul>

FRUIT CROPS			
Citrus	Maturity/ Harvesting		<ul style="list-style-type: none"> <li>• Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plant</li> <li>• Go for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is over</li> <li>• During dry weather, spray Imagalil 34WP @ 1g/lit against blue &amp; green mauld in Khasi mandarin when the harvesting is over</li> <li>• Remove all water sprouts from the base of the plant</li> </ul>
Banana			<ul style="list-style-type: none"> <li>• Timely/regular removal of extra suckers, dry leaves &amp; pseudo-stem</li> </ul>
Apple/Kiwi			<ul style="list-style-type: none"> <li>• Traning &amp; Pruning on priority to remove diseased and crisscross branches to maintain health and shape of plants</li> </ul>
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Pigs	All stages		<ul style="list-style-type: none"> <li>• Provide proper bedding materials for newly born piglets to protect them from cold</li> <li>• Vaccination as per the schedule with proper consultation with vet.</li> <li>• Deworm at regular interval</li> <li>• During scarcity of fodder provide hay/silage/feedblocks.</li> </ul>
Poultry	All stages		<ul style="list-style-type: none"> <li>• Provide sufficient bedding materials to keep them warm</li> <li>• Provide adequate clean drinking water+electrolyte</li> <li>• Vaccination as per the schedule with proper consultation with vet.</li> <li>• Provide anti-coccidial drugs</li> <li>• Remove wet litter/provide artificial heating and light</li> </ul>
Fishery			<ul style="list-style-type: none"> <li>• Use soil-water probiotics in fish pond for making disease free environment</li> </ul>
<b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247</b>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit  
ICAR, BASAR

# Agro-met Advisory Bulletin

(Period: 5<sup>th</sup> to 9<sup>th</sup> February, 2016)



## GRAMIN KRISHI MAUSAM SEWA

(Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region  
Arunachal Pradesh Centre, Basar-791101



**District: Lohit**



**No: 11/2015-16/Fri** **Dated: 9<sup>th</sup> February, 2016**



Weather summery of the previous week	Weather forecast valid up to 9 <sup>th</sup> February, 2016
Rainfall = 0.0 mm.	Partly cloudy sky with possibility of light rain. Max Temp=21°C and Min Temp=13°C, RH=19-81%, Wind Speed=4-5 kmph, Wind direction: E

### Weather Based Agro-meteorological Advisories Bulletin

• Obtain seeds from reputed/certified/reliable source			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
VEGETABES			
Tomato	Vegetative		<ul style="list-style-type: none"><li>Cover the crop with polyethene/cloths to protect from frost</li><li>Place 2-3 inch crop residue mulch around the base of the plant</li></ul>
Pea/Cauliflow er/Cabbage	Vegetative	Leaf eating insects 	<ul style="list-style-type: none"><li>On clear weather, spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in cabbage/cauliflower,/pea &amp; other vegetables is advocated</li></ul>

Potato	Harvesting	Ants 	<ul style="list-style-type: none"><li>• Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants.</li></ul>
FRUIT CROPS			
Citrus	Maturity/ Harvesting		<ul style="list-style-type: none"><li>• Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plant</li><li>• Go for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is over</li><li>• On dry weather, spray Imagalil 34WP @ 1g/lit against blue &amp; green mauld in Khasi mandarin when the harvesting is over</li><li>• Remove all water sprouts from the base of the plant</li></ul>
Banana			<ul style="list-style-type: none"><li>• Timely/regular removal of extra suckers, dry leaves &amp; pseudo-stem</li></ul>
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Pigs	All stages		<ul style="list-style-type: none"><li>• Provide proper bedding materials for newly born piglets to protect them from cold</li><li>• Vaccination as per the schedule with proper consultation with vet.</li><li>• Deworm at regular interval</li><li>• During scarcity of fodder provide hay/silage/feedblocks.</li></ul>
Poultry	All stages		<ul style="list-style-type: none"><li>• Provide sufficient bedding materials to keep them warm</li><li>• Provide adequate clean drinking water+electrolyte</li><li>• Vaccination as per the schedule with proper consultation with vet.</li><li>• Provide anti-coccidial drugs</li><li>• Remove wet litter/provide artificial heating and light</li></ul>
Fishery			<ul style="list-style-type: none"><li>• Use soil-water probiotics in fish pond for making disease free environment</li></ul>
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit  
ICAR, BASAR