

Agro-met Advisory Bulletin

(Period: 4th to 8th March, 2016)



GRAMIN KRISHI MAUSAM SEWA (Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101




District: Longding


No: 19/2015-16/Fri

Dated: 4th March, 2016

Weather summary of the previous week	Weather forecast valid up to 8 th March, 2016
Rainfall = NA	Partially cloudy sky with possibility of light rain.

Weather Based Agro-meteorological Advisories Bulletin

<ul style="list-style-type: none">• Right time for the construction of jalkund• Obtain seeds from reputed/certified/reliable source• Use fertilizers/pesticides as per recommended doses for the location• Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Maize/ cowpea			<ul style="list-style-type: none">• Right time for sowing of local/early maize and Cowpea• Seed treatment of early maize/cowpea with Imidichloropid @ 1-2g/10kg is recommended before sowing
Toria/mustard			<ul style="list-style-type: none">• On sunny days, go for harvesting of toria/mustard if mature
Maize (winter)			<ul style="list-style-type: none">• Right time for harvesting
VEGETABLES			
Summer Vegetables			<ul style="list-style-type: none">• Right time for the sowing of summer vegetables like cucumber, sponge gourd, bottle gourd, ladies finger etc.• Get seeds/planting materials of preferred or recommended varieties from right sources
Pea/Cauliflower/Cabbage	Vegetative	Leaf eating insects 	<ul style="list-style-type: none">• On clear weather, spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in cabbage/cauliflower/pea & other vegetables is advocated
Potato	Harvesting	Ants 	<ul style="list-style-type: none">• Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants.• Store potatoes in a cool, well ventilated place• Don't wash potatoes before storing as dampness promotes early spoilage.

FRUIT CROPS			
Citrus	Maturity/ Harvesting		<ul style="list-style-type: none"> • Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plant • Go for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is over • Spray Imagalil 34WP @ 1g/lit against blue & green mauld in Khasi mandarin when the harvesting is over • Remove all water sprouts from the base of the plant
Banana			<ul style="list-style-type: none"> • Timely/regular removal of extra suckers, dry leaves & pseudo-stem
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Pigs	All stages		<ul style="list-style-type: none"> • Provide concentrate feed and multi-vitamins • Provide proper bedding materials for newly born piglets to protect them from cold • Vaccination as per the schedule with proper consultation with vet. • Deworm at regular interval • During scarcity of fodder provide hay/silage/feedblocks.
Poultry	All stages 		<ul style="list-style-type: none"> • Avoid rearing of Day-Old-Chicks • Provide sufficient bedding materials to keep them warm • Provide adequate clean drinking water+electrolyte • Vaccination as per the schedule with proper consultation with vet. • Provide anti-coccidial drugs • Remove wet litter/provide artificial heating and light
Fishery			<ul style="list-style-type: none"> • Use soil-water probiotics in fish pond for making disease free environment
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit
ICAR, BASAR

Agro-met Advisory Bulletin

(Period: 4th to 8th March, 2016)




GRAMIN KRISHI MAUSAM SEWA (Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Lower Dibang Valley





No: 19/2015-16/Fri Dated: 4th March, 2016



Weather summery of the previous week	Weather forecast valid up to 8 th March, 2016
Rainfall = 0.0 mm	Partial cloudy weather is forecasted for next 2-3 days with light rain in few places. Max Temp=17°C and Min Temp=3°C, RH=19-71%, Wind Speed=6-8 kmph, Wind direction: E
Warning:	Non

Weather Based Agro-meteorological Advisories Bulletin

<ul style="list-style-type: none">Right time for digging pond for construction of <i>Jalkunds</i>.Go for low cost polyhouses for year round cultivation of vegetablesObtain seeds from reputed/certified/reliable sourceUse fertilizers/pesticides as per recommended doses for the locationPreference must be given on use of organic manures/bio-fertilizers & bio-pesticidesICAR Basar recommend poultry variety VANARAJA for backyard rearing			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Maize/ cowpea			<ul style="list-style-type: none">Right time for sowing of local/early maize and CowpeaSeed treatment of early maize/cowpea with Imidichloropid @ 1-2g/10kg is recommended before sowing
Toria/mustard			<ul style="list-style-type: none">On sunny days, go for harvesting of toria/mustard if mature

Maize (winter)			<ul style="list-style-type: none"> Right time for harvesting
Pea		Powdery Mildew 	<ul style="list-style-type: none"> Need based spray of Carbendazim 50WP @ 1g/lit or Difenoconazole 5EC @ 0.5 ml/lit against powdery mildew.
VEGETABLES			
Summer Vegetables			<ul style="list-style-type: none"> Right time for the sowing of summer vegetables like cucumber, sponge gourd, bottle gourd, ladies finger etc. Get seeds/planting materials of preferred or recommended varieties from right sources
Pea/Cauliflower/Cabbage	Vegetative	Leaf eating insects 	<ul style="list-style-type: none"> On clear weather, need based spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in cabbage/cauliflower/pea & other vegetables is advocated using stickers
Potato	Harvesting	Ants 	<ul style="list-style-type: none"> Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants. Store potatoes in a cool, well ventilated place Don't wash potatoes before storing as dampness promotes early spoilage.
FRUIT CROPS			
Citrus	Maturity/Harvesting		<ul style="list-style-type: none"> Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plant Go for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is over Spray Imagalil 34WP @ 1g/lit against blue & green mauld in Khasi mandarin when the harvesting is over Remove all water sprouts from the base of the plant Need based spray of Carbendazim 50WP @ 1g/lit or Difenoconazole 5EC @ 0.5 ml/lit against powdery mildew
Banana			<ul style="list-style-type: none"> Timely/regular removal of extra suckers, dry leaves & pseudo-stem
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Pigs	All stages		<ul style="list-style-type: none"> Provide concentrate feed and multi-vitamins Provide proper bedding materials for newly born piglets to protect them from cold Vaccination as per the schedule with proper consultation with vet. Deworm at regular interval During scarcity of fodder provide hay/silage/feedblocks.
Poultry	All stages 		<ul style="list-style-type: none"> Provide sufficient bedding materials to keep them warm Provide adequate clean drinking water+electrolyte Vaccination as per the schedule with proper consultation with vet. Provide anti-coccidial drugs Remove wet litter/provide artificial heating and light Quarantine the newly procured birds for one month before introduction with old birds.
Fishery			<ul style="list-style-type: none"> Use soil-water probiotics in fish pond for making disease free environment
Dog		Canine Parvovirus	<ul style="list-style-type: none"> Vaccinate as per consultation with vet

Cattle		FMD	• Vaccination is advocated
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit
ICAR. BASAR

Agro-met Advisory Bulletin

(Period: 4th to 8th March, 2016)



GRAMIN KRISHI MAUSAM SEWA (Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Lower Subansiri




No: 19/2015-16/Fri Dated: 4th March, 2016




Weather summary of the previous week	Weather forecast valid up to 8 th March, 2016
Rainfall = 0.0	Partially cloudy weather with light rain is forecasted for next 2-3 days. Max Temp=8°C and Min Temp=2°C, RH=46-83%, Wind Speed=4-6 kmph, Wind direction: WSW
Warning:	Non

Weather Based Agro-meteorological Advisories Bulletin

<ul style="list-style-type: none">Right time for digging pond for construction of <i>Jalkunds</i>.Go for low cost polyhouses for year round cultivation of vegetablesObtain seeds from reputed/certified/reliable sourceUse fertilizers/pesticides as per recommended doses for the locationPreference must be given on use of organic manures/bio-fertilizers & bio-pesticidesICAR Basar recommend poultry variety VANARAJA for backyard rearing			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Maize/ cowpea			<ul style="list-style-type: none">Right time for sowing of local/early maize and CowpeaSeed treatment of early maize/cowpea with Imidichloropid @ 1-2g/10kg is recommended before sowing
Toria/mustard			<ul style="list-style-type: none">On sunny days, go for harvesting of toria/mustard if mature
Maize (winter)			<ul style="list-style-type: none">Right time for harvesting

Pea		Powdery Mildew 	<ul style="list-style-type: none"> Need based spray of Carbendazim 50WP @ 1g/lit or Difenconazole 5EC @ 0.5 ml/lit against powdery mildew.
VEGETABLES			
Summer Vegetables			<ul style="list-style-type: none"> Right time for the sowing of summer vegetables like cucumber, sponge gourd, bottle gourd, ladies finger etc. Get seeds/planting materials of preferred or recommended varieties from right sources
Pea/Cauliflower/Cabbage	Vegetative	Leaf eating insects 	<ul style="list-style-type: none"> On clear weather, need based spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in cabbage/cauliflower/pea & other vegetables is advocated.
Potato	Harvesting	Ants 	<ul style="list-style-type: none"> Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants. Store potatoes in a cool, well ventilated place Don't wash potatoes before storing as dampness promotes early spoilage.
FRUIT CROPS			
Citrus	Maturity/Harvesting		<ul style="list-style-type: none"> Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plant Go for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is over Spray Imagalil 34WP @ 1g/lit against blue & green mauld in Khasi mandarin when the harvesting is over Remove all water sprouts from the base of the plant Need based spray of Carbendazim 50WP @ 1g/lit or Difenconazole 5EC @ 0.5 ml/lit against powdery mildew.
Banana			<ul style="list-style-type: none"> Timely/regular removal of extra suckers, dry leaves & pseudo-stem
Apple/Kiwi			<ul style="list-style-type: none"> Traning & Pruning on priority to remove diseased and crisscross branches to maintain health and shape of plants Best time to apply fertilizers to the apple trees containing potassium, calcium, Boron and nitrogen Fertilizers should be applied a foot away from the base of the plants
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Pigs	All stages		<ul style="list-style-type: none"> Provide concentrate feed and multi-vitamins Provide proper bedding materials for newly born piglets to protect them from cold Vaccination as per the schedule with proper consultation with vet. Deworm at regular interval During scarcity of fodder provide hay/silage/feedblocks.
Poultry	All stages		<ul style="list-style-type: none"> Avoid rearing of Day-Old-Chicks Provide sufficient bedding materials to keep them warm Provide adequate clean drinking water+electrolyte Vaccination as per the schedule with proper consultation with vet.

			<ul style="list-style-type: none">• Provide anti-coccidial drugs• Remove wet litter/provide artificial heating and light• Quarantine the newly procured birds for one month before introduction with old birds.
Fishery			<ul style="list-style-type: none">• Use soil-water probiotics in fish pond for making disease free environment
Dog		Canine Parvovirus	<ul style="list-style-type: none">• Vaccinate as per consultation with vet
Cattle		FMD	<ul style="list-style-type: none">• Vaccination is advocated
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit
ICAR, BASAR

Agro-met Advisory Bulletin

(Period: 4th to 8th March, 2016)



GRAMIN KRISHI MAUSAM SEWA (Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Papum Pare


No: 19/2015-16/Fri





Dated: 4th March, 2016



Weather summery of the previous week	Weather forecast valid up to 8 th March, 2016
Rainfall: 0.0 mm; T.Max: 32.5°C; T.Min:15.9°C ; RH: 42-62 %	Partially cloudy weather with possibility of light rain in next 2-3 days. Max Temp=30°C and Min Temp=17°C, RH=30-74%, Wind Speed=5-6 kmph, Wind direction: E
Warning:	Non

Weather Based Agro-meteorological Advisories Bulletin

<ul style="list-style-type: none">Right time for digging pond for construction of <i>Jalkunds</i>.Go for low cost polyhouses for year round cultivation of vegetablesObtain seeds from reputed/certified/reliable sourceUse fertilizers/pesticides as per recommended doses for the locationPreference must be given on use of organic manures/bio-fertilizers & bio-pesticidesICAR Basar recommend poultry variety VANARAJA for backyard rearing			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Maize/ cowpea			<ul style="list-style-type: none">Right time for sowing of local/early maize and CowpeaSeed treatment of early maize/cowpea with Imidichloropid @ 1-2g/10kg is recommended before sowing
Toria/mustard			<ul style="list-style-type: none">On sunny days, go for harvesting of toria/mustard if mature
Maize (winter)			<ul style="list-style-type: none">Right time for harvesting

Pea		Powdery Mildew 	<ul style="list-style-type: none"> Need based spray of Carbendazim 50WP @ 1g/lit or Difenconazole 5EC @ 0.5 ml/lit against powdery mildew.
VEGETABLES			
Summer Vegetables			<ul style="list-style-type: none"> Right time for the sowing of summer vegetables like cucumber, sponge gourd, bottle gourd, ladies finger etc. Get seeds/planting materials of preferred or recommended varieties from right sources
Pea/Cauliflower/Cabbage	Vegetative	Leaf eating insects 	<ul style="list-style-type: none"> On clear weather, need based spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in cabbage/cauliflower/pea & other vegetables is advocated.
Potato	Harvesting	Ants 	<ul style="list-style-type: none"> Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants. Store potatoes in a cool, well ventilated place Don't wash potatoes before storing as dampness promotes early spoilage.
FRUIT CROPS			
Citrus	Maturity/Harvesting		<ul style="list-style-type: none"> Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plant Go for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is over On dry weather, spray Imagilil 34WP @ 1g/lit against blue & green mould in Khasi mandarin when the harvesting is over Remove all water sprouts from the base of the plant Need based spray of Carbendazim 50WP @ 1g/lit or Difenconazole 5EC @ 0.5 ml/lit against powdery mildew
Banana			<ul style="list-style-type: none"> Timely/regular removal of extra suckers, dry leaves & pseudo-stem
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Pigs	All stages		<ul style="list-style-type: none"> Provide concentrate feed and multi-vitamins Provide proper bedding materials for newly born piglets to protect them from cold Vaccination as per the schedule with proper consultation with vet. Deworm at regular interval During scarcity of fodder provide hay/silage/feedblocks.
Poultry	All stages 		<ul style="list-style-type: none"> Avoid rearing of Day-Old-Chicks Provide sufficient bedding materials to keep them warm Provide adequate clean drinking water+electrolyte Vaccination as per the schedule with proper consultation with vet. Provide anti-coccidial drugs Remove wet litter/provide artificial heating and light Quarantine the newly procured birds for one month before

			introduction with old birds
Fishery			<ul style="list-style-type: none">• Use soil-water probiotics in fish pond for making disease free environment
Dog		Canine Parvovirus	<ul style="list-style-type: none">• Vaccinate as per consultation with vet
Cattle		FMD	<ul style="list-style-type: none">• Vaccination is advocated
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit
ICAR, BASAR

Agro-met Advisory Bulletin

(Period: 4th to 8th March, 2016)



GRAMIN KRISHI MAUSAM SEWA (Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Tawang




No: 19/2015-16/Fri Dated: 4th March, 2016



Weather summery of the previous week	Weather forecast valid up to 8 th March, 2016
Rainfall = 0.0 mm.	Cloudy weather is forecasted for next 2-3 days, with light rain in few isolated places. Max Temp=11°C and Min Temp=3°C, RH=30-72%, Wind Speed=6-9 kmph, Wind direction: ENE
Warning:	Non

Weather Based Agro-meteorological Advisories Bulletin

<ul style="list-style-type: none">Right time for digging pond for construction of <i>Jalkunds</i>.Go for low cost polyhouses for year round cultivation of vegetablesObtain seeds from reputed/certified/reliable sourceUse fertilizers/pesticides as per recommended doses for the locationPreference must be given on use of organic manures/bio-fertilizers & bio-pesticidesICAR Basar recommend poultry variety VANARAJA for backyard rearing			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Maize/ cowpea			<ul style="list-style-type: none">Right time for sowing of local/early maize and CowpeaSeed treatment of early maize/cowpea with Imidichloropid @ 1-2g/10kg is recommended before sowing
Toria/mustard			<ul style="list-style-type: none">On sunny days, go for harvesting of toria/mustard if mature
Maize (winter)			<ul style="list-style-type: none">Right time for harvesting
VEGETABLES			
Pea/Cauliflower/Cabbage	Vegetative	Leaf eating insects	<ul style="list-style-type: none">On clear weather, need based spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in

			cabbage/cauliflower,pea & other vegetables is advocated.
Potato	Harvesting	Ants 	<ul style="list-style-type: none"> • Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants. • Store potatoes in a cool, well ventilated place • Don't wash potatoes before storing as dampness promotes early spoilage.
FRUIT CROPS			
Citrus	Maturity/ Harvesting		<ul style="list-style-type: none"> • Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plant • Go for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is over • Spray Imagalil 34WP @ 1g/lit against blue & green mauld in Khasi mandarin when the harvesting is over • Remove all water sprouts from the base of the plant • Need based spray of Carbendazim 50WP @ 1g/lit or Difenconazole 5EC @ 0.5 ml/lit against powdery mildew
Banana			<ul style="list-style-type: none"> • Timely/regular removal of extra suckers, dry leaves & pseudo-stem
Apple/Kiwi			<ul style="list-style-type: none"> • Traning & Pruning on priority to remove diseased and crisscross branches to maintain health and shape of plants • Best time to apply fertilizers to the apple trees containing potassium, calcium, Boron and nitrogen • Fertilizers should be applied a foot away from the base of the plants
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Pigs	All stages		<ul style="list-style-type: none"> • Provide concentrate feed and multi-vitamins • Provide proper bedding materials for newly born piglets to protect them from cold • Vaccination as per the schedule with proper consultation with vet. • Deworm at regular interval • During scarcity of fodder provide hay/silage/feedblocks.
Poultry	All stages 		<ul style="list-style-type: none"> • Avoid rearing of Day-Old-Chicks • Provide sufficient bedding materials to keep them warm • Provide adequate clean drinking water+electrolyte • Vaccination as per the schedule with proper consultation with vet. • Provide anti-coccidial drugs • Remove wet litter/provide artificial heating and light • Quarantine the newly procured birds for one month before introduction with old birds.
Fishery			<ul style="list-style-type: none"> • Use soil-water probiotics in fish pond for making disease free environment
Dog		Canine Parvovirus	<ul style="list-style-type: none"> • Vaccinate as per consultation with vet
Cattle		FMD	<ul style="list-style-type: none"> • Vaccination is advocated
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

Agro-met Advisory Bulletin

(Period: 4th to 8th March, 2016)



GRAMIN KRISHI MAUSAM SEWA (Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Tirap





No: 19/2015-16/Fri Dated: 4th March, 2016



Weather summery of the previous week	Weather forecast valid up to 8 th March, 2016
Rainfall = 0.0 mm	Partly cloudy sky with possibility of no rain in next 2-3 days. Max Temp=30°C and Min Temp=17°C, RH=45-89%, Wind Speed=2 kmph, Wind direction: ESE
Warning:	Non

Weather Based Agro-meteorological Advisories Bulletin

<ul style="list-style-type: none">Right time for digging pond for construction of <i>Jalkunds</i>.Go for low cost polyhouses for year round cultivation of vegetablesObtain seeds from reputed/certified/reliable sourceUse fertilizers/pesticides as per recommended doses for the locationPreference must be given on use of organic manures/bio-fertilizers & bio-pesticidesICAR Basar recommend poultry variety VANARAJA for backyard rearing			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Maize/ cowpea			<ul style="list-style-type: none">Right time for sowing of local/early maize and CowpeaSeed treatment of early maize/cowpea with Imidichloropid @ 1-2g/10kg is recommended before sowing
Toria/mustard			<ul style="list-style-type: none">On sunny days, go for harvesting of toria/mustard if mature
Maize (winter)			<ul style="list-style-type: none">Right time for harvesting

Pea		Powdery Mildew 	<ul style="list-style-type: none"> Need based spray of Carbendazim 50WP @ 1g/lit or Difenonazole 5EC @ 0.5 ml/lit against powdery mildew.
VEGETABLES			
Summer Vegetables			<ul style="list-style-type: none"> Right time for the sowing of summer vegetables like cucumber, sponge gourd, bottle gourd, ladies finger etc. Get seeds/planting materials of preferred or recommended varieties from right sources
Pea/Cauliflower/Cabbage	Vegetative	Leaf eating insects 	<ul style="list-style-type: none"> On clear weather, spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in cabbage/cauliflower, pea & other vegetables is advocated
Potato	Harvesting	Ants 	<ul style="list-style-type: none"> Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants. Store potatoes in a cool, well ventilated place Don't wash potatoes before storing as dampness promotes early spoilage.
FRUIT CROPS			
Citrus	Maturity/Harvesting		<ul style="list-style-type: none"> Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plant Go for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is over Spray Imagalil 34WP @ 1g/lit against blue & green mould in Khasi mandarin when the harvesting is over Remove all water sprouts from the base of the plant Need based spray of Carbendazim 50WP @ 1g/lit or Difenonazole 5EC @ 0.5 ml/lit against powdery mildew
Banana			<ul style="list-style-type: none"> Timely/regular removal of extra suckers, dry leaves & pseudo-stem
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Pigs	All stages		<ul style="list-style-type: none"> Provide concentrate feed and multi-vitamins Provide proper bedding materials for newly born piglets to protect them from cold Vaccination as per the schedule with proper consultation with vet. Deworm at regular interval During scarcity of fodder provide hay/silage/feedblocks.
Poultry	All stages 		<ul style="list-style-type: none"> Avoid rearing of Day-Old-Chicks Provide sufficient bedding materials to keep them warm Provide adequate clean drinking water+electrolyte Vaccination as per the schedule with proper consultation with vet. Provide anti-coccidial drugs Remove wet litter/provide artificial heating and light Quarantine the newly procured birds for one month before

			introduction with old birds.
Fishery			<ul style="list-style-type: none">• Use soil-water probiotics in fish pond for making disease free environment
Dog		Canine Parvovirus	<ul style="list-style-type: none">• Vaccinate as per consultation with vet
Cattle		FMD	<ul style="list-style-type: none">• Vaccination is advocated
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit
ICAR, BASAR

Agro-met Advisory Bulletin

(Period: 4th to 8th March, 2016)



GRAMIN KRISHI MAUSAM SEWA (Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Upper Siang






No: 19/2015-16/Fri **Dated: 4th March, 2016**




Weather summery of the previous week	Weather forecast valid up to 8 th March, 2016
Rainfall = 0.0 mm.	Moderate cloud cover with light/moderate rain is forecasted for next 2-3 days. Max Temp=10°C and Min Temp=2°C, RH=21-68%, Wind Speed=7-9 kmph, Wind direction: SSE
Warning:	Heavy rainfall is likely in few places on 7th

Weather Based Agro-meteorological Advisories Bulletin

<ul style="list-style-type: none">Right time for digging pond for construction of <i>Jalkunds</i>.Go for low cost polyhouses for year round cultivation of vegetablesObtain seeds from reputed/certified/reliable sourceUse fertilizers/pesticides as per recommended doses for the locationPreference must be given on use of organic manures/bio-fertilizers & bio-pesticidesICAR Basar recommend poultry variety VANARAJA for backyard rearing			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Maize/ cowpea			<ul style="list-style-type: none">Right time for sowing of local/early maize and CowpeaSeed treatment of early maize/cowpea with Imidichloropid @ 1-2g/10kg is recommended before sowing

Toria/mustard			<ul style="list-style-type: none"> On sunny days, go for harvesting of toria/mustard if mature
Maize (winter)			<ul style="list-style-type: none"> Right time for harvesting
Pea		Powdery Mildew 	<ul style="list-style-type: none"> Need based spray of Carbendazim 50WP @ 1g/lit or Difenoconazole 5EC @ 0.5 ml/lit against powdery mildew.
VEGETABLES			
Summer Vegetables			<ul style="list-style-type: none"> Right time for the sowing of summer vegetables like cucumber, sponge gourd, bottle gourd, ladies finger etc. Get seeds/planting materials of preferred or recommended varieties from right sources
Pea/Cauliflower/Cabbage	Vegetative	Leaf eating insects 	<ul style="list-style-type: none"> On clear weather, need based spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in cabbage/cauliflower/pea & other vegetables is advocated with stickers.
Potato	Harvesting	Ants 	<ul style="list-style-type: none"> Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants. Store potatoes in a cool, well ventilated place Don't wash potatoes before storing as dampness promotes early spoilage.
FRUIT CROPS			
Citrus	Maturity/Harvesting		<ul style="list-style-type: none"> Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plant Go for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is over on clear day On dry weather, spray Imagalil 34WP @ 1g/lit against blue & green mauld in Khasi mandarin when the harvesting is over Remove all water sprouts from the base of the plant on clear day using stickers Need based spray of Carbendazim 50WP @ 1g/lit or Difenoconazole 5EC @ 0.5 ml/lit against powdery mildew.
Banana			<ul style="list-style-type: none"> Timely/regular removal of extra suckers, dry leaves & pseudo-stem
Apple /Kiwi			<ul style="list-style-type: none"> Traning & Pruning on priority to remove diseased and crisscross branches to maintain health and shape of plants Best time to apply fertilizers to the apple trees containing potassium, calcium, Boron and nitrogen Fertilizers should be applied a foot away from the base of the plants
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Pigs	All stages		<ul style="list-style-type: none"> Provide concentrate feed and multi-vitamins Provide proper bedding materials for newly born piglets to protect them from cold Vaccination as per the schedule with proper consultation with vet.

			<ul style="list-style-type: none">• Deworm at regular interval• During scarcity of fodder provide hay/silage/feedblocks.
Poultry	All stages		<ul style="list-style-type: none">• Avoid rearing of Day-Old-Chicks• Provide sufficient bedding materials to keep them warm• Provide adequate clean drinking water+electrolyte• Vaccination as per the schedule with proper consultation with vet.• Provide anti-coccidial drugs• Remove wet litter/provide artificial heating and light• Quarantine the newly procured birds for one month before introduction with old birds.
Fishery			<ul style="list-style-type: none">• Use soil-water probiotics in fish pond for making disease free environment
Dog		Canine Parvovirus	<ul style="list-style-type: none">• Vaccinate as per consultation with vet
Cattle		FMD	<ul style="list-style-type: none">• Vaccination is advocated
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit
ICAR, BASAR

Agro-met Advisory Bulletin

(Period: 4th to 8th March, 2016)



GRAMIN KRISHI MAUSAM SEWA (Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Upper Subansiri





No: 19/2015-16/Fri Dated: 4th March, 2016



Weather summery of the previous week	Weather forecast valid up to 8 th March, 2016
Rainfall = 0.0 mm	Partial cloud cover with light rain is forecasted for next 2-3 days. Max Temp=10°C and Min Temp=2°C, RH=47-85%, Wind Speed=5-6 kmph, Wind direction: WNW

Weather Based Agro-meteorological Advisories Bulletin

<ul style="list-style-type: none">Right time for digging pond for construction of <i>Jalkunds</i>.Go for low cost polyhouses for year round cultivation of vegetablesObtain seeds from reputed/certified/reliable sourceUse fertilizers/pesticides as per recommended doses for the locationPreference must be given on use of organic manures/bio-fertilizers & bio-pesticidesICAR Basar recommend poultry variety VANARAJA for backyard rearing			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Maize/ cowpea			<ul style="list-style-type: none">Right time for sowing of local/early maize and CowpeaSeed treatment of early maize/cowpea with Imidichloropid @ 1-2g/10kg is recommended before sowing
Toria/mustard			<ul style="list-style-type: none">On sunny days, go for harvesting of toria/mustard if mature

Maize (winter)			<ul style="list-style-type: none"> Right time for harvesting
Pea		Powdery Mildew 	<ul style="list-style-type: none"> Need based spray of Carbendazim 50WP @ 1g/lit or Difenoconazole 5EC @ 0.5 ml/lit against powdery mildew.
VEGETABLES			
Summer Vegetables			<ul style="list-style-type: none"> Right time for the sowing of summer vegetables like cucumber, sponge gourd, bottle gourd, ladies finger etc. Get seeds/planting materials of preferred or recommended varieties from right sources
Pea/Cauliflower/Cabbage	Vegetative	Leaf eating insects 	<ul style="list-style-type: none"> On clear weather, need based spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in cabbage/cauliflower,/pea & other vegetables is advocated with sticker.
Potato	Harvesting	Ants 	<ul style="list-style-type: none"> Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants. Best time to apply fertilizers to the apple trees containing potassium, calcium, Boron and nitrogen Fertilizers should be applied a foot away from the base of the plants
FRUIT CROPS			
Citrus	Maturity/Harvesting		<ul style="list-style-type: none"> Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plant Go for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is over Spray Imagalil 34WP @ 1g/lit against blue & green mauld in Khasi mandarin when the harvesting is over Remove all water sprouts from the base of the plant Need based spray of Carbendazim 50WP @ 1g/lit or Difenoconazole 5EC @ 0.5 ml/lit against powdery mildew
Banana			<ul style="list-style-type: none"> Timely/regular removal of extra suckers, dry leaves & pseudo-stem
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Pigs	All stages		<ul style="list-style-type: none"> Provide concentrate feed and multi-vitamins Provide proper bedding materials for newly born piglets to protect them from cold Vaccination as per the schedule with proper consultation with vet. Deworm at regular interval During scarcity of fodder provide hay/silage/feedblocks.
Poultry	All stages		<ul style="list-style-type: none"> Avoid rearing of Day-Old-Chicks Provide sufficient bedding materials to keep them warm Provide adequate clean drinking water+electrolyte Vaccination as per the schedule with proper consultation with vet. Provide anti-coccidial drugs Remove wet litter/provide artificial heating and light Quarantine the newly procured birds for one month before

			introduction with old birds.
Fishery			<ul style="list-style-type: none">• Use soil-water probiotics in fish pond for making disease free environment
Dog		Canine Parvovirus	<ul style="list-style-type: none">• Vaccinate as per consultation with vet
Cattle		FMD	<ul style="list-style-type: none">• Vaccination is advocated
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit
ICAR, BASAR

Agro-met Advisory Bulletin

(Period: 4th to 8th March, 2016)



GRAMIN KRISHI MAUSAM SEWA (Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: West Kameng






No: 19/2015-16/Fri Dated: 4th March, 2016




Weather summery of the previous week	Weather forecast valid up to 8 th March, 2016
Rainfall = 0.0.	Moderate cloud cover with light rain is forecasted for next 2-3 days. Max Temp=15°C and Min Temp=6°C, RH=37-69%, Wind Speed=6-7 kmph, Wind direction: ENE
Warning:	Non

Weather Based Agro-meteorological Advisories Bulletin

<ul style="list-style-type: none">Right time for digging pond for construction of <i>Jalkunds</i>.Go for low cost polyhouses for year round cultivation of vegetablesObtain seeds from reputed/certified/reliable sourceUse fertilizers/pesticides as per recommended doses for the locationPreference must be given on use of organic manures/bio-fertilizers & bio-pesticidesICAR Basar recommend poultry variety VANARAJA for backyard rearing			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Maize/ cowpea			<ul style="list-style-type: none">Right time for sowing of local/early maize and CowpeaSeed treatment of early maize/cowpea with Imidichloropid @ 1-2g/10kg is recommended before sowing

Toria/mustard			<ul style="list-style-type: none"> On sunny days, go for harvesting of toria/mustard if mature
Maize (winter)			<ul style="list-style-type: none"> Right time for harvesting
Pea		Powdery Mildew 	<ul style="list-style-type: none"> Need based spray of Carbendazim 50WP @ 1g/lit or Difenoconazole 5EC @ 0.5 ml/lit against powdery mildew.
VEGETABLES			
Pea/Cauliflower/Cabbage	Vegetative	Leaf eating insects 	<ul style="list-style-type: none"> On clear weather, need based spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in cabbage/cauliflower/pea & other vegetables is advocated.
Potato	Harvesting	Ants 	<ul style="list-style-type: none"> Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants. Store potatoes in a cool, well ventilated place Don't wash potatoes before storing as dampness promotes early spoilage.
Summer Vegetables			<ul style="list-style-type: none"> Right time for the sowing of summer vegetables like cucumber, sponge gourd, bottle gourd, ladies finger etc. Get seeds/planting materials of preferred or recommended varieties from right sources
FRUIT CROPS			
Citrus	Maturity/Harvesting		<ul style="list-style-type: none"> Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plant Go for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is over Spray Imagalil 34WP @ 1g/lit against blue & green mauld in Khasi mandarin when the harvesting is over Remove all water sprouts from the base of the plant Need based spray of Carbendazim 50WP @ 1g/lit or Difenoconazole 5EC @ 0.5 ml/lit against powdery mildew
Banana			<ul style="list-style-type: none"> Timely/regular removal of extra suckers, dry leaves & pseudo-stem
Apple/Kiwi			<ul style="list-style-type: none"> Traning & Pruning on priority to remove diseased and crisscross branches to maintain health and shape of plants Best time to apply fertilizers to the apple trees containing potassium, calcium, Boron and nitrogen Fertilizers should be applied a foot away from the base of the plants
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Pigs	All stages		<ul style="list-style-type: none"> Provide concentrate feed and multi-vitamins Provide proper bedding materials for newly born piglets to protect them from cold Vaccination as per the schedule with proper consultation with vet. Deworm at regular interval

			<ul style="list-style-type: none"> During scarcity of fodder provide hay/silage/feedblocks.
Poultry	All stages		<ul style="list-style-type: none"> Avoid rearing of Day-Old-Chicks Provide sufficient bedding materials to keep them warm Provide adequate clean drinking water+electrolyte Vaccination as per the schedule with proper consultation with vet. Provide anti-coccidial drugs Remove wet litter/provide artificial heating and light Need based spray of Carbendazim 50WP @ 1g/lit or Difenoconazole 5EC @ 0.5 ml/lit against powdery mildew
Fishery			<ul style="list-style-type: none"> Use soil-water probiotics in fish pond for making disease free environment
Dog		Canine Parvovirus	<ul style="list-style-type: none"> Vaccinate as per consultation with vet
Cattle		FMD	<ul style="list-style-type: none"> Vaccination is advocated
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
 Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit
 ICAR. BASAR

Agro-met Advisory Bulletin

(Period: 4th to 8th March, 2016)




GRAMIN KRISHI MAUSAM SEWA (Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: West Siang

No: 19/2015-16/Fri Dated: 4th March, 2016









Weather summery of the previous week	Weather forecast valid up to 8 th March, 2016
Rainfall = 0.0 mm. T. Max: 26.3°C, T. Min: 9.8°C, RH: 31-78 %, Wind Speed: 2 kmph, Wind Direction: NE	Moderate cloud cover with possibility of rain in few places. Max Temp=28°C and Min Temp=10°C, RH=24-73%, Wind Speed=6-7 kmph, Wind direction: WNW
Warning:	Heavy rainfall is likely in few places on 7th March

Weather Based Agro-meteorological Advisories Bulletin

- Right time for digging pond for construction of *Jalkunds*.
- Go for low cost polyhouses for year round cultivation of vegetables
- Obtain seeds from reputed/certified/reliable source
- Use fertilizers/pesticides as per recommended doses for the location
- Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides


- **ICAR Basar recommend poultry variety VANARAJA for backyard rearing**

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Maize/ cowpea			<ul style="list-style-type: none"> • Right time for sowing of local/early maize and Cowpea • Seed treatment of early maize/cowpea with Imidichloropid @ 1-2g/10kg is recommended before sowing
Toria/mustard			<ul style="list-style-type: none"> • On sunny days, go for harvesting of toria/mustard if mature
Maize (winter)			<ul style="list-style-type: none"> • Right time for harvesting
Pea		Powdery Mildew 	<ul style="list-style-type: none"> • Need based foliar spray of Carbendazim 50WP @ 1g/lit or Difenconazole 5EC @ 0.5 ml/lit against powdery mildew.
VEGETABLES			
Tomato	Vegetative		<ul style="list-style-type: none"> • Cover the crop with polyethene/cloths to protect from frost • Place 2-3 inch crop residue mulch around the base of the plant
Summer Vegetables			<ul style="list-style-type: none"> • Right time for the sowing of summer vegetables like cucumber, sponge gourd, bottle gourd, ladies finger etc. • Get seeds/planting materials of preferred or recommended varieties from right sources
Pea/Cauliflower/Cabbage	Vegetative	Leaf eating insects 	<ul style="list-style-type: none"> • On clear weather, need based spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in cabbage/cauliflower/pea & other vegetables is advocated.
Potato	Harvesting	Ants 	<ul style="list-style-type: none"> • Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants. • Store potatoes in a cool, well ventilated place • Don't wash potatoes before storing as dampness promotes early spoilage.
Mustard			<ul style="list-style-type: none"> • Provide provision for drainage in the crop in WRC field to protect in case of heavy rainfall
FRUIT CROPS			
Citrus	Maturity/Harvesting	Powdery Mildew 	<ul style="list-style-type: none"> • Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plant • Go for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is over • Spray Imagalil 34WP @ 1g/lit against blue & green mauld in Khasi mandarin when the harvesting is over • Remove all water sprouts from the base of the plant • Need based foliar spray of Carbendazim 50WP @ 1g/lit or Difenconazole 5EC @ 0.5 ml/lit against powdery mildew.
Banana			<ul style="list-style-type: none"> • Timely/regular removal of extra suckers, dry leaves & pseudo-stem
Apple/Kiwi			<ul style="list-style-type: none"> • Traning & Pruning on priority to remove diseased and crisscross branches to maintain health and shape of plants



- Store potatoes in a cool, well ventilated place
- Don't wash potatoes before storing as dampness promotes early spoilage.

LIVESTOCKS

Livestock	Stage	Disease	Precautionary Practices to be followed
Pigs	All stages		<ul style="list-style-type: none"> • Provide concentrate feed and multi-vitamins • Provide proper bedding materials for newly born piglets to protect them from cold • Vaccination as per the schedule with proper consultation with vet. • Deworm at regular interval • During scarcity of fodder provide hay/silage/feedblocks.
Poultry	All stages		<ul style="list-style-type: none"> • Avoid rearing of Day-Old-Chicks • Provide sufficient bedding materials to keep them warm • Provide adequate clean drinking water+electrolyte • Vaccination as per the schedule with proper consultation with vet. • Provide anti-coccidial drugs • Remove wet litter/provide artificial heating and light • Quarantine the newly procured birds for one month before introduction with old birds.
Fishery			<ul style="list-style-type: none"> • Use soil-water probiotics in fish pond for making disease free environment
Dog		Canine Parvovirus	<ul style="list-style-type: none"> • Vaccinate as per consultation with vet
Cattle		FMD	<ul style="list-style-type: none"> • Vaccination is advocated

Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
 Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit
 ICAR, BASAR**

Agro-met Advisory Bulletin

(Period: 4th to 8th March, 2016)



GRAMIN KRISHI MAUSAM SEWA


(Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Anjaw






No: 19/2015-16/Fri Dated: 4th March, 2016




Weather summery of the previous week	Weather forecast valid up to 8 th March, 2016
Rainfall = 0.0	Partially cloudy sky with possibility of rain in few places. Max Temp=11°C and Min Temp=2°C, RH=16-76%, Wind Speed=9-11 kmph, Wind direction: E
Warning:	Heavy rain on 7th in few isolated places

Weather Based Agro-meteorological Advisories Bulletin

<ul style="list-style-type: none">Right time for digging pond for construction of <i>Jalkunds</i>.Go for low cost polyhouses for year round cultivation of vegetablesObtain seeds from reputed/certified/reliable sourceUse fertilizers/pesticides as per recommended doses for the locationPreference must be given on use of organic manures/bio-fertilizers & bio-pesticidesICAR Basar recommend poultry variety VANARAJA for backyard rearing			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Maize/ cowpea			<ul style="list-style-type: none">Right time for sowing of local/early maize and CowpeaSeed treatment of early maize/cowpea with Imidichloropid @ 1-2g/10kg is recommended before sowing
Toria/mustard			<ul style="list-style-type: none">On sunny days, go for harvesting of toria/mustard if mature
Maize (winter)			<ul style="list-style-type: none">Right time for harvesting

Pea		Powdery Mildew 	<ul style="list-style-type: none"> On clear/dry weather, need based spray of Carbendazim 50WP @ 1g/lit or Difenoconazole 5EC @ 0.5 ml/lit against powdery mildew.
VEGETABLES			
Summer Vegetables			<ul style="list-style-type: none"> Right time for the sowing of summer vegetables like cucumber, sponge gourd, bottle gourd, ladies finger etc. Get seeds/planting materials of preferred or recommended varieties from right sources
Pea/Cauliflower/Cabbage	Vegetative	Leaf eating insects 	<ul style="list-style-type: none"> Need based spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in cabbage/cauliflower, pea & other vegetables is advocated using sticker
Potato	Harvesting	Ants 	<ul style="list-style-type: none"> Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants. Store potatoes in a cool, well ventilated place Don't wash potatoes before storing as dampness promotes early spoilage.
FRUIT CROPS			
Citrus	Maturity/Harvesting	Powdery Mildew 	<ul style="list-style-type: none"> Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plant Go for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is over On clear day, spray Imagalil 34WP @ 1g/lit against blue & green mauld in Khasi mandarin when the harvesting is over Remove all water sprouts from the base of the plant Need based spray of Carbendazim 50WP @ 1g/lit or Difenoconazole 5EC @ 0.5 ml/lit against powdery mildew
Banana			<ul style="list-style-type: none"> Timely/regular removal of extra suckers, dry leaves & pseudo-stem
Apple/Kiwi			<ul style="list-style-type: none"> Traning & Pruning on priority to remove diseased and crisscross branches to maintain health and shape of plants Best time to apply fertilizers to the apple trees containing potassium, calcium, Boron and nitrogen Fertilizers should be applied a foot away from the base of the plants
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Pigs	All stages		<ul style="list-style-type: none"> Provide concentrate feed and multi-vitamins Provide proper bedding materials for newly born piglets to protect them from cold Vaccination as per the schedule with proper consultation with vet. Deworm at regular interval During scarcity of fodder provide hay/silage/feedblocks.
Poultry	All stages		<ul style="list-style-type: none"> Avoid rearing of Day-Old-Chicks Provide sufficient bedding materials to keep them warm Provide adequate clean drinking water+electrolyte and multi-vitamins Vaccination as per the schedule with proper consultation

			<div>with vet.</div> <ul style="list-style-type: none">• Provide anti-coccidial drugs• Remove wet litter/provide artificial heating and light• Quarantine the newly procured birds for one month before introduction with old birds.
Fishery			<ul style="list-style-type: none">• Use soil-water probiotics in fish pond for making disease free environment
Dog		Canine Parvovirus	<ul style="list-style-type: none">• Vaccinate as per consultation with vet
Cattle		FMD	<ul style="list-style-type: none">• Vaccination is advocated
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit
ICAR, BASAR

Agro-met Advisory Bulletin

(Period: 4th to 8th March, 2016)



GRAMIN KRISHI MAUSAM SEWA (Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Changlang






No: 19/2015-16/Fri Dated: 4th March, 2016



Weather summary of the previous week	Weather forecast valid up to 8 th March, 2016
Rainfall = 0.0 mm.	Moderately cloudy sky with possibility of light rain. Max Temp=21°C and Min Temp=10°C, RH=41-83%, Wind Speed=4-6 kmph, Wind direction: E
Warning	Non

Weather Based Agro-meteorological Advisories Bulletin

<ul style="list-style-type: none">Right time for digging ponds for construction of jalkundsObtain seeds from reputed/certified/reliable sourceUse fertilizers/pesticides as per recommended doses for the locationPreference must be given on use of organic manures/bio-fertilizers & bio-pesticidesICAR Basar recommend poultry variety VANARAJA for backyard rearing			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Maize/ cowpea			<ul style="list-style-type: none">Right time for sowing of local/early maize and CowpeaSeed treatment of early maize/cowpea with Imidichloropid @ 1-2g/10kg is recommended before sowing
Toria/mustard			<ul style="list-style-type: none">On sunny days, go for harvesting of toria/mustard if mature
Maize (winter)			<ul style="list-style-type: none">Right time for harvesting

Pea		<div>Powdery Mildew</div> 	<ul style="list-style-type: none">Need based spray of Carbendazim 50WP @ 1g/lit or Difenconazole 5EC @ 0.5 ml/lit against powdery mildew.
VEGETABLES			
Pea/Cauliflower/Cabbage	Vegetative	<div>Leaf eating insects</div> 	<ul style="list-style-type: none">On clear weather, need based spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in cabbage/cauliflower/pea & other vegetables is advocated using stickers.
Summer Vegetables			<ul style="list-style-type: none">Right time for the sowing of summer vegetables like cucumber, sponge gourd, bottle gourd, ladies finger etc.Get seeds/planting materials of preferred or recommended varieties from right sources
Potato	Harvesting	<div>Ants</div> 	<ul style="list-style-type: none">Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants.Store potatoes in a cool, well ventilated placeDon't wash potatoes before storing as dampness promotes early spoilage.
FRUIT CROPS			
Citrus	Maturity/Harvesting	<div>Powdery Mildew</div> 	<ul style="list-style-type: none">Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plantGo for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is overOn dry weather, spray Imagalil 34WP @ 1g/lit against blue & green mauld in Khasi mandarin when the harvesting is overRemove all water sprouts from the base of the plantNeed based spray of Carbendazim 50WP @ 1g/lit or Difenconazole 5EC @ 0.5 ml/lit against powdery mildew.
Banana			<ul style="list-style-type: none">Timely/regular removal of extra suckers, dry leaves & pseudo-stem
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Pigs	All stages		<ul style="list-style-type: none">Provide concentrate feed and multi-vitaminsProvide proper bedding materials for newly born piglets to protect them from coldVaccination as per the schedule with proper consultation with vet.Deworm at regular intervalDuring scarcity of fodder provide hay/silage/feedblocks.
Poultry	<div>All stages</div> 		<ul style="list-style-type: none">Avoid rearing of Day-Old-ChicksProvide sufficient bedding materials to keep them warmProvide adequate clean drinking water+electrolyteVaccination as per the schedule with proper consultation with vet.Provide anti-coccidial drugsRemove wet litter/provide artificial heating and light
Fishery			<ul style="list-style-type: none">Use soil-water probiotics in fish pond for making disease free environment
Dog		<div>Canine Parvovirus</div>	<ul style="list-style-type: none">Vaccinate as per consultation with vet
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit
ICAR, BASAR

Agro-met Advisory Bulletin

(Period: 4th to 8th March, 2016)



GRAMIN KRISHI MAUSAM SEWA

(Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Dibang Valley






No: 19/2015-16/Fri Dated: 4th March, 2016



Weather summery of the previous week	Weather forecast valid up to 8 th March, 2016
Rainfall = 3.0 mm.	Moderate cloud cover with possibility of very light rain. Max Temp=14°C and Min Temp=5°C, RH=20-81%, Wind Speed=6-11 kmph, Wind direction: ENE
Warning:	Non

Weather Based Agro-meteorological Advisories Bulletin

<ul style="list-style-type: none">Right time for digging pond for construction of <i>Jalkunds</i>.Go for low cost polyhouses for year round cultivation of vegetablesObtain seeds from reputed/certified/reliable sourceUse fertilizers/pesticides as per recommended doses for the locationPreference must be given on use of organic manures/bio-fertilizers & bio-pesticidesICAR Basar recommend poultry variety VANARAJA for backyard rearing			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Maize/ cowpea			<ul style="list-style-type: none">Right time for sowing of local/early maize and CowpeaSeed treatment of early maize/cowpea with Imidichloropid @ 1-2g/10kg is recommended before sowing
Toria/mustard			<ul style="list-style-type: none">On sunny days, go for harvesting of toria/mustard if mature
Maize (winter)			<ul style="list-style-type: none">Right time for harvesting

Pea		Powdery Mildew 	<ul style="list-style-type: none"> Need based spray of Carbendazim 50WP @ 1g/lit or Difenconazole 5EC @ 0.5 ml/lit against powdery mildew.
VEGETABLES			
Tomato	Vegetative		<ul style="list-style-type: none"> Cover the crop with polyethene/cloths to protect from frost/rain Place 2-3 inch crop residue mulch around the base of the plant
Summer Vegetables			<ul style="list-style-type: none"> Right time for the sowing of summer vegetables like cucumber, sponge gourd, bottle gourd, ladies finger etc. Get seeds/planting materials of preferred or recommended varieties from right sources
Pea/Cauliflower/Cabbage	Vegetative	Leaf eating insects 	<ul style="list-style-type: none"> On clear weather, spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in cabbage/cauliflower/pea & other vegetables is advocated using stickers.
Potato	Harvesting	Ants 	<ul style="list-style-type: none"> Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants. Store potatoes in a cool, well ventilated place Don't wash potatoes before storing as dampness promotes early spoilage.
FRUIT CROPS			
Citrus	Maturity/Harvesting		<ul style="list-style-type: none"> Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plant Go for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is over Remove all water sprouts from the base of the plant Need based spray of Carbendazim 50WP @ 1g/lit or Difenconazole 5EC @ 0.5 ml/lit against powdery mildew
Banana			<ul style="list-style-type: none"> Timely/regular removal of extra suckers, dry leaves & pseudo-stem
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Pigs	All stages 		<ul style="list-style-type: none"> Provide concentrate feed and multi-vitamins Provide proper bedding materials for newly born piglets to protect them from cold Vaccination as per the schedule with proper consultation with vet. Deworm at regular interval During scarcity of fodder provide hay/silage/feedblocks.
Poultry	All stages 		<ul style="list-style-type: none"> Avoid rearing of Day-Old-Chicks Provide sufficient bedding materials to keep them warm Provide adequate clean drinking water+electrolyte Vaccination as per the schedule with proper consultation with vet. Provide anti-coccidial drugs Remove wet litter/provide artificial heating and light Quarantine the newly procured birds for one month before introduction with old birds
Fishery			<ul style="list-style-type: none"> Use soil-water probiotics in fish pond for making disease free environment

Dog		Canine Parvovirus	• Vaccinate as per consultation with vet
Cattle		FMD	• Vaccination is advocated
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit
ICAR, BASAR

Agro-met Advisory Bulletin

(Period: 4th to 8th March, 2016)



GRAMIN KRISHI MAUSAM SEWA

(Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: East Kameng






No: 19/2015-16/Fri Dated: 4th March, 2016



Weather summary of the previous week	Weather forecast valid up to 8 th March, 2016
Rainfall = 0.0 mm.	Clear/dry weather is forecasted for next 2-3 days. Max Temp=11°C and Min Temp=3°C, RH=36-72%, Wind Speed=5-6 kmph, Wind direction: ESE
Warning:	Non

Weather Based Agro-meteorological Advisories Bulletin

<ul style="list-style-type: none">• Right time for digging pond for construction of <i>Jalkunds</i>.• Go for low cost polyhouses for year round cultivation of vegetables• Obtain seeds from reputed/certified/reliable source• Use fertilizers/pesticides as per recommended doses for the location• Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides• ICAR Basar recommend poultry variety VANARAJA for backyard rearing			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Maize/ cowpea			<ul style="list-style-type: none">• Right time for sowing of local/early maize and Cowpea• Seed treatment of early maize/cowpea with Imidichloropid @ 1-2g/10kg is recommended before sowing• Need-based irrigation if crops sown

Toria/mustard			<ul style="list-style-type: none"> On sunny days, go for harvesting of toria/mustard if mature
Maize (winter)			<ul style="list-style-type: none"> Right time for harvesting
Pea		Powdery Mildew 	<ul style="list-style-type: none"> Need based spray of Carbendazim 50WP @ 1g/lit or Difenoconazole 5EC @ 0.5 ml/lit against powdery mildew.
VEGETABLES			
Pea/Cauliflower/Cabbage	Vegetative	Leaf eating insects 	<ul style="list-style-type: none"> On clear weather, spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in cabbage/cauliflower, pea & other vegetables is advocated.
Potato	Harvesting	Ants 	<ul style="list-style-type: none"> Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants. Store potatoes in a cool, well ventilated place Don't wash potatoes before storing as dampness promotes early spoilage.
Summer Vegetables			<ul style="list-style-type: none"> Right time for the sowing of summer vegetables like cucumber, sponge gourd, bottle gourd, ladies finger etc. Get seeds/planting materials of preferred or recommended varieties from right sources
FRUIT CROPS			
Citrus	Maturity/Harvesting		<ul style="list-style-type: none"> Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plant Go for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is over Spray Imagalil 34WP @ 1g/lit against blue & green mauld in Khasi mandarin when the harvesting is over Remove all water sprouts from the base of the plant Need based spray of Carbendazim 50WP @ 1g/lit or Difenoconazole 5EC @ 0.5 ml/lit against powdery mildew.
Banana			<ul style="list-style-type: none"> Timely/regular removal of extra suckers, dry leaves & pseudo-stem
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Pigs	All stages		<ul style="list-style-type: none"> Provide concentrate feed and multi-vitamins Provide proper bedding materials for newly born piglets to protect them from cold Vaccination as per the schedule with proper consultation with vet. Deworm at regular interval During scarcity of fodder provide hay/silage/feedblocks.
Poultry	All stages 		<ul style="list-style-type: none"> Avoid rearing of Day-Old-Chicks Provide sufficient bedding materials to keep them warm Provide adequate clean drinking water+electrolyte Vaccination as per the schedule with proper consultation with vet. Provide anti-coccidial drugs Remove wet litter/provide artificial heating and light

			<ul style="list-style-type: none">Quarantine the newly procured birds for one month before introduction with old birds.
Fishery			<ul style="list-style-type: none">Use soil-water probiotics in fish pond for making disease free environment
Dog		Canine Parvovirus	<ul style="list-style-type: none">Vaccinate as per consultation with vet
Cattle		FMD	<ul style="list-style-type: none">Vaccination is advocated
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit
ICAR, BASAR

Agro-met Advisory Bulletin

(Period: 4th to 8th March, 2016)



GRAMIN KRISHI MAUSAM SEWA (Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: East Siang

No: 19/2015-16/Fri





Dated: 4th March, 2016



Weather summary of the previous week	Weather forecast valid up to 8 th March, 2016
Rainfall: 0.0 mm; T. Max: 30.0°C; T. Min: 17.1°C; RH: 54-64%	Partially cloudy sky with possibility of very light rain in few isolated places. Max Temp=29°C and Min Temp=18°C, RH=23-76%, Wind Speed=6 kmph, Wind direction: W
Warning:	Heavy rainfall is forecasted on 7th in some places

Weather Based Agro-meteorological Advisories Bulletin

<ul style="list-style-type: none">Right time for digging pond for construction of <i>Jalkunds</i>.Go for low cost polyhouses for year round cultivation of vegetablesObtain seeds from reputed/certified/reliable sourceUse fertilizers/pesticides as per recommended doses for the locationPreference must be given on use of organic manures/bio-fertilizers & bio-pesticidesICAR Basar recommend poultry variety VANARAJA for backyard rearing			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Maize/ cowpea			<ul style="list-style-type: none">Right time for sowing of local/early maize and CowpeaSeed treatment of early maize/cowpea with Imidichloropid @ 1-2g/10kg is recommended before sowing
Toria/mustard			<ul style="list-style-type: none">On sunny days, go for harvesting of toria/mustard if mature

Maize (winter)			<ul style="list-style-type: none"> Right time for harvesting
Pea		Powdery Mildew 	<ul style="list-style-type: none"> Need based spray of Carbendazim 50WP @ 1g/lit or Difenoconazole 5EC @ 0.5 ml/lit against powdery mildew.
VEGETABLES			
Summer Vegetables			<ul style="list-style-type: none"> Right time for the sowing of summer vegetables like cucumber, sponge gourd, bottle gourd, ladies finger etc. Get seeds/planting materials of preferred or recommended varieties from right sources
Pea/Cauliflower/Cabbage	Vegetative	Leaf eating insects 	<ul style="list-style-type: none"> On clear weather, spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in cabbage/cauliflower/pea & other vegetables is advocated using stickers.
Potato	Harvesting	Ants 	<ul style="list-style-type: none"> Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants. Store potatoes in a cool, well ventilated place Don't wash potatoes before storing as dampness promotes early spoilage.
FRUIT CROPS			
Citrus	Maturity/Harvesting		<ul style="list-style-type: none"> Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plant Go for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is over Spray Imagalil 34WP @ 1g/lit against blue & green mauld in Khasi mandarin when the harvesting is over Remove all water sprouts from the base of the plant Need based spray of Carbendazim 50WP @ 1g/lit or Difenoconazole 5EC @ 0.5 ml/lit against powdery mildew.
Banana			<ul style="list-style-type: none"> Timely/regular removal of extra suckers, dry leaves & pseudo-stem
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Pigs	All stages		<ul style="list-style-type: none"> Provide concentrate feed and multi-vitamins Provide proper bedding materials for newly born piglets to protect them from cold Vaccination as per the schedule with proper consultation with vet. Deworm at regular interval During scarcity of fodder provide hay/silage/feedblocks.
Poultry	All stages		<ul style="list-style-type: none"> Provide sufficient bedding materials to keep them warm Provide adequate clean drinking water+electrolyte Vaccination as per the schedule with proper consultation with vet. Provide anti-coccidial drugs Remove wet litter Quarantine the newly procured birds for one month before introduction with old birds.
Fishery			<ul style="list-style-type: none"> Use soil-water probiotics in fish pond for making disease free environment
Dog		Canine Parvovirus	<ul style="list-style-type: none"> Vaccinate as per consultation with vet

Cattle		FMD	• Vaccination is advocated
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit
ICAR. BASAR

Agro-met Advisory Bulletin

(Period: 4th to 8th March, 2016)



GRAMIN KRISHI MAUSAM SEWA

(Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Kurung Kumey





No: 19/2015-16/Fri Dated: 4th March, 2016




Weather summery of the previous week	Weather forecast valid up to 8 th March, 2016
Rainfall = 0.0	Partially cloudy sky with light rain is forecasted for next 2-3 days. Max Temp=8°C and Min Temp=2°C, RH=47-85%, Wind Speed=4-6 kmph, Wind direction: WNW

Weather Based Agro-meteorological Advisories Bulletin

<ul style="list-style-type: none">Right time for digging pond for construction of <i>Jalkunds</i>.Go for low cost polyhouses for year round cultivation of vegetablesObtain seeds from reputed/certified/reliable sourceUse fertilizers/pesticides as per recommended doses for the locationPreference must be given on use of organic manures/bio-fertilizers & bio-pesticidesICAR Basar recommend poultry variety VANARAJA for backyard rearing			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Maize/ cowpea			<ul style="list-style-type: none">Right time for sowing of local/early maize and CowpeaSeed treatment of early maize/cowpea with Imidichloropid @ 1-2g/10kg is recommended before sowing
Toria/mustard			<ul style="list-style-type: none">On sunny days, go for harvesting of toria/mustard if mature
Maize (winter)			<ul style="list-style-type: none">Right time for harvesting

Pea		Powdery Mildew 	<ul style="list-style-type: none"> Need based spray of Carbendazim 50WP @ 1g/lit or Difenconazole 5EC @ 0.5 ml/lit against powdery mildew.
VEGETABLES			
Summer Vegetables			<ul style="list-style-type: none"> Right time for the sowing of summer vegetables like cucumber, sponge gourd, bottle gourd, ladies finger etc. Get seeds/planting materials of preferred or recommended varieties from right sources
Pea/Cauliflower/Cabbage	Vegetative	Leaf eating insects 	<ul style="list-style-type: none"> On clear weather, spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in cabbage/cauliflower/pea & other vegetables is advocated.
Potato	Harvesting	Ants 	<ul style="list-style-type: none"> Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants. Store potatoes in a cool, well ventilated place Don't wash potatoes before storing as dampness promotes early spoilage.
FRUIT CROPS			
Citrus	Maturity/Harvesting		<ul style="list-style-type: none"> Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plant Go for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is over During dry weather, spray Imagalil 34WP @ 1g/lit against blue & green mould in Khasi mandarin when the harvesting is over Remove all water sprouts from the base of the plant Need based spray of Carbendazim 50WP @ 1g/lit or Difenconazole 5EC @ 0.5 ml/lit against powdery mildew.
Banana			<ul style="list-style-type: none"> Timely/regular removal of extra suckers, dry leaves & pseudo-stem
Apple/Kiwi			<ul style="list-style-type: none"> Traning & Pruning on priority to remove diseased and crisscross branches to maintain health and shape of plants Best time to apply fertilizers to the apple trees containing potassium, calcium, Boron and nitrogen Fertilizers should be applied a foot away from the base of the plants
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Pigs	All stages		<ul style="list-style-type: none"> Provide concentrate feed and multi-vitamins Provide proper bedding materials for newly born piglets to protect them from cold Vaccination as per the schedule with proper consultation with vet. Deworm at regular interval During scarcity of fodder provide hay/silage/feedblocks.
Poultry	All stages		<ul style="list-style-type: none"> Avoid rearing of Day-Old-Chicks Provide sufficient bedding materials to keep them warm Provide adequate clean drinking water+electrolyte Vaccination as per the schedule with proper consultation with vet.

			<ul style="list-style-type: none">• Provide anti-coccidial drugs• Remove wet litter/provide artificial heating and light• Quarantine the newly procured birds for one month before introduction with old birds.
Fishery			<ul style="list-style-type: none">• Use soil-water probiotics in fish pond for making disease free environment
Dog		Canine Parvovirus	<ul style="list-style-type: none">• Vaccinate as per consultation with vet
Cattle		FMD	<ul style="list-style-type: none">• Vaccination is advocated
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit
ICAR, BASAR

Agro-met Advisory Bulletin

(Period: 4th to 8th March, 2016)



GRAMIN KRISHI MAUSAM SEWA

(Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Lohit





No: 19/2015-16/Fri **Dated: 4th March, 2016**



Weather summery of the previous week	Weather forecast valid up to 8 th March, 2016
Rainfall = 0.0	Moderate cloud cover with possibility of rain in few places. Max Temp=18°C and Min Temp=10°C, RH=28-82%, Wind Speed=6-7 kmph, Wind direction: E
Warning:	Non

Weather Based Agro-meteorological Advisories Bulletin

<ul style="list-style-type: none">Right time for digging pond for construction of <i>Jalkunds</i>.Go for low cost polyhouses for year round cultivation of vegetablesObtain seeds from reputed/certified/reliable sourceUse fertilizers/pesticides as per recommended doses for the locationPreference must be given on use of organic manures/bio-fertilizers & bio-pesticidesICAR Basar recommend poultry variety VANARAJA for backyard rearing			
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Maize/ cowpea			<ul style="list-style-type: none">Right time for sowing of local/early maize and CowpeaSeed treatment of early maize/cowpea with Imidichloropid @ 1-2g/10kg is recommended before sowing
Toria/mustard			<ul style="list-style-type: none">On sunny days, go for harvesting of toria/mustard if mature
Maize (winter)			<ul style="list-style-type: none">Right time for harvesting

Pea		Powdery Mildew 	<ul style="list-style-type: none"> Need based spray of Carbendazim 50WP @ 1g/lit or Difenconazole 5EC @ 0.5 ml/lit against powdery mildew.
VEGETABLES			
Summer Vegetables			<ul style="list-style-type: none"> Right time for the sowing of summer vegetables like cucumber, sponge gourd, bottle gourd, ladies finger etc. Get seeds/planting materials of preferred or recommended varieties from right sources
Pea/Cauliflower/Cabbage	Vegetative	Leaf eating insects 	<ul style="list-style-type: none"> On clear weather, spray of Azadirachtin 0.15% EC @ 6ml/lit against leaf eating insects in cabbage/cauliflower/pea & other vegetables is advocated using stickers.
Potato	Harvesting	Ants 	<ul style="list-style-type: none"> Apply Neem-based granular insecticides in soil against soil inhabiting insect pests/ants. Store potatoes in a cool, well ventilated place Don't wash potatoes before storing as dampness promotes early spoilage.
FRUIT CROPS			
Citrus	Maturity/Harvesting		<ul style="list-style-type: none"> Prune away all diseased portion of the citrus plant after harvest of the fruit to maintain the health of the plant Go for drenching with Dichlorvos 76 EC @ 0.05% against trunk borer in citrus trees only if the harvesting is over On dry weather, spray Imagalil 34WP @ 1g/lit against blue & green mould in Khasi mandarin when the harvesting is over Remove all water sprouts from the base of the plant Need based spray of Carbendazim 50WP @ 1g/lit or Difenconazole 5EC @ 0.5 ml/lit against powdery mildew.
Banana			<ul style="list-style-type: none"> Timely/regular removal of extra suckers, dry leaves & pseudo-stem
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Pigs	All stages		<ul style="list-style-type: none"> Provide concentrate feed and multi-vitamins Provide proper bedding materials for newly born piglets to protect them from cold Vaccination as per the schedule with proper consultation with vet. Deworm at regular interval During scarcity of fodder provide hay/silage/feedblocks.
Poultry	All stages		<ul style="list-style-type: none"> Avoid rearing of Day-Old-Chicks Provide sufficient bedding materials to keep them warm Provide adequate clean drinking water+electrolyte Vaccination as per the schedule with proper consultation with vet. Provide anti-coccidial drugs Remove wet litter/provide artificial heating and light Quarantine the newly procured birds for one month before introduction with old birds.
Fishery			<ul style="list-style-type: none"> Use soil-water probiotics in fish pond for making disease free environment

Dog		Canine Parvovirus	• Vaccinate as per consultation with vet
Cattle		FMD	• Vaccination is advocated
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit
ICAR, BASAR