

कृषि मौसम सलाहकार सेवा पत्रिका

Agro-met Advisory Bulletin

(Period: 24th to 27th February 2018)

ग्रामीण कृषि मौसम सेवा



GRAMIN KRISHI MAUSAM SEWA
(Integrated Agro-Meteorological Advisory Services)



AMFU, ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Anjaw

No: AMFU-Basar/2018/16

Dated: Fri, 23rd February, 2018



Hawai: Latitude: 28.2179994/ Longitude: 94.7277528/Altitude: 1296 m amsl

Weather summary of the previous week	Weather forecast valid up to 28 th February 2018
No weather summary available	Sunny/partly cloudy with possibility of light/moderate rainfall on 24th and 27th Feb. Daily maximum and minimum temperature for the following period is predicted to be 23-24°C and 10-11°C respectively. Expected morning and evening relative humidity will be 64-89% and 43-62% respectively. Wind direction is mainly SSE and E with average wind speed range of 7-9 kmph.
Warning:	• Nil

Weather Based Agro-meteorological Advisories Bulletin

General Recommendations

- **Spraying operation should be done only on dry/clear weather**
- **Obtain seeds from reputed/certified/reliable source**
- **Use fertilizers/pesticides as per recommended doses for the location**
- **Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides**
- **Warning: Spraying of DDT in crops is lethal**

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Pea	Pod formation	Powdery mildew/ leaf eating insects	<ul style="list-style-type: none"> Need based foliar spray of Dinocap (0.2%) or Carbendazim 50WP (0.1%) or Propiconazole (0.1%) on clear/dry days or using sticker in case of heavy infestation
Toria/ Mustard	Seed formation	Aphids	<ul style="list-style-type: none"> In case of heavy infestation, Spray Dimethoate 30EC @2ml/litre against aphids in Rapeseed/Mustard using sticker.
VEGETABLES			
Ginger/ Turmeric	Maturity		<ul style="list-style-type: none"> Harvest the mature crop on clear day, Cure should be taken while harvest so that the rhizome is not damaged,
Potato	Vegetative stage/ Harvesting stage	Late blight/ Red ant	<ul style="list-style-type: none"> Avoid harvesting Store already harvested crops in a cool, well ventilated place/ do not wash before storing as dampness promotes early spoilage. Regularly monitor against the disease symptoms. Prominent symptoms include small, dark, circular to irregular shaped lesion in the leaves Prophylactic spraying of Mancozeb @ 2 gram per 4 litres of water is advised using stickers in case of rainfall..
Cabbage/ Cauliflower	Harvesting		<ul style="list-style-type: none"> Harvest mature crop. Constantly monitor against attack of insects pests/mole.
HORTICULTURAL CROPS			
Kiwi	Pruning/Training		<ul style="list-style-type: none"> Diseased and senile branches should be removed. Pruning should be done retaining 8-10 buds per branch for enhancing new flush and flowering. Pruned healthy hard wood of pencil thickness size having 4-5 buds can be used as cutting for propagation
Banana	Bunching	Cold	<ul style="list-style-type: none"> Cover newly emerged bunches with locally available sack to protect from cold/frost Mulching with straw/crop residue at root zone to avoid frost injury in banana.
Pineapple	Harvesting		<ul style="list-style-type: none"> After winter harvest, collection/Selection of suckers & slip to be done for next season planting
Citrus	Maturity/ harvesting	Bark eating caterpillar, Trunk borer, Blue & Green mould	<ul style="list-style-type: none"> Orange (only if harvesting is over): Go for drenching with Dichlorovos 76 EC@0.05% against trunk-borer on dry days. On clear weather or using sticker, against Blue & Green mould, spray Imazalil 34 WP @ 1g/lit. Prune away the diseased portion of the plant. In case of Bark eating caterpillar, remove the wooden frass present in between the joints of the trunk and inject Dichlorovos (DDVP) 76 EC 0.1 percent @ 5ml in the larval tunnel with the help of disposable syringe and push cotton swab soaked in Dichlorovos solution in the larval tunnel.
Temperate fruits	After harvesting		<ul style="list-style-type: none"> Training & pruning of the temperate crops (apple/pears/peach) is recommended In case of flowering, stop spraying insecticides/pesticides
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Poultry	All stages		<ul style="list-style-type: none"> Provide shelter to the backyard/scavenging poultry. Feed grains in the morning/evening to supplement scavenging. Day old chick-Maintain warmth by lighting, In present weather conditions protect the birds from cold by using gunny bags/plastic sheets around the poultry shed during night, Provide heat by lighting 100W bulb @ 1 bulb per 20 newly born chicks.
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.) Swine Fever	<ul style="list-style-type: none"> Avoid dampness and cover weak and sick animals with sack cloth. The bedding is always dry and that there is plenty. If possible wooden sleeping box should be provided. Animals should be given lukewarm feed and water.

			<ul style="list-style-type: none"> • Right time to deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight.
Fishery			<ul style="list-style-type: none"> • Keep fish stoking density thin @1200/ha
Cattle	All stages	FMD	<ul style="list-style-type: none"> • Animals should be given lukewarm feed and water. • Right time to deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight. • Provide proper bedding materials especially to new-born and lukewarm food & water • Provide feed supplements
Mithun		FMD	<ul style="list-style-type: none"> • Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route. Pregnant animals above 7 months of pregnancy should be avoided from vaccination.
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

कृषि मौसम सलाहकार सेवा पत्रिका

Agro-met Advisory Bulletin

(Period: 24th to 27th February 2018)

ग्रामीण कृषि मौसम सेवा



GRAMIN KRISHI MAUSAM SEWA
(Integrated Agro-Meteorological Advisory Services)




AMFU, ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Changlang

No: AMFU-Basar/2018/16

Dated: Fri, 23rd February, 2018



Changlang (HQ): Latitude: 27.1305856/Longitude: 95.7342373/Altitude: 580 masl

Weather summary of the previous week	Weather forecast valid up to 28 th February 2018
Rainfall = Changlang: 0.0 mm Miao: 4.0 mm	Mostly/partly cloudy with possibility of light/moderate rainfall. Daily maximum and minimum temperature for the following period is predicted to be 28-29°C and 15-16°C respectively. Expected morning and evening relative humidity will be 69-78% and 29-47% respectively. Wind direction is mainly ESE with speed range of 5-6 kmph.
Warning	• Nil

Weather Based Agro-meteorological Advisories Bulletin

General Recommendations

- **Spraying operation should be done only on dry/clear weather**
- **Go for low cost polyhouses for year round cultivation of vegetables**
- **Obtain seeds from reputed/certified/reliable source**
- **Use fertilizers/pesticides as per recommended doses for the location**
- **Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides**
- **ICAR Basar recommend poultry variety VANARAJA for backyard rearing**

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Pea	Pod mature	Birds	<ul style="list-style-type: none"> • Harvest of green pods, • Use of scare crow and tying of bird scare tape around the crop field,
Toria/ Mustard	Vegetative/ Flowering	Aphid	<ul style="list-style-type: none"> • Harvesting of mature crop, • Drying of harvested crop to protect against disease development, • Spraying of dimethoate @ 2 ml per litre of water is advise during clear/dry days or using stickers
Maize	Siliqua maturation and harvest	Rats	<ul style="list-style-type: none"> • Protect the crop against attack of rats using locally available traps, • When rat menace has occurred application of multi dose rodenticide like bromadiolone with proper consultation with experts is advised.
VEGETABLES			
Potato	Vegetative stage/ Harvesting stage	Late blight/ Red ant	<ul style="list-style-type: none"> • Avoid harvesting • If already harvested, store in a cool, well ventilated place/ do not wash before storing as dampness promotes early spoilage. • Regularly monitor against the disease symptoms. Prominent symptoms include small, dark, circular to irregular shaped lesion in the leaves • Prophylactic spraying of Mancozeb @ 2 gram per 4 litres of water is advised using stickers in case of rainfall..
Cabbage/ Cauliflower	Harvesting		<ul style="list-style-type: none"> • Harvest mature crop. • Constantly monitor against attack of insects pests/mole.
Ginger/ Turmeric	Maturity		<ul style="list-style-type: none"> • Harvest the mature crop, • Cure should be taken while harvest so that the rhizome is not damaged,
HORTICULTURAL CROPS			
Citrus	Maturity/ harvesting	Bark eating caterpillar, Trunk borer, Blue & Green mould	<ul style="list-style-type: none"> • Orange (only if harvesting is over): Go for drenching with Dichlorovos 76 EC@0.05% against trunk-borer. • Against Blue & Green mould, spray Imazalil 34 WP @ 1g/lit. Prune away the diseased portion of the plant using sticker in case of rainfall. • In case of Bark eating caterpillar, remove the wooden frass present in between the joints of the trunk and inject Dichlorovos (DDVP) 76 EC 0.1 percent @ 5ml in the larval tunnel with the help of disposable syringe and push cotton swab soaked in Dichlorovos solution in the larval tunnel.
Banana	Bunching	Cold	<ul style="list-style-type: none"> • Cover newly emerged bunches with locally available sack to protect from cold/frost • Mulching with straw/crop residue at root zone to avoid frost injury in banana.
Honey Bees			
Honey bees			<ul style="list-style-type: none"> • Box must be covered with straw/dry leaves to provide warmth • Provide sugar syrup
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
General			
<ul style="list-style-type: none"> • Damaged grain (not fungus contaminated) to be diverted for Poultry/livestock feeding • Clean drinking water should be ensured round the clock and water troughs/feeders should be regularly cleaned, 			
Poultry	All stages		<ul style="list-style-type: none"> • Provide shelter to the backyard/scavenging poultry. Feed grains in the morning/evening to supplement scavenging. • Day old chick-Maintain warmth by lighting, • In present weather conditions protect the birds from cold by using gunny bags/plastic sheets around the poultry shed during night, • Provide heat by lighting 100W bulb @ 1 bulb per 20 newly born chicks.
Pigs	All stages	FMD (high fever and appearance	<ul style="list-style-type: none"> • Avoid dampness and cover weak and sick animals with

		of mucous membrane of the mouth and foot.) Swine Fever	<p>sack cloth.</p> <ul style="list-style-type: none"> • The bedding is always dry and that there is plenty. • If possible wooden sleeping box should be provided. • Animals should be given lukewarm feed and water. • Right time to deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight.
Goat	All stages		<ul style="list-style-type: none"> • Deworming for gastrointestinal parasite
Cattle	All stages	FMD	<ul style="list-style-type: none"> • Animals should be given lukewarm feed and water. • Right time to deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight. • Provide proper bedding materials especially to new-born and lukewarm food & water • Provide feed supplements
<p>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247</p>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit
ICAR, BASAR**

कृषि मौसम सलाहकार सेवा पत्रिका

Agro-met Advisory Bulletin

(Period: 24th to 27th February 2018)

ग्रामीण कृषि मौसम सेवा



GRAMIN KRISHI MAUSAM SEWA
(Integrated Agro-Meteorological Advisory Services)




AMFU, ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Dibang Valley

No: AMFU-Basar/2018/16

Dated: Fri, 23rd February, 2018



Anini: Latitude: 28.2179994/Longitude: 94.7277528/Altitude: 1968 masl

Weather summary of the previous week	Weather forecast valid up to 28 th February 2018
Rainfall = 27.0 mm (Anini)	Partly cloudy with possibility of moderate rainfall on 24 th and 27 th Feb. Daily maximum and minimum temperature for the following period is predicted to be 20-21°C and 08-09°C respectively. Expected morning and evening relative humidity will be 82-96% and 42-54% respectively. Wind direction is mainly SSE with speed range of 6-9 kmph.
Warning:	• Nil

Weather Based Agro-meteorological Advisories Bulletin

General recommendations

- Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated
- Go for spray operation only on dry/clear weather.
- Go for low cost polyhouses for year round cultivation of vegetables
- Obtain seeds from reputed/certified/reliable source
- Use fertilizers/pesticides as per recommended doses for the location
- Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides
- ICAR Basar recommend poultry variety VANARAJA for backyard rearing

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Pea	Pod formation	Powdery mildew/ leaf eating insects	<ul style="list-style-type: none"> In case of heavy infestation, need based foliar spray of Dinocap (0.2%) or Carbendazim 50WP (0.1%) or Propiconazole (0.1%) during clear/dry days or using appropriate sticker.
Toria/ Mustard	Seed formation	Aphids	<ul style="list-style-type: none"> In case of heavy infestation, on clear/dry days or using sticker, spray Dimethoate 30EC @2ml/litre against aphids in Rapeseed/Mustard.
VEGETABLES			
Potato	Maturity/ Harvesting		<ul style="list-style-type: none"> Avoid harvesting Store already harvested potatoes in a cool, well ventilated place/ do not wash before storing as dampness promotes early spoilage.
Cabbage/ Cauliflower	Harvesting		<ul style="list-style-type: none"> Harvest mature crop. Constantly monitor against attack of insects pests/mole.
HORTICULTURAL CROPS			
Kiwi	Pruning/Training		<ul style="list-style-type: none"> Diseased and senile branches should be removed. Pruning should be done retaining 8-10 buds per branch for enhancing new flush and flowering. Pruned healthy hard wood of pencil thickness size having 4-5 buds can be used as cutting for propagation
Citrus	Maturity/ harvesting	Bark eating caterpillar, Trunk borer, Blue & Green mould	<ul style="list-style-type: none"> Orange (only if harvesting is over): Go for drenching with Dichlorovos 76 EC@0.05% against trunk-borer on dry days. On clear weather or using sticker, against Blue & Green mould, spray Imazalil 34 WP @ 1g/lit. Prune away the diseased portion of the plant. In case of Bark eating caterpillar, remove the wooden frass present in between the joints of the trunk and inject Dichlorovos (DDVP) 76 EC 0.1 percent @ 5ml in the larval tunnel with the help of disposable syringe and push cotton swab soaked in Dichlorovos solution in the larval tunnel.
Banana	Bunching	Cold	<ul style="list-style-type: none"> Cover newly emerged bunches with locally available sack to protect from cold/frost Mulching with straw/crop residue at root zone to avoid frost injury in banana.
Temperate fruits	After harvesting		<ul style="list-style-type: none"> Training & pruning of the temperate crops (apple/pears/peach) is recommended
Honey Bees			
Honey bees			<ul style="list-style-type: none"> Cover the boxes with locally available dry grass or straw to provide warmth Provide sugar syrup in a bowl in front of the box in case of heavy rainfall
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
General			
<ul style="list-style-type: none"> Damaged grain (not fungus contaminated) to be diverted for Poultry/livestock feeding Clean drinking water should be ensured round the clock and water troughs/feeders should be regularly cleaned, 			
Poultry	All stages		<ul style="list-style-type: none"> Provide shelter to the backyard/scavenging poultry. Feed grains in the morning/evening to supplement scavenging. Day old chick-Maintain warmth by lighting, In present weather conditions protect the birds from cold by using gunny bags/plastic sheets around the poultry shed during night, Provide heat by lighting 100W bulb @ 1 bulb per 20 newly born chicks.
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.) Swine Fever	<ul style="list-style-type: none"> Avoid dampness and cover weak and sick animals with sack cloth. The bedding is always dry and that there is plenty. If possible wooden sleeping box should be provided. Animals should be given lukewarm feed and water.

			<ul style="list-style-type: none"> • Right time to deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight.
Cattle	All stages	FMD	<ul style="list-style-type: none"> • Animals should be given lukewarm feed and water. • Right time to deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight. • Provide proper bedding materials especially to new-born and lukewarm food & water • Heavy mortality in cloven footed animals has been reported. • Supportive treatment of infected animals under supervision of veterinary department officials, • Avoid mixing of infected and healthy animals, • Keep infected animals separated from the healthy ones, • Vaccinate the healthy animals against FMD (Through State Veterinary Dept.) • Provide feed supplements
Mithun		FMD	<ul style="list-style-type: none"> • Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route. • Pregnant animals above 7 months of pregnancy should be avoided.
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit
ICAR, BASAR**

कृषि मौसम सलाहकार सेवा पत्रिका

Agro-met Advisory Bulletin

(Period: 24th to 27th February 2018)

ग्रामीण कृषि मौसम सेवा



GRAMIN KRISHI MAUSAM SEWA
(Integrated Agro-Meteorological Advisory Services)



भा.क.अनु.प. - पू.प.सं.
ICAR - RCNEH

AMFU, ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: East Kameng

No: AMFU-Basar/2018/16

Dated: Fri, 23rd February, 2018



Weather summary of the previous week	Weather forecast valid up to 28 th February 2018
Rainfall = 0.0 mm	Mostly/partly cloudy with possibility light/moderate rain. Daily maximum and minimum temperature for the following period is predicted to be 24-25°C and 12-13°C respectively. Expected morning and evening relative humidity will be 77-87% and 30-47% respectively. Expected wind direction is mainly SE & S with speed range of 5 kmph.
Warning:	<ul style="list-style-type: none"> Rain in few isolated places

Weather Based Agro-meteorological Advisories Bulletin

General Recommendations

- Go for low cost polyhouses for year round cultivation of vegetables
- Obtain seeds from reputed/certified/reliable source
- Use fertilizers/pesticides as per recommended doses for the location
- Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides
- ICAR Basar recommend poultry variety VANARAJA for backyard rearing
- Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Pea	Pod mature	Birds	<ul style="list-style-type: none"> • Harvest of green pods, • Use of scare crow and tying of bird scare tape around the crop field,
VEGETABLES			
Potato	Vegetative stage/ Harvesting stage	Late blight/ Red ant	<ul style="list-style-type: none"> • Avoid harvesting • Regularly monitor against the disease symptoms. Prominent symptoms include small, dark, circular to irregular shaped lesion in the leaves • Prophylactic spraying of Mancozeb @ 2 gram per 4 litres of water is advised using stickers in case of rainfall..
Cabbage/ Cauliflower	Harvesting		<ul style="list-style-type: none"> • Harvest mature crop. • Constantly monitor against attack of insects pests/mole.
HORTICULTURAL CROPS			
Citrus	Maturity/ harvesting	Bark eating caterpillar, Trunk borer, Blue & Green mould	<ul style="list-style-type: none"> • Orange (only if harvesting is over): Go for drenching with Dichlorovos 76 EC@0.05% against trunk-borer. • Against Blue & Green mould, spray Imazalil 34 WP @ 1g/lit on clear/dry day. Prune away the diseased portion of the plant. • In case of Bark eating caterpillar, remove the wooden frass present in between the joints of the trunk and inject Dichlorovos (DDVP) 76 EC 0.1 percent @ 5ml in the larval tunnel with the help of disposable syringe and push cotton swab soaked in Dichlorovos solution in the larval tunnel.
Banana	Bunching	Cold	<ul style="list-style-type: none"> • Cover newly emerged bunches with locally available sack to protect from cold/frost • Mulching with straw/crop residue at root zone to avoid frost injury in banana.
Large Cardamom	Vegetative/ flowering stage		<ul style="list-style-type: none"> • Provide proper drainage to avoid water stagnation in the base of the plants.
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
General			
<ul style="list-style-type: none"> • Damaged grain (not fungus contaminated) to be diverted for Poultry/livestock feeding • Clean drinking water should be ensured round the clock and water troughs/feeders should be regularly cleaned, 			
Poultry	All stages		<ul style="list-style-type: none"> • Provide shelter to the backyard/scavenging poultry. Feed grains in the morning/evening to supplement scavenging. • Day old chick-Maintain warmth by lighting, • In present weather conditions protect the birds from cold by using gunny bags/plastic sheets around the poultry shed during night, • Provide heat by lighting 100W bulb @ 1 bulb per 20 newly born chicks.
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.) Swine Fever	<ul style="list-style-type: none"> • Avoid dampness and cover weak and sick animals with sack cloth. • The bedding is always dry and that there is plenty. • If possible wooden sleeping box should be provided. • Animals should be given lukewarm feed and water. • Right time to deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight.
Cattle	All stages	FMD	<ul style="list-style-type: none"> • Animals should be given lukewarm feed and water. • Right time to deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight. • Provide proper bedding materials especially to new-born and lukewarm food & water • Provide feed supplements
Fishery			<ul style="list-style-type: none"> • Application of quick lime is highly recommended in order to rectify water pH. • Application of fish feed (mustard oil cake) is essential for achieving higher yield
Mithun		FMD	<ul style="list-style-type: none"> • Polyvalent FMD vaccine with serotype O, A and Asia at 6 months

			of age and repeat after that 6 months interval by sub-cutaneous route. <ul style="list-style-type: none">• Pregnant animals above 7 months of pregnancy should be avoided.
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit
ICAR, BASAR**

कृषि मौसम सलाहकार सेवा पत्रिका

Agro-met Advisory Bulletin

(Period: 24th to 27th February 2018)

ग्रामीण कृषि मौसम सेवा



GRAMIN KRISHI MAUSAM SEWA
(Integrated Agro-Meteorological Advisory Services)



भा.क.अनु.प. - पू.प.सं.
ICAR - RCNEH

AMFU, ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: East Siang

No: AMFU-Basar/2018/16

Dated: Fri, 23rd February, 2018



Pasighat: Latitude: 28.0700/Longitude: 95.3300/Altitude: 153 masl

Weather summary of the previous week	Weather forecast valid up to 28 th February 2018
Rainfall: 0.0 mm; T. Max: 28.0°C; T. Min: 15.0°C ; RH: 54-69%	Mostly/partly cloudy with possibility of light/moderate rainfall. Daily maximum and minimum temperature for the following period is predicted to be 26-28°C and 16-17°C respectively. Expected morning and evening relative humidity will be 69-86% and 22-41% respectively. Expected wind direction is mainly S with speed range of 6 kmph.
Warning:	• Nil

Weather Based Agro-meteorological Advisories Bulletin

General Recommendations

- Go for low cost polyhouses for year round cultivation of vegetables
- Obtain seeds from reputed/certified/reliable source
- Use fertilizers/pesticides as per recommended doses for the location
- Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides
- ICAR Basar recommend poultry variety VANARAJA for backyard rearing

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Pea	Pod mature	Birds	<ul style="list-style-type: none"> • Harvest of green pods, • Use of scare crow and tying of bird scare tape around the crop field,
Toria/ Mustard	Vegetative/ Flowering	Aphid	<ul style="list-style-type: none"> • Harvesting of mature crop, • If only infestation is high, Spraying of dimethoate @ 2 ml per litre of water is advised using proper sticker,
VEGETABLES			
Potato	Vegetative stage/ Harvesting stage	Red ant	<ul style="list-style-type: none"> • Avoid harvesting • If already harvested, Store in a cool, well ventilated place/ do not wash before storing as dampness promotes early spoilage.
Cabbage/ Cauliflower	Harvesting		<ul style="list-style-type: none"> • Harvest mature crop. • Constantly monitor against attack of insects pests/mole.
Ginger/ Turmeric	Maturity		<ul style="list-style-type: none"> • Harvest the mature crop, • Cure should be taken while harvest so that the rhizome is not damaged,
HORTICULTURAL CROPS			
Kiwi	Pruning/Training		<ul style="list-style-type: none"> • Diseased and senile branches should be removed. • Pruning should be done retaining 8-10 buds per branch for enhancing new flush and flowering. • Pruned healthy hard wood of pencil thickness size having 4-5 buds can be used as cutting for propagation
Citrus	Maturity/ harvesting	Bark eating caterpillar, Trunk borer, Blue & Green mould	<ul style="list-style-type: none"> • Orange (only if harvesting is over): Go for drenching with Dichlorovos 76 EC@0.05% against trunk-borer on dry days. • On clear weather or using sticker, against Blue & Green mould, spray Imazalil 34 WP @ 1g/lit. Prune away the diseased portion of the plant. • In case of Bark eating caterpillar, remove the wooden frass present in between the joints of the trunk and inject Dichlorovos (DDVP) 76 EC 0.1 percent @ 5ml in the larval tunnel with the help of disposable syringe and push cotton swab soaked in Dichlorovos solution in the larval tunnel.
Honey Bees			
Honey bees			<ul style="list-style-type: none"> • Cover the boxes with locally available dry grass or straw to provide warmth
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
General			
<ul style="list-style-type: none"> • Damaged grain (not fungus contaminated) to be diverted for Poultry/livestock feeding • Clean drinking water should be ensured round the clock and water troughs/feeders should be regularly cleaned, 			
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.) Swine Fever	<ul style="list-style-type: none"> • Avoid dampness and cover weak and sick animals with sack cloth. • Animals should be given lukewarm feed and water. • Right time to deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight. • Provide proper bedding materials especially to new-born piglets and lukewarm food & water
Cattle	All stages	FMD	<ul style="list-style-type: none"> • Animals should be given lukewarm feed and water. • Right time to deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight. • Provide proper bedding materials especially to new-born and lukewarm food & water • Heavy mortality in cloven footed animals has been reported. • Supportive treatment of infected animals under supervision of veterinary department officials, • Avoid mixing of infected and healthy animals, • Keep infected animals separated from the healthy ones, • Vaccinate the healthy animals against FMD (Through State Veterinary Dept.)

			<ul style="list-style-type: none"> • Provide feed supplements
Poultry	All stages		<ul style="list-style-type: none"> • Day old chick-Maintain warmth by lighting, • In present weather conditions protect the birds from cold by using gunny bags/plastic sheets around the poultry shed during night, • Provide heat by lighting 100W bulb @ 1 bulb per 20 newly born chicks.
Mithun		FMD	<ul style="list-style-type: none"> • Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by subcutaneous route. • Pregnant animals above 7 months of pregnancy should be avoided.
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit
ICAR, BASAR**

कृषि मौसम सलाहकार सेवा पत्रिका

Agro-met Advisory Bulletin

(Period: 24th to 27th February 2018)

ग्रामीण कृषि मौसम सेवा



GRAMIN KRISHI MAUSAM SEWA

(Integrated Agro-Meteorological Advisory Services)



भा.क.अनु.प. - पू.प.सं.
ICAR - RCNEH

AMFU, ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Kurung Kumey

No: AMFU-Basar/2018/16

Dated: Fri, 23rd February, 2018



Koloriang: Latitude: 27.9167°/Longitude: 93.3500°/Altitude: 1040 masl

Weather summary of the previous week	Weather forecast valid up to 28 th February 2018
Past weather summary not available	Mostly cloudy with possibility of light/moderate rain. Daily maximum and minimum temperature for the following period is predicted to be 19-20°C and 07-09°C respectively. Expected morning and evening relative humidity will be 58-89% and 24-54% respectively. Wind direction is mainly variable with wind speed range of 4-18 kmph.
Warning	• Nil

Weather Based Agro-meteorological Advisories Bulletin

General Recommendations

- **Spray operations should be done only on dry/clear weather or in extreme case use stickers**
- **Go for low cost polyhouses for year round cultivation of vegetables**
- **Obtain seeds from reputed/certified/reliable source**
- **Use fertilizers/pesticides as per recommended doses for the location**
- **Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides**
- **ICAR Basar recommend poultry variety VANARAJA for backyard rearing**
- **Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated**

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Pea	Pod formation	Powdery mildew/ leaf eating insects	<ul style="list-style-type: none"> Need based foliar spray of Dinocap (0.2%) or Carbendazim 50WP (0.1%) or Propiconazole (0.1%) in case of heavy infestation using proper stickers.
Toria/ Mustard	Seed formation	Aphids	<ul style="list-style-type: none"> Avoid spray operation In case of heavy infestation, Spray Dimethoate 30EC @2ml/litre against aphids in Rapeseed/Mustard using appropriate stickers@ 1ml/lit.
VEGETABLES			
Potato	Vegetative stage/ Harvesting stage	Late blight/ Red ant	<ul style="list-style-type: none"> Avoid harvesting Regularly monitor against the disease symptoms. Prominent symptoms include small, dark, circular to irregular shaped lesion in the leaves Prophylactic spraying of Mancozeb @ 2 gram per 4 litres of water is advised using stickers in case of rainfall..
Cabbage/ Cauliflower	Harvesting		<ul style="list-style-type: none"> Harvest mature crop. Constantly monitor against attack of insects pests/mole.
HORTICULTURAL CROPS			
Temperate fruits	After harvesting		<ul style="list-style-type: none"> Training & pruning of the temperate crops (apple/pears/peach) is recommended In case of flowering, stop spraying insecticides/pesticides
Citrus	Maturity/ harvesting	Bark eating caterpillar, Trunk borer, Blue & Green mould	<ul style="list-style-type: none"> Orange (only if harvesting is over): Go for drenching with Dichlorovos 76 EC@0.05% against trunk-borer. Against Blue & Green mould, spray Imazalil 34 WP @ 1g/lit using sticker in case of rainfall. Prune away the diseased portion of the plant. In case of Bark eating caterpillar, remove the wooden frass present in between the joints of the trunk and inject Dichlorovos (DDVP) 76 EC 0.1 percent @ 5ml in the larval tunnel with the help of disposable syringe and push cotton swab soaked in Dichlorovos solution in the larval tunnel.
Pineapple	Harvesting		<ul style="list-style-type: none"> After winter harvest, collection/Selection of suckers & slip to be done for next season planting
Banana	Bunching	Cold	<ul style="list-style-type: none"> Cover newly emerged bunches with locally available sack to protect from cold/frost Mulching with straw/crop residue at root zone to avoid frost injury in banana.
Kiwi	Pruning/Training		<ul style="list-style-type: none"> Diseased and senile branches should be removed. Pruning should be done retaining 8-10 buds per branch for enhancing new flush and flowering. Pruned healthy hard wood of pencil thickness size having 4-5 buds can be used as cutting for propagation
Honey Bees			
			<ul style="list-style-type: none"> Box must be covered with straw/dry leaves to provide warmth
LIVESTOCKS			
General			
<ul style="list-style-type: none"> Damaged grain (not fungus contaminated) to be diverted for Poultry/livestock feeding Clean drinking water should be ensured round the clock and water troughs/feeders should be regularly cleaned, 			
Livestock	Stage	Disease	Precautionary Practices to be followed
Cattle	All stages	FMD	<ul style="list-style-type: none"> Animals should be given lukewarm feed and water. Right time to deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight. Provide proper bedding materials especially to new-born and lukewarm food & water Provide feed supplements
Poultry	All stages		<ul style="list-style-type: none"> Provide shelter to the backyard/scavenging poultry. Feed grains in the morning/evening to supplement scavenging. Day old chick-Maintain warmth by lighting, In present weather conditions protect the birds from cold by

			<p>using gunny bags/plastic sheets around the poultry shed during night,</p> <ul style="list-style-type: none"> • Provide heat by lighting 100W bulb @ 1 bulb per 20 newly born chicks.
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.) Swine Fever	<ul style="list-style-type: none"> • Avoid dampness and cover weak and sick animals with sack cloth. • The bedding is always dry and that there is plenty. • If possible wooden sleeping box should be provided. • Animals should be given lukewarm feed and water. • Right time to deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight.
Mithun		FMD	<ul style="list-style-type: none"> • Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route. • Pregnant animals above 7 months of pregnancy should be avoided.
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit
ICAR, BASAR**

कृषि मौसम सलाहकार सेवा पत्रिका

Agro-met Advisory Bulletin

(Period: 24th to 27th February 2018)

ग्रामीण कृषि मौसम सेवा



GRAMIN KRISHI MAUSAM SEWA

(Integrated Agro-Meteorological Advisory Services)



भा.क.अनु.प. - पू.प.सं.
ICAR - RCNEH

AMFU, ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Lohit

No: AMFU-Basar/2018/16

Dated: Fri, 23rd February, 2018



Weather summary of the previous week	Weather forecast valid up to 28 th February 2018
Rainfall = 0.0 mm	Mostly/partly cloudy with possibility of moderate rainfall. Daily maximum and minimum temperature for the following period is predicted to be 27-28°C and 14-15°C respectively. Expected morning and evening relative humidity will be 84-92% and 37-58% respectively. Wind direction is mainly ENE with speed range of 6-7 kmph.
Warning:	• Nil

Weather Based Agro-meteorological Advisories Bulletin

General Recommendations

- Spraying operation should be done only on dry/clear weather
- Go for low cost poly-houses for year round cultivation of vegetables
- Obtain seeds from reputed/certified/reliable source
- Use fertilizers/pesticides as per recommended doses for the location
- Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides
- ICAR Basar recommend poultry variety VANARAJA for backyard rearing.

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Pea	Pod mature	Birds	<ul style="list-style-type: none"> • Harvest of green pods, • Use of scare crow and tying of bird scare tape around the crop field,
Toria/ Mustard	Vegetative/ Flowering	Aphid	<ul style="list-style-type: none"> • Harvest mature crops on clear/dry days • In case of heavy infestation, Spraying of dimethoate @ 2 ml per litre of water is advised using proper stickers@ 1ml/lit,
Maize	Vegetative/ Maturity/ Harvesting	Rats	<ul style="list-style-type: none"> • Protect the mature crop against attack of rats using locally available traps, • When rat menace has occurred application of multi dose rodenticide like bromadiolone with proper consultation with experts is advised.
VEGETABLES			
Potato	Vegetative stage/ Harvesting stage	Late blight/ Red ant	<ul style="list-style-type: none"> • Avoid harvesting • Regularly monitor against the disease symptoms. Prominent symptoms include small, dark, circular to irregular shaped lesion in the leaves • Prophylactic spraying of Mancozeb @ 2 gram per 4 litres of water is advised using stickers in case of rainfall..
Cabbage/ Cauliflower	Harvesting		<ul style="list-style-type: none"> • Harvest mature crop. • Constantly monitor against attack of insects pests/mole.
Ginger/ Turmeric	Maturity		<ul style="list-style-type: none"> • Harvest the mature crop, • Cure should be taken while harvest so that the rhizome is not damaged,
HORTICULTURAL CROPS			
Kiwi	Pruning/Training		<ul style="list-style-type: none"> • Diseased and senile branches should be removed. • Pruning should be done retaining 8-10 buds per branch for enhancing new flush and flowering. • Pruned healthy hard wood of pencil thickness size having 4-5 buds can be used as cutting for propagation
Citrus	Maturity/ harvesting	Bark eating caterpillar, Trunk borer, Blue & Green mould	<ul style="list-style-type: none"> • Orange (only if harvesting is over): Go for drenching with Dichlorovos 76 <u>EC@0.05%</u> against trunk-borer on dry days. • On clear weather or using sticker, against Blue & Green mould, spray Imazalil 34 WP @ 1g/lit. Prune away the diseased portion of the plant. • In case of Bark eating caterpillar, remove the wooden frass present in between the joints of the trunk and inject Dichlorovos (DDVP) 76 EC 0.1 percent @ 5ml in the larval tunnel with the help of disposable syringe and push cotton swab soaked in Dichlorovos solution in the larval tunnel.
Large Cardamom	Fruiting	Chirkey Disease	<ul style="list-style-type: none"> • Remove dry/diseased/old tillers & branches & spray Bio NPK@ 2ml/lit water for proper growth on clear/dry days.
Banana	All stages	Sigatoka Leaf spot	<ul style="list-style-type: none"> • One sword sucker may be allowed after shooting (Flowering), • Periodical removal of side suckers by cutting them above the ground level, scoop the core and pour 2 ml of kerosene in the core, • Foliar spray of Propiconazole (e.g., Tilt) @ 1 ml per litre of wate using stickers
Honey Bees			
Honey bees			<ul style="list-style-type: none"> • Cover the boxes with locally available dry grass or straw to provide warmth
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
General			
<ul style="list-style-type: none"> • Damaged grain (not fungus contaminated) to be diverted for Poultry/livestock feeding • Clean drinking water should be ensured round the clock and water troughs/feeders should be regularly cleaned, 			
Poultry	All stages		<ul style="list-style-type: none"> • Provide shelter to the backyard/scavenging poultry. Feed grains in the morning/evening to supplement scavenging. • Day old chick-Maintain warmth by lighting,

			<ul style="list-style-type: none"> • In present weather conditions protect the birds from cold by using gunny bags/plastic sheets around the poultry shed during night, • Provide heat by lighting 100W bulb @ 1 bulb per 20 newly born chicks.
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.) Swine Fever	<ul style="list-style-type: none"> • Avoid dampness and cover weak and sick animals with sack cloth. • The bedding is always dry and that there is plenty. • If possible wooden sleeping box should be provided. • Animals should be given lukewarm feed and water. • Right time to deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight.
Cattle	All stages	FMD	<ul style="list-style-type: none"> • Animals should be given lukewarm feed and water. • Right time to deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight. • Provide proper bedding materials especially to new-born and lukewarm food & water • Heavy mortality in cloven footed animals has been reported. • Supportive treatment of infected animals under supervision of veterinary department officials, • Avoid mixing of infected and healthy animals, • Keep infected animals separated from the healthy ones, • Vaccinate the healthy animals against FMD (Through State Veterinary Dept.) • Provide feed supplements
Mithun		FMD	<ul style="list-style-type: none"> • Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route. • Pregnant animals above 7 months of pregnancy should be avoided.

Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit
ICAR, BASAR**

कृषि मौसम सलाहकार सेवा पत्रिका

Agro-met Advisory Bulletin

(Period: 24th to 27th February 2018)

ग्रामीण कृषि मौसम सेवा



GRAMIN KRISHI MAUSAM SEWA

(Integrated Agro-Meteorological Advisory Services)



भा.क.अनु.पू.प.सं.
ICAR-RCNEH

AMFU, ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Lower Dibang Valley

No: AMFU-Basar/2018/16

Dated: Fri, 23rd February, 2018



Weather summary of the previous week	Weather forecast valid up to 28 th February 2018
Rainfall = 0.0 mm (Roing)	Mostly cloudy with possibility of light/moderate rain. Daily maximum and minimum temperature for the following period is predicted to be 24-25°C and 12-13°C respectively. Expected morning and evening relative humidity will be 81-87% and 30-47% respectively. Wind direction is mainly SSE with speed range of 6-7 kmph.
Warning:	• Nil

Weather Based Agro-meteorological Advisories Bulletin

General Recommendations

- Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated
- Spraying operation should be done only on dry/clear weather
- Go for low cost polyhouses for year round cultivation of vegetables
- Obtain seeds from reputed/certified/reliable source
- Use fertilizers/pesticides as per recommended doses for the location
- Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides
- ICAR Basar recommend poultry variety VANARAJA for backyard rearing

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Pea	Pod mature	Birds	<ul style="list-style-type: none"> • Harvest of green pods, • Use of scare crow and tying of bird scare tape around the crop field,
Toria/ Mustard	Vegetative/ Flowering	Aphid	<ul style="list-style-type: none"> • In case of heavy infestation, Spraying of dimethoate @ 2 ml per litre of water is advised using appropriate stickers,
Maize	Vegetative/ Maturity/ Harvesting	Rats	<ul style="list-style-type: none"> • Protect the mature crop against attack of rats using locally available traps, • When rat menace has occurred application of multi dose rodenticide like bromadiolone with proper consultation with experts is advised.
VEGETABLES			
Potato	Vegetative stage/ Harvesting stage	Late blight/ Red ant	<ul style="list-style-type: none"> • Avoid harvesting • Regularly monitor against the disease symptoms. Prominent symptoms include small, dark, circular to irregular shaped lesion in the leaves • Prophylactic spraying of Mancozeb @ 2 gram per 4 litres of water is advised using stickers in case of rainfall..
Cabbage/ Cauliflower	Harvesting		<ul style="list-style-type: none"> • Harvest mature crop. • Constantly monitor against attack of insects pests/mole.
HORTICULTURAL CROPS			
Citrus	Maturity/ harvesting	Bark eating caterpillar, Trunk borer, Blue & Green mould	<ul style="list-style-type: none"> • Orange (only if harvesting is over): Go for drenching with Dichlorovos 76 EC@0.05% against trunk-borer on dry days. • On clear weather or using sticker, against Blue & Green mould, spray Imazalil 34 WP @ 1g/lit. Prune away the diseased portion of the plant. • In case of Bark eating caterpillar, remove the wooden frass present in between the joints of the trunk and inject Dichlorovos (DDVP) 76 EC 0.1 percent @ 5ml in the larval tunnel with the help of disposable syringe and push cotton swab soaked in Dichlorovos solution in the larval tunnel.
Large Cardamom	Fruiting	Chirkey Disease	<ul style="list-style-type: none"> • Remove dry/diseased/old tillers & branches & spray Bio NPK@ 2ml/lit water for proper growth on clear/dry days.
Banana	Multiple stages	Sigatoka Leaf spot	<ul style="list-style-type: none"> • One sword sucker may be allowed after shooting (Flowering), • Periodical removal of side suckers by cutting them above the ground level, scoop the core and pour 2 ml of kerosene in the core, • Foliar spray of Propiconazole (e.g., Tilt) @ 1 ml per litre of wate using stickers in case of rainfall.
Honey Bees			
Honey bees			<ul style="list-style-type: none"> • Box must be covered with straw/dry leaves to provide warmth
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
General			
<ul style="list-style-type: none"> • Damaged grain (not fungus contaminated) to be diverted for Poultry/livestock feeding • Clean drinking water should be ensured round the clock and water troughs/feeders should be regularly cleaned, 			
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.) Swine Fever	<ul style="list-style-type: none"> • Avoid dampness and cover weak and sick animals with sack cloth. • Animals should be given lukewarm feed and water. • Right time to deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight. • Provide proper bedding materials especially to new-born piglets and lukewarm food & water
Cattle	All stages	FMD	<ul style="list-style-type: none"> • Animals should be given lukewarm feed and water. • Right time to deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight. • Provide proper bedding materials especially to new-born and lukewarm food & water • Heavy mortality in cloven footed animals has been reported.

			<ul style="list-style-type: none"> • Supportive treatment of infected animals under supervision of veterinary department officials, • Avoid mixing of infected and healthy animals, • Keep infected animals separated from the healthy ones, • Vaccinate the healthy animals against FMD (Through State Veterinary Dept.) • Provide feed supplements
Poultry	All stages		<ul style="list-style-type: none"> • Day old chick-Maintain warmth by lighting, • In present weather conditions protect the birds from cold by using gunny bags/plastic sheets around the poultry shed during night, • Provide heat by lighting 100W bulb @ 1 bulb per 20 newly born chicks.
Goat	All stages		<ul style="list-style-type: none"> • Deworming for gastrointestinal parasite
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit
ICAR, BASAR**

कृषि मौसम सलाहकार सेवा पत्रिका

Agro-met Advisory Bulletin

(Period: 24th to 27th February 2018)

ग्रामीण कृषि मौसम सेवा



GRAMIN KRISHI MAUSAM SEWA

(Integrated Agro-Meteorological Advisory Services)



AMFU, ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Lower Subansiri

No: AMFU-Basar/2018/16

Dated: Fri, 23rd February, 2018



Weather summary of the previous week	Weather forecast valid up to 28 th February 2018
Rainfall = NA	Mostly/Partly cloudy with possibility of moderate rainfall. Daily maximum and minimum temperature for the following period is predicted to be 27-28°C and 14-15°C respectively. Expected morning and evening relative humidity will be 84-92% and 27-58% respectively. Wind direction is mainly SSW & SW with speed range of 4-8 kmph.
Warning:	• Nil

Weather Based Agro-meteorological Advisories Bulletin

- **Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated**
- **Spraying operation should be done only on dry/clear weather**
- **Go for low cost polyhouses for year round cultivation of vegetables**
- **Obtain seeds from reputed/certified/reliable source**
- **Use fertilizers/pesticides as per recommended doses for the location**
- **Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides**
- **ICAR Basar recommend poultry variety VANARAJA for backyard rearing**

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Pea	Pod formation	Powdery mildew/ leaf eating insects	<ul style="list-style-type: none"> In case of heavy infestation, need based foliar spray of Dinocap (0.2%) or Carbendazim 50WP (0.1%) or Propiconazole (0.1%) using stickers.
Toria/ Mustard	Seed formation	Aphids	<ul style="list-style-type: none"> Spray Dimethoate 30EC @2ml/litre against aphids in Rapeseed/Mustard on clear/dry days.
VEGETABLES			
Potato	Vegetative stage/ Harvesting stage	Late blight/ Red ant	<ul style="list-style-type: none"> Avoid harvesting under current weather condition. Regularly monitor against the disease symptoms. Prominent symptoms include small, dark, circular to irregular shaped lesion in the leaves Prophylactic spraying of Mancozeb @ 2 gram per 4 litres of water is advised using stickers in case of rainfall..
Cabbage/ Cauliflower	Harvesting		<ul style="list-style-type: none"> Harvest mature crop. Constantly monitor against attack of insects pests/mole.
HORTICULTURAL CROPS			
Kiwi	Pruning/Training		<ul style="list-style-type: none"> Diseased and senile branches should be removed. Pruning should be done retaining 8-10 buds per branch for enhancing new flush and flowering. Pruned healthy hard wood of pencil thickness size having 4-5 buds can be used as cutting for propagation
Citrus	Maturity/ harvesting	Bark eating caterpillar, Trunk borer, Blue & Green mould	<ul style="list-style-type: none"> Orange (only if harvesting is over): Go for drenching with Dichlorovos 76 EC@0.05% against trunk-borer on dry days. On clear weather or using sticker, against Blue & Green mould, spray Imazalil 34 WP @ 1g/lit. Prune away the diseased portion of the plant. In case of Bark eating caterpillar, remove the wooden frass present in between the joints of the trunk and inject Dichlorovos (DDVP) 76 EC 0.1 percent @ 5ml in the larval tunnel with the help of disposable syringe and push cotton swab soaked in Dichlorovos solution in the larval tunnel.
Temperate fruits	After harvesting		<ul style="list-style-type: none"> Training & pruning of the temperate crops (apple/pears/peach) is recommended In case of flowering, stop spraying insecticides/pesticides
Banana	Bunching	Cold	<ul style="list-style-type: none"> Cover newly emerged bunches with locally available sack to protect from cold/frost Mulching with straw/crop residue at root zone to avoid frost injury in banana.
Agroforestry			
Toko palm	Nursery		Filling of nursery polybags for raising of takopalm seedlings
Honey Bees			
Honey bees			<ul style="list-style-type: none"> Box must be covered with straw/dry leaves to provide warmth
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
General			
<ul style="list-style-type: none"> Damaged grain (not fungus contaminated) to be diverted for Poultry/livestock feeding Clean drinking water should be ensured round the clock and water troughs/feeders should be regularly cleaned, 			
Poultry	All stages		<ul style="list-style-type: none"> Provide shelter to the backyard/scavenging poultry. Feed grains in the morning/evening to supplement scavenging. Day old chick-Maintain warmth by lighting, In present weather conditions protect the birds from cold by using gunny bags/plastic sheets around the poultry shed during night, Provide heat by lighting 100W bulb @ 1 bulb per 20 newly born chicks.
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the	<ul style="list-style-type: none"> Avoid dampness and cover weak and sick animals with sack cloth. The bedding is always dry and that there is plenty.

		mouth and foot.) Swine Fever	<ul style="list-style-type: none"> • If possible wooden sleeping box should be provided. • Animals should be given lukewarm feed and water. • Right time to deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight.
Cattle	All stages	FMD	<ul style="list-style-type: none"> • Animals should be given lukewarm feed and water. • Right time to deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight. • Provide proper bedding materials especially to new-born and lukewarm food & water • Provide feed supplements
Mithun		FMD	<ul style="list-style-type: none"> • Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route. • Pregnant animals above 7 months of pregnancy should be avoided.
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit
ICAR, BASAR**

कृषि मौसम सलाहकार सेवा पत्रिका

Agro-met Advisory Bulletin

(Period: 24th to 27th February 2018)

ग्रामीण कृषि मौसम सेवा



GRAMIN KRISHI MAUSAM SEWA
(Integrated Agro-Meteorological Advisory Services)



भा.क.अनु.प.सं.
ICAR-RCNEH

AMFU, ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Papum Pare

No: AMFU-Basar/2018/16

Dated: Fri, 23rd February, 2018



Weather summary of the previous week	Weather forecast valid up to 28 th February 2018
Rainfall: 0.0 mm; T. Max: 30.0°C; T. Min: 15.0°C ; RH: 55-70%	Mostly cloudy with possibility of moderate rain. Daily maximum and minimum temperature for the following period is predicted to be 27-29°C and 16-17°C respectively. Expected morning and evening relative humidity will be 76-86% and 40-51% respectively. Expected wind direction mainly WSW with speed range of 5 kmph.
Warning:	<ul style="list-style-type: none">Moderate to dense fog likely to occur at isolated places

Weather Based Agro-meteorological Advisories Bulletin

General Recommendations

- Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated
- Spraying operation should be done only on dry/clear weather
- Go for low cost polyhouses for year round cultivation of vegetables
- Obtain seeds from reputed/certified/reliable source
- Use fertilizers/pesticides as per recommended doses for the location
- Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides
- ICAR Basar recommend poultry variety VANARAJA for backyard rearing

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Pea	Pod mature	Birds	<ul style="list-style-type: none"> • Harvest of green pods, • Use of scare crow and tying of bird scare tape around the crop field,
Toria/ Mustard	Vegetative/ Flowering	Aphid	<ul style="list-style-type: none"> • Harvesting of mature crop, • Drying of harvested crop to protect against disease development, • Spraying of dimethoate @ 2 ml per litre of water is advised using sticker if the infestation is high,
VEGETABLES			
Potato	Maturity/ Harvesting stage	Late blight/ Red ant	<ul style="list-style-type: none"> • Avoid harvesting • Regularly monitor against the disease symptoms. Prominent symptoms include small, dark, circular to irregular shaped lesion in the leaves • Prophylactic spraying of Mancozeb @ 2 gram per 4 litres of water is advised using stickers in case of rainfall..
Cabbage/ Cauliflower	Harvesting		<ul style="list-style-type: none"> • Harvest mature crop. • Constantly monitor against attack of insects pests/mole.
Ginger/ Turmeric	Maturity		<ul style="list-style-type: none"> • Harvest the mature crop during clear/dry days, • Cure should be taken while harvest so that the rhizome is not damaged,
HORTICULTURAL CROPS			
Citrus	Maturity/ harvesting	Bark eating caterpillar, Gummosis	<ul style="list-style-type: none"> • In case of Bark eating caterpillar, remove the wooden frass present in between the joints of the trunk and inject Dichlorovos (DDVP) 76 EC 0.1 percent @ 5ml in the larval tunnel with the help of disposable syringe and push cotton swab soaked in Dichlorovos solution in the larval tunnel. • Collect dropped fruits & dispose by putting in airtight bags or bury deep in ground. • Harvest the mature fruit with stalk and store in dry places. • Dead wood pruning may be done after harvesting of crop followed by spraying of fungicide, Carbendazim @ 1gm/lit of water. • Breakdown the soil collected around the tree trunk.
Large Cardamom	Vegetative		<ul style="list-style-type: none"> • Provide proper drainage to avoid water stagnation in the base of the plant.
Banana	Multiple stages	Sigatoka Leaf spot	<ul style="list-style-type: none"> • One sword sucker may be allowed after shooting (Flowering), • Periodical removal of side suckers by cutting them above the ground level, scoop the core and pour 2 ml of kerosene in the core, • Foliar spray of Propiconazole (e.g., Tilt) @ 1 ml per litre of water using stickers in case of rainfall
Honey Bees			
Honey bees			<ul style="list-style-type: none"> • Box must be covered with straw/dry leaves to provide warmth
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
General			
<ul style="list-style-type: none"> • Damaged grain (not fungus contaminated) to be diverted for Poultry/livestock feeding • Clean drinking water should be ensured round the clock and water troughs/feeders should be regularly cleaned, 			
Goat	All stages		<ul style="list-style-type: none"> • Deworming for gastrointestinal parasite
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.) Swine Fever	<ul style="list-style-type: none"> • Avoid dampness and cover weak and sick animals with sack cloth. • Animals should be given lukewarm feed and water. • Right time to deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight. • Provide proper bedding materials especially to new-born piglets and lukewarm food & water
Fishery			<ul style="list-style-type: none"> • Keep fish stoking density thin @ 1200/ha
Cattle	All stages	FMD	<ul style="list-style-type: none"> • Animals should be given lukewarm feed and water. • Right time to deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30

			<p>kg body weight.</p> <ul style="list-style-type: none"> • Provide proper bedding materials especially to new-born and lukewarm food & water • Provide feed supplements
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.) Swine Fever	<ul style="list-style-type: none"> • Iron injection within 10 days of birth is advised to newly born piglets if they are kept in confined system • Deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight <p>In present weather conditions vaccinate against swine fever (Vaccines available in State Veterinary Depts),</p>
<p>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247</p>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit
ICAR, BASAR

कृषि मौसम सलाहकार सेवा पत्रिका

Agro-met Advisory Bulletin

(Period: 24th to 27th February 2018)

ग्रामीण कृषि मौसम सेवा



GRAMIN KRISHI MAUSAM SEWA

(Integrated Agro-Meteorological Advisory Services)



भा.क.अनु.प.सं.
ICAR-RCNEH

AMFU, ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Tawang

No: AMFU-Basar/2018/16

Dated: Fri, 23rd February, 2018



Weather summary of the previous week	Weather forecast valid up to 28 th February 2018
Rainfall = 10.0 mm	Mostly/partly cloudy with possibility light/moderate rain. Daily maximum and minimum temperature for the following period is predicted to be 12-13°C and 2.0-3.0°C respectively. Expected morning and evening relative humidity will be 74-87% and 26-39% respectively. Wind direction is mainly SE with probable speed range of 4-13 kmph.
Warning:	•

Weather Based Agro-meteorological Advisories Bulletin

General Recommendations

- Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated
- All spraying operation should be done only on dry/clear weather
- Go for low cost polyhouses for year round cultivation of vegetables
- Obtain seeds from reputed/certified/reliable source
- Use fertilizers/pesticides as per recommended doses for the location
- Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides
- ICAR Basar recommend poultry variety VANARAJA for backyard rearing

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Pea	Pod formation	Powdery mildew/ leaf eating insects	<ul style="list-style-type: none"> In case of heavy infestation, spray of Dinocap (0.2%) or Carbendazim 50WP (0.1%) or Propiconazole (0.1%) using sticker.
VEGETABLES			
Potato	Vegetative stage/ Harvesting stage	Late blight/ Red ant	<ul style="list-style-type: none"> Avoid harvesting in present weather. Regularly monitor against the disease symptoms. Prominent symptoms include small, dark, circular to irregular shaped lesion in the leaves Prophylactic spraying of Mancozeb @ 2 gram per 4 litres of water is advised using stickers in case of rainfall.
Cabbage/ Cauliflower	Harvesting		<ul style="list-style-type: none"> Harvest mature crop. Constantly monitor against attack of insects pests/mole.
HORTICULTURAL CROPS			
Kiwi	Pruning/Training		<ul style="list-style-type: none"> Diseased and senile branches should be removed. Pruning should be done retaining 8-10 buds per branch for enhancing new flush and flowering. Pruned healthy hard wood of pencil thickness size having 4-5 buds can be used as cutting for propagation
Temperate fruits	After harvesting		<ul style="list-style-type: none"> Training & pruning of the temperate crops (apple/pears/peach) is recommended In case of flowering, stop spraying insecticides/pesticides
Banana	Bunching	Cold	<ul style="list-style-type: none"> Cover newly emerged bunches with locally available sack to protect from cold/frost Mulching with straw/crop residue at root zone to avoid frost injury in banana.
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
General			
<ul style="list-style-type: none"> Damaged grain (not fungus contaminated) to be diverted for Poultry/livestock feeding Clean drinking water should be ensured round the clock and water troughs/feeders should be regularly cleaned, 			
Poultry	All stages		<ul style="list-style-type: none"> Provide shelter to the backyard/scavenging poultry. Feed grains in the morning/evening to supplement scavenging. Day old chick-Maintain warmth by lighting, In present weather conditions protect the birds from cold by using gunny bags/plastic sheets around the poultry shed during night, Provide heat by lighting 100W bulb @ 1 bulb per 20 newly born chicks.
Yak	All stages		<ul style="list-style-type: none"> Feeding of concentrate feed along with mineral mixture and salt. Hay feeding is recommended.
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.) Swine Fever	<ul style="list-style-type: none"> Avoid dampness and cover weak and sick animals with sack cloth. The bedding is always dry and that there is plenty. If possible wooden sleeping box should be provided. Animals should be given lukewarm feed and water. Right time to deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight.
Cattle	All stages	FMD	<ul style="list-style-type: none"> Animals should be given lukewarm feed and water. Right time to deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight. Provide proper bedding materials especially to new-born and lukewarm food & water Provide feed supplements
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

**Agrometeorological Field Unit
ICAR, BASAR**

कृषि मौसम सलाहकार सेवा पत्रिका

Agro-met Advisory Bulletin

(Period: 24th to 27th February 2018)

ग्रामीण कृषि मौसम सेवा



GRAMIN KRISHI MAUSAM SEWA

(Integrated Agro-Meteorological Advisory Services)




भा.क.अनु.प. - पू.प.सं.
ICAR - RCNEH

AMFU, ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Tirap

No: AMFU-Basar/2018/16

Dated: Fri, 23rd February, 2018



Weather summary of the previous week	Weather forecast valid up to 28 th February 2018
Rainfall = 0.0 mm	Mostly cloudy with possibility of light/moderate rain on 24 th and 27 th Feb. Daily maximum and minimum temperature for the following period is predicted to be 27-29°C and 16-17°C respectively. Expected morning and evening relative humidity will be 76-86 % and 29-43% respectively. Wind direction is mainly ESE & SE with speed range of 2-6 kmph.
Warning:	• Nil

Weather Based Agro-meteorological Advisories Bulletin

- Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated.
- Go for low cost polyhouses for year round cultivation of vegetables
- Obtain seeds from reputed/certified/reliable source
- Use fertilizers/pesticides as per recommended doses for the location
- Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides
- ICAR Basar recommend poultry variety VANARAJA for backyard rearing

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Maize	Vegetative/ Maturity/ Harvesting	Rats	<ul style="list-style-type: none"> Protect the crop against attack of rats using locally available traps, When rat menace has occurred application of multi dose rodenticide like bromadiolone with proper consultation with experts is advised.
Pea	Pod mature	Birds	<ul style="list-style-type: none"> Harvest of green pods during clear days, Use of scare crow and tying of bird scare tape around the crop field,
Toria/ Mustard	Vegetative/ Flowering	Aphid	<ul style="list-style-type: none"> Harvesting of mature crop during sunny/dry days, Drying of harvested crop to protect against disease development, Spraying of dimethoate @ 2 ml per litre of water is advised using sticker in case of heavy infestation,
VEGETABLES			
Potato	Vegetative stage/ Harvesting stage	Late blight/ Red ant	<ul style="list-style-type: none"> Avoid Harvesting If already harvested, Store in a cool, well ventilated place/ do not wash before storing as dampness promotes early spoilage. Regularly monitor against the disease symptoms. Prominent symptoms include small, dark, circular to irregular shaped lesion in the leaves Prophylactic spraying of Mancozeb @ 2 gram per 4 litres of water is advised using stickers in case of rainfall..
Cabbage/ Cauliflower	Harvesting		<ul style="list-style-type: none"> Harvest mature crop. Constantly monitor against attack of insects pests/mole.
Ginger/ Turmeric	Maturity		<ul style="list-style-type: none"> Harvest the mature crop, Cure should be taken while harvest so that the rhizome is not damaged,
HORTICULTURAL CROPS			
Citrus	Maturity/ harvesting	Bark eating caterpillar, Trunk borer, Blue & Green mould	<ul style="list-style-type: none"> Orange (only if harvesting is over): Go for drenching with Dichlorovos 76 EC@0.05% against trunk-borer. Against Blue & Green mould, spray Imazalil 34 WP @ 1g/lit using stickers in case of rainfall. Prune away the diseased portion of the plant. In case of Bark eating caterpillar, remove the wooden frass present in between the joints of the trunk and inject Dichlorovos (DDVP) 76 EC 0.1 percent @ 5ml in the larval tunnel with the help of disposable syringe and push cotton swab soaked in Dichlorovos solution in the larval tunnel.
Banana	All stages	Sigatoka Leaf spot	<ul style="list-style-type: none"> One sword sucker may be allowed after shooting (Flowering), Periodical removal of side suckers by cutting them above the ground level, scoop the core and pour 2 ml of kerosene in the core, Foliar spray of Propiconazole (e.g., Tilt) @ 1 ml per litre of water using stickers in case of rainfall
Honey Bees			
			<ul style="list-style-type: none"> Box must be covered with straw/dry leaves to provide warmth
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
General			
<ul style="list-style-type: none"> Damaged grain (not fungus contaminated) to be diverted for Poultry/livestock feeding Clean drinking water should be ensured round the clock and water troughs/feeders should be regularly cleaned, 			
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.) Swine Fever	<ul style="list-style-type: none"> Avoid dampness and cover weak and sick animals with sack cloth. Animals should be given lukewarm feed and water. Right time to deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight. Provide proper bedding materials especially to new-born piglets and lukewarm food & water
Cattle	All stages	FMD	<ul style="list-style-type: none"> Animals should be given lukewarm feed and water.

			<ul style="list-style-type: none"> • Right time to deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight. • Provide proper bedding materials especially to new-born and lukewarm food & water • Provide feed supplements
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.) Swine Fever	<ul style="list-style-type: none"> • Iron injection within 10 days of birth is advised to newly born piglets if they are kept in confined system • Deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight • In present weather conditions vaccinate against swine fever (Vaccines available in State Veterinary Depts),
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
 Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit
 ICAR, BASAR**

कृषि मौसम सलाहकार सेवा पत्रिका

Agro-met Advisory Bulletin

(Period: 24th to 27th February 2018)

ग्रामीण कृषि मौसम सेवा



GRAMIN KRISHI MAUSAM SEWA

(Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Upper Siang

No: AMFU-Basar/2018/16

Dated: Fri, 23rd February, 2018



Weather summary of the previous week	Weather forecast valid up to 28 th February 2018
Rainfall = 0.0 mm	Partly cloudy with possibility of /moderate rainfall on 24 th and 27 th Feb. Daily maximum and minimum temperature for the following period is predicted to be 23-24°C and 10-11°C respectively. Expected morning and evening relative humidity will be 67-87% and 27-57% respectively. Wind direction is mainly variable with speed range of 5-10 kmph.
Warning:	• Nil

Weather Based Agro-meteorological Advisories Bulletin

General Recommendations

- Disinfecting the animal/bird sheds with lime powder/bleaching powder/4%NaOH at regular interval is advocated
- Spraying operation should be done only on dry/clear weather
- Go for low cost poly-houses for year round cultivation of vegetables
- Obtain seeds from reputed/certified/reliable source
- Use fertilizers/pesticides as per recommended doses for the location
- Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides
- ICAR Basar recommend poultry variety VANARAJA for backyard rearing

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Pea	Vegetative	Powdery mildew/ leaf eating insects	<ul style="list-style-type: none"> • Avoid/suspend all spray operations and application of fertilizers. • In case of heavy infestation, spray of Dinocap (0.2%) or Carbendazim 50WP (0.1%) or Propiconazole (0.1%) on clear/dry days or using sticker advocated.
Toria/ Mustard	Vegetative/ Flowering		<ul style="list-style-type: none"> • Provide proper drainage facilities for the toria grown in the WRC field to protect them against heavy rainfall.
Maize	Vegetative/ Maturity/ Harvesting	Rats	<ul style="list-style-type: none"> • Prepare field for sowing of crop, • Protect the crop against attack of rats using locally available traps, • When rat menace has occurred application of multi dose rodenticide like bromadiolone with proper consultation with experts is advised.
VEGETABLES			
Potato	Vegetative stage/ Harvesting stage	Late blight/ Red ant	<ul style="list-style-type: none"> • Avoid harvesting • Regularly monitor against the disease symptoms. Prominent symptoms include small, dark, circular to irregular shaped lesion in the leaves • Prophylactic spraying of Mancozeb @ 2 gram per 4 litres of water is advised using stickers in case of rainfall..
Cabbage/ Cauliflower	Harvesting		<ul style="list-style-type: none"> • Harvest mature crop. • Constantly monitor against attack of insects pests/mole.
Ginger/ Turmeric	Maturity		<ul style="list-style-type: none"> • Harvest the mature crop, • Cure should be taken while harvest so that the rhizome is not damaged,
HORTICULTURAL CROPS			
Kiwi	Pruning/Training		<ul style="list-style-type: none"> • Diseased and senile branches should be removed. • Pruning should be done retaining 8-10 buds per branch for enhancing new flush and flowering. • Pruned healthy hard wood of pencil thickness size having 4-5 buds can be used as cutting for propagation
Citrus	Maturity/ harvesting	Bark eating caterpillar, Trunk borer, Blue & Green mould	<ul style="list-style-type: none"> • Orange (only if harvesting is over): Go for drenching with Dichlorovos 76 EC@0.05% against trunk-borer on dry days. • On clear weather or using sticker, against Blue & Green mould, spray Imazalil 34 WP @ 1g/lit. Prune away the diseased portion of the plant. • In case of Bark eating caterpillar, remove the wooden frass present in between the joints of the trunk and inject Dichlorovos (DDVP) 76 EC 0.1 percent @ 5ml in the larval tunnel with the help of disposable syringe and push cotton swab soaked in Dichlorovos solution in the larval tunnel.
Temperate fruits	After harvesting		<ul style="list-style-type: none"> • Training & pruning of the temperate crops (apple/pears/peach) is recommended • In case of flowering, stop spraying insecticides/pesticides
Banana	Bunching	Cold	<ul style="list-style-type: none"> • Cover newly emerged bunches with locally available sack to protect from cold/frost • Mulching with straw/crop residue at root zone to avoid frost injury in banana.
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
General			
<ul style="list-style-type: none"> • Damaged grain (not fungus contaminated) to be diverted for Poultry/livestock feeding • Clean drinking water should be ensured round the clock and water troughs/feeders should be regularly cleaned, 			
Poultry	All stages		<ul style="list-style-type: none"> • Provide shelter to the backyard/scavenging poultry. Feed grains in the morning/evening to supplement scavenging. • Day old chick-Maintain warmth by lighting, • In present weather conditions protect the birds from cold by using gunny bags/plastic sheets around the poultry shed during night,

			<ul style="list-style-type: none"> • Provide heat by lighting 100W bulb @ 1 bulb per 20 newly born chicks.
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.) Swine Fever	<ul style="list-style-type: none"> • Avoid dampness and cover weak and sick animals with sack cloth. • The bedding is always dry and that there is plenty. • If possible wooden sleeping box should be provided. • Animals should be given lukewarm feed and water. • Right time to deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight.
Cattle	All stages	FMD	<ul style="list-style-type: none"> • Animals should be given lukewarm feed and water. • Right time to deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight. • Provide proper bedding materials especially to new-born and lukewarm food & water • Provide feed supplements
Fishery			<ul style="list-style-type: none"> • Application of quick lime is highly recommended in order to rectify water pH. • Application of fish feed (mustard oil cake) is essential for achieving higher yield
Mithun		FMD	<ul style="list-style-type: none"> • Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route. • Pregnant animals above 7 months of pregnancy should be avoided.
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit
ICAR, BASAR**

कृषि मौसम सलाहकार सेवा पत्रिका

Agro-met Advisory Bulletin

(Period: 24th to 27th February 2018)

ग्रामीण कृषि मौसम सेवा



GRAMIN KRISHI MAUSAM SEWA
(Integrated Agro-Meteorological Advisory Services)




भा.क.अनु.प. - पू.प.सं.
ICAR - RCNEH

AMFU, ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Upper Subansiri

No: AMFU-Basar/2018/16

Dated: Fri, 23rd February, 2018



Weather summary of the previous week	Weather forecast valid up to 28 th February 2018
Daporijo: 7.2 mm	Mostly/Partly cloudy with possibility of moderate rainfall. Daily maximum and minimum temperature for the following period is predicted to be 27-28°C and 14-15°C respectively. Expected morning and evening relative humidity will be 84-92% and 23-57% respectively. Expected wind direction is mainly S with speed range of 3-12 kmph.
Warning	• Rain in few isolated places

Weather Based Agro-meteorological Advisories Bulletin

- Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated
- Avoid spray of insecticides during flowering stage as it may hinder pollination by honey bees and other insects.
- Spraying operation should be done only on dry/clear weather
- Go for low cost polyhouses for year round cultivation of vegetables
- Obtain seeds from reputed/certified/reliable source
- Use fertilizers/pesticides as per recommended doses for the location
- Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides
- ICAR Basar recommend poultry variety VANARAJA for backyard rearing

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Pea	Pod formation	Powdery mildew/ leaf eating insects	<ul style="list-style-type: none"> In case of heavy infestation, foliar spray of Dinocap (0.2%) or Carbendazim 50WP (0.1%) or Propiconazole (0.1%) using appropriate sticker advocated.
Toria/ Mustard	Seed formation	Aphids	<ul style="list-style-type: none"> Spray Dimethoate 30EC @2ml/litre against aphids in Rapeseed/Mustard using stickers.
Maize	Vegetative/ Maturity/ Harvesting	Rats	<ul style="list-style-type: none"> Prepare field for sowing of crop, Protect the crop against attack of rats using locally available traps, When rat menace has occurred application of multi dose rodenticide like bromadiolone with proper consultation with experts is advised.
VEGETABLES			
Potato	Vegetative stage/ Harvesting stage	Late blight/ Red ant	<ul style="list-style-type: none"> Avoid harvesting Regularly monitor against the disease symptoms. Prominent symptoms include small, dark, circular to irregular shaped lesion in the leaves Prophylactic spraying of Mancozeb @ 2 gram per 4 litres of water is advised using stickers in case of rainfall..
Cabbage/ Cauliflower	Harvesting		<ul style="list-style-type: none"> Harvest mature crop. Constantly monitor against attack of insects pests/mole.
HORTICULTURAL CROPS			
Kiwi	Pruning/Training		<ul style="list-style-type: none"> Diseased and senile branches should be removed. Pruning should be done retaining 8-10 buds per branch for enhancing new flush and flowering. Pruned healthy hard wood of pencil thickness size having 4-5 buds can be used as cutting for propagation
Citrus	Maturity/ harvesting	Bark eating caterpillar, Trunk borer, Blue & Green mould	<ul style="list-style-type: none"> Orange (only if harvesting is over): Go for drenching with Dichlorovos 76 EC@0.05% against trunk-borer on dry days. On clear weather or using sticker, against Blue & Green mould, spray Imazalil 34 WP @ 1g/lit. Prune away the diseased portion of the plant. In case of Bark eating caterpillar, remove the wooden frass present in between the joints of the trunk and inject Dichlorovos (DDVP) 76 EC 0.1 percent @ 5ml in the larval tunnel with the help of disposable syringe and push cotton swab soaked in Dichlorovos solution in the larval tunnel.
Temperate fruits	After harvesting		<ul style="list-style-type: none"> Training & pruning of the temperate crops (apple/pears/peach) is recommended
Large Cardamom	Fruiting	Chirkey Disease	<ul style="list-style-type: none"> Remove dry/diseased/old tillers & branches & spray Bio NPK@ 2ml/lit water for proper growth using stickers in case of rainfall.
Banana	Bunching	Cold	<ul style="list-style-type: none"> Cover newly emerged bunches with locally available sack to protect from cold/frost
Orchards			<ul style="list-style-type: none"> Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour. Follow proper drainage and earthing up practices in low land/plain areas to avoid water stagnation in fruit orchards
Honey Bees			
Honey bees			<ul style="list-style-type: none"> Provide sugar syrup near the box Add insulation and ventilation systems
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
General			
<ul style="list-style-type: none"> Damaged grain (not fungus contaminated) to be diverted for Poultry/livestock feeding Clean drinking water should be ensured round the clock and water troughs/feeders should be regularly cleaned, 			
Poultry	All stages		<ul style="list-style-type: none"> Provide shelter to the backyard/scavenging poultry. Feed grains in the morning/evening to supplement scavenging. Day old chick-Maintain warmth by lighting, In present weather conditions protect the birds from cold by

			<p>using gunny bags/plastic sheets around the poultry shed during night,</p> <ul style="list-style-type: none"> • Provide heat by lighting 100W bulb @ 1 bulb per 20 newly born chicks.
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.) Swine Fever	<ul style="list-style-type: none"> • Avoid dampness and cover weak and sick animals with sack cloth. • The bedding is always dry and that there is plenty. • If possible wooden sleeping box should be provided. • Animals should be given lukewarm feed and water. • Right time to deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight.
Cattle	All stages	FMD	<ul style="list-style-type: none"> • Animals should be given lukewarm feed and water. • Right time to deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight. • Provide proper bedding materials especially to new-born and lukewarm food & water • Provide feed supplements
Mithun		FMD	<ul style="list-style-type: none"> • Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route. • Pregnant animals above 7 months of pregnancy should be avoided.

Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit
ICAR, BASAR**

कृषि मौसम सलाहकार सेवा पत्रिका

Agro-met Advisory Bulletin

(Period: 24th to 27th February 2018)

ग्रामीण कृषि मौसम सेवा



GRAMIN KRISHI MAUSAM SEWA
(Integrated Agro-Meteorological Advisory Services)



AMFU, ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: West Kameng

No: AMFU-Basar/2018/16

Dated: Fri, 23rd February, 2018



Weather summary of the previous week	Weather forecast valid up to 28 th February 2018
Rainfall = Bomdila: 0.0 ; Bhalukpong: 0.0 mm	Mostly/Partly cloudy with possibility light/moderate rain. Daily maximum and minimum temperature for the following period is predicted to be 22-23°C and 07-08°C respectively. Expected morning and evening relative humidity will be 86-90% and 28-48% respectively. Wind direction is mainly NNW with speed range of 5-7 kmph.
Warning:	• Moderate to dense fog likely to occur at isolated places

Weather Based Agro-meteorological Advisories Bulletin

General Recommendations

- **Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated**
- **Spraying operation should be done only on dry/clear weather**
- **Go for low cost polyhouses for year round cultivation of vegetables**
- **Obtain seeds from reputed/certified/reliable source**
- **Use fertilizers/pesticides as per recommended doses for the location**
- **Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides**
- **ICAR Basar recommend poultry variety VANARAJA for backyard rearing**

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Pea	Pod formation	Powdery mildew/ leaf eating insects	<ul style="list-style-type: none"> Need based foliar spray of Dinocap (0.2%) or Carbendazim 50WP (0.1%) or Propiconazole (0.1%) using sticker if the infestation is high.
Toria/ Mustard	Seed formation	Aphids	<ul style="list-style-type: none"> If the infestation is high, Spray Dimethoate 30EC @2ml/litre against aphids in Rapeseed/Mustard using sticker or during clear/dry days.
VEGETABLES			
Potato	Vegetative stage/ Harvesting stage	Late blight/ Red ant	<ul style="list-style-type: none"> Avoid harvesting during rain. Regularly monitor against the disease symptoms. Prominent symptoms include small, dark, circular to irregular shaped lesion in the leaves Prophylactic spraying of Mancozeb @ 2 gram per 4 litres of water is advised using stickers in case of rainfall..
Cabbage/ Cauliflower	Harvesting		<ul style="list-style-type: none"> Harvest mature crop. Constantly monitor against attack of insects pests/mole.
HORTICULTURAL CROPS			
Kiwi	Pruning/Training		<ul style="list-style-type: none"> Diseased and senile branches should be removed. Pruning should be done retaining 8-10 buds per branch for enhancing new flush and flowering. Pruned healthy hard wood of pencil thickness size having 4-5 buds can be used as cutting for propagation
Temperate fruits	After harvesting		<ul style="list-style-type: none"> Training & pruning of the temperate crops (apple/pears/peach) is recommended In case of flowering, stop spraying insecticides/pesticides
Banana	Bunching	Cold	<ul style="list-style-type: none"> Cover newly emerged bunches with locally available sack to protect from cold/frost
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
General			
<ul style="list-style-type: none"> Damaged grain (not fungus contaminated) to be diverted for Poultry/livestock feeding Clean drinking water should be ensured round the clock and water troughs/feeders should be regularly cleaned, 			
Yak	All stages		<ul style="list-style-type: none"> Feeding of concentrate feed along with mineral mixture and salt. Hay feeding is recommended.
Fishery			<ul style="list-style-type: none"> Keep fish stoking density thin @1200/ha
Poultry	All stages		<ul style="list-style-type: none"> Provide shelter to the backyard/scavenging poultry. Feed grains in the morning/evening to supplement scavenging. Day old chick-Maintain warmth by lighting, In present weather conditions protect the birds from cold by using gunny bags/plastic sheets around the poultry shed during night, Provide heat by lighting 100W bulb @ 1 bulb per 20 newly born chicks.
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.) Swine Fever	<ul style="list-style-type: none"> Avoid dampness and cover weak and sick animals with sack cloth. The bedding is always dry and that there is plenty. If possible wooden sleeping box should be provided. Animals should be given lukewarm feed and water. Right time to deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight.
Cattle	All stages	FMD	<ul style="list-style-type: none"> Animals should be given lukewarm feed and water. Right time to deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight. Provide proper bedding materials especially to new-born and lukewarm food & water Heavy mortality in cloven footed animals has been reported. Supportive treatment of infected animals under supervision of veterinary department officials, Avoid mixing of infected and healthy animals,

			<ul style="list-style-type: none"> • Keep infected animals separated from the healthy ones, • Vaccinate the healthy animals against FMD (Through State Veterinary Dept.) • Provide feed supplements
Mithun		FMD	<ul style="list-style-type: none"> • Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route. • Pregnant animals above 7 months of pregnancy should be avoided.

Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
 Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit
 ICAR, BASAR**

कृषि मौसम सलाहकार सेवा पत्रिका Agro-met Advisory Bulletin

(Period: 24th to 27th February 2018)

ग्रामीण कृषि मौसम सेवा



GRAMIN KRISHI MAUSAM SEWA
(Integrated Agro-Meteorological Advisory Services)




AMFU, ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: West Siang

No: AMFU-Basar/2018/16

Dated: Fri, 23rd February, 2018



Basar: Latitude: 27°59.53/Longitude: 94.41.269/Altitude: 616 amsl

Weather summary of the previous week	Weather forecast valid up to 28 th February 2018
Basar: Rainfall = 1.3 mm. T. Max: 22.6°C, T. Min: 7.0°C, RH: 44-97%, Wind Speed: 6-8 kmph, Wind Direction: ENE & NNE	Mostly/partly cloudy with possibility of light/moderate rain. Daily maximum and minimum temperature for the following period is predicted to be 28-29°C and 15-16°C respectively. Expected morning and evening relative humidity will be 62-78% and 29-47% respectively. Wind direction is mainly W & SW with speed range of 5-6 kmph.
Warning:	• Nil

Weather Based Agro-meteorological Advisories Bulletin

General Recommendations

- Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated
- Go for low cost polyhouses for year round cultivation of vegetables
- Obtain seeds from reputed/certified/reliable source
- Use fertilizers/pesticides as per recommended doses for the location
- Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides
- ICAR Basar recommend poultry variety VANARAJA for backyard rearing

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Pea	Pod mature	Birds	<ul style="list-style-type: none"> • Harvest of green pods, • Use of scare crow and tying of bird scare tape around the crop field,
Toria/ Mustard	Vegetative/ Flowering	Aphid	<ul style="list-style-type: none"> • Spraying of dimethoate @ 2 ml per litre of water is advised on clear/dry days or using sticker in case of heavy infestation.
Maize	Vegetative/ Maturity/ Harvesting	Rats	<ul style="list-style-type: none"> • Protect the crop against attack of rats using locally available traps, • When rat menace has occurred application of multi dose rodenticide like bromadiolone with proper consultation with experts is advised.
VEGETABLES			
Potato	Vegetative stage/ Harvesting stage	Late blight/ Red ant	<ul style="list-style-type: none"> • Avoid harvesting • Regularly monitor against the disease symptoms. Prominent symptoms include small, dark, circular to irregular shaped lesion in the leaves • Prophylactic spraying of Mancozeb @ 2 gram per 4 litres of water is advised using stickers in case of rainfall..
Cabbage/ Cauliflower	Harvesting		<ul style="list-style-type: none"> • Harvest mature crop. • Constantly monitor against attack of insects pests/mole.
Ginger/ Turmeric	Maturity		<ul style="list-style-type: none"> • Harvest the mature crop, • Cure should be taken while harvest so that the rhizome is not damaged,
HORTICULTURAL CROPS			
Kiwi	Pruning/Training		<ul style="list-style-type: none"> • Diseased and senile branches should be removed. • Pruning should be done retaining 8-10 buds per branch for enhancing new flush and flowering. • Pruned healthy hard wood of pencil thickness size having 4-5 buds can be used as cutting for propagation
Citrus	Maturity/ harvesting	Bark eating caterpillar, Trunk borer, Blue & Green mould	<ul style="list-style-type: none"> • Orange (only if harvesting is over): Go for drenching with Dichlorovos 76 EC@0.05% against trunk-borer on dry days. • On clear weather or using sticker, against Blue & Green mould, spray Imazalil 34 WP @ 1g/lit. Prune away the diseased portion of the plant. • In case of Bark eating caterpillar, remove the wooden frass present in between the joints of the trunk and inject Dichlorovos (DDVP) 76 EC 0.1 percent @ 5ml in the larval tunnel with the help of disposable syringe and push cotton swab soaked in Dichlorovos solution in the larval tunnel.
Temperate fruits	After harvesting		<ul style="list-style-type: none"> • Training & pruning of the temperate crops (apple/pears/peach) is recommended • In case of flowering, stop spraying insecticides/pesticides
Banana	Bunching	Cold	<ul style="list-style-type: none"> • Cover newly emerged bunches with locally available sack to protect from cold/frost • Mulching with straw/crop residue at root zone to avoid frost injury in banana.
Large Cardamom	Flowering/ Fruiting	Chirkey Disease	<ul style="list-style-type: none"> • Remove dry/diseased/old tillers & branches & spray Bio NPK@ 2ml/lit water for proper growth using stickers in case of rainfall.
Honey Bees			
Honey bees			<ul style="list-style-type: none"> • Cover the boxes with locally available dry grass or straw to provide warmth
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
General			
<ul style="list-style-type: none"> • Damaged grain (not fungus contaminated) to be diverted for Poultry/livestock feeding • Clean drinking water should be ensured round the clock and water troughs/feeders should be regularly cleaned, 			
Poultry	All stages		<ul style="list-style-type: none"> • Provide shelter to the backyard/scavenging poultry. Feed grains in the morning/evening to supplement scavenging. • Day old chick-Maintain warmth by lighting,

			<ul style="list-style-type: none"> • In present weather conditions protect the birds from cold by using gunny bags/plastic sheets around the poultry shed during night, • Provide heat by lighting 100W bulb @ 1 bulb per 20 newly born chicks.
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.) Swine Fever	<ul style="list-style-type: none"> • Avoid dampness and cover weak and sick animals with sack cloth. • The bedding is always dry and that there is plenty. • If possible wooden sleeping box should be provided. • Animals should be given lukewarm feed and water. • Right time to deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight.
Cattle	All stages	FMD	<ul style="list-style-type: none"> • Animals should be given lukewarm feed and water. • Right time to deworm with anti helmetic drugs (particularly Tapeworms) viz. Fento Plus @ 1 tablet per 30 kg body weight. • Provide proper bedding materials especially to new-born and lukewarm food & water • Provide feed supplements
Mithun		FMD	<ul style="list-style-type: none"> • Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route. • Pregnant animals above 7 months of pregnancy should be avoided.
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)