





Dated: Friday, 8th February 2019



Gramin Krishi Mausam Sewa ICAR Research Complex for NEH Region, Arunachal Pradesh Centre, Basar-791101

#### **ADVISORY FOR BLOCK: BASAR**

{Basar: Latitude: 27°59.53/Longitude: 94.41.269/Altitude: 616 amsl}

Weather summery of the previous week	Weather forecast valid up to 13 <sup>th</sup> February 2019
Rainfall: 0.0 mm, Max. T: 24.5°C, Min. T: 4.3°C, Wind Speed: 3-6 km/h, RH: 40-96%, Wind Direction: SSW	<b>Partly cloudy with possibility of light/moderate.</b> Daily maximum and minimum temperature for the following period is predicted to be 17-29°C and 10-14°C respectively. Expected morning and evening relative humidity will be 63-97% and 26-92% respectively. Wind direction is mainly NNW with average wind speed range of 3-4 kmph.
Warning:	•

#### **GENERAL RECOMMENDATIONS**

- Monitor for possible attack of insects/pests
- Soil moisture conservation techniques like mulching with crop residue and provide life saving irrigation to the crops recommended.
- Application of fertilizers/manure can be done during clear/dry days

Check of	f vaccination sche			
DETAILS OF ADVISORIES FOR THE PERIOD				
Crop	Stage	Pest/Disease		Precautionary Practices to be followed
FIELD CROPS				
Pea	Pod Formation			Regularly harvest mature pods,
	/mature			Regular monitoring of the crop against possible attack of blight.
	/harvesting			Provide support to the crop,
				Provide proper drainage to prevent water stagnation
Maize	Land			Procure good quality certified seeds,
	preparation		*	Seed treatment of early maize with Imidichloropid @ 1-2g/10kg
				is recommended before sowing.
			*	Provide proper drainage in the field
			*	Avoid sowing till fair weather forecasted
VEGETABLES				
Early Kharif	Sowing		*	Nursery of early kharif vegetables like Capsicum, tomato &
vegetables				other vegetables can be done
_			*	Procure seed from certified agencies
				Carry out seed treatment with Thiram/Carbendazim @ 2g/kg
			·	seed before sowing
			**	Transplanting of the healthy seedling to main field may be done
			•	during clear/dry weather
			*	Provide proper drainage in the field
Potato	Vegetative			Avoid harvesting during rainy days
1 otato	stage/			
	Ü		*	Provide proper drainage in the field
HODELGIU EUD	harvesting			
HORTICULTUR				
Khasi Mandarin	After harvesting/	Lichens/mosses	**	In present weather conditions remove lichens/mosses from the
	transplanting		*	affected trees using gunny bags smoothly, Transplantation of new seedling in the main field may be done
LIVESTOCKS			**	Transplantation of new seeding in the main field may be done
Livestock	Stago	Disease		Duccoutionary Ducctions to be followed
	Stage	Disease	.♦.	Precautionary Practices to be followed
Cattle	All Growth			Provide less ventilation to avoid heat loss
			*	Keep cattle clean and dry so that they stay warm,
Pig	Pigs/ Piglets		*	The bedding is always dry and that there is plenty.
			*	If possible wooden sleeping box should be provided.
Cattle	All stages	FMD	*	Vaccination – of healthy animals
			*	Treatment – sick animals,
		Respiratory	*	Intra muscular injection of Oxi-tetracycline @ 20ml/400 kg
				body weight animal (Avoid injecting to Pregnant animals)





### **Block Level Experimental Agro-met Advisory Bulletin**

(Period: 9th to 13th February 2019)

**No: AMFU-Basar/2019/12** 

Dated: Friday, 8th February 2019



Gramin Krishi Mausam Sewa ICAR Research Complex for NEH Region, Arunachal Pradesh Centre, Basar-791101

### **ADVISORY FOR BLOCK: DARING**

{Latitude: 28°11.651'/ Longitude: 94°45.712'/Altitude: 350-450 amsl}

Weather summery of the previous week	Weather forecast valid up to 13 <sup>th</sup> February 2019
NA	Partly cloudy with possibility of light/moderate rain. Daily maximum and
	minimum temperature for the following period is predicted to be 15-28°C
	and 10-14°C respectively. Expected morning and evening relative humidity
	will be 66-98% and 26-96% respectively. Wind direction is mainly NNW
	with average wind speed range of 3-4 kmph.
Warning:	•

#### **GENERAL RECOMMENDATIONS**

- Monitor for possible attack of insects/pests
- Soil moisture conservation techniques like mulching with crop residue and provide life saving irrigation to the crops recommended
- Application of fertilizers/manure can be done during clear/dry days

Proper drainage to all crop to avoid runoff			
Check of vaccination schedule in poultry/livestock			
		DETAILS OF A	ADVISORIES FOR THE PERIOD
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Pea	Pod Formation /mature /harvesting		<ul> <li>Regularly harvest mature pods,</li> <li>Regular monitoring of the crop against possible attack of blight.</li> <li>Provide support to the crop,</li> <li>Provide proper drainage to prevent water stagnation</li> </ul>
Maize	Land		❖ Procure good quality certified seeds,
	preparation		<ul> <li>Seed treatment of early maize with Imidichloropid @ 1-2g/10kg is recommended before sowing.</li> <li>Provide proper drainage in the field</li> <li>Avoid sowing till fair weather forecasted</li> </ul>
VEGETABLES			
Early Kharif vegetables  Potato  HORTICULTUR Khasi Mandarin		Lichens/mosses	<ul> <li>Nursery of early kharif vegetables like Capsicum, tomato &amp; other vegetables can be done</li> <li>Procure seed from certified agencies</li> <li>Carry out seed treatment with Thiram/Carbendazim @ 2g/kg seed before sowing</li> <li>Transplanting of the healthy seedling to main field may be done during clear/dry weather</li> <li>Provide proper drainage in the field</li> <li>Avoid harvesting during rainy days</li> <li>Provide proper drainage in the field</li> <li>In present weather conditions remove lichens/mosses from the affected trees using gunny bags smoothly,</li> </ul>
			<ul> <li>Transplantation of new seedling in the main field may be done</li> </ul>
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Cattle	All Growth		<ul> <li>Provide less ventilation to avoid heat loss</li> <li>Keep cattle clean and dry so that they stay warm,</li> </ul>
Pig	Pigs/ Piglets		<ul> <li>The bedding is always dry and that there is plenty.</li> <li>If possible wooden sleeping box should be provided.</li> </ul>
Cattle	All stages	FMD Respiratory	<ul> <li>Vaccination – of healthy animals</li> <li>Treatment – sick animals,</li> <li>Intra muscular injection of Oxi-tetracycline @ 20ml/400 kg body weight animal (Avoid injecting to Pregnant animals)</li> </ul> K staff for measures against adverse conditions or call at 03795-226237/226247





### **Block Level Experimental Agro-met Advisory Bulletin**

(Period: 9th to 13th February 2019)

No: AMFU-Basar/2019/12

Dated: Friday, 8th February 2019



Gramin Krishi Mausam Sewa ICAR Research Complex for NEH Region, Arunachal Pradesh Centre, Basar-791101

### **ADVISORY FOR BLOCK: GENSI**

{Latitude: 27°48.044'/ Longitude: 94°43.409'/Altitude: 750-850 amsl}

Weather summery of the previous week	Weather forecast valid up to 13 <sup>th</sup> February 2019
NA	Partly cloudy with possibility of light/moderate rain. Daily maximum and
	minimum temperature for the following period is predicted to be 16-29°C
	and 10-15°C respectively. Expected morning and evening relative humidity
	will be 634-98% and 28-89% respectively. Wind direction is mainly NNW
	with average wind speed range of 4-5 kmph.
Warning:	•

#### **GENERAL RECOMMENDATIONS**

GENER	GENERAL RECUMINIENDATIONS				
Monitor for possible attack of insects/pests					
Application of fertilizers/manure can be done during clear/dry days					
	Proper drainage to all crop to avoid runoff				
_		•			
• CHECK O	Check of vaccination schedule in poultry/livestock     DETAILS OF ADVISORIES FOR THE PERIOD				
C	Gt.		AD VISC		
Crop	Stage	Pest/Disease		Precautionary Practices to be followed	
FIELD CROPS	771				
Pea	Flowering/		*	8 p p, p	
	Maturity			stimulate generation of new flowers	
				Regular monitoring against insect attack	
				Provide proper drainage in the field	
				Avoid any spray operation/ application of fertilizers	
Maize	Land		*	Procure good quality certified seeds,	
	preparation		*		
				is recommended before sowing.	
			*	Avoid sowing till forecast of clear weather is issued	
			*	Provide proper drainage to the already sown crop	
<b>EGETABLES</b>					
Early Kharif	Sowing		*	Nursery of early kharif vegetables like Capsicum, tomato &	
vegetables				other vegetables can be done	
			*	Procure seed from certified agencies	
			*	Carry out seed treatment with Thiram/Carbendazim @ 2g/kg	
				seed before sowing	
			*	Avoid transplanting to main field	
Potato/	Tuber			Avoid harvesting during rainy days	
Tomato	formation/			Do not wash the tubers before storage	
	Maturity/			Washing may lead to early spoilage	
	Harvest		Ť	The state of the s	
HORTICULT	URAL CROPS				
Citrus/ Oranges	After harvesting		*	Provide proper drainage	
				Avoid application of manure/fertilizers	
Pineapple	Maturing/	Rhinocerous		Install light trap to capture and kill adult,	
	Harvesting	beetle		Use of iron rods to pull out the adults from the lower part of the	
				crop	
			*	•	
				stagnation	
			*	During clear/dry days, immediately harvest mature crops	
LIVESTOCKS					
Livestock	Stage	Disease		Precautionary Practices to be followed	
Pig	Pigs/ Piglets		*	The bedding is always dry and that there is plenty.	
	0 1 10100			If possible wooden sleeping box should be provided.	
Goat	All			Deworming provide dry bedding	
Fish	-			Harvesting and pond repairing	
1 1511	_			mai vosting and pond repairing	

Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247





### **Block Level Experimental Agro-met Advisory Bulletin**

(Period: 9th to 13th February 2019)

No: AMFU-Basar/2019/12

Dated: Friday, 8th February 2019



Gramin Krishi Mausam Sewa ICAR Research Complex for NEH Region, Arunachal Pradesh Centre, Basar-791101

### **ADVISORY FOR BLOCK: LIKABALI-KANGKU**

{Latitude: 27°59.56'/ Longitude: 94°76.79'/Altitude: 150-250 amsl}

Weather summery of the previous week	Weather forecast valid up to 13 <sup>th</sup> February 2019
NA	Fair to partly cloudy with possibility of light/moderate rain. Daily
	maximum and minimum temperature for the following period is predicted to
	be 19-29°C and 09-15°C respectively. Expected morning and evening
	relative humidity will be 58-92% and 25-77% respectively. Wind direction
	is mainly NNW with average wind speed range of 4-5 kmph.
Warning:	•

	AL RECOMMEN		
• Do not g	for possible attac o for fertilizer/ma	nure applications	S
-	drainage to all cro		
Check of	f vaccination sche		
	a. I		ADVISORIES FOR THE PERIOD
Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS	F1 . /		• F 1 C 1.10
Pea	Flowering/ Maturity		❖ For green pods of pea, pick/harvest at 4-5 days interval so as to stimulate generation of new flowers
			Regular monitoring against insect attack
			<ul> <li>Provide proper drainage in the field</li> <li>Avoid any spray operation/ application of fertilizers</li> </ul>
Maize	Land		<ul> <li>Avoid any spray operation application of fertilizers</li> <li>Procure good quality certified seeds,</li> </ul>
Widize	preparation		<ul> <li>Seed treatment of early maize with Imidichloropid @ 1-2g/10kg</li> </ul>
	FF		is recommended before sowing.
			<b>Avoid sowing till forecast of clear weather is issued</b>
			Provide proper drainage to the already sown crop
<b>EGETABLES</b>			
Early Kharif	Sowing		Nursery of early kharif vegetables like Capsicum, tomato &
vegetables			other vegetables can be done
			<ul> <li>Procure seed from certified agencies</li> </ul>
			Carry out seed treatment with Thiram/Carbendazim @ 2g/kg
			seed before sowing
Potato/	Tuber		Avoid transplanting to main field     Avoid howard a device a device a device and a second
Tomato	formation/		<ul> <li>Avoid harvesting during rainy days</li> <li>Do not wash the tubers before storage</li> </ul>
Tomato	Maturity/		<ul> <li>Washing may lead to early spoilage</li> </ul>
	Harvest		v vasining may read to early sponage
HORTICULT	URAL CROPS		
Citrus/ Oranges	After harvesting		❖ Provide proper drainage
			Avoid application of manure/fertilizers
Pineapple	Maturing/	Rhinocerous	Install light trap to capture and kill adult,
	Harvesting	beetle	Use of iron rods to pull out the adults from the lower part of the
			crop
			Provide proper drainage in the field to avoid water
			stagnation
LIVESTOCKS			<b>❖</b> During clear/dry days, immediately harvest mature crops
Livestock	Stage	Disease	Precautionary Practices to be followed
	Pigs/ Piglets	Discase	<u> </u>
Pig	rigs/ rigiets		<ul> <li>The bedding is always dry and that there is plenty.</li> <li>If possible wooden sleeping box should be provided.</li> </ul>
Goat	All		Deworming provide dry bedding
Fish	7 111		Harvesting and pond repairing
	- (ICLP)	. A 1 Cl4 . 00 TZTTT	staff for measures against adverse conditions or call at 03795-226237/226247