



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

**MANIPUR BULLETIN**

**28<sup>th</sup> October, 2014**

**Churachandpur**

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 29<sup>th</sup> October to 2<sup>nd</sup> November, 2014 as given below:

**Significant past weather for the preceding bi-weekly of Imphal - West (24<sup>th</sup> to 27<sup>th</sup> October, 2014)**

During past four days sky was mainly clear to partly cloudy but mainly cloudy to cloudy on 27<sup>th</sup> Oct. **5.2 mm** rainfall was recorded during past fourdays. Maximum temperature range observed was 21.1 – 29.7 °C, range of minimum temperature observed was 11.0 – 16.4 °C, mean relative humidity observed in morning and afternoon were 89.0 % and 66.5 % respectively. The average wind speed range was 0.9 – 1.4 km/hr.

**Weather forecast for next five days of Churachandpur (29<sup>th</sup> October to 2<sup>nd</sup> November, 2014)**

There may be no rain upto Sunday. Sky may be mainly clear to partly cloudy with maximum temp. of 26.0 – 29.0 °C & minimum temp. 12.0 – 14.0 °C. Relative humidity may reach upto 99 % (maximum RH) & 61 % (minimum RH). The average wind speed may be 3-4 km/ hr during next five days.

Name of the crop	Stage	Pest and disease infestation	Agro-meteorological advisories
<b>Field Crops</b>			
<b>Kharif Rice</b>	Maturity & harvesting		<ul style="list-style-type: none"> <li>- Drain out water before harvesting.</li> <li>- As there is likely to have no rain during next five days, harvesting of rice may be done if 80 % of the spikelet is yellow if facility for threshing is available.</li> <li>- Stacking of the harvested panicles for 5-7 days by turning the heap of panicles can help in better threshing.</li> <li>- Clean and sundry seeds properly before storage.</li> </ul>
<b>Rapeseed/ mustard</b>	Sowing		- In foot-hills and upland area zero tillage rape seed may be sown just after harvesting rice to make full utilization of the residual moisture.
<b>Field pea</b>	Land preparation to sowing		Land preparation may be done. Sowing may be done at a spacing of 30cm X 10-15cm. Recommended variety - Rachana. Seed rate is 75 kg/ hectare.
<b>Potato</b>	Land preparation to sowing		Land preparation may be done. Sowing may be done at a spacing of 60cm X 15-20cm. Recommended varieties are Kufri Megha and Kufri Jyoti. Seed rate is 25-30 quintals/ hectare.
<b>Groundnut</b>	Maturity to harvesting		<ul style="list-style-type: none"> <li>- Harvest the crop when 125 days old (var. ICGS-76) as well as dark tint inside the shell appears.</li> <li>- Make bundles and dry the crops in shade for 2-3 days for curing. Dry it for 4-5 days before storage. Select good quality pods for seeds.</li> </ul>
<b>Blackgram</b>	Maturity & harvesting		Harvest mature crops.
<b>Storage</b>			- Clean and store the seeds in RC-seed bins after proper drying. Check for germination of seeds or fungal growth. After proper drying, store grains/ seeds in well ventilated godowns.
<b>Horticulture Crops</b>			
<b>* As there is likely to have no rain in next five days, sowing/ planting of crops may be done to utilize the soil moisture for germination and vegetative growth.</b>			
<b>Coriander</b>	Land preparation & sowing		<ul style="list-style-type: none"> <li>- Select area with bright sunshine rich in organic manure for preparing beds. Treat seeds using Dichlorovos @ 1ml/ litre of water 2-3 days before sowing to protect seeds from ant and other insects.</li> <li>- Apply 200-250 kg of FYM (@ ½ cart load) in 100 sq. m and mix thoroughly to improve soil fertility.</li> <li>- Seed rate is 100g seeds for 100 sq. m.</li> </ul>
<b>Spinach beet (Palak)</b>	Land preparation &		<ul style="list-style-type: none"> <li>- Well drained soil is best for growing spinach beet.</li> <li>- Avoid damping off by adding Captan 50WP in the soil of bed.</li> </ul>



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

	sowing		Spaning may be maintained at 10cm x 20-25cm apart. - Seed rate is 100-120g/ 100 sq m. If broadcasting is done, double the dose is required.
<b>Onion</b>	Nursery raising		- Apply Captan 50 WP during final land preparation as preventive measures against soil borne diseases. - For 25 sq m area we need ½ kg seeds. - check that the new sprouts/ young plants in nursery beds are not damaged. - 45-60 days old seedlings may be transplanted.
<b>Cabbage/ Cauliflower</b>	Nursery		- Sowing of late season crop may be done. - Care should be taken for the young seedlings below 20 days old.

**Animal Sciences**

Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
<b>Poultry</b>	All susceptible age groups (as and when required)	Coccidiosis	Give preventive dose of Coximar @ 1g/ lit of drinking water for 7 days along with glucose.
		Deworming	Use Ventrimizole @ 1g in 1 litre of drinking water for 3 days.
		Litter management	Open the hessian cloth hanging on the wall and cross ventilation must be provided. Stir the litter material regularly.
<b>Sheep/ Goat</b>	Above 3 months of age (as and when required)	Tetanus	In pregnant animals, inject Tetanus Toxoid @ 0.5ml at 3.5 months of pregnancy.
<b>Pig</b>	Above 3 months of age (as and when required)	Skin infection	Give Ivermectin ( Indectin tab.) @ 10mg/33kg body weight.
		Coccidial infection (Diarrhea)	Ofloxacin/ Ornidazole (O <sub>2</sub> ) tab twice daily for 5 days.
		Deworming	- Deworm using Piperazine citrate @ 30 ml per adult animal once.
		Ticks	Add Butox liquid @ 2 ml/ lit of water and use the solution twice a week while brushing the animal.
<b>Cattle / Buffalo/ Mithun</b>	Above 3 months of age (as and when required)	Fasciolosis	- Give Tolzan-F @30 ml/ 100 kg body weight.
		Deworming	Give Nilzan/ Tolzan- F @ 30ml/ 100 kg body weight.
		Ticks/ Wounds	Topicure Spray may be used twice daily on the affected parts.

**Fishery:**

Type of Farming	Stage & Management	Agro-metrological advisories
<b>Carp/ Catfish culture</b>	Harvesting and stock maintenance	- Fish harvesting can be started. Also stock fish for next year and for this practices mentioned below may be done in small scale. - Ponds should be free from unwanted vegetation and insects and water depth should be maintained at 1.0-1.5m. - Feeds should be provided 1-2 times a day @ 3 % body weight per day. - Well balanced feed fortified with vitamins and minerals should be provided. - Apply Green AQ @ 4litre/ ha or lime @ 200kg/ ha. Maintain good quality of pond water.
<b>Paddy cum fish culture</b>	Harvesting	- Fish harvesting can be started. - Maintained 1.0 m water depth of the trenches in paddy field. - Provide rice bran and mustard oil cake (1:1) as supplementary feed. Avoid insecticide/pesticide application.

(I. Meghachandra Singh)  
Scientist i/c, GKMS



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

**MANIPUR BULLETIN**

**28<sup>th</sup> October, 2014**

**Imphal-East**

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 29<sup>th</sup> October to 2<sup>nd</sup> November, 2014 as given below:

**Significant past weather for the preceding bi-weekly of Imphal - West (24<sup>th</sup> to 27<sup>th</sup> October, 2014)**

During past four days sky was mainly clear to partly cloudy but mainly cloudy to cloudy on 27<sup>th</sup> Oct. **5.2 mm** rainfall was recorded during past fourdays. Maximum temperature range observed was 21.1 – 29.7 °C, range of minimum temperature observed was 11.0 – 16.4 °C, mean relative humidity observed in morning and afternoon were 89.0 % and 66.5 % respectively. The average wind speed range was 0.9 – 1.4 km/hr.

**Weather forecast for next five days of Imphal-East (29<sup>th</sup> October to 2<sup>nd</sup> November, 2014)**

There may be no rain upto Sunday. Sky may be mainly clear to partly cloudy with maximum temp. of 26.0 – 29.0 °C & minimum temp. 13.0 – 15.0 °C. Relative humidity may reach upto 100 % (maximum RH) & 54 % (minimum RH). The average wind speed may be 2-4 km/ hr during next five days.

Name of the crop	Stage	Pest and disease infestation	Agro-meteorological advisories
<b>Field Crops</b>			
<b>Kharif Rice</b>	Maturity & harvesting		<ul style="list-style-type: none"> <li>- Drain out water before harvesting.</li> <li>- As there is likely to have no rain during next five days, harvesting of rice may be done if 80 % of the spikelet is yellow if facility for threshing is available.</li> <li>- Stacking of the harvested panicles for 5-7 days by turning the heap of panicles can help in better threshing.</li> <li>- Clean and sundry seeds properly before storage.</li> </ul>
<b>Rapeseed/ mustard</b>	Sowing		- In foot-hills and upland area zero tillage rape seed may be sown just after harvesting rice to make full utilization of the residual moisture.
<b>Field pea</b>	Land preparation to sowing		Land preparation may be done. Sowing may be done at a spacing of 30cm X 10-15cm. Recommended variety - Rachana. Seed rate is 75 kg/ hectare.
<b>Potato</b>	Land preparation to sowing		Land preparation may be done. Sowing may be done at a spacing of 60cm X 15-20cm. Recommended varieties are Kufri Megha and Kufri Jyoti. Seed rate is 25-30 quintals/ hectare.
<b>Groundnut</b>	Maturity to harvesting		<ul style="list-style-type: none"> <li>- Harvest the crop when 125 days old (var. ICGS-76) as well as dark tint inside the shell appears.</li> <li>- Make bundles and dry the crops in shade for 2-3 days for curing. Dry it for 4-5 days before storage. Select good quality pods for seeds.</li> </ul>
<b>Blackgram</b>	Maturity & harvesting		Harvest mature crops.
<b>Storage</b>			- Clean and store the seeds in RC-seed bins after proper drying. Check for germination of seeds or fungal growth. After proper drying, store grains/ seeds in well ventilated godowns.
<b>Horticulture Crops</b>			
<b>* As there is likely to have no rain in next five days, sowing/ planting of crops may be done to utilize the soil moisture for germination and vegetative growth.</b>			
<b>Coriander</b>	Land preparation & sowing		<ul style="list-style-type: none"> <li>- Select area with bright sunshine rich in organic manure for preparing beds. Treat seeds using Dichlorovos @ 1ml/ litre of water 2-3 days before sowing to protect seeds from ant and other insects.</li> <li>- Apply 200-250 kg of FYM (@ ½ cart load) in 100 sq. m and mix thoroughly to improve soil fertility.</li> <li>- Seed rate is 100g seeds for 100 sq. m.</li> </ul>
<b>Spinach beet (Palak)</b>	Land preparation &		<ul style="list-style-type: none"> <li>- Well drained soil is best for growing spinach beet.</li> <li>- Avoid damping off by adding Captan 50WP in the soil of bed.</li> </ul>



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

	sowing		Spaning may be maintained at 10cm x 20-25cm apart. - Seed rate is 100-120g/ 100 sq m. If broadcasting is done, double the dose is required.
<b>Onion</b>	Nursery raising		- Apply Captan 50 WP during final land preparation as preventive measures against soil borne diseases. - For 25 sq m area we need ½ kg seeds. - check that the new sprouts/ young plants in nursery beds are not damaged. - 45-60 days old seedlings may be transplanted.
<b>Cabbage/ Cauliflower</b>	Nursery		- Sowing of late season crop may be done. - Care should be taken for the young seedlings below 20 days old.

**Animal Sciences**

Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
<b>Poultry</b>	All susceptible age groups (as and when required)	Coccidiosis	Give preventive dose of Coximar @ 1g/ lit of drinking water for 7 days along with glucose.
		Deworming	Use Ventrimizole @ 1g in 1 litre of drinking water for 3 days.
		Litter management	Open the hessian cloth hanging on the wall and cross ventilation must be provided. Stir the litter material regularly.
<b>Sheep/ Goat</b>	Above 3 months of age (as and when required)	Tetanus	In pregnant animals, inject Tetanus Toxoid @ 0.5ml at 3.5 months of pregnancy.
<b>Pig</b>	Above 3 months of age (as and when required)	Skin infection	Give Ivermectin ( Indectin tab.) @ 10mg/33kg body weight.
		Coccidial infection (Diarrhea)	Ofloxacin/ Ornidazole (O <sub>2</sub> ) tab twice daily for 5 days.
		Deworming	- Deworm using Piperazine citrate @ 30 ml per adult animal once.
		Ticks	Add Butox liquid @ 2 ml/ lit of water and use the solution twice a week while brushing the animal.
<b>Cattle / Buffalo/ Mithun</b>	Above 3 months of age (as and when required)	Fasciolosis	- Give Tolzan-F @30 ml/ 100 kg body weight.
		Deworming	Give Nilzan/ Tolzan- F @ 30ml/ 100 kg body weight.
		Ticks/ Wounds	Topicure Spray may be used twice daily on the affected parts.

**Fishery:**

Type of Farming	Stage & Management	Agro-metrological advisories
<b>Carp/ Catfish culture</b>	Harvesting and stock maintenance	- Fish harvesting can be started. Also stock fish for next year and for this practices mentioned below may be done in small scale. - Ponds should be free from unwanted vegetation and insects and water depth should be maintained at 1.0-1.5m. - Feeds should be provided 1-2 times a day @ 3 % body weight per day. - Well balanced feed fortified with vitamins and minerals should be provided. - Apply Green AQ @ 4litre/ ha or lime @ 200kg/ ha. Maintain good quality of pond water.
<b>Paddy cum fish culture</b>	Harvesting	- Fish harvesting can be started. - Maintained 1.0 m water depth of the trenches in paddy field. - Provide rice bran and mustard oil cake (1:1) as supplementary feed. Avoid insecticide/pesticide application.

(I. Meghachandra Singh)  
Scientist i/c, GKMS



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

**MANIPUR BULLETIN**

**28<sup>th</sup> October, 2014**

**Imphal-West**

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 29<sup>th</sup> October to 2<sup>nd</sup> November, 2014 as given below:

**Significant past weather for the preceding bi-weekly of Imphal - West (24<sup>th</sup> to 27<sup>th</sup> October, 2014)**

During past four days sky was mainly clear to partly cloudy but mainly cloudy to cloudy on 27<sup>th</sup> Oct. **5.2 mm** rainfall was recorded during past fourdays. Maximum temperature range observed was 21.1 – 29.7 °C, range of minimum temperature observed was 11.0 – 16.4 °C, mean relative humidity observed in morning and afternoon were 89.0 % and 66.5 % respectively. The average wind speed range was 0.9 – 1.4 km/hr.

**Weather forecast for next five days of Imphal-West (29<sup>th</sup> October to 2<sup>nd</sup> November, 2014)**

There may be no rain from Wednesday. Sky may be mainly clear to partly cloudy but mainly cloudy on 31<sup>st</sup> Oct with maximum temp. of 26.0 – 29.0 °C & minimum temp. 13.0 – 15.0 °C. Relative humidity may reach upto 98 % (maximum RH) & 51 % (minimum RH). The average wind speed may be 0-3 km/ hr during next five days.

Name of the crop	Stage	Pest and disease infestation	Agro-meteorological advisories
<b>Field Crops</b>			
<b>Kharif Rice</b>	Maturity & harvesting		<ul style="list-style-type: none"> <li>- Drain out water before harvesting.</li> <li>- As there is likely to have no rain during next five days, harvesting of rice may be done if 80 % of the spikelet is yellow if facility for threshing is available.</li> <li>- Stacking of the harvested panicles for 5-7 days by turning the heap of panicles can help in better threshing.</li> <li>- Clean and sundry seeds properly before storage.</li> </ul>
<b>Rapeseed/ mustard</b>	Sowing		- In foot-hills and upland area zero tillage rape seed may be sown just after harvesting rice to make full utilization of the residual moisture.
<b>Field pea</b>	Land preparation to sowing		Land preparation may be done. Sowing may be done at a spacing of 30cm X 10-15cm. Recommended variety - Rachana. Seed rate is 75 kg/ hectare.
<b>Potato</b>	Land preparation to sowing		Land preparation may be done. Sowing may be done at a spacing of 60cm X 15-20cm. Recommended varieties are Kufri Megha and Kufri Jyoti. Seed rate is 25-30 quintals/ hectare.
<b>Groundnut</b>	Maturity to harvesting		<ul style="list-style-type: none"> <li>- Harvest the crop when 125 days old (var. ICGS-76) as well as dark tint inside the shell appears.</li> <li>- Make bundles and dry the crops in shade for 2-3 days for curing. Dry it for 4-5 days before storage. Select good quality pods for seeds.</li> </ul>
<b>Blackgram</b>	Maturity & harvesting		Harvest mature crops.
<b>Storage</b>			- Clean and store the seeds in RC-seed bins after proper drying. Check for germination of seeds or fungal growth. After proper drying, store grains/ seeds in well ventilated godowns.
<b>Horticulture Crops</b>			
<b>* As there is likely to have no rain in next five days, sowing/ planting of crops may be done to utilize the soil moisture for germination and vegetative growth.</b>			
<b>Coriander</b>	Land preparation & sowing		<ul style="list-style-type: none"> <li>- Select area with bright sunshine rich in organic manure for preparing beds. Treat seeds using Dichlorovos @ 1ml/ litre of water 2-3 days before sowing to protect seeds from ant and other insects.</li> <li>- Apply 200-250 kg of FYM (@ ½ cart load) in 100 sq. m and mix thoroughly to improve soil fertility.</li> <li>- Seed rate is 100g seeds for 100 sq. m.</li> </ul>
<b>Spinach beet (Palak)</b>	Land preparation &		<ul style="list-style-type: none"> <li>- Well drained soil is best for growing spinach beet.</li> <li>- Avoid damping off by adding Captan 50WP in the soil of bed.</li> </ul>



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

	sowing		Spaning may be maintained at 10cm x 20-25cm apart. - Seed rate is 100-120g/ 100 sq m. If broadcasting is done, double the dose is required.
<b>Onion</b>	Nursery raising		- Apply Captan 50 WP during final land preparation as preventive measures against soil borne diseases. - For 25 sq m area we need ½ kg seeds. - check that the new sprouts/ young plants in nursery beds are not damaged. - 45-60 days old seedlings may be transplanted.
<b>Cabbage/ Cauliflower</b>	Nursery		- Sowing of late season crop may be done. - Care should be taken for the young seedlings below 20 days old.

**Animal Sciences**

Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
<b>Poultry</b>	All susceptible age groups (as and when required)	Coccidiosis	Give preventive dose of Coximar @ 1g/ lit of drinking water for 7 days along with glucose.
		Deworming	Use Ventrimizole @ 1g in 1 litre of drinking water for 3 days.
		Litter management	Open the hessian cloth hanging on the wall and cross ventilation must be provided. Stir the litter material regularly.
<b>Sheep/ Goat</b>	Above 3 months of age (as and when required)	Tetanus	In pregnant animals, inject Tetanus Toxoid @ 0.5ml at 3.5 months of pregnancy.
<b>Pig</b>	Above 3 months of age (as and when required)	Skin infection	Give Ivermectin ( Indectin tab.) @ 10mg/33kg body weight.
		Coccidial infection (Diarrhea)	Ofloxacin/ Ornidazole (O <sub>2</sub> ) tab twice daily for 5 days.
		Deworming	- Deworm using Piperazine citrate @ 30 ml per adult animal once.
		Ticks	Add Butox liquid @ 2 ml/ lit of water and use the solution twice a week while brushing the animal.
<b>Cattle / Buffalo/ Mithun</b>	Above 3 months of age (as and when required)	Fasciolosis	- Give Tolzan-F @30 ml/ 100 kg body weight.
		Deworming	Give Nilzan/ Tolzan- F @ 30ml/ 100 kg body weight.
		Ticks/ Wounds	Topicure Spray may be used twice daily on the affected parts.

**Fishery:**

Type of Farming	Stage & Management	Agro-metrological advisories
<b>Carp/ Catfish culture</b>	Harvesting and stock maintenance	- Fish harvesting can be started. Also stock fish for next year and for this practices mentioned below may be done in small scale. - Ponds should be free from unwanted vegetation and insects and water depth should be maintained at 1.0-1.5m. - Feeds should be provided 1-2 times a day @ 3 % body weight per day. - Well balanced feed fortified with vitamins and minerals should be provided. - Apply Green AQ @ 4litre/ ha or lime @ 200kg/ ha. Maintain good quality of pond water.
<b>Paddy cum fish culture</b>	Harvesting	- Fish harvesting can be started. - Maintained 1.0 m water depth of the trenches in paddy field. - Provide rice bran and mustard oil cake (1:1) as supplementary feed. Avoid insecticide/pesticide application.

(I. Meghachandra Singh)  
Scientist i/c, GKMS



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

**MANIPUR BULLETIN**

**28<sup>th</sup> October, 2014**

**Senapati**

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 29<sup>th</sup> October to 2<sup>nd</sup> November, 2014 as given below:

**Significant past weather for the preceding bi-weekly of Imphal - West (24<sup>th</sup> to 27<sup>th</sup> October, 2014)**

During past four days sky was mainly clear to partly cloudy but mainly cloudy to cloudy on 27<sup>th</sup> Oct. **5.2 mm** rainfall was recorded during past fourdays. Maximum temperature range observed was 21.1 – 29.7 °C, range of minimum temperature observed was 11.0 – 16.4 °C, mean relative humidity observed in morning and afternoon were 89.0 % and 66.5 % respectively. The average wind speed range was 0.9 – 1.4 km/hr.

**Weather forecast for next five days of Senapati (29<sup>th</sup> October to 2<sup>nd</sup> November, 2014)**

There may be no rain upto Sunday. Sky may be mainly clear to partly cloudy with maximum temp. of 25.0 – 27.0 °C & minimum temp. 9.0 – 11.0 °C. Relative humidity may reach upto 99 % (maximum RH) & 50 % (minimum RH). The average wind speed may be 2-4 km/ hr during next five days.

Name of the crop	Stage	Pest and disease infestation	Agro-meteorological advisories
<b>Field Crops</b>			
<b>Kharif Rice</b>	Maturity & harvesting		<ul style="list-style-type: none"> <li>- Drain out water before harvesting.</li> <li>- As there is likely to have no rain during next five days, harvesting of rice may be done if 80 % of the spikelet is yellow if facility for threshing is available.</li> <li>- Stacking of the harvested panicles for 5-7 days by turning the heap of panicles can help in better threshing.</li> <li>- Clean and sundry seeds properly before storage.</li> </ul>
<b>Rapeseed/ mustard</b>	Sowing		- In foot-hills and upland area zero tillage rape seed may be sown just after harvesting rice to make full utilization of the residual moisture.
<b>Field pea</b>	Land preparation to sowing		Land preparation may be done. Sowing may be done at a spacing of 30cm X 10-15cm. Recommended variety - Rachana. Seed rate is 75 kg/ hectare.
<b>Groundnut</b>	Maturity to harvesting		<ul style="list-style-type: none"> <li>- Harvest the crop when 125 days old (var. ICGS-76) as well as dark tint inside the shell appears.</li> <li>- Make bundles and dry the crops in shade for 2-3 days for curing. Dry it for 4-5 days before storage. Select good quality pods for seeds.</li> </ul>
<b>Blackgram</b>	Maturity & harvesting		Harvest mature crops.
<b>Storage</b>			- Clean and store the seeds in RC-seed bins after proper drying. Check for germination of seeds or fungal growth. After proper drying, store grains/ seeds in well ventilated godowns.
<b>Horticulture Crops</b>			
<b>* As there is likely to have no rain in next five days, sowing/ planting of crops may be done to utilize the soil moisture for germination and vegetative growth.</b>			
<b>Coriander</b>	Land preparation & sowing		<ul style="list-style-type: none"> <li>- Select area with bright sunshine rich in organic manure for preparing beds. Treat seeds using Dichlorovos @ 1ml/ litre of water 2-3 days before sowing to protect seeds from ant and other insects.</li> <li>- Apply 200-250 kg of FYM (@ ½ cart load) in 100 sq. m and mix thoroughly to improve soil fertility.</li> <li>- Seed rate is 100g seeds for 100 sq. m.</li> </ul>
<b>Spinach beet (Palak)</b>	Land preparation & sowing		<ul style="list-style-type: none"> <li>- Well drained soil is best for growing spinach beet.</li> <li>- Avoid damping off by adding Captan 50WP in the soil of bed. Spaning may be maintained at 10cm x 20-25cm apart.</li> <li>- Seed rate is 100-120g/ 100 sq m. If broadcasting is done, double the dose is required.</li> </ul>



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

<b>Onion</b>	Nursery raising		<ul style="list-style-type: none"> <li>- Apply Captan 50 WP during final land preparation as preventive measures against soil borne diseases.</li> <li>- For 25 sq m area we need ½ kg seeds.</li> <li>- check that the new sprouts/ young plants in nursery beds are not damaged.</li> <li>- 45-60 days old seedlings may be transplanted.</li> </ul>
<b>Cabbage/ Cauliflower</b>	Nursery		<ul style="list-style-type: none"> <li>- Sowing of late season crop may be done.</li> <li>- Care should be taken for the young seedlings below 20 days old.</li> </ul>

**Animal Sciences**

Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
<b>Poultry</b>	All susceptible age groups (as and when required)	Coccidiosis	Give preventive dose of Coximar @ 1g/ lit of drinking water for 7 days along with glucose.
		Deworming	Use Ventrimizole @ 1g in 1 litre of drinking water for 3 days.
		Litter management	Open the hessian cloth hanging on the wall and cross ventilation must be provided. Stir the litter material regularly.
<b>Sheep/ Goat</b>	Above 3 months of age (as and when required)	Tetanus	In pregnant animals, inject Tetanus Toxoid @ 0.5ml at 3.5 months of pregnancy.
<b>Pig</b>	Above 3 months of age (as and when required)	Skin infection	Give Ivermectin ( Indectin tab.) @ 10mg/33kg body weight.
		Coccidial infection (Diarrhea)	Ofloxacin/ Ornidazole (O <sub>2</sub> ) tab twice daily for 5 days.
		Deworming	- Deworm using Piperazine citrate @ 30 ml per adult animal once.
		Ticks	Add Butox liquid @ 2 ml/ lit of water and use the solution twice a week while brushing the animal.
<b>Cattle / Buffalo/ Mithun</b>	Above 3 months of age (as and when required)	Fasciolosis	- Give Tolzan-F @ 30 ml/ 100 kg body weight.
		Deworming	Give Nilzan/ Tolzan- F @ 30ml/ 100 kg body weight.
		Ticks/ Wounds	Topicure Spray may be used twice daily on the affected parts.

**Fishery:**

Type of Farming	Stage & Management	Agro-metrological advisories
<b>Carp/ Catfish culture</b>	Harvesting and stock maintenance	<ul style="list-style-type: none"> <li>- Fish harvesting can be started. Also stock fish for next year and for this practices mentioned below may be done in small scale.</li> <li>- Ponds should be free from unwanted vegetation and insects and water depth should be maintained at 1.0-1.5m.</li> <li>- Feeds should be provided 1-2 times a day @ 3 % body weight per day.</li> <li>- Well balanced feed fortified with vitamins and minerals should be provided.</li> <li>- Apply Green AQ @ 4litre/ ha or lime @ 200kg/ ha. Maintain good quality of pond water.</li> </ul>
<b>Paddy cum fish culture</b>	Harvesting	<ul style="list-style-type: none"> <li>- Fish harvesting can be started.</li> <li>- Maintained 1.0 m water depth of the trenches in paddy field.</li> <li>- Provide rice bran and mustard oil cake (1:1) as supplementary feed. Avoid insecticide/pesticide application.</li> </ul>

(I. Meghachandra Singh)  
Scientist i/c, GKMS



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

**MANIPUR BULLETIN**

**28<sup>th</sup> October, 2014**

**Tamenglong**

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 29<sup>th</sup> October to 2<sup>nd</sup> November, 2014 as given below:

**Significant past weather for the preceding bi-weekly of Imphal - West (24<sup>th</sup> to 27<sup>th</sup> October, 2014)**

During past four days sky was mainly clear to partly cloudy but mainly cloudy to cloudy on 27<sup>th</sup> Oct. **5.2 mm** rainfall was recorded during past fourdays. Maximum temperature range observed was 21.1 – 29.7 °C, range of minimum temperature observed was 11.0 – 16.4 °C, mean relative humidity observed in morning and afternoon were 89.0 % and 66.5 % respectively. The average wind speed range was 0.9 – 1.4 km/hr.

**Weather forecast for next five days of Tamenglong (29<sup>th</sup> October to 2<sup>nd</sup> November, 2014)**

There may be no rain upto Sunday. Sky may be mainly clear to partly cloudy with maximum temp. of 27.0 – 30.0 °C & minimum temp. 13.0 – 15.0 °C. Relative humidity may reach upto 97 % (maximum RH) & 51 % (minimum RH). The average wind speed may be 3-4 km/ hr during next five days.

Name of the crop	Stage	Pest and disease infestation	Agro-meteorological advisories
<b>Field Crops</b>			
<b>Kharif Rice</b>	Maturity & harvesting		<ul style="list-style-type: none"> <li>- Drain out water before harvesting.</li> <li>- As there is likely to have no rain during next five days, harvesting of rice may be done if 80 % of the spikelet is yellow if facility for threshing is available.</li> <li>- Stacking of the harvested panicles for 5-7 days by turning the heap of panicles can help in better threshing.</li> <li>- Clean and sundry seeds properly before storage.</li> </ul>
<b>Rapeseed/ mustard</b>	Sowing		- In foot-hills and upland area zero tillage rape seed may be sown just after harvesting rice to make full utilization of the residual moisture.
<b>Field pea</b>	Land preparation to sowing		Land preparation may be done. Sowing may be done at a spacing of 30cm X 10-15cm. Recommended variety - Rachana. Seed rate is 75 kg/ hectare.
<b>Potato</b>	Land preparation to sowing		Land preparation may be done. Sowing may be done at a spacing of 60cm X 15-20cm. Recommended varieties are Kufri Megha and Kufri Jyoti. Seed rate is 25-30 quintals/ hectare.
<b>Groundnut</b>	Maturity to harvesting		<ul style="list-style-type: none"> <li>- Harvest the crop when 125 days old (var. ICGS-76) as well as dark tint inside the shell appears.</li> <li>- Make bundles and dry the crops in shade for 2-3 days for curing. Dry it for 4-5 days before storage. Select good quality pods for seeds.</li> </ul>
<b>Blackgram</b>	Maturity & harvesting		Harvest mature crops.
<b>Storage</b>			- Clean and store the seeds in RC-seed bins after proper drying. Check for germination of seeds or fungal growth. After proper drying, store grains/ seeds in well ventilated godowns.
<b>Horticulture Crops</b>			
<b>* As there is likely to have no rain in next five days, sowing/ planting of crops may be done to utilize the soil moisture for germination and vegetative growth.</b>			
<b>Coriander</b>	Land preparation & sowing		<ul style="list-style-type: none"> <li>- Select area with bright sunshine rich in organic manure for preparing beds. Treat seeds using Dichlorovos @ 1ml/ litre of water 2-3 days before sowing to protect seeds from ant and other insects.</li> <li>- Apply 200-250 kg of FYM (@ ½ cart load) in 100 sq. m and mix thoroughly to improve soil fertility.</li> <li>- Seed rate is 100g seeds for 100 sq. m.</li> </ul>
<b>Spinach beet (Palak)</b>	Land preparation &		<ul style="list-style-type: none"> <li>- Well drained soil is best for growing spinach beet.</li> <li>- Avoid damping off by adding Captan 50WP in the soil of bed.</li> </ul>



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

	sowing		Spaning may be maintained at 10cm x 20-25cm apart. - Seed rate is 100-120g/ 100 sq m. If broadcasting is done, double the dose is required.
<b>Onion</b>	Nursery raising		- Apply Captan 50 WP during final land preparation as preventive measures against soil borne diseases. - For 25 sq m area we need ½ kg seeds. - check that the new sprouts/ young plants in nursery beds are not damaged. - 45-60 days old seedlings may be transplanted.
<b>Cabbage/ Cauliflower</b>	Nursery		- Sowing of late season crop may be done. - Care should be taken for the young seedlings below 20 days old.

**Animal Sciences**

Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
<b>Poultry</b>	All susceptible age groups (as and when required)	Coccidiosis	Give preventive dose of Coximar @ 1g/ lit of drinking water for 7 days along with glucose.
		Deworming	Use Ventrimizole @ 1g in 1 litre of drinking water for 3 days.
		Litter management	Open the hessian cloth hanging on the wall and cross ventilation must be provided. Stir the litter material regularly.
<b>Sheep/ Goat</b>	Above 3 months of age (as and when required)	Tetanus	In pregnant animals, inject Tetanus Toxoid @ 0.5ml at 3.5 months of pregnancy.
<b>Pig</b>	Above 3 months of age (as and when required)	Skin infection	Give Ivermectin ( Indectin tab.) @ 10mg/33kg body weight.
		Coccidial infection (Diarrhea)	Ofloxacin/ Ornidazole (O <sub>2</sub> ) tab twice daily for 5 days.
		Deworming	- Deworm using Piperazine citrate @ 30 ml per adult animal once.
		Ticks	Add Butox liquid @ 2 ml/ lit of water and use the solution twice a week while brushing the animal.
<b>Cattle / Buffalo/ Mithun</b>	Above 3 months of age (as and when required)	Fasciolosis	- Give Tolzan-F @30 ml/ 100 kg body weight.
		Deworming	Give Nilzan/ Tolzan- F @ 30ml/ 100 kg body weight.
		Ticks/ Wounds	Topicure Spray may be used twice daily on the affected parts.

**Fishery:**

Type of Farming	Stage & Management	Agro-metrological advisories
<b>Carp/ Catfish culture</b>	Harvesting and stock maintenance	- Fish harvesting can be started. Also stock fish for next year and for this practices mentioned below may be done in small scale. - Ponds should be free from unwanted vegetation and insects and water depth should be maintained at 1.0-1.5m. - Feeds should be provided 1-2 times a day @ 3 % body weight per day. - Well balanced feed fortified with vitamins and minerals should be provided. - Apply Green AQ @ 4litre/ ha or lime @ 200kg/ ha. Maintain good quality of pond water.
<b>Paddy cum fish culture</b>	Harvesting	- Fish harvesting can be started. - Maintained 1.0 m water depth of the trenches in paddy field. - Provide rice bran and mustard oil cake (1:1) as supplementary feed. Avoid insecticide/pesticide application.

(I. Meghachandra Singh)  
Scientist i/c, GKMS



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

**MANIPUR BULLETIN**

**28<sup>th</sup> October, 2014**

**Thoubal**

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 29<sup>th</sup> October to 2<sup>nd</sup> November, 2014 as given below:

**Significant past weather for the preceding bi-weekly of Imphal - West (24<sup>th</sup> to 27<sup>th</sup> October, 2014)**

During past four days sky was mainly clear to partly cloudy but mainly cloudy to cloudy on 27<sup>th</sup> Oct. **5.2 mm** rainfall was recorded during past fourdays. Maximum temperature range observed was 21.1 – 29.7 °C, range of minimum temperature observed was 11.0 – 16.4 °C, mean relative humidity observed in morning and afternoon were 89.0 % and 66.5 % respectively. The average wind speed range was 0.9 – 1.4 km/hr.

**Weather forecast for next five days of Thoubal (29<sup>th</sup> October to 2<sup>nd</sup> November, 2014)**

There may be no rain upto Sunday. Sky may be mainly clear to partly cloudy with maximum temp. of 27.0 – 28.0 °C & minimum temp. 13.0 – 14.0 °C. Relative humidity may reach upto 100 % (maximum RH) & 73 % (minimum RH). The average wind speed may be 2 km/ hr during next five days.

Name of the crop	Stage	Pest and disease infestation	Agro-meteorological advisories
<b>Field Crops</b>			
<b>Kharif Rice</b>	Maturity & harvesting		<ul style="list-style-type: none"> <li>- Drain out water before harvesting.</li> <li>- As there is likely to have no rain during next five days, harvesting of rice may be done if 80 % of the spikelet is yellow if facility for threshing is available.</li> <li>- Stacking of the harvested panicles for 5-7 days by turning the heap of panicles can help in better threshing.</li> <li>- Clean and sundry seeds properly before storage.</li> </ul>
<b>Rapeseed/ mustard</b>	Sowing		- In foot-hills and upland area zero tillage rape seed may be sown just after harvesting rice to make full utilization of the residual moisture.
<b>Field pea</b>	Land preparation to sowing		Land preparation may be done. Sowing may be done at a spacing of 30cm X 10-15cm. Recommended variety - Rachana. Seed rate is 75 kg/ hectare.
<b>Potato</b>	Land preparation to sowing		Land preparation may be done. Sowing may be done at a spacing of 60cm X 15-20cm. Recommended varieties are Kufri Megha and Kufri Jyoti. Seed rate is 25-30 quintals/ hectare.
<b>Groundnut</b>	Maturity to harvesting		<ul style="list-style-type: none"> <li>- Harvest the crop when 125 days old (var. ICGS-76) as well as dark tint inside the shell appears.</li> <li>- Make bundles and dry the crops in shade for 2-3 days for curing. Dry it for 4-5 days before storage. Select good quality pods for seeds.</li> </ul>
<b>Blackgram</b>	Maturity & harvesting		Harvest mature crops.
<b>Storage</b>			- Clean and store the seeds in RC-seed bins after proper drying. Check for germination of seeds or fungal growth. After proper drying, store grains/ seeds in well ventilated godowns.
<b>Horticulture Crops</b>			
<b>* As there is likely to have no rain in next five days, sowing/ planting of crops may be done to utilize the soil moisture for germination and vegetative growth.</b>			
<b>Coriander</b>	Land preparation & sowing		<ul style="list-style-type: none"> <li>- Select area with bright sunshine rich in organic manure for preparing beds. Treat seeds using Dichlorovos @ 1ml/ litre of water 2-3 days before sowing to protect seeds from ant and other insects.</li> <li>- Apply 200-250 kg of FYM (@ ½ cart load) in 100 sq. m and mix thoroughly to improve soil fertility.</li> <li>- Seed rate is 100g seeds for 100 sq. m.</li> </ul>
<b>Spinach beet (Palak)</b>	Land preparation &		<ul style="list-style-type: none"> <li>- Well drained soil is best for growing spinach beet.</li> <li>- Avoid damping off by adding Captan 50WP in the soil of bed.</li> </ul>



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

	sowing		Spaning may be maintained at 10cm x 20-25cm apart. - Seed rate is 100-120g/ 100 sq m. If broadcasting is done, double the dose is required.
<b>Onion</b>	Nursery raising		- Apply Captan 50 WP during final land preparation as preventive measures against soil borne diseases. - For 25 sq m area we need ½ kg seeds. - check that the new sprouts/ young plants in nursery beds are not damaged. - 45-60 days old seedlings may be transplanted.
<b>Cabbage/ Cauliflower</b>	Nursery		- Sowing of late season crop may be done. - Care should be taken for the young seedlings below 20 days old.

**Animal Sciences**

Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
<b>Poultry</b>	All susceptible age groups (as and when required)	Coccidiosis	Give preventive dose of Coximar @ 1g/ lit of drinking water for 7 days along with glucose.
		Deworming	Use Ventrimizole @ 1g in 1 litre of drinking water for 3 days.
		Litter management	Open the hessian cloth hanging on the wall and cross ventilation must be provided. Stir the litter material regularly.
<b>Sheep/ Goat</b>	Above 3 months of age (as and when required)	Tetanus	In pregnant animals, inject Tetanus Toxoid @ 0.5ml at 3.5 months of pregnancy.
<b>Pig</b>	Above 3 months of age (as and when required)	Skin infection	Give Ivermectin ( Indectin tab.) @ 10mg/33kg body weight.
		Coccidial infection (Diarrhea)	Ofloxacin/ Ornidazole (O <sub>2</sub> ) tab twice daily for 5 days.
		Deworming	- Deworm using Piperazine citrate @ 30 ml per adult animal once.
		Ticks	Add Butox liquid @ 2 ml/ lit of water and use the solution twice a week while brushing the animal.
<b>Cattle / Buffalo/ Mithun</b>	Above 3 months of age (as and when required)	Fasciolosis	- Give Tolzan-F @30 ml/ 100 kg body weight.
		Deworming	Give Nilzan/ Tolzan- F @ 30ml/ 100 kg body weight.
		Ticks/ Wounds	Topicure Spray may be used twice daily on the affected parts.

**Fishery:**

Type of Farming	Stage & Management	Agro-metrological advisories
<b>Carp/ Catfish culture</b>	Harvesting and stock maintenance	- Fish harvesting can be started. Also stock fish for next year and for this practices mentioned below may be done in small scale. - Ponds should be free from unwanted vegetation and insects and water depth should be maintained at 1.0-1.5m. - Feeds should be provided 1-2 times a day @ 3 % body weight per day. - Well balanced feed fortified with vitamins and minerals should be provided. - Apply Green AQ @ 4litre/ ha or lime @ 200kg/ ha. Maintain good quality of pond water.
<b>Paddy cum fish culture</b>	Harvesting	- Fish harvesting can be started. - Maintained 1.0 m water depth of the trenches in paddy field. - Provide rice bran and mustard oil cake (1:1) as supplementary feed. Avoid insecticide/pesticide application.

(I. Meghachandra Singh)  
Scientist i/c, GKMS



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

**MANIPUR BULLETIN**

**28<sup>th</sup> October, 2014**

**Ukhrul**

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 29<sup>th</sup> October to 2<sup>nd</sup> November, 2014 as given below:

**Significant past weather for the preceding bi-weekly of Imphal - West (24<sup>th</sup> to 27<sup>th</sup> October, 2014)**

During past four days sky was mainly clear to partly cloudy but mainly cloudy to cloudy on 27<sup>th</sup> Oct. **5.2 mm** rainfall was recorded during past fourdays. Maximum temperature range observed was 21.1 – 29.7 °C, range of minimum temperature observed was 11.0 – 16.4 °C, mean relative humidity observed in morning and afternoon were 89.0 % and 66.5 % respectively. The average wind speed range was 0.9 – 1.4 km/hr.

**Weather forecast for next five days of Ukhrul (29<sup>th</sup> October to 2<sup>nd</sup> November, 2014)**

There may be no rain upto Sunday. Sky may be mainly clear to partly cloudy with maximum temp. of 25.0 – 27.0 °C & minimum temp. 9.0 – 11.0 °C. Relative humidity may reach upto 100 % (maximum RH) & 73 % (minimum RH). The average wind speed may be 3 km/ hr during next five days.

Name of the crop	Stage	Pest and disease infestation	Agro-meteorological advisories
<b>Field Crops</b>			
<b>Kharif Rice</b>	Maturity & harvesting		<ul style="list-style-type: none"> <li>- Drain out water before harvesting.</li> <li>- As there is likely to have no rain during next five days, harvesting of rice may be done if 80 % of the spikelet is yellow if facility for threshing is available.</li> <li>- Stacking of the harvested panicles for 5-7 days by turning the heap of panicles can help in better threshing.</li> <li>- Clean and sundry seeds properly before storage.</li> </ul>
<b>Rapeseed/ mustard</b>	Sowing		- In foot-hills and upland area zero tillage rape seed may be sown just after harvesting rice to make full utilization of the residual moisture.
<b>Field pea</b>	Land preparation to sowing		Land preparation may be done. Sowing may be done at a spacing of 30cm X 10-15cm. Recommended variety - Rachana. Seed rate is 75 kg/ hectare.
<b>Groundnut</b>	Maturity to harvesting		<ul style="list-style-type: none"> <li>- Harvest the crop when 125 days old (var. ICGS-76) as well as dark tint inside the shell appears.</li> <li>- Make bundles and dry the crops in shade for 2-3 days for curing. Dry it for 4-5 days before storage. Select good quality pods for seeds.</li> </ul>
<b>Blackgram</b>	Maturity & harvesting		Harvest mature crops.
<b>Storage</b>			- Clean and store the seeds in RC-seed bins after proper drying. Check for germination of seeds or fungal growth. After proper drying, store grains/ seeds in well ventilated godowns.
<b>Horticulture Crops</b>			
<b>* As there is likely to have no rain in next five days, sowing/ planting of crops may be done to utilize the soil moisture for germination and vegetative growth.</b>			
<b>Coriander</b>	Land preparation & sowing		<ul style="list-style-type: none"> <li>- Select area with bright sunshine rich in organic manure for preparing beds. Treat seeds using Dichlorovos @ 1ml/ litre of water 2-3 days before sowing to protect seeds from ant and other insects.</li> <li>- Apply 200-250 kg of FYM (@ ½ cart load) in 100 sq. m and mix thoroughly to improve soil fertility.</li> <li>- Seed rate is 100g seeds for 100 sq. m.</li> </ul>
<b>Spinach beet (Palak)</b>	Land preparation & sowing		<ul style="list-style-type: none"> <li>- Well drained soil is best for growing spinach beet.</li> <li>- Avoid damping off by adding Captan 50WP in the soil of bed. Spaning may be maintained at 10cm x 20-25cm apart.</li> <li>- Seed rate is 100-120g/ 100 sq m. If broadcasting is done, double the dose is required.</li> </ul>



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

<b>Onion</b>	Nursery raising		<ul style="list-style-type: none"> <li>- Apply Captan 50 WP during final land preparation as preventive measures against soil borne diseases.</li> <li>- For 25 sq m area we need ½ kg seeds.</li> <li>- check that the new sprouts/ young plants in nursery beds are not damaged.</li> <li>- 45-60 days old seedlings may be transplanted.</li> </ul>
<b>Cabbage/ Cauliflower</b>	Nursery		<ul style="list-style-type: none"> <li>- Sowing of late season crop may be done.</li> <li>- Care should be taken for the young seedlings below 20 days old.</li> </ul>

**Animal Sciences**

Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
<b>Poultry</b>	All susceptible age groups (as and when required)	Coccidiosis	Give preventive dose of Coximar @ 1g/ lit of drinking water for 7 days along with glucose.
		Deworming	Use Ventrimizole @ 1g in 1 litre of drinking water for 3 days.
		Litter management	Open the hessian cloth hanging on the wall and cross ventilation must be provided. Stir the litter material regularly.
<b>Sheep/ Goat</b>	Above 3 months of age (as and when required)	Tetanus	In pregnant animals, inject Tetanus Toxoid @ 0.5ml at 3.5 months of pregnancy.
<b>Pig</b>	Above 3 months of age (as and when required)	Skin infection	Give Ivermectin ( Indectin tab.) @ 10mg/33kg body weight.
		Coccidial infection (Diarrhea)	Ofloxacin/ Ornidazole (O <sub>2</sub> ) tab twice daily for 5 days.
		Deworming	- Deworm using Piperazine citrate @ 30 ml per adult animal once.
		Ticks	Add Butox liquid @ 2 ml/ lit of water and use the solution twice a week while brushing the animal.
<b>Cattle / Buffalo/ Mithun</b>	Above 3 months of age (as and when required)	Fasciolosis	- Give Tolzan-F @ 30 ml/ 100 kg body weight.
		Deworming	Give Nilzan/ Tolzan- F @ 30ml/ 100 kg body weight.
		Ticks/ Wounds	Topicure Spray may be used twice daily on the affected parts.

**Fishery:**

Type of Farming	Stage & Management	Agro-metrological advisories
<b>Carp/ Catfish culture</b>	Harvesting and stock maintenance	<ul style="list-style-type: none"> <li>- Fish harvesting can be started. Also stock fish for next year and for this practices mentioned below may be done in small scale.</li> <li>- Ponds should be free from unwanted vegetation and insects and water depth should be maintained at 1.0-1.5m.</li> <li>- Feeds should be provided 1-2 times a day @ 3 % body weight per day.</li> <li>- Well balanced feed fortified with vitamins and minerals should be provided.</li> <li>- Apply Green AQ @ 4litre/ ha or lime @ 200kg/ ha. Maintain good quality of pond water.</li> </ul>
<b>Paddy cum fish culture</b>	Harvesting	<ul style="list-style-type: none"> <li>- Fish harvesting can be started.</li> <li>- Maintained 1.0 m water depth of the trenches in paddy field.</li> <li>- Provide rice bran and mustard oil cake (1:1) as supplementary feed. Avoid insecticide/pesticide application.</li> </ul>

(I. Meghachandra Singh)  
Scientist i/c, GKMS



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

**MANIPUR BULLETIN**

**28<sup>th</sup> October, 2014**

**Bishnupur**

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 29<sup>th</sup> October to 2<sup>nd</sup> November, 2014 as given below:

**Significant past weather for the preceding bi-weekly of Imphal - West (24<sup>th</sup> to 27<sup>th</sup> October, 2014)**

During past four days sky was mainly clear to partly cloudy but mainly cloudy to cloudy on 27<sup>th</sup> Oct. **5.2 mm** rainfall was recorded during past fourdays. Maximum temperature range observed was 21.1 – 29.7 °C, range of minimum temperature observed was 11.0 – 16.4 °C, mean relative humidity observed in morning and afternoon were 89.0 % and 66.5 % respectively. The average wind speed range was 0.9 – 1.4 km/hr.

**Weather forecast for next five days of Bishnupur (29<sup>th</sup> October to 2<sup>nd</sup> November, 2014)**

There may be no rain upto Sunday. Sky may be mainly clear to partly cloudy with maximum temp. of 27.0 – 29.0 °C & minimum temp. 12.0 – 14.0 °C. Relative humidity may reach upto 99 % (maximum RH) & 66 % (minimum RH). The average wind speed may be 2-4 km/ hr during next five days.

Name of the crop	Stage	Pest and disease infestation	Agro-meteorological advisories
<b>Field Crops</b>			
<b>Kharif Rice</b>	Maturity & harvesting		<ul style="list-style-type: none"> <li>- Drain out water before harvesting.</li> <li>- As there is likely to have no rain during next five days, harvesting of rice may be done if 80 % of the spikelet is yellow if facility for threshing is available.</li> <li>- Stacking of the harvested panicles for 5-7 days by turning the heap of panicles can help in better threshing.</li> <li>- Clean and sundry seeds properly before storage.</li> </ul>
<b>Rapeseed/ mustard</b>	Sowing		- In foot-hills and upland area zero tillage rape seed may be sown just after harvesting rice to make full utilization of the residual moisture.
<b>Field pea</b>	Land preparation to sowing		Land preparation may be done. Sowing may be done at a spacing of 30cm X 10-15cm. Recommended variety - Rachana. Seed rate is 75 kg/ hectare.
<b>Potato</b>	Land preparation to sowing		Land preparation may be done. Sowing may be done at a spacing of 60cm X 15-20cm. Recommended varieties are Kufri Megha and Kufri Jyoti. Seed rate is 25-30 quintals/ hectare.
<b>Groundnut</b>	Maturity to harvesting		<ul style="list-style-type: none"> <li>- Harvest the crop when 125 days old (var. ICGS-76) as well as dark tint inside the shell appears.</li> <li>- Make bundles and dry the crops in shade for 2-3 days for curing. Dry it for 4-5 days before storage. Select good quality pods for seeds.</li> </ul>
<b>Blackgram</b>	Maturity & harvesting		Harvest mature crops.
<b>Storage</b>			- Clean and store the seeds in RC-seed bins after proper drying. Check for germination of seeds or fungal growth. After proper drying, store grains/ seeds in well ventilated godowns.
<b>Horticulture Crops</b>			
<b>* As there is likely to have no rain in next five days, sowing/ planting of crops may be done to utilize the soil moisture for germination and vegetative growth.</b>			
<b>Coriander</b>	Land preparation & sowing		<ul style="list-style-type: none"> <li>- Select area with bright sunshine rich in organic manure for preparing beds. Treat seeds using Dichlorovos @ 1ml/ litre of water 2-3 days before sowing to protect seeds from ant and other insects.</li> <li>- Apply 200-250 kg of FYM (@ ½ cart load) in 100 sq. m and mix thoroughly to improve soil fertility.</li> <li>- Seed rate is 100g seeds for 100 sq. m.</li> </ul>
<b>Spinach beet (Palak)</b>	Land preparation &		<ul style="list-style-type: none"> <li>- Well drained soil is best for growing spinach beet.</li> <li>- Avoid damping off by adding Captan 50WP in the soil of bed.</li> </ul>



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

	sowing		Spaning may be maintained at 10cm x 20-25cm apart. - Seed rate is 100-120g/ 100 sq m. If broadcasting is done, double the dose is required.
<b>Onion</b>	Nursery raising		- Apply Captan 50 WP during final land preparation as preventive measures against soil borne diseases. - For 25 sq m area we need ½ kg seeds. - check that the new sprouts/ young plants in nursery beds are not damaged. - 45-60 days old seedlings may be transplanted.
<b>Cabbage/ Cauliflower</b>	Nursery		- Sowing of late season crop may be done. - Care should be taken for the young seedlings below 20 days old.

**Animal Sciences**

Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
<b>Poultry</b>	All susceptible age groups (as and when required)	Coccidiosis	Give preventive dose of Coximar @ 1g/ lit of drinking water for 7 days along with glucose.
		Deworming	Use Ventrimizole @ 1g in 1 litre of drinking water for 3 days.
		Litter management	Open the hessian cloth hanging on the wall and cross ventilation must be provided. Stir the litter material regularly.
<b>Sheep/ Goat</b>	Above 3 months of age (as and when required)	Tetanus	In pregnant animals, inject Tetanus Toxoid @ 0.5ml at 3.5 months of pregnancy.
<b>Pig</b>	Above 3 months of age (as and when required)	Skin infection	Give Ivermectin ( Indectin tab.) @ 10mg/33kg body weight.
		Coccidial infection (Diarrhea)	Ofloxacin/ Ornidazole (O <sub>2</sub> ) tab twice daily for 5 days.
		Deworming	- Deworm using Piperazine citrate @ 30 ml per adult animal once.
		Ticks	Add Butox liquid @ 2 ml/ lit of water and use the solution twice a week while brushing the animal.
<b>Cattle / Buffalo/ Mithun</b>	Above 3 months of age (as and when required)	Fasciolosis	- Give Tolzan-F @30 ml/ 100 kg body weight.
		Deworming	Give Nilzan/ Tolzan- F @ 30ml/ 100 kg body weight.
		Ticks/ Wounds	Topicure Spray may be used twice daily on the affected parts.

**Fishery:**

Type of Farming	Stage & Management	Agro-metrological advisories
<b>Carp/ Catfish culture</b>	Harvesting and stock maintenance	- Fish harvesting can be started. Also stock fish for next year and for this practices mentioned below may be done in small scale. - Ponds should be free from unwanted vegetation and insects and water depth should be maintained at 1.0-1.5m. - Feeds should be provided 1-2 times a day @ 3 % body weight per day. - Well balanced feed fortified with vitamins and minerals should be provided. - Apply Green AQ @ 4litre/ ha or lime @ 200kg/ ha. Maintain good quality of pond water.
<b>Paddy cum fish culture</b>	Harvesting	- Fish harvesting can be started. - Maintained 1.0 m water depth of the trenches in paddy field. - Provide rice bran and mustard oil cake (1:1) as supplementary feed. Avoid insecticide/pesticide application.

(I. Meghachandra Singh)  
Scientist i/c, GKMS



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

**MANIPUR BULLETIN**

**28<sup>th</sup> October, 2014**

**Chandel**

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 29<sup>th</sup> October to 2<sup>nd</sup> November, 2014 as given below:

**Significant past weather for the preceding bi-weekly of Imphal - West (24<sup>th</sup> to 27<sup>th</sup> October, 2014)**

During past four days sky was mainly clear to partly cloudy but mainly cloudy to cloudy on 27<sup>th</sup> Oct. **5.2 mm** rainfall was recorded during past fourdays. Maximum temperature range observed was 21.1 – 29.7 °C, range of minimum temperature observed was 11.0 – 16.4 °C, mean relative humidity observed in morning and afternoon were 89.0 % and 66.5 % respectively. The average wind speed range was 0.9 – 1.4 km/hr.

**Weather forecast for next five days of Chandel (29<sup>th</sup> October to 2<sup>nd</sup> November, 2014)**

There may be no rain upto Sunday. Sky may be mainly clear to partly cloudy with maximum temp. of 27.0 – 28.0 °C & minimum temp. 13.0 – 14.0 °C. Relative humidity may reach upto 100 % (maximum RH) & 73 % (minimum RH). The average wind speed may be 2 km/ hr during next five days.

Name of the crop	Stage	Pest and disease infestation	Agro-meteorological advisories
<b>Field Crops</b>			
<b>Kharif Rice</b>	Maturity & harvesting		<ul style="list-style-type: none"> <li>- Drain out water before harvesting.</li> <li>- As there is likely to have no rain during next five days, harvesting of rice may be done if 80 % of the spikelet is yellow if facility for threshing is available.</li> <li>- Stacking of the harvested panicles for 5-7 days by turning the heap of panicles can help in better threshing.</li> <li>- Clean and sundry seeds properly before storage.</li> </ul>
<b>Rapeseed/ mustard</b>	Sowing		- In foot-hills and upland area zero tillage rape seed may be sown just after harvesting rice to make full utilization of the residual moisture.
<b>Field pea</b>	Land preparation to sowing		Land preparation may be done. Sowing may be done at a spacing of 30cm X 10-15cm. Recommended variety - Rachana. Seed rate is 75 kg/ hectare.
<b>Potato</b>	Land preparation to sowing		Land preparation may be done. Sowing may be done at a spacing of 60cm X 15-20cm. Recommended varieties are Kufri Megha and Kufri Jyoti. Seed rate is 25-30 quintals/ hectare.
<b>Groundnut</b>	Maturity to harvesting		<ul style="list-style-type: none"> <li>- Harvest the crop when 125 days old (var. ICGS-76) as well as dark tint inside the shell appears.</li> <li>- Make bundles and dry the crops in shade for 2-3 days for curing. Dry it for 4-5 days before storage. Select good quality pods for seeds.</li> </ul>
<b>Blackgram</b>	Maturity & harvesting		Harvest mature crops.
<b>Storage</b>			- Clean and store the seeds in RC-seed bins after proper drying. Check for germination of seeds or fungal growth. After proper drying, store grains/ seeds in well ventilated godowns.
<b>Horticulture Crops</b>			
<b>* As there is likely to have no rain in next five days, sowing/ planting of crops may be done to utilize the soil moisture for germination and vegetative growth.</b>			
<b>Coriander</b>	Land preparation & sowing		<ul style="list-style-type: none"> <li>- Select area with bright sunshine rich in organic manure for preparing beds. Treat seeds using Dichlorovos @ 1ml/ litre of water 2-3 days before sowing to protect seeds from ant and other insects.</li> <li>- Apply 200-250 kg of FYM (@ ½ cart load) in 100 sq. m and mix thoroughly to improve soil fertility.</li> <li>- Seed rate is 100g seeds for 100 sq. m.</li> </ul>
<b>Spinach beet (Palak)</b>	Land preparation &		<ul style="list-style-type: none"> <li>- Well drained soil is best for growing spinach beet.</li> <li>- Avoid damping off by adding Captan 50WP in the soil of bed.</li> </ul>



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

	sowing		Spaning may be maintained at 10cm x 20-25cm apart. - Seed rate is 100-120g/ 100 sq m. If broadcasting is done, double the dose is required.
<b>Onion</b>	Nursery raising		- Apply Captan 50 WP during final land preparation as preventive measures against soil borne diseases. - For 25 sq m area we need ½ kg seeds. - check that the new sprouts/ young plants in nursery beds are not damaged. - 45-60 days old seedlings may be transplanted.
<b>Cabbage/ Cauliflower</b>	Nursery		- Sowing of late season crop may be done. - Care should be taken for the young seedlings below 20 days old.

**Animal Sciences**

Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
<b>Poultry</b>	All susceptible age groups (as and when required)	Coccidiosis	Give preventive dose of Coximar @ 1g/ lit of drinking water for 7 days along with glucose.
		Deworming	Use Ventrimizole @ 1g in 1 litre of drinking water for 3 days.
		Litter management	Open the hessian cloth hanging on the wall and cross ventilation must be provided. Stir the litter material regularly.
<b>Sheep/ Goat</b>	Above 3 months of age (as and when required)	Tetanus	In pregnant animals, inject Tetanus Toxoid @ 0.5ml at 3.5 months of pregnancy.
<b>Pig</b>	Above 3 months of age (as and when required)	Skin infection	Give Ivermectin ( Indectin tab.) @ 10mg/33kg body weight.
		Coccidial infection (Diarrhea)	Ofloxacin/ Ornidazole (O <sub>2</sub> ) tab twice daily for 5 days.
		Deworming	- Deworm using Piperazine citrate @ 30 ml per adult animal once.
		Ticks	Add Butox liquid @ 2 ml/ lit of water and use the solution twice a week while brushing the animal.
<b>Cattle / Buffalo/ Mithun</b>	Above 3 months of age (as and when required)	Fasciolosis	- Give Tolzan-F @30 ml/ 100 kg body weight.
		Deworming	Give Nilzan/ Tolzan- F @ 30ml/ 100 kg body weight.
		Ticks/ Wounds	Topicure Spray may be used twice daily on the affected parts.

**Fishery:**

Type of Farming	Stage & Management	Agro-metrological advisories
<b>Carp/ Catfish culture</b>	Harvesting and stock maintenance	- Fish harvesting can be started. Also stock fish for next year and for this practices mentioned below may be done in small scale. - Ponds should be free from unwanted vegetation and insects and water depth should be maintained at 1.0-1.5m. - Feeds should be provided 1-2 times a day @ 3 % body weight per day. - Well balanced feed fortified with vitamins and minerals should be provided. - Apply Green AQ @ 4litre/ ha or lime @ 200kg/ ha. Maintain good quality of pond water.
<b>Paddy cum fish culture</b>	Harvesting	- Fish harvesting can be started. - Maintained 1.0 m water depth of the trenches in paddy field. - Provide rice bran and mustard oil cake (1:1) as supplementary feed. Avoid insecticide/pesticide application.

(I. Meghachandra Singh)  
Scientist i/c, GKMS