



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Integrated Agromet Advisory Services”

MANIPUR BULLETIN

17th February, 2015

Imphal-East

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 18th to 22nd February, 2015 as below:

Significant past weather for the preceding bi-weekly of Imphal - West (13th to 16th February, 2015)

During past four days sky was mainly clear to partly cloudy but mainly cloudy during afternoon hours of 10th & 11th February. No rainfall was recorded during past four days. Maximum temperature range observed was 22.4 – 23.1 °C, range of minimum temperature observed was 4.5 – 6.0 °C, mean relative humidity observed in morning and afternoon were 80.0 % and 49.8 % respectively. The average wind speed range was 2.1 – 5.0 km/hr.

Weather forecast for next five days of Imphal- East (18th to 22nd February, 2015)

There may be rains from Thursday. Sky will be mainly clear to partly cloudy with maximum temp. of 23.0 – 24.0 °C & minimum temp. 5.0 – 7.0 °C. Relative humidity may reach upto 98 % (max RH) & 36 % (min RH). The average wind speed may be 2-5 km/ hr during next five days.

Name of the crop	Stage	Pest and disease infestation	Agro-meteorological advisories
Field Crops			
Pre-kharif Rice	Land Preparation for nursery		<ul style="list-style-type: none"> - Prepare nursery bed of 700 m² area, apply well decomposed FYM/ compost @ 1 t/ 700 m² at final leveling. Seed rate is 60-65kg for a hectare. - Soaking and incubation of seeds may be started before sowing when minimum temperature reaches 10-12 °C.
Rabi Maize	Vegetative to pre-flowering		<ul style="list-style-type: none"> - If the soil is dry, give light irrigation to maintain soil moisture status. - Weeding may be done.
		Stem borer	- To control the infestation of stem borer, apply few granules of Carbofuran 3G at maize whorl.
Rapeseed/ mustard	Siliqua formation to maturity	Pieris brassica	If incidence of Pieris brassica is observed (which bore the pods of rapeseeds of mustard) then spray Coragen @ 2ml/15 lit of water.
			Irrigation is essential at this stage for sustaining favorable condition for seed formation.
Pulses (Field pea/ Lentil)	Fruiting to maturity and harvesting		Well filled up green pods should be harvested. Retain well filled disease free mature pods for seeds.
Storage			<ul style="list-style-type: none"> - Dry, clean and store the seeds in RC-seed bins after proper drying of seeds and charcoal. Check for moisture content or weight of seeds. After proper drying, store grains/ seeds in well ventilated godowns.
Horticulture Crops			
Bhindi	Sowing		Land preparation for sowing bhindi may be done. Sowing of bhindi/ladies finger may be done at the spacing of 30cm x 45cm. Apply 25kg urea, 33kg DAP and 25kg MOP per sangam at final stage of land preparation before sowing. Recommended varieties are - Arka Anamika and Pusa Sawani
Tree bean/ Lime/ Lemon/ Khasi Mandarin/ Banana	Land preparation and collection of planting material		<ul style="list-style-type: none"> - Preparation of pits or field for planting tree bean/ lime/ lemon/ khasi mandarin/ banana may be started. - Collect good planting material.
Tomato	Transplanting		<ul style="list-style-type: none"> - 25 to 30 days old tomato seedlings may be transplanted in prepared land incorporated with well decomposed FYM at a spacing of 40-45cm x 50-60cm during evening. Apply Sevin dust @ 2.5 – 5.0 g/ hill at the time of transplanting to protect the seedlings from caterpillar/ cutworm. - A light irrigation should be given regularly for establishment.



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Integrated Agromet Advisory Services”

Fruit Crops			Life saving irrigation to passion fruit, pineapple and other fruit crops may be applied and mulch with dry grass at the basin to conserve the soil moisture.
Animal Sciences			
Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
Poultry	All susceptible age groups (as and when required)	Coccidiosis	Give Coximar @ 2g/lit of drinking water for 5-7 days.
		Ranikhet disease	Birds must be vaccinated as per recommended schedule mentioned below as the probability of occurrence of the disease is high during March- April months. 1. F-1 Strain may be first administered within 7 days of age and second dose may be given at 30days for all birds. 2. R ₂ B vaccine of 8-9 weeks of age group birds (layers). 3. Killed ND virus vaccine of 16-18 weeks of age group birds (layers).
		Gumboro	Vaccinate using one shot IBD vaccine against Gumboro.
		Farm premises management	Keep the farm premises neat and clean using quality disinfectants.
Sheep/ Goat	Above 3 months of age (as and when required)	Skin infections	- Use Butox liq. @ 2ml/ litres of water and brush the animal using the solution at 15 days interval.
		F.M.D.	- Vaccinate against F.M.D.
Pig	Above 3 months of age (as and when required)	Hog cholera	- Vaccinate against swine fever after the age of 3 months.
		Skin infection	Give Ivermectin (Indectin tab.) @ 200mg/ kg body weight once only.
		F.M.D.	- Vaccinate against F.M.D.
Cattle / Buffalo/ Mithun	Above 3 months of age (as and when required)	Skin infections	- Use Butox liq. @ 2ml/ litres of water and brush the animal using the solution at 15 days interval. * For Mithun, give Ivermectin @ 200 mg per kg body weight at the time of salt feeding.
		Black Quarter/ Haemorrhagic	Vaccinate against B.Q./ H.S. using proven vaccine of the diseases mentioned.
		F.M.D.	- Vaccinate against F.M.D.

(I. Meghachandra Singh)
Scientist i/c, GKMS



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Integrated Agromet Advisory Services”

MANIPUR BULLETIN

17th February, 2015

Imphal-West

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 18th to 22nd February, 2015 as below:

Significant past weather for the preceding bi-weekly of Imphal - West (13th to 16th February, 2015)

During past four days sky was mainly clear to partly cloudy but mainly cloudy during afternoon hours of 10th & 11th February. No rainfall was recorded during past four days. Maximum temperature range observed was 22.4 – 23.1 °C, range of minimum temperature observed was 4.5 – 6.0 °C, mean relative humidity observed in morning and afternoon were 80.0 % and 49.8 % respectively. The average wind speed range was 2.1 – 5.0 km/hr.

Weather forecast for next five days of Imphal- West (18th to 22nd February, 2015)

There may be rains from Thursday. Sky will be mainly clear to partly cloudy with maximum temp. of 24.0 – 25.0 °C & minimum temp. 7.0 – 8.0 °C. Relative humidity may reach upto 95 % (max RH) & 35 % (min RH). The average wind speed may be 2-5 km/ hr during next five days.

Name of the crop	Stage	Pest and disease infestation	Agro-meteorological advisories
Field Crops			
Pre-kharif Rice	Land Preparation for nursery		<ul style="list-style-type: none"> - Prepare nursery bed of 700 m² area, apply well decomposed FYM/ compost @ 1 t/ 700 m² at final leveling. Seed rate is 60-65kg for a hectare. - Soaking and incubation of seeds may be started before sowing when minimum temperature reaches 10-12 °C.
Rabi Maize	Vegetative to pre-flowering		<ul style="list-style-type: none"> - If the soil is dry, give light irrigation to maintain soil moisture status. - Weeding may be done.
		Stem borer	- To control the infestation of stem borer, apply few granules of Carbofuran 3G at maize whorl.
Rapeseed/ mustard	Silique formation to maturity	Pieris brassica	If incidence of Pieris brassica is observed (which bore the pods of rapeseeds of mustard) then spray Coragen @ 2ml/15 lit of water.
			Irrigation is essential at this stage for sustaining favorable condition for seed formation.
Pulses (Field pea/ Lentil)	Fruiting to maturity and harvesting		Well filled up green pods should be harvested. Retain well filled disease free mature pods for seeds.
Storage			<ul style="list-style-type: none"> - Dry, clean and store the seeds in RC-seed bins after proper drying of seeds and charcoal. Check for moisture content or weight of seeds. After proper drying, store grains/ seeds in well ventilated godowns.
Horticulture Crops			
Bhindi	Sowing		Land preparation for sowing bhindi may be done. Sowing of bhindi/ladies finger may be done at the spacing of 30cm x 45cm. Apply 25kg urea, 33kg DAP and 25kg MOP per sangam at final stage of land preparation before sowing. Recommended varieties are - Arka Anamika and Pusa Sawani
Tree bean/ Lime/ Lemon/ Khasi Mandarin/ Banana	Land preparation and collection of planting material		<ul style="list-style-type: none"> - Preparation of pits or field for planting tree bean/ lime/ lemon/ khasi mandarin/ banana may be started. - Collect good planting material.
Tomato	Transplanting		<ul style="list-style-type: none"> - 25 to 30 days old tomato seedlings may be transplanted in prepared land incorporated with well decomposed FYM at a spacing of 40-45cm x 50-60cm during evening. Apply Sevin dust @ 2.5 – 5.0 g/ hill at the time of transplanting to protect the seedlings from caterpillar/ cutworm. - A light irrigation should be given regularly for establishment.



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Integrated Agromet Advisory Services”

Fruit Crops			Life saving irrigation to passion fruit, pineapple and other fruit crops may be applied and mulch with dry grass at the basin to conserve the soil moisture.
Animal Sciences			
Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
Poultry	All susceptible age groups (as and when required)	Coccidiosis	Give Coximar @ 2g/lit of drinking water for 5-7 days.
		Ranikhet disease	Birds must be vaccinated as per recommended schedule mentioned below as the probability of occurrence of the disease is high during March- April months. 1. F-1 Strain may be first administered within 7 days of age and second dose may be given at 30days for all birds. 2. R ₂ B vaccine of 8-9 weeks of age group birds (layers). 3. Killed ND virus vaccine of 16-18 weeks of age group birds (layers).
		Gumboro	Vaccinate using one shot IBD vaccine against Gumboro.
		Farm premises management	Keep the farm premises neat and clean using quality disinfectants.
Sheep/ Goat	Above 3 months of age (as and when required)	Skin infections	- Use Butox liq. @ 2ml/ litres of water and brush the animal using the solution at 15 days interval.
		F.M.D.	- Vaccinate against F.M.D.
Pig	Above 3 months of age (as and when required)	Hog cholera	- Vaccinate against swine fever after the age of 3 months.
		Skin infection	Give Ivermectin (Indectin tab.) @ 200mg/ kg body weight once only.
		F.M.D.	- Vaccinate against F.M.D.
Cattle / Buffalo/ Mithun	Above 3 months of age (as and when required)	Skin infections	- Use Butox liq. @ 2ml/ litres of water and brush the animal using the solution at 15 days interval. * For Mithun, give Ivermectin @ 200 mg per kg body weight at the time of salt feeding.
		Black Quarter/ Haemorrhagic	Vaccinate against B.Q./ H.S. using proven vaccine of the diseases mentioned.
		F.M.D.	- Vaccinate against F.M.D.

(I. Meghachandra Singh)
Scientist i/c, GKMS



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Integrated Agromet Advisory Services”

MANIPUR BULLETIN

17th February, 2015

Senapati

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 18th to 22nd February, 2015 as below:

Significant past weather for the preceding bi-weekly of Imphal - West (13th to 16th February, 2015)

During past four days sky was mainly clear to partly cloudy but mainly cloudy during afternoon hours of 10th & 11th February. No rainfall was recorded during past four days. Maximum temperature range observed was 22.4 – 23.1 °C, range of minimum temperature observed was 4.5 – 6.0 °C, mean relative humidity observed in morning and afternoon were 80.0 % and 49.8 % respectively. The average wind speed range was 2.1 – 5.0 km/hr.

Weather forecast for next five days of Senapati (18th to 22nd February, 2015)

There may be rains from Thursday. Sky will be mainly clear to partly cloudy with maximum temp. of 23.0 – 24.0 °C & minimum temp. 4.0 – 6.0 °C. Relative humidity may reach upto 99 % (max RH) & 34 % (min RH). The average wind speed may be 2-4 km/ hr during next five days.

Name of the crop	Stage	Pest and disease infestation	Agro-meteorological advisories
Field Crops			
Pre-kharif Rice	Land Preparation for nursery		<ul style="list-style-type: none"> - Prepare nursery bed of 700 m² area, apply well decomposed FYM/ compost @ 1 t/ 700 m² at final leveling. Seed rate is 60-65kg for a hectare. - Soaking and incubation of seeds may be started before sowing when minimum temperature reaches 10-12 °C.
Rabi Maize	Vegetative to pre-flowering		<ul style="list-style-type: none"> - If the soil is dry, give light irrigation to maintain soil moisture status. - Weeding may be done.
		Stem borer	- To control the infestation of stem borer, apply few granules of Carbofuran 3G at maize whorl.
Rapeseed/ mustard	Silique formation to maturity	Pieris brassica	If incidence of Pieris brassica is observed (which bore the pods of rapeseeds of mustard) then spray Coragen @ 2ml/15 lit of water.
			Irrigation is essential at this stage for sustaining favorable condition for seed formation.
Pulses (Field pea/ Lentil)	Fruiting to maturity and harvesting		Well filled up green pods should be harvested. Retain well filled disease free mature pods for seeds.
Storage			<ul style="list-style-type: none"> - Dry, clean and store the seeds in RC-seed bins after proper drying of seeds and charcoal. Check for moisture content or weight of seeds. After proper drying, store grains/ seeds in well ventilated godowns.
Horticulture Crops			
Bhindi	Sowing		Land preparation for sowing bhindi may be done. Sowing of bhindi/ladies finger may be done at the spacing of 30cm x 45cm. Apply 25kg urea, 33kg DAP and 25kg MOP per sangam at final stage of land preparation before sowing. Recommended varieties are - Arka Anamika and Pusa Sawani
Tree bean/ Lime/ Lemon/ Khasi Mandarin/ Banana	Land preparation and collection of planting material		<ul style="list-style-type: none"> - Preparation of pits or field for planting tree bean/ lime/ lemon/ khasi mandarin/ banana may be started. - Collect good planting material.
Tomato	Transplanting		<ul style="list-style-type: none"> - 25 to 30 days old tomato seedlings may be transplanted in prepared land incorporated with well decomposed FYM at a spacing of 40-45cm x 50-60cm during evening. Apply Sevin dust @ 2.5 – 5.0 g/ hill at the time of transplanting to protect the seedlings from caterpillar/ cutworm. - A light irrigation should be given regularly for establishment.



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Integrated Agromet Advisory Services”

Fruit Crops			Life saving irrigation to passion fruit, pineapple and other fruit crops may be applied and mulch with dry grass at the basin to conserve the soil moisture.
Animal Sciences			
Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
Poultry	All susceptible age groups (as and when required)	Coccidiosis	Give Coximar @ 2g/lit of drinking water for 5-7 days.
		Ranikhet disease	Birds must be vaccinated as per recommended schedule mentioned below as the probability of occurrence of the disease is high during March- April months. 1. F-1 Strain may be first administered within 7 days of age and second dose may be given at 30days for all birds. 2. R ₂ B vaccine of 8-9 weeks of age group birds (layers). 3. Killed ND virus vaccine of 16-18 weeks of age group birds (layers).
		Gumboro	Vaccinate using one shot IBD vaccine against Gumboro.
		Farm premises management	Keep the farm premises neat and clean using quality disinfectants.
Sheep/ Goat	Above 3 months of age (as and when required)	Skin infections	- Use Butox liq. @ 2ml/ litres of water and brush the animal using the solution at 15 days interval.
		F.M.D.	- Vaccinate against F.M.D.
Pig	Above 3 months of age (as and when required)	Hog cholera	- Vaccinate against swine fever after the age of 3 months.
		Skin infection	Give Ivermectin (Indectin tab.) @ 200mg/ kg body weight once only.
		F.M.D.	- Vaccinate against F.M.D.
Cattle / Buffalo/ Mithun	Above 3 months of age (as and when required)	Skin infections	- Use Butox liq. @ 2ml/ litres of water and brush the animal using the solution at 15 days interval. * For Mithun, give Ivermectin @ 200 mg per kg body weight at the time of salt feeding.
		Black Quarter/ Haemorrhagic	Vaccinate against B.Q./ H.S. using proven vaccine of the diseases mentioned.
		F.M.D.	- Vaccinate against F.M.D.

(I. Meghachandra Singh)
Scientist i/c, GKMS



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Integrated Agromet Advisory Services”

MANIPUR BULLETIN

17th February, 2015

Tamenglong

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 18th to 22nd February, 2015 as below:

Significant past weather for the preceding bi-weekly of Imphal - West (13th to 16th February, 2015)

During past four days sky was mainly clear to partly cloudy but mainly cloudy during afternoon hours of 10th & 11th February. No rainfall was recorded during past four days. Maximum temperature range observed was 22.4 – 23.1 °C, range of minimum temperature observed was 4.5 – 6.0 °C, mean relative humidity observed in morning and afternoon were 80.0 % and 49.8 % respectively. The average wind speed range was 2.1 – 5.0 km/hr.

Weather forecast for next five days of Tamenglong (18th to 22nd February, 2015)

There may be rains from Thursday. Sky will be mainly clear to partly cloudy with maximum temp. of 24.0 – 25.0 °C & minimum temp. 7.0 – 8.0 °C. Relative humidity may reach upto 96 % (max RH) & 36 % (min RH). The average wind speed may be 3-5 km/ hr during next five days.

Name of the crop	Stage	Pest and disease infestation	Agro-meteorological advisories
Field Crops			
Pre-kharif Rice	Land Preparation for nursery		<ul style="list-style-type: none"> - Prepare nursery bed of 700 m² area, apply well decomposed FYM/ compost @ 1 t/ 700 m² at final leveling. Seed rate is 60-65kg for a hectare. - Soaking and incubation of seeds may be started before sowing when minimum temperature reaches 10-12 °C.
Rabi Maize	Vegetative to pre-flowering		<ul style="list-style-type: none"> - If the soil is dry, give light irrigation to maintain soil moisture status. - Weeding may be done.
		Stem borer	- To control the infestation of stem borer, apply few granules of Carbofuran 3G at maize whorl.
Rapeseed/ mustard	Siliqua formation to maturity	Pieris brassica	If incidence of Pieris brassica is observed (which bore the pods of rapeseeds of mustard) then spray Coragen @ 2ml/15 lit of water.
			Irrigation is essential at this stage for sustaining favorable condition for seed formation.
Pulses (Field pea/ Lentil)	Fruiting to maturity and harvesting		Well filled up green pods should be harvested. Retain well filled disease free mature pods for seeds.
Storage			<ul style="list-style-type: none"> - Dry, clean and store the seeds in RC-seed bins after proper drying of seeds and charcoal. Check for moisture content or weight of seeds. After proper drying, store grains/ seeds in well ventilated godowns.
Horticulture Crops			
Bhindi	Sowing		Land preparation for sowing bhindi may be done. Sowing of bhindi/ladies finger may be done at the spacing of 30cm x 45cm. Apply 25kg urea, 33kg DAP and 25kg MOP per sangam at final stage of land preparation before sowing. Recommended varieties are - Arka Anamika and Pusa Sawani
Tree bean/ Lime/ Lemon/ Khasi Mandarin/ Banana	Land preparation and collection of planting material		<ul style="list-style-type: none"> - Preparation of pits or field for planting tree bean/ lime/ lemon/ khasi mandarin/ banana may be started. - Collect good planting material.
Tomato	Transplanting		<ul style="list-style-type: none"> - 25 to 30 days old tomato seedlings may be transplanted in prepared land incorporated with well decomposed FYM at a spacing of 40-45cm x 50-60cm during evening. Apply Sevin dust @ 2.5 – 5.0 g/ hill at the time of transplanting to protect the seedlings from caterpillar/ cutworm. - A light irrigation should be given regularly for establishment.



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Integrated Agromet Advisory Services”

Fruit Crops			Life saving irrigation to passion fruit, pineapple and other fruit crops may be applied and mulch with dry grass at the basin to conserve the soil moisture.
Animal Sciences			
Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
Poultry	All susceptible age groups (as and when required)	Coccidiosis	Give Coximar @ 2g/lit of drinking water for 5-7 days.
		Ranikhet disease	Birds must be vaccinated as per recommended schedule mentioned below as the probability of occurrence of the disease is high during March- April months. 1. F-1 Strain may be first administered within 7 days of age and second dose may be given at 30days for all birds. 2. R ₂ B vaccine of 8-9 weeks of age group birds (layers). 3. Killed ND virus vaccine of 16-18 weeks of age group birds (layers).
		Gumboro	Vaccinate using one shot IBD vaccine against Gumboro.
		Farm premises management	Keep the farm premises neat and clean using quality disinfectants.
Sheep/ Goat	Above 3 months of age (as and when required)	Skin infections	- Use Butox liq. @ 2ml/ litres of water and brush the animal using the solution at 15 days interval.
		F.M.D.	- Vaccinate against F.M.D.
Pig	Above 3 months of age (as and when required)	Hog cholera	- Vaccinate against swine fever after the age of 3 months.
		Skin infection	Give Ivermectin (Indectin tab.) @ 200mg/ kg body weight once only.
		F.M.D.	- Vaccinate against F.M.D.
Cattle / Buffalo/ Mithun	Above 3 months of age (as and when required)	Skin infections	- Use Butox liq. @ 2ml/ litres of water and brush the animal using the solution at 15 days interval. * For Mithun, give Ivermectin @ 200 mg per kg body weight at the time of salt feeding.
		Black Quarter/ Haemorrhagic	Vaccinate against B.Q./ H.S. using proven vaccine of the diseases mentioned.
		F.M.D.	- Vaccinate against F.M.D.

(I. Meghachandra Singh)
Scientist i/c, GKMS



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Integrated Agromet Advisory Services”

MANIPUR BULLETIN

17th February, 2015

Thoubal

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 18th to 22nd February, 2015 as below:

Significant past weather for the preceding bi-weekly of Imphal - West (13th to 16th February, 2015)

During past four days sky was mainly clear to partly cloudy but mainly cloudy during afternoon hours of 10th & 11th February. No rainfall was recorded during past four days. Maximum temperature range observed was 22.4 – 23.1 °C, range of minimum temperature observed was 4.5 – 6.0 °C, mean relative humidity observed in morning and afternoon were 80.0 % and 49.8 % respectively. The average wind speed range was 2.1 – 5.0 km/hr.

Weather forecast for next five days of Thoubal (18th to 22nd February, 2015)

There may be rains from Thursday. Sky will be mainly clear to partly cloudy with maximum temp. of 22.0 – 24.0 °C & minimum temp. 6.0 – 9.0 °C. Relative humidity may reach upto 94 % (max RH) & 44 % (min RH). The average wind speed may be 4-6 km/ hr during next five days.

Name of the crop	Stage	Pest and disease infestation	Agro-meteorological advisories
Field Crops			
Pre-kharif Rice	Land Preparation for nursery		<ul style="list-style-type: none"> - Prepare nursery bed of 700 m² area, apply well decomposed FYM/ compost @ 1 t/ 700 m² at final leveling. Seed rate is 60-65kg for a hectare. - Soaking and incubation of seeds may be started before sowing when minimum temperature reaches 10-12 °C.
Rabi Maize	Vegetative to pre-flowering		<ul style="list-style-type: none"> - If the soil is dry, give light irrigation to maintain soil moisture status. - Weeding may be done.
		Stem borer	- To control the infestation of stem borer, apply few granules of Carbofuran 3G at maize whorl.
Rapeseed/ mustard	Siliqua formation to maturity	Pieris brassica	If incidence of Pieris brassica is observed (which bore the pods of rapeseeds of mustard) then spray Coragen @ 2ml/15 lit of water.
			Irrigation is essential at this stage for sustaining favorable condition for seed formation.
Pulses (Field pea/ Lentil)	Fruiting to maturity and harvesting		Well filled up green pods should be harvested. Retain well filled disease free mature pods for seeds.
Storage			<ul style="list-style-type: none"> - Dry, clean and store the seeds in RC-seed bins after proper drying of seeds and charcoal. Check for moisture content or weight of seeds. After proper drying, store grains/ seeds in well ventilated godowns.
Horticulture Crops			
Bhindi	Sowing		Land preparation for sowing bhindi may be done. Sowing of bhindi/ladies finger may be done at the spacing of 30cm x 45cm. Apply 25kg urea, 33kg DAP and 25kg MOP per sangam at final stage of land preparation before sowing. Recommended varieties are - Arka Anamika and Pusa Sawani
Tree bean/ Lime/ Lemon/ Khasi Mandarin/ Banana	Land preparation and collection of planting material		<ul style="list-style-type: none"> - Preparation of pits or field for planting tree bean/ lime/ lemon/ khasi mandarin/ banana may be started. - Collect good planting material.
Tomato	Transplanting		<ul style="list-style-type: none"> - 25 to 30 days old tomato seedlings may be transplanted in prepared land incorporated with well decomposed FYM at a spacing of 40-45cm x 50-60cm during evening. Apply Sevin dust @ 2.5 – 5.0 g/ hill at the time of transplanting to protect the seedlings from caterpillar/ cutworm. - A light irrigation should be given regularly for establishment.



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Integrated Agromet Advisory Services”

Fruit Crops			Life saving irrigation to passion fruit, pineapple and other fruit crops may be applied and mulch with dry grass at the basin to conserve the soil moisture.
Animal Sciences			
Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
Poultry	All susceptible age groups (as and when required)	Coccidiosis	Give Coximar @ 2g/lit of drinking water for 5-7 days.
		Ranikhet disease	Birds must be vaccinated as per recommended schedule mentioned below as the probability of occurrence of the disease is high during March- April months. 1. F-1 Strain may be first administered within 7 days of age and second dose may be given at 30days for all birds. 2. R ₂ B vaccine of 8-9 weeks of age group birds (layers). 3. Killed ND virus vaccine of 16-18 weeks of age group birds (layers).
		Gumboro	Vaccinate using one shot IBD vaccine against Gumboro.
		Farm premises management	Keep the farm premises neat and clean using quality disinfectants.
Sheep/ Goat	Above 3 months of age (as and when required)	Skin infections	- Use Butox liq. @ 2ml/ litres of water and brush the animal using the solution at 15 days interval.
		F.M.D.	- Vaccinate against F.M.D.
Pig	Above 3 months of age (as and when required)	Hog cholera	- Vaccinate against swine fever after the age of 3 months.
		Skin infection	Give Ivermectin (Indectin tab.) @ 200mg/ kg body weight once only.
		F.M.D.	- Vaccinate against F.M.D.
Cattle / Buffalo/ Mithun	Above 3 months of age (as and when required)	Skin infections	- Use Butox liq. @ 2ml/ litres of water and brush the animal using the solution at 15 days interval. * For Mithun, give Ivermectin @ 200 mg per kg body weight at the time of salt feeding.
		Black Quarter/ Haemorrhagic	Vaccinate against B.Q./ H.S. using proven vaccine of the diseases mentioned.
		F.M.D.	- Vaccinate against F.M.D.

(I. Meghachandra Singh)
Scientist i/c, GKMS



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Integrated Agromet Advisory Services”

MANIPUR BULLETIN

17th February, 2015

Ukhrul

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 18th to 22nd February, 2015 as below:

Significant past weather for the preceding bi-weekly of Imphal - West (13th to 16th February, 2015)

During past four days sky was mainly clear to partly cloudy but mainly cloudy during afternoon hours of 10th & 11th February. No rainfall was recorded during past four days. Maximum temperature range observed was 22.4 – 23.1 °C, range of minimum temperature observed was 4.5 – 6.0 °C, mean relative humidity observed in morning and afternoon were 80.0 % and 49.8 % respectively. The average wind speed range was 2.1 – 5.0 km/hr.

Weather forecast for next five days of Ukhrul (18th to 22nd February, 2015)

There may be rains from Thursday. Sky will be mainly clear to partly cloudy with maximum temp. of 23.0 – 24.0 °C & minimum temp. 4.0 – 6.0 °C. Relative humidity may reach upto 93 % (max RH) & 38 % (min RH). The average wind speed may be 3-5 km/ hr during next five days.

Name of the crop	Stage	Pest and disease infestation	Agro-meteorological advisories
Field Crops			
Pre-kharif Rice	Land Preparation for nursery		<ul style="list-style-type: none"> - Prepare nursery bed of 700 m² area, apply well decomposed FYM/ compost @ 1 t/ 700 m² at final leveling. Seed rate is 60-65kg for a hectare. - Soaking and incubation of seeds may be started before sowing when minimum temperature reaches 10-12 °C.
Rabi Maize	Vegetative to pre-flowering		<ul style="list-style-type: none"> - If the soil is dry, give light irrigation to maintain soil moisture status. - Weeding may be done.
		Stem borer	- To control the infestation of stem borer, apply few granules of Carbofuran 3G at maize whorl.
Rapeseed/ mustard	Siliqua formation to maturity	Pieris brassica	If incidence of Pieris brassica is observed (which bore the pods of rapeseeds of mustard) then spray Coragen @ 2ml/15 lit of water.
			Irrigation is essential at this stage for sustaining favorable condition for seed formation.
Pulses (Field pea/ Lentil)	Fruiting to maturity and harvesting		Well filled up green pods should be harvested. Retain well filled disease free mature pods for seeds.
Storage			<ul style="list-style-type: none"> - Dry, clean and store the seeds in RC-seed bins after proper drying of seeds and charcoal. Check for moisture content or weight of seeds. After proper drying, store grains/ seeds in well ventilated godowns.
Horticulture Crops			
Bhindi	Sowing		Land preparation for sowing bhindi may be done. Sowing of bhindi/ladies finger may be done at the spacing of 30cm x 45cm. Apply 25kg urea, 33kg DAP and 25kg MOP per sangam at final stage of land preparation before sowing. Recommended varieties are - Arka Anamika and Pusa Sawani
Tree bean/ Lime/ Lemon/ Khasi Mandarin/ Banana	Land preparation and collection of planting material		<ul style="list-style-type: none"> - Preparation of pits or field for planting tree bean/ lime/ lemon/ khasi mandarin/ banana may be started. - Collect good planting material.
Tomato	Transplanting		<ul style="list-style-type: none"> - 25 to 30 days old tomato seedlings may be transplanted in prepared land incorporated with well decomposed FYM at a spacing of 40-45cm x 50-60cm during evening. Apply Sevin dust @ 2.5 – 5.0 g/ hill at the time of transplanting to protect the seedlings from caterpillar/ cutworm. - A light irrigation should be given regularly for establishment.



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Integrated Agromet Advisory Services”

Fruit Crops			Life saving irrigation to passion fruit, pineapple and other fruit crops may be applied and mulch with dry grass at the basin to conserve the soil moisture.
Animal Sciences			
Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
Poultry	All susceptible age groups (as and when required)	Coccidiosis	Give Coximar @ 2g/lit of drinking water for 5-7 days.
		Ranikhet disease	Birds must be vaccinated as per recommended schedule mentioned below as the probability of occurrence of the disease is high during March- April months. 1. F-1 Strain may be first administered within 7 days of age and second dose may be given at 30days for all birds. 2. R ₂ B vaccine of 8-9 weeks of age group birds (layers). 3. Killed ND virus vaccine of 16-18 weeks of age group birds (layers).
		Gumboro	Vaccinate using one shot IBD vaccine against Gumboro.
		Farm premises management	Keep the farm premises neat and clean using quality disinfectants.
Sheep/ Goat	Above 3 months of age (as and when required)	Skin infections	- Use Butox liq. @ 2ml/ litres of water and brush the animal using the solution at 15 days interval.
		F.M.D.	- Vaccinate against F.M.D.
Pig	Above 3 months of age (as and when required)	Hog cholera	- Vaccinate against swine fever after the age of 3 months.
		Skin infection	Give Ivermectin (Indectin tab.) @ 200mg/ kg body weight once only.
		F.M.D.	- Vaccinate against F.M.D.
Cattle / Buffalo/ Mithun	Above 3 months of age (as and when required)	Skin infections	- Use Butox liq. @ 2ml/ litres of water and brush the animal using the solution at 15 days interval. * For Mithun, give Ivermectin @ 200 mg per kg body weight at the time of salt feeding.
		Black Quarter/ Haemorrhagic	Vaccinate against B.Q./ H.S. using proven vaccine of the diseases mentioned.
		F.M.D.	- Vaccinate against F.M.D.

(I. Meghachandra Singh)
Scientist i/c, GKMS



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Integrated Agromet Advisory Services”

MANIPUR BULLETIN

17th February, 2015

Bishnupur

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 18th to 22nd February, 2015 as below:

Significant past weather for the preceding bi-weekly of Imphal - West (13th to 16th February, 2015)

During past four days sky was mainly clear to partly cloudy but mainly cloudy during afternoon hours of 10th & 11th February. No rainfall was recorded during past four days. Maximum temperature range observed was 22.4 – 23.1 °C, range of minimum temperature observed was 4.5 – 6.0 °C, mean relative humidity observed in morning and afternoon were 80.0 % and 49.8 % respectively. The average wind speed range was 2.1 – 5.0 km/hr.

Weather forecast for next five days of Bishnupur (18th to 22nd February, 2015)

There may be rains from Thursday. Sky will be mainly clear to partly cloudy with maximum temp. of 24.0 – 25.0 °C & minimum temp. 7.0 – 10.0 °C. Relative humidity may reach upto 95 % (max RH) & 41 % (min RH). The average wind speed may be 4-6 km/ hr during next five days.

Name of the crop	Stage	Pest and disease infestation	Agro-meteorological advisories
Field Crops			
Pre-kharif Rice	Land Preparation for nursery		<ul style="list-style-type: none"> - Prepare nursery bed of 700 m² area, apply well decomposed FYM/ compost @ 1 t/ 700 m² at final leveling. Seed rate is 60-65kg for a hectare. - Soaking and incubation of seeds may be started before sowing when minimum temperature reaches 10-12 °C.
Rabi Maize	Vegetative to pre-flowering		<ul style="list-style-type: none"> - If the soil is dry, give light irrigation to maintain soil moisture status. - Weeding may be done.
		Stem borer	- To control the infestation of stem borer, apply few granules of Carbofuran 3G at maize whorl.
Rapeseed/ mustard	Siliquea formation to maturity	Pieris brassica	If incidence of Pieris brassica is observed (which bore the pods of rapeseeds of mustard) then spray Coragen @ 2ml/15 lit of water.
			Irrigation is essential at this stage for sustaining favorable condition for seed formation.
Pulses (Field pea/ Lentil)	Fruiting to maturity and harvesting		Well filled up green pods should be harvested. Retain well filled disease free mature pods for seeds.
Storage			<ul style="list-style-type: none"> - Dry, clean and store the seeds in RC-seed bins after proper drying of seeds and charcoal. Check for moisture content or weight of seeds. After proper drying, store grains/ seeds in well ventilated godowns.
Horticulture Crops			
Bhindi	Sowing		Land preparation for sowing bhindi may be done. Sowing of bhindi/ladies finger may be done at the spacing of 30cm x 45cm. Apply 25kg urea, 33kg DAP and 25kg MOP per sangam at final stage of land preparation before sowing. Recommended varieties are - Arka Anamika and Pusa Sawani
Tree bean/ Lime/ Lemon/ Khasi Mandarin/ Banana	Land preparation and collection of planting material		<ul style="list-style-type: none"> - Preparation of pits or field for planting tree bean/ lime/ lemon/ khasi mandarin/ banana may be started. - Collect good planting material.
Tomato	Transplanting		<ul style="list-style-type: none"> - 25 to 30 days old tomato seedlings may be transplanted in prepared land incorporated with well decomposed FYM at a spacing of 40-45cm x 50-60cm during evening. Apply Sevin dust @ 2.5 – 5.0 g/ hill at the time of transplanting to protect the seedlings from caterpillar/ cutworm. - A light irrigation should be given regularly for establishment.



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Integrated Agromet Advisory Services”

Fruit Crops			Life saving irrigation to passion fruit, pineapple and other fruit crops may be applied and mulch with dry grass at the basin to conserve the soil moisture.
Animal Sciences			
Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
Poultry	All susceptible age groups (as and when required)	Coccidiosis	Give Coximar @ 2g/lit of drinking water for 5-7 days.
		Ranikhet disease	Birds must be vaccinated as per recommended schedule mentioned below as the probability of occurrence of the disease is high during March- April months. 1. F-1 Strain may be first administered within 7 days of age and second dose may be given at 30days for all birds. 2. R ₂ B vaccine of 8-9 weeks of age group birds (layers). 3. Killed ND virus vaccine of 16-18 weeks of age group birds (layers).
		Gumboro	Vaccinate using one shot IBD vaccine against Gumboro.
		Farm premises management	Keep the farm premises neat and clean using quality disinfectants.
Sheep/ Goat	Above 3 months of age (as and when required)	Skin infections	- Use Butox liq. @ 2ml/ litres of water and brush the animal using the solution at 15 days interval.
		F.M.D.	- Vaccinate against F.M.D.
Pig	Above 3 months of age (as and when required)	Hog cholera	- Vaccinate against swine fever after the age of 3 months.
		Skin infection	Give Ivermectin (Indectin tab.) @ 200mg/ kg body weight once only.
		F.M.D.	- Vaccinate against F.M.D.
Cattle / Buffalo/ Mithun	Above 3 months of age (as and when required)	Skin infections	- Use Butox liq. @ 2ml/ litres of water and brush the animal using the solution at 15 days interval. * For Mithun, give Ivermectin @ 200 mg per kg body weight at the time of salt feeding.
		Black Quarter/ Haemorrhagic	Vaccinate against B.Q./ H.S. using proven vaccine of the diseases mentioned.
		F.M.D.	- Vaccinate against F.M.D.

(I. Meghachandra Singh)
Scientist i/c, GKMS



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Integrated Agromet Advisory Services”

MANIPUR BULLETIN

17th February, 2015

Chandel

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 18th to 22nd February, 2015 as below:

Significant past weather for the preceding bi-weekly of Imphal - West (13th to 16th February, 2015)

During past four days sky was mainly clear to partly cloudy but mainly cloudy during afternoon hours of 10th & 11th February. No rainfall was recorded during past four days. Maximum temperature range observed was 22.4 – 23.1 °C, range of minimum temperature observed was 4.5 – 6.0 °C, mean relative humidity observed in morning and afternoon were 80.0 % and 49.8 % respectively. The average wind speed range was 2.1 – 5.0 km/hr.

Weather forecast for next five days of Chandel (18th to 22nd February, 2015)

There may be rains from Thursday. Sky will be mainly clear to partly cloudy with maximum temp. of 21.0 – 25.0 °C & minimum temp. 5.0 – 9.0 °C. Relative humidity may reach upto 90 % (max RH) & 44 % (min RH). The average wind speed may be 5 km/ hr during next five days.

Name of the crop	Stage	Pest and disease infestation	Agro-meteorological advisories
Field Crops			
Pre-kharif Rice	Land Preparation for nursery		<ul style="list-style-type: none"> - Prepare nursery bed of 700 m² area, apply well decomposed FYM/ compost @ 1 t/ 700 m² at final leveling. Seed rate is 60-65kg for a hectare. - Soaking and incubation of seeds may be started before sowing when minimum temperature reaches 10-12 °C.
Rabi Maize	Vegetative to pre-flowering		<ul style="list-style-type: none"> - If the soil is dry, give light irrigation to maintain soil moisture status. - Weeding may be done.
		Stem borer	- To control the infestation of stem borer, apply few granules of Carbofuran 3G at maize whorl.
Rapeseed/ mustard	Silique formation to maturity	Pieris brassica	If incidence of Pieris brassica is observed (which bore the pods of rapeseeds of mustard) then spray Coragen @ 2ml/15 lit of water.
			Irrigation is essential at this stage for sustaining favorable condition for seed formation.
Pulses (Field pea/ Lentil)	Fruiting to maturity and harvesting		Well filled up green pods should be harvested. Retain well filled disease free mature pods for seeds.
Storage			<ul style="list-style-type: none"> - Dry, clean and store the seeds in RC-seed bins after proper drying of seeds and charcoal. Check for moisture content or weight of seeds. After proper drying, store grains/ seeds in well ventilated godowns.
Horticulture Crops			
Bhindi	Sowing		Land preparation for sowing bhindi may be done. Sowing of bhindi/ladies finger may be done at the spacing of 30cm x 45cm. Apply 25kg urea, 33kg DAP and 25kg MOP per sangam at final stage of land preparation before sowing. Recommended varieties are - Arka Anamika and Pusa Sawani
Tree bean/ Lime/ Lemon/ Khasi Mandarin/ Banana	Land preparation and collection of planting material		<ul style="list-style-type: none"> - Preparation of pits or field for planting tree bean/ lime/ lemon/ khasi mandarin/ banana may be started. - Collect good planting material.
Tomato	Transplanting		<ul style="list-style-type: none"> - 25 to 30 days old tomato seedlings may be transplanted in prepared land incorporated with well decomposed FYM at a spacing of 40-45cm x 50-60cm during evening. Apply Sevin dust @ 2.5 – 5.0 g/ hill at the time of transplanting to protect the seedlings from caterpillar/ cutworm. - A light irrigation should be given regularly for establishment.



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Integrated Agromet Advisory Services”

Fruit Crops			Life saving irrigation to passion fruit, pineapple and other fruit crops may be applied and mulch with dry grass at the basin to conserve the soil moisture.
Animal Sciences			
Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
Poultry	All susceptible age groups (as and when required)	Coccidiosis	Give Coximar @ 2g/lit of drinking water for 5-7 days.
		Ranikhet disease	Birds must be vaccinated as per recommended schedule mentioned below as the probability of occurrence of the disease is high during March- April months. 1. F-1 Strain may be first administered within 7 days of age and second dose may be given at 30days for all birds. 2. R ₂ B vaccine of 8-9 weeks of age group birds (layers). 3. Killed ND virus vaccine of 16-18 weeks of age group birds (layers).
		Gumboro	Vaccinate using one shot IBD vaccine against Gumboro.
		Farm premises management	Keep the farm premises neat and clean using quality disinfectants.
Sheep/ Goat	Above 3 months of age (as and when required)	Skin infections	- Use Butox liq. @ 2ml/ litres of water and brush the animal using the solution at 15 days interval.
		F.M.D.	- Vaccinate against F.M.D.
Pig	Above 3 months of age (as and when required)	Hog cholera	- Vaccinate against swine fever after the age of 3 months.
		Skin infection	Give Ivermectin (Indectin tab.) @ 200mg/ kg body weight once only.
		F.M.D.	- Vaccinate against F.M.D.
Cattle / Buffalo/ Mithun	Above 3 months of age (as and when required)	Skin infections	- Use Butox liq. @ 2ml/ litres of water and brush the animal using the solution at 15 days interval. * For Mithun, give Ivermectin @ 200 mg per kg body weight at the time of salt feeding.
		Black Quarter/ Haemorrhagic	Vaccinate against B.Q./ H.S. using proven vaccine of the diseases mentioned.
		F.M.D.	- Vaccinate against F.M.D.

(I. Meghachandra Singh)
Scientist i/c, GKMS



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Integrated Agromet Advisory Services”

MANIPUR BULLETIN

17th February, 2015

Churachandpur

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 18th to 22nd February, 2015 as below:

Significant past weather for the preceding bi-weekly of Imphal - West (13th to 16th February, 2015)

During past four days sky was mainly clear to partly cloudy but mainly cloudy during afternoon hours of 10th & 11th February. No rainfall was recorded during past four days. Maximum temperature range observed was 22.4 – 23.1 °C, range of minimum temperature observed was 4.5 – 6.0 °C, mean relative humidity observed in morning and afternoon were 80.0 % and 49.8 % respectively. The average wind speed range was 2.1 – 5.0 km/hr.

Weather forecast for next five days of Churachandpur (18th to 22nd February, 2015)

There may be rains from Thursday. Sky will be mainly clear to partly cloudy with maximum temp. of 25.0 – 26.0 °C & minimum temp. 7.0 – 11.0 °C. Relative humidity may reach upto 94 % (max RH) & 34 % (min RH). The average wind speed may be 4-6 km/ hr during next five days.

Name of the crop	Stage	Pest and disease infestation	Agro-meteorological advisories
Field Crops			
Pre-kharif Rice	Land Preparation for nursery		<ul style="list-style-type: none"> - Prepare nursery bed of 700 m² area, apply well decomposed FYM/ compost @ 1 t/ 700 m² at final leveling. Seed rate is 60-65kg for a hectare. - Soaking and incubation of seeds may be started before sowing when minimum temperature reaches 10-12 °C.
Rabi Maize	Vegetative to pre-flowering		<ul style="list-style-type: none"> - If the soil is dry, give light irrigation to maintain soil moisture status. - Weeding may be done.
		Stem borer	- To control the infestation of stem borer, apply few granules of Carbofuran 3G at maize whorl.
Rapeseed/ mustard	Silique formation to maturity	Pieris brassica	If incidence of Pieris brassica is observed (which bore the pods of rapeseeds of mustard) then spray Coragen @ 2ml/15 lit of water.
			Irrigation is essential at this stage for sustaining favorable condition for seed formation.
Pulses (Field pea/ Lentil)	Fruiting to maturity and harvesting		Well filled up green pods should be harvested. Retain well filled disease free mature pods for seeds.
Storage			<ul style="list-style-type: none"> - Dry, clean and store the seeds in RC-seed bins after proper drying of seeds and charcoal. Check for moisture content or weight of seeds. After proper drying, store grains/ seeds in well ventilated godowns.
Horticulture Crops			
Bhindi	Sowing		Land preparation for sowing bhindi may be done. Sowing of bhindi/ladies finger may be done at the spacing of 30cm x 45cm. Apply 25kg urea, 33kg DAP and 25kg MOP per sangam at final stage of land preparation before sowing. Recommended varieties are - Arka Anamika and Pusa Sawani
Tree bean/ Lime/ Lemon/ Khasi Mandarin/ Banana	Land preparation and collection of planting material		<ul style="list-style-type: none"> - Preparation of pits or field for planting tree bean/ lime/ lemon/ khasi mandarin/ banana may be started. - Collect good planting material.
Tomato	Transplanting		<ul style="list-style-type: none"> - 25 to 30 days old tomato seedlings may be transplanted in prepared land incorporated with well decomposed FYM at a spacing of 40-45cm x 50-60cm during evening. Apply Sevin dust @ 2.5 – 5.0 g/ hill at the time of transplanting to protect the seedlings from caterpillar/ cutworm. - A light irrigation should be given regularly for establishment.



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Integrated Agromet Advisory Services”

Fruit Crops			Life saving irrigation to passion fruit, pineapple and other fruit crops may be applied and mulch with dry grass at the basin to conserve the soil moisture.
Animal Sciences			
Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
Poultry	All susceptible age groups (as and when required)	Coccidiosis	Give Coximar @ 2g/lit of drinking water for 5-7 days.
		Ranikhet disease	Birds must be vaccinated as per recommended schedule mentioned below as the probability of occurrence of the disease is high during March- April months. 1. F-1 Strain may be first administered within 7 days of age and second dose may be given at 30days for all birds. 2. R ₂ B vaccine of 8-9 weeks of age group birds (layers). 3. Killed ND virus vaccine of 16-18 weeks of age group birds (layers).
		Gumboro	Vaccinate using one shot IBD vaccine against Gumboro.
		Farm premises management	Keep the farm premises neat and clean using quality disinfectants.
Sheep/ Goat	Above 3 months of age (as and when required)	Skin infections	- Use Butox liq. @ 2ml/ litres of water and brush the animal using the solution at 15 days interval.
		F.M.D.	- Vaccinate against F.M.D.
Pig	Above 3 months of age (as and when required)	Hog cholera	- Vaccinate against swine fever after the age of 3 months.
		Skin infection	Give Ivermectin (Indectin tab.) @ 200mg/ kg body weight once only.
		F.M.D.	- Vaccinate against F.M.D.
Cattle / Buffalo/ Mithun	Above 3 months of age (as and when required)	Skin infections	- Use Butox liq. @ 2ml/ litres of water and brush the animal using the solution at 15 days interval. * For Mithun, give Ivermectin @ 200 mg per kg body weight at the time of salt feeding.
		Black Quarter/ Haemorrhagic	Vaccinate against B.Q./ H.S. using proven vaccine of the diseases mentioned.
		F.M.D.	- Vaccinate against F.M.D.

(I. Meghachandra Singh)
Scientist i/c, GKMS