



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Gramin Krishi Mausam Seva”

MANIPUR BULLETIN

22nd January, 2016

Imphal-West

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 23rd to 27th January, 2016 as below:

Significant past weather for the preceding bi-weekly of Imphal - West (19th to 21st January, 2016)

During last three days the sky was mainly clear to partly cloudy. 0.2 mm rainfall was recorded on 21st Jan. 2016, with maximum temperature range observed was 22.0 – 24.6°C, range of minimum temperature observed was 2.9 –9.0 °C, mean relative humidity observed in morning and afternoon were 93.67% and 51.67% respectively. The average wind speed range was 2.6 – 3.4 km/hr.

Weather forecast for next five days of Imphal-West (23rd to 27th January, 2016)

There may be no rain on Thursday. The Sky will remain mainly clear to partly foggy during morning hours, with maximum temp. of 22.0 – 23.0 °C & minimum temp. 5.0 – 6.0 °C. Relative humidity may reach upto 99 % (max RH) & 28 % (min RH). The average wind speed may be 3-4 km/ hr during next five days. The Wind direction may be north easterly to south south-westerly during next five days.

Weather Forecast for next 5 days * upto 0830 hours IST of 27th January

Moderate to dense fog very likely at isolated places over Manipur during next 2 days

Weather Outlook for subsequent 2 days from 27th to 29th January 2016

Rain/thundershowers is likely at isolated places over northeast India

Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Field Crops			
Rabi Maize	Early vegetative		- Light irrigation may be given to keep soil moist if the soil is dry. - Thinning may be done in vegetative stage for mustard or pea intercropped with maize.
Rapeseed/ Mustard	Vegetative to flowering		- Apply foliar application of urea {5 pumps/ sangam @ 10g/ pump} or top dress with ½ N (@ 66kg Urea/ ha) just after irrigation. - If flowering has started, then four bee boxes may be kept for one hectare crop area. Avoid spray of insecticides in such area.
		Aphid	- Spray Monocrotophos @ 1.5 ml/lit of water to the infected patches only or use yellow plates filled with water.
Field Pea/ Lentil	Vegetative	Powdery Mildew	- Spray Bayleton (Triaclemefon 25% WP) @ 7.5 g/ pump or Dinocap (Karathane 46% EC) @ 7.5ml/ pump.
		Aphid	- Spray Cypermethrin @2ml/lit of water or Nuvan @ 0.5ml/ lit of water against aphid. - For organic control against aphid, sprays that consist of Azadirachtin/ neem oil or pyrethrin may be used.
Storage (seed)			- Clean and store the seeds in RC-seed bins after proper drying of the desiccant charcoal. Check for moisture content or weight of seeds. For grains, store in well ventilated godowns with regular check for moisture.
Horticulture Crops			
Turmeric	Harvesting		-Already mature plants whose leaves are more than 70 % dried may be harvested. -Harvested rhizome should be spread in the shed for at least 4- 5 days and screen for seed purposes. -Curing process should be started within one week of harvest.
French bean			-land preparation for sowing of French bean may be started.
Potato	Vegetative		- Spray Mencozeb 2gm/litre of water at least twice at an interval of 7-10 days to prevent spreading of early blight as the temperature rises. - If aphid is observed in small patches, wash off it with plain water but if it is severe, spray Cypermethrin 20 E.C. @ 1ml/ litre of water or Nuvan @ 0.5ml/litre of water. - Give light irrigation to attain optimum soil moisture.
Cucurbits	Nursery		- Nurseries of bitter gourd, cucumber, sponge gourd, bottle gourd, pumpkin, etc. may be done under polyhouse.
Tomato	Transplanting		- Proper field preparation for transplanting of 4 leaf stage should be done. -Due consideration for removal of potentially infected material such as



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Gramin Krishi Mausam Seva”

			vines and stubbles for early blight should be done while ploughing.
	Flowering		- prophylactic measures against tomato fruit borer should be done.
Animal Sciences			
Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
Poultry (Chicken/ Turkey/ Duck)	All age groups	Care & management	- To protect birds of all age groups from inclement weather e.g. frost, snowfall etc., there should be adequate arrangements for artificial lighting and heating. - Needed to increase energy giving ingredients of feed (i.e. Maize, Wheat, Bajra etc.) and can be added 5% more.
		Coccidiosis	- After routine preventive and prophylaxis medication for the said disease, if still there is any defaecation of reddish colour faeces, somnolence, drooping of wings, offed etc., Amprolium agent can be used in feed and water. Add vitamin k, vit. A and electrolytes etc. also add mineral mixture in the feeds and water.
Sheep/ Goat	All age groups excepts the young ones	PPR (Peste Des Petits Ruminants)	- If the animals have not yet been vaccinated, give vaccine now for PPR (Peste Des Petits Ruminants).
		Enterotoxemia	- If the Lambs and Kids in particular should be given the vaccination to prevent Enterotoxaemia.
		Ectoparasitism	- Keep the sheds clean. Hang Bouquets of Basil (Ocimum sanctum) or Lemon Grass (Cymbopogon citrates) in the animals sheds to keep away the ecto-parasites. Neem oil based disinfectants can be sprayed. Butox solution (2 C.C. in 1 litre of water) can also be used.
Pig	All age groups excepts the young ones	Care & management for newly born one	-Artificial lighting and heating to the piglets. -Provide adequate bedding materials.
		Deworming	-Give Morantel citrate tablet @ 15 mg/kg body weight.
		Ectoparasitism	- Keep the sheds clean. Hang Bouquets of Basil (Ocimum sanctum) or Lemon Grass (Cymbopogon citrates) in the animals sheds to keep away the ecto-parasites. Neem oil based disinfectants can be sprayed. Butox solution (2 C.C. in 1 litre of water) can also be used.
Cattle	All age groups excepts the young ones	B.Q./ H.S/FMD	-If the animals have not yet been vaccinated, give vaccine, now for Black Quarter, Haemorrhagic septicaemia, Foot & mouth Disease.
		Ectoparasitism	- Keep the sheds clean. Hang Bouquets of Basil (Ocimum sanctum) or Lemon Grass (Cymbopogon citrates) in the animals sheds to keep away the ecto-parasites. Neem oil based disinfectants can be sprayed. Butox solution (2 C.C. in 1 litre of water) can also be used.
		Fodder cultivation	-Irrigate the crops like Berseem Clover every 20-30 days. -At every 20-22 days for Oat.
Fishery:			
Type of Farming	Stage	Management	Agro-metrological advisories
Carp culture	Fries/ fingerlings/ brooders	Pond and water	Remove aquatic weeds. Maintain minimum depth of 1.5 – 2.0 m. Apply Agri-Lime in brooder pond at small quantity at an interval of 30 days regularly.
		Feed	Provide feed 1-2 times @ 2 % body weight per day.
		Fish	- Male and female brooders should be stock in separate ponds for early breeding programs. - harvesting of fish may be done in area where water availability is low.

(I. Meghachandra Singh)
Scientist i/c, IAAS



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Gramin Krishi Mausam Seva”

MANIPUR BULLETIN

22nd January, 2016

Senapati

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 23rd to 27th January, 2016 as below:

<p>Significant past weather for the preceding bi-weekly of Imphal - West (19th to 21st January, 2016)</p> <p>During last three days the sky was mainly clear to partly cloudy. 0.2 mm rainfall was recorded on 21st Jan. 2016, with maximum temperature range observed was 22.0 – 24.6°C, range of minimum temperature observed was 2.9 –9.0 °C, mean relative humidity observed in morning and afternoon were 93.67% and 51.67% respectively. The average wind speed range was 2.6 – 3.4 km/hr.</p>	<p>Weather forecast for next five days of Senapati (23rd to 27th January, 2016)</p> <p>There may be no rain on Thursday. The Sky will remain mainly clear to partly foggy during morning hours, with maximum temp. of 18.0 – 20.0 °C & minimum temp.4.0 – 5.0 °C. Relative humidity may reach upto 97 % (max RH) & 24 % (min RH). The average wind speed may be 3-4 km/ hr during next five days. The Wind direction may be north east easterly to north west westerly during next five days.</p>
--	--

Weather Forecast for next 5 days *upto 0830 hours IST of 27th January

Moderate to dense fog very likely at isolated places over Manipur during next 2 days

Weather Outlook for subsequent 2 days from 27th to 29th January 2016

Rain/thundershowers is likely at isolated places over northeast India

Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Field Crops			
Rabi Maize	Early vegetative		- Light irrigation may be given to keep soil moist if the soil is dry. - Thinning may be done in vegetative stage for mustard or pea intercropped with maize.
Rapeseed/ Mustard	Vegetative to flowering		- Apply foliar application of urea {5 pumps/ sangam @ 10g/ pump} or top dress with ½ N (@ 66kg Urea/ ha) just after irrigation. - If flowering has started, then four bee boxes may be kept for one hectare crop area. Avoid spray of insecticides in such area.
		Aphid	- Spray Monocrotophos @ 1.5 ml/lit of water to the infected patches only or use yellow plates filled with water.
Field Pea/ Lentil	Vegetative	Powdery Mildew	- Spray Bayleton (Triaclemefon 25% WP) @ 7.5 g/ pump or Dinocap (Karathane 46% EC) @ 7.5ml/ pump.
		Aphid	- Spray Cypermethrin @2ml/lit of water or Nuvan @ 0.5ml/ lit of water against aphid. - For organic control against aphid, sprays that consist of Azadirachtin/ neem oil or pyrethrin may be used.
Storage (seed)			- Clean and store the seeds in RC-seed bins after proper drying of the desiccant charcoal. Check for moisture content or weight of seeds. For grains, store in well ventilated godowns with regular check for moisture.
Horticulture Crops			
Turmeric	Harvesting		-Already mature plants whose leaves are more than 70 % dried may be harvested. -Harvested rhizome should be spread in the shed for at least 4- 5 days and screen for seed purposes. -Curing process should be started within one week of harvest.
French bean			-land preparation for sowing of French bean may be started.
Potato	Vegetative		- Spray Mencozeb 2gm/litre of water at least twice at an interval of 7-10 days to prevent spreading of early blight as the temperature rises. - If aphid is observed in small patches, wash off it with plain water but if it is severe, spray Cypermethrin 20 E.C. @ 1ml/ litre of water or Nuvan @ 0.5ml/litre of water. - Give light irrigation to attain optimum soil moisture.
Cucurbits	Nursery		- Nurseries of bitter gourd, cucumber, sponge gourd, bottle gourd, pumpkin, etc. may be done under polyhouse.
Tomato	Transplanting		- Proper field preparation for transplanting of 4 leaf stage should be done. -Due consideration for removal of potentially infected material such as



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Gramin Krishi Mausam Seva”

			vines and stubbles for early blight should be done while ploughing.
	Flowering		- prophylactic measures against tomato fruit borer should be done.
Animal Sciences			
Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
Poultry (Chicken/ Turkey/ Duck)	All age groups	Care & management	- To protect birds of all age groups from inclement weather e.g. frost, snowfall etc., there should be adequate arrangements for artificial lighting and heating. - Needed to increase energy giving ingredients of feed (i.e. Maize, Wheat, Bajra etc.) and can be added 5% more.
		Coccidiosis	- After routine preventive and prophylaxis medication for the said disease, if still there is any defaecation of reddish colour faeces, somnolence, drooping of wings, offed etc., Amprolium agent can be used in feed and water. Add vitamin k, vit. A and electrolytes etc. also add mineral mixture in the feeds and water.
Sheep/ Goat	All age groups excepts the young ones	PPR (Peste Des Petits Ruminants)	- If the animals have not yet been vaccinated, give vaccine now for PPR (Peste Des Petits Ruminants).
		Enterotoxemia	- If the Lambs and Kids in particular should be given the vaccination to prevent Enterotoxaemia.
		Ectoparasitism	- Keep the sheds clean. Hang Bouquets of Basil (Ocimum sanctum) or Lemon Grass (Cymbopogon citrates) in the animals sheds to keep away the ecto-parasites. Neem oil based disinfectants can be sprayed. Butox solution (2 C.C. in 1 litre of water) can also be used.
Pig	All age groups excepts the young ones	Care & management for newly born one	-Artificial lighting and heating to the piglets. -Provide adequate bedding materials.
		Deworming	-Give Morantel citrate tablet @ 15 mg/kg body weight.
		Ectoparasitism	- Keep the sheds clean. Hang Bouquets of Basil (Ocimum sanctum) or Lemon Grass (Cymbopogon citrates) in the animals sheds to keep away the ecto-parasites. Neem oil based disinfectants can be sprayed. Butox solution (2 C.C. in 1 litre of water) can also be used.
Cattle	All age groups excepts the young ones	B.Q./ H.S/FMD	-If the animals have not yet been vaccinated, give vaccine, now for Black Quarter, Haemorrhagic septicaemia, Foot & mouth Disease.
		Ectoparasitism	- Keep the sheds clean. Hang Bouquets of Basil (Ocimum sanctum) or Lemon Grass (Cymbopogon citrates) in the animals sheds to keep away the ecto-parasites. Neem oil based disinfectants can be sprayed. Butox solution (2 C.C. in 1 litre of water) can also be used.
		Fodder cultivation	-Irrigate the crops like Berseem Clover every 20-30 days. -At every 20-22 days for Oat.
Fishery:			
Type of Farming	Stage	Management	Agro-metrological advisories
Carp culture	Fries/ fingerlings/ brooders	Pond and water	Remove aquatic weeds. Maintain minimum depth of 1.5 – 2.0 m. Apply Agri-Lime in brooder pond at small quantity at an interval of 30 days regularly.
		Feed	Provide feed 1-2 times @ 2 % body weight per day.
		Fish	- Male and female brooders should be stock in separate ponds for early breeding programs. - harvesting of fish may be done in area where water availability is low.

(I. Meghachandra Singh)
Scientist i/c, IAAS



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Gramin Krishi Mausam Seva”

MANIPUR BULLETIN

22nd January, 2016

Tamenglong

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 23rd to 27th January, 2016 as below:

<p>Significant past weather for the preceding bi-weekly of Imphal - West (19th to 21st January, 2016)</p> <p>During last three days the sky was mainly clear to partly cloudy. 0.2 mm rainfall was recorded on 21st Jan. 2016, with maximum temperature range observed was 22.0 – 24.6°C, range of minimum temperature observed was 2.9 –9.0 °C, mean relative humidity observed in morning and afternoon were 93.67% and 51.67% respectively. The average wind speed range was 2.6 – 3.4 km/hr.</p>	<p>Weather forecast for next five days of Tamenglong (23rd to 27th January, 2016)</p> <p>There may be no rain on Thursday. The Sky will remain mainly clear to partly foggy during morning hours, with maximum temp. of 19.0 – 21.0 °C & minimum temp.4.0 – 5.0 °C. Relative humidity may reach upto 98 % (max RH) & 27 % (min RH). The average wind speed may be 3-4 km/ hr during next five days. The Wind direction may be north east easterly to southerly during next five days.</p>
--	--

Weather Forecast for next 5 days *upto 0830 hours IST of 27th January

Moderate to dense fog very likely at isolated places over Manipur during next 2 days

Weather Outlook for subsequent 2 days from 27th to 29th January 2016

Rain/thundershowers is likely at isolated places over northeast India

Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Field Crops			
Rabi Maize	Early vegetative		- Light irrigation may be given to keep soil moist if the soil is dry. - Thinning may be done in vegetative stage for mustard or pea intercropped with maize.
Rapeseed/ Mustard	Vegetative to flowering		- Apply foliar application of urea {5 pumps/ sangam @ 10g/ pump} or top dress with ½ N (@ 66kg Urea/ ha) just after irrigation. - If flowering has started, then four bee boxes may be kept for one hectare crop area. Avoid spray of insecticides in such area.
		Aphid	- Spray Monocrotophos @ 1.5 ml/lit of water to the infected patches only or use yellow plates filled with water.
Field Pea/ Lentil	Vegetative	Powdery Mildew	- Spray Bayleton (Triaclemofon 25% WP) @ 7.5 g/ pump or Dinocap (Karathane 46% EC) @ 7.5ml/ pump.
		Aphid	- Spray Cypermethrin @2ml/lit of water or Nuvan @ 0.5ml/ lit of water against aphid. - For organic control against aphid, sprays that consist of Azadirachtin/ neem oil or pyrethrin may be used.
Storage (seed)			- Clean and store the seeds in RC-seed bins after proper drying of the desiccant charcoal. Check for moisture content or weight of seeds. For grains, store in well ventilated godowns with regular check for moisture.
Horticulture Crops			
Turmeric	Harvesting		-Already mature plants whose leaves are more than 70 % dried may be harvested. -Harvested rhizome should be spread in the shed for at least 4- 5 days and screen for seed purposes. -Curing process should be started within one week of harvest.
French bean			-land preparation for sowing of French bean may be started.
Potato	Vegetative		- Spray Mencozeb 2gm/litre of water at least twice at an interval of 7-10 days to prevent spreading of early blight as the temperature rises. - If aphid is observed in small patches, wash off it with plain water but if it is severe, spray Cypermethrin 20 E.C. @ 1ml/ litre of water or Nuvan @ 0.5ml/litre of water. - Give light irrigation to attain optimum soil moisture.
Cucurbits	Nursery		- Nurseries of bitter gourd, cucumber, sponge gourd, bottle gourd, pumpkin, etc. may be done under polyhouse.
Tomato	Transplanting		- Proper field preparation for transplanting of 4 leaf stage should be done. -Due consideration for removal of potentially infected material such as



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Gramin Krishi Mausam Seva”

			vines and stubbles for early blight should be done while ploughing.
	Flowering		- prophylactic measures against tomato fruit borer should be done.
Animal Sciences			
Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
Poultry (Chicken/ Turkey/ Duck)	All age groups	Care & management	- To protect birds of all age groups from inclement weather e.g. frost, snowfall etc., there should be adequate arrangements for artificial lighting and heating. - Needed to increase energy giving ingredients of feed (i.e. Maize, Wheat, Bajra etc.) and can be added 5% more.
		Coccidiosis	- After routine preventive and prophylaxis medication for the said disease, if still there is any defaecation of reddish colour faeces, somnolence, drooping of wings, offed etc., Amprolium agent can be used in feed and water. Add vitamin k, vit. A and electrolytes etc. also add mineral mixture in the feeds and water.
Sheep/ Goat	All age groups excepts the young ones	PPR (Peste Des Petits Ruminants)	- If the animals have not yet been vaccinated, give vaccine now for PPR (Peste Des Petits Ruminants).
		Enterotoxemia	- If the Lambs and Kids in particular should be given the vaccination to prevent Enterotoxaemia.
		Ectoparasitism	- Keep the sheds clean. Hang Bouquets of Basil (Ocimum sanctum) or Lemon Grass (Cymbopogon citrates) in the animals sheds to keep away the ecto-parasites. Neem oil based disinfectants can be sprayed. Butox solution (2 C.C. in 1 litre of water) can also be used.
Pig	All age groups excepts the young ones	Care & management for newly born one	-Artificial lighting and heating to the piglets. -Provide adequate bedding materials.
		Deworming	-Give Morantel citrate tablet @ 15 mg/kg body weight.
		Ectoparasitism	- Keep the sheds clean. Hang Bouquets of Basil (Ocimum sanctum) or Lemon Grass (Cymbopogon citrates) in the animals sheds to keep away the ecto-parasites. Neem oil based disinfectants can be sprayed. Butox solution (2 C.C. in 1 litre of water) can also be used.
Cattle	All age groups excepts the young ones	B.Q./ H.S/FMD	-If the animals have not yet been vaccinated, give vaccine, now for Black Quarter, Haemorrhagic septicaemia, Foot & mouth Disease.
		Ectoparasitism	- Keep the sheds clean. Hang Bouquets of Basil (Ocimum sanctum) or Lemon Grass (Cymbopogon citrates) in the animals sheds to keep away the ecto-parasites. Neem oil based disinfectants can be sprayed. Butox solution (2 C.C. in 1 litre of water) can also be used.
		Fodder cultivation	-Irrigate the crops like Berseem Clover every 20-30 days. -At every 20-22 days for Oat.
Fishery:			
Type of Farming	Stage	Management	Agro-metrological advisories
Carp culture	Fries/ fingerlings/ brooders	Pond and water	Remove aquatic weeds. Maintain minimum depth of 1.5 – 2.0 m. Apply Agri-Lime in brooder pond at small quantity at an interval of 30 days regularly.
		Feed	Provide feed 1-2 times @ 2 % body weight per day.
		Fish	- Male and female brooders should be stock in separate ponds for early breeding programs. - harvesting of fish may be done in area where water availability is low.

(I. Meghachandra Singh)
Scientist i/c, IAAS



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Gramin Krishi Mausam Seva”

MANIPUR BULLETIN

22nd January, 2016

Thoubal

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 23rd to 27th January, 2016 as below:

Significant past weather for the preceding bi-weekly of Imphal - West (19th to 21st January, 2016)

During last three days the sky was mainly clear to partly cloudy. 0.2 mm rainfall was recorded on 21st Jan. 2016, with maximum temperature range observed was 22.0 – 24.6°C, range of minimum temperature observed was 2.9 –9.0 °C, mean relative humidity observed in morning and afternoon were 93.67% and 51.67% respectively. The average wind speed range was 2.6 – 3.4 km/hr.

Weather forecast for next five days of Thoubal (23rd to 27th January, 2016)

There may be no rain on Thursday. The Sky will remain mainly clear to partly foggy during morning hours, with maximum temp. of 19.0 – 20.0 °C & minimum temp. 5.0 – 6.0 °C. Relative humidity may reach upto 98 % (max RH) & 27 % (min RH). The average wind speed may be 4-5 km/ hr during next five days. The Wind direction may be south south easterly to north west westerly during next five days.

Weather Forecast for next 5 days *upto 0830 hours IST of 27th January

Moderate to dense fog very likely at isolated places over Manipur during next 2 days

Weather Outlook for subsequent 2 days from 27th to 29th January 2016

Rain/thundershowers is likely at isolated places over northeast India

Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Field Crops			
Rabi Maize	Early vegetative		- Light irrigation may be given to keep soil moist if the soil is dry. - Thinning may be done in vegetative stage for mustard or pea intercropped with maize.
Rapeseed/ Mustard	Vegetative to flowering		- Apply foliar application of urea {5 pumps/ sangam @ 10g/ pump} or top dress with ½ N (@ 66kg Urea/ ha) just after irrigation. - If flowering has started, then four bee boxes may be kept for one hectare crop area. Avoid spray of insecticides in such area.
		Aphid	- Spray Monocrotophos @ 1.5 ml/lit of water to the infected patches only or use yellow plates filled with water.
Field Pea/ Lentil	Vegetative	Powdery Mildew	- Spray Bayleton (Triaclemofon 25% WP) @ 7.5 g/ pump or Dinocap (Karathane 46% EC) @ 7.5ml/ pump.
		Aphid	- Spray Cypermethrin @2ml/lit of water or Nuvan @ 0.5ml/ lit of water against aphid. - For organic control against aphid, sprays that consist of Azadirachtin/ neem oil or pyrethrin may be used.
Storage (seed)			- Clean and store the seeds in RC-seed bins after proper drying of the desiccant charcoal. Check for moisture content or weight of seeds. For grains, store in well ventilated godowns with regular check for moisture.
Horticulture Crops			
Turmeric	Harvesting		-Already mature plants whose leaves are more than 70 % dried may be harvested. -Harvested rhizome should be spread in the shed for at least 4- 5 days and screen for seed purposes. -Curing process should be started within one week of harvest.
French bean			-land preparation for sowing of French bean may be started.
Potato	Vegetative		- Spray Mencozeb 2gm/litre of water at least twice at an interval of 7-10 days to prevent spreading of early blight as the temperature rises. - If aphid is observed in small patches, wash off it with plain water but if it is severe, spray Cypermethrin 20 E.C. @ 1ml/ litre of water or Nuvan @ 0.5ml/litre of water. - Give light irrigation to attain optimum soil moisture.
Cucurbits	Nursery		- Nurseries of bitter gourd, cucumber, sponge gourd, bottle gourd, pumpkin, etc. may be done under polyhouse.
Tomato	Transplanting		- Proper field preparation for transplanting of 4 leaf stage should be done. -Due consideration for removal of potentially infected material such as



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Gramin Krishi Mausam Seva”

			vines and stubbles for early blight should be done while ploughing.
	Flowering		- prophylactic measures against tomato fruit borer should be done.
Animal Sciences			
Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
Poultry (Chicken/ Turkey/ Duck)	All age groups	Care & management	- To protect birds of all age groups from inclement weather e.g. frost, snowfall etc., there should be adequate arrangements for artificial lighting and heating. - Needed to increase energy giving ingredients of feed (i.e. Maize, Wheat, Bajra etc.) and can be added 5% more.
		Coccidiosis	- After routine preventive and prophylaxis medication for the said disease, if still there is any defaecation of reddish colour faeces, somnolence, drooping of wings, offed etc., Amprolium agent can be used in feed and water. Add vitamin k, vit. A and electrolytes etc. also add mineral mixture in the feeds and water.
Sheep/ Goat	All age groups excepts the young ones	PPR (Peste Des Petits Ruminants)	- If the animals have not yet been vaccinated, give vaccine now for PPR (Peste Des Petits Ruminants).
		Enterotoxemia	- If the Lambs and Kids in particular should be given the vaccination to prevent Enterotoxaemia.
		Ectoparasitism	- Keep the sheds clean. Hang Bouquets of Basil (Ocimum sanctum) or Lemon Grass (Cymbopogon citrates) in the animals sheds to keep away the ecto-parasites. Neem oil based disinfectants can be sprayed. Butox solution (2 C.C. in 1 litre of water) can also be used.
Pig	All age groups excepts the young ones	Care & management for newly born one	-Artificial lighting and heating to the piglets. -Provide adequate bedding materials.
		Deworming	-Give Morantel citrate tablet @ 15 mg/kg body weight.
		Ectoparasitism	- Keep the sheds clean. Hang Bouquets of Basil (Ocimum sanctum) or Lemon Grass (Cymbopogon citrates) in the animals sheds to keep away the ecto-parasites. Neem oil based disinfectants can be sprayed. Butox solution (2 C.C. in 1 litre of water) can also be used.
Cattle	All age groups excepts the young ones	B.Q./ H.S/FMD	-If the animals have not yet been vaccinated, give vaccine, now for Black Quarter, Haemorrhagic septicaemia, Foot & mouth Disease.
		Ectoparasitism	- Keep the sheds clean. Hang Bouquets of Basil (Ocimum sanctum) or Lemon Grass (Cymbopogon citrates) in the animals sheds to keep away the ecto-parasites. Neem oil based disinfectants can be sprayed. Butox solution (2 C.C. in 1 litre of water) can also be used.
		Fodder cultivation	-Irrigate the crops like Berseem Clover every 20-30 days. -At every 20-22 days for Oat.
Fishery:			
Type of Farming	Stage	Management	Agro-metrological advisories
Carp culture	Fries/ fingerlings/ brooders	Pond and water	Remove aquatic weeds. Maintain minimum depth of 1.5 – 2.0 m. Apply Agri-Lime in brooder pond at small quantity at an interval of 30 days regularly.
		Feed	Provide feed 1-2 times @ 2 % body weight per day.
		Fish	- Male and female brooders should be stock in separate ponds for early breeding programs. - harvesting of fish may be done in area where water availability is low.

(I. Meghachandra Singh)
Scientist i/c, IAAS



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Gramin Krishi Mausam Seva”

MANIPUR BULLETIN

22nd January, 2016

Ukhrul

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 23rd to 27th January, 2016 as below:

Significant past weather for the preceding bi-weekly of Imphal - West (19th to 21st January, 2016)

During last three days the sky was mainly clear to partly cloudy. 0.2 mm rainfall was recorded on 21st Jan. 2016, with maximum temperature range observed was 22.0 – 24.6°C, range of minimum temperature observed was 2.9 –9.0 °C, mean relative humidity observed in morning and afternoon were 93.67% and 51.67% respectively. The average wind speed range was 2.6 – 3.4 km/hr.

Weather forecast for next five days of Ukhrul (23rd to 27th January, 2016)

There may be no rain on Thursday. The Sky will remain mainly clear to partly foggy during morning hours, with maximum temp. of 20.0 – 21.0 °C & minimum temp. 3.0 – 4.0 °C. Relative humidity may reach upto 95 % (max RH) & 30 % (min RH). The average wind speed may be 3-4 km/ hr during next five days. The Wind direction may be south westerly to north westerly during next five days.

Weather Forecast for next 5 days *upto 0830 hours IST of 27th January

Moderate to dense fog very likely at isolated places over Manipur during next 2 days

Weather Outlook for subsequent 2 days from 27th to 29th January 2016

Rain/thundershowers is likely at isolated places over northeast India

Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Field Crops			
Rabi Maize	Early vegetative		- Light irrigation may be given to keep soil moist if the soil is dry. - Thinning may be done in vegetative stage for mustard or pea intercropped with maize.
Rapeseed/ Mustard	Vegetative to flowering		- Apply foliar application of urea {5 pumps/ sangam @ 10g/ pump} or top dress with ½ N (@ 66kg Urea/ ha) just after irrigation. - If flowering has started, then four bee boxes may be kept for one hectare crop area. Avoid spray of insecticides in such area.
		Aphid	- Spray Monocrotophos @ 1.5 ml/lit of water to the infected patches only or use yellow plates filled with water.
Field Pea/ Lentil	Vegetative	Powdery Mildew	- Spray Bayleton (Triaclemefon 25% WP) @ 7.5 g/ pump or Dinocap (Karathane 46% EC) @ 7.5ml/ pump.
		Aphid	- Spray Cypermethrin @2ml/lit of water or Nuvan @ 0.5ml/ lit of water against aphid. - For organic control against aphid, sprays that consist of Azadirachtin/ neem oil or pyrethrin may be used.
Storage (seed)			- Clean and store the seeds in RC-seed bins after proper drying of the desiccant charcoal. Check for moisture content or weight of seeds. For grains, store in well ventilated godowns with regular check for moisture.
Horticulture Crops			
Turmeric	Harvesting		-Already mature plants whose leaves are more than 70 % dried may be harvested. -Harvested rhizome should be spread in the shed for at least 4- 5 days and screen for seed purposes. -Curing process should be started within one week of harvest.
French bean			-land preparation for sowing of French bean may be started.
Potato	Vegetative		- Spray Mencozeb 2gm/litre of water at least twice at an interval of 7-10 days to prevent spreading of early blight as the temperature rises. - If aphid is observed in small patches, wash off it with plain water but if it is severe, spray Cypermethrin 20 E.C. @ 1ml/ litre of water or Nuvan @ 0.5ml/litre of water. - Give light irrigation to attain optimum soil moisture.
Cucurbits	Nursery		- Nurseries of bitter gourd, cucumber, sponge gourd, bottle gourd, pumpkin, etc. may be done under polyhouse.
Tomato	Transplanting		- Proper field preparation for transplanting of 4 leaf stage should be done. -Due consideration for removal of potentially infected material such as



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Gramin Krishi Mausam Seva”

			vines and stubbles for early blight should be done while ploughing.
	Flowering		- prophylactic measures against tomato fruit borer should be done.
Animal Sciences			
Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
Poultry (Chicken/ Turkey/ Duck)	All age groups	Care & management	- To protect birds of all age groups from inclement weather e.g. frost, snowfall etc., there should be adequate arrangements for artificial lighting and heating. - Needed to increase energy giving ingredients of feed (i.e. Maize, Wheat, Bajra etc.) and can be added 5% more.
		Coccidiosis	- After routine preventive and prophylaxis medication for the said disease, if still there is any defaecation of reddish colour faeces, somnolence, drooping of wings, offed etc., Amprolium agent can be used in feed and water. Add vitamin k, vit. A and electrolytes etc. also add mineral mixture in the feeds and water.
Sheep/ Goat	All age groups excepts the young ones	PPR (Peste Des Petits Ruminants)	- If the animals have not yet been vaccinated, give vaccine now for PPR (Peste Des Petits Ruminants).
		Enterotoxemia	- If the Lambs and Kids in particular should be given the vaccination to prevent Enterotoxaemia.
		Ectoparasitism	- Keep the sheds clean. Hang Bouquets of Basil (Ocimum sactum) or Lemon Grass (Cymbopogon citrates) in the animals sheds to keep away the ecto-parasites. Neem oil based disinfectants can be sprayed. Butox solution (2 C.C. in 1 litre of water) can also be used.
Pig	All age groups excepts the young ones	Care & management for newly born one	-Artificial lighting and heating to the piglets. -Provide adequate bedding materials.
		Deworming	-Give Morantel citrate tablet @ 15 mg/kg body weight.
		Ectoparasitism	- Keep the sheds clean. Hang Bouquets of Basil (Ocimum sactum) or Lemon Grass (Cymbopogon citrates) in the animals sheds to keep away the ecto-parasites. Neem oil based disinfectants can be sprayed. Butox solution (2 C.C. in 1 litre of water) can also be used.
Cattle	All age groups excepts the young ones	B.Q./ H.S/FMD	-If the animals have not yet been vaccinated, give vaccine, now for Black Quarter, Haemorrhagic septicaemia, Foot & mouth Disease.
		Ectoparasitism	- Keep the sheds clean. Hang Bouquets of Basil (Ocimum sactum) or Lemon Grass (Cymbopogon citrates) in the animals sheds to keep away the ecto-parasites. Neem oil based disinfectants can be sprayed. Butox solution (2 C.C. in 1 litre of water) can also be used.
		Fodder cultivation	-Irrigate the crops like Berseem Clover every 20-30 days. -At every 20-22 days for Oat.
Fishery:			
Type of Farming	Stage	Management	Agro-metrological advisories
Carp culture	Fries/ fingerlings/ brooders	Pond and water	Remove aquatic weeds. Maintain minimum depth of 1.5 – 2.0 m. Apply Agri-Lime in brooder pond at small quantity at an interval of 30 days regularly.
		Feed	Provide feed 1-2 times @ 2 % body weight per day.
		Fish	- Male and female brooders should be stock in separate ponds for early breeding programs. - harvesting of fish may be done in area where water availability is low.

(I. Meghachandra Singh)
Scientist i/c, IAAS



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Gramin Krishi Mausam Seva”

MANIPUR BULLETIN

22nd January, 2016

Bishnupur

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 23rd to 27th January, 2016 as below:

Significant past weather for the preceding bi-weekly of Imphal - West (19th to 21st January, 2016)

During last three days the sky was mainly clear to partly cloudy. 0.2 mm rainfall was recorded on 21st Jan. 2016, with maximum temperature range observed was 22.0 – 24.6°C, range of minimum temperature observed was 2.9 –9.0 °C, mean relative humidity observed in morning and afternoon were 93.67% and 51.67% respectively. The average wind speed range was 2.6 – 3.4 km/hr.

Weather forecast for next five days of Bishnupur (23rd to 27th January, 2016)

There may be no rain on Thursday. The Sky will remain mainly clear to partly foggy during morning hours, with maximum temp. of 20.0 – 21.0 °C & minimum temp. 4.0 – 5.0 °C. Relative humidity may reach upto 97 % (max RH) & 29 % (min RH). The average wind speed may be 4-5 km/ hr during next five days. The Wind direction may be south easterly to south westerly during next five days.

Weather Forecast for next 5 days *upto 0830 hours IST of 27th January

Moderate to dense fog very likely at isolated places over Manipur during next 2 days

Weather Outlook for subsequent 2 days from 27th to 29th January 2016

Rain/thundershowers is likely at isolated places over northeast India

Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Field Crops			
Rabi Maize	Early vegetative		- Light irrigation may be given to keep soil moist if the soil is dry. - Thinning may be done in vegetative stage for mustard or pea intercropped with maize.
Rapeseed/ Mustard	Vegetative to flowering		- Apply foliar application of urea {5 pumps/ sangam @ 10g/ pump} or top dress with ½ N (@ 66kg Urea/ ha) just after irrigation. - If flowering has started, then four bee boxes may be kept for one hectare crop area. Avoid spray of insecticides in such area.
		Aphid	- Spray Monocrotophos @ 1.5 ml/lit of water to the infected patches only or use yellow plates filled with water.
Field Pea/ Lentil	Vegetative	Powdery Mildew	- Spray Bayleton (Triaclemefon 25% WP) @ 7.5 g/ pump or Dinocap (Karathane 46% EC) @ 7.5ml/ pump.
		Aphid	- Spray Cypermethrin @2ml/lit of water or Nuvan @ 0.5ml/ lit of water against aphid. - For organic control against aphid, sprays that consist of Azadirachtin/ neem oil or pyrethrin may be used.
Storage (seed)			- Clean and store the seeds in RC-seed bins after proper drying of the desiccant charcoal. Check for moisture content or weight of seeds. For grains, store in well ventilated godowns with regular check for moisture.
Horticulture Crops			
Turmeric	Harvesting		-Already mature plants whose leaves are more than 70 % dried may be harvested. -Harvested rhizome should be spread in the shed for at least 4- 5 days and screen for seed purposes. -Curing process should be started within one week of harvest.
French bean			-land preparation for sowing of French bean may be started.
Potato	Vegetative		- Spray Mencozeb 2gm/litre of water at least twice at an interval of 7-10 days to prevent spreading of early blight as the temperature rises. - If aphid is observed in small patches, wash off it with plain water but if it is severe, spray Cypermethrin 20 E.C. @ 1ml/ litre of water or Nuvan @ 0.5ml/litre of water. - Give light irrigation to attain optimum soil moisture.
Cucurbits	Nursery		- Nurseries of bitter gourd, cucumber, sponge gourd, bottle gourd, pumpkin, etc. may be done under polyhouse.
Tomato	Transplanting		- Proper field preparation for transplanting of 4 leaf stage should be done. -Due consideration for removal of potentially infected material such as



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Gramin Krishi Mausam Seva”

			vines and stubbles for early blight should be done while ploughing.
	Flowering		- prophylactic measures against tomato fruit borer should be done.
Animal Sciences			
Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
Poultry (Chicken/ Turkey/ Duck)	All age groups	Care & management	- To protect birds of all age groups from inclement weather e.g. frost, snowfall etc., there should be adequate arrangements for artificial lighting and heating. - Needed to increase energy giving ingredients of feed (i.e. Maize, Wheat, Bajra etc.) and can be added 5% more.
		Coccidiosis	- After routine preventive and prophylaxis medication for the said disease, if still there is any defaecation of reddish colour faeces, somnolence, drooping of wings, offed etc., Amprolium agent can be used in feed and water. Add vitamin k, vit. A and electrolytes etc. also add mineral mixture in the feeds and water.
Sheep/ Goat	All age groups excepts the young ones	PPR (Peste Des Petits Ruminants)	- If the animals have not yet been vaccinated, give vaccine now for PPR (Peste Des Petits Ruminants).
		Enterotoxemia	- If the Lambs and Kids in particular should be given the vaccination to prevent Enterotoxaemia.
		Ectoparasitism	- Keep the sheds clean. Hang Bouquets of Basil (Ocimum sanctum) or Lemon Grass (Cymbopogon citrates) in the animals sheds to keep away the ecto-parasites. Neem oil based disinfectants can be sprayed. Butox solution (2 C.C. in 1 litre of water) can also be used.
Pig	All age groups excepts the young ones	Care & management for newly born one	-Artificial lighting and heating to the piglets. -Provide adequate bedding materials.
		Deworming	-Give Morantel citrate tablet @ 15 mg/kg body weight.
		Ectoparasitism	- Keep the sheds clean. Hang Bouquets of Basil (Ocimum sanctum) or Lemon Grass (Cymbopogon citrates) in the animals sheds to keep away the ecto-parasites. Neem oil based disinfectants can be sprayed. Butox solution (2 C.C. in 1 litre of water) can also be used.
Cattle	All age groups excepts the young ones	B.Q./ H.S/FMD	-If the animals have not yet been vaccinated, give vaccine, now for Black Quarter, Haemorrhagic septicaemia, Foot & mouth Disease.
		Ectoparasitism	- Keep the sheds clean. Hang Bouquets of Basil (Ocimum sanctum) or Lemon Grass (Cymbopogon citrates) in the animals sheds to keep away the ecto-parasites. Neem oil based disinfectants can be sprayed. Butox solution (2 C.C. in 1 litre of water) can also be used.
		Fodder cultivation	-Irrigate the crops like Berseem Clover every 20-30 days. -At every 20-22 days for Oat.
Fishery:			
Type of Farming	Stage	Management	Agro-metrological advisories
Carp culture	Fries/ fingerlings/ brooders	Pond and water	Remove aquatic weeds. Maintain minimum depth of 1.5 – 2.0 m. Apply Agri-Lime in brooder pond at small quantity at an interval of 30 days regularly.
		Feed	Provide feed 1-2 times @ 2 % body weight per day.
		Fish	- Male and female brooders should be stock in separate ponds for early breeding programs. - harvesting of fish may be done in area where water availability is low.

(I. Meghachandra Singh)
Scientist i/c, IAAS



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Gramin Krishi Mausam Seva”

MANIPUR BULLETIN

22nd January, 2016

Chandel

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 23rd to 27th January, 2016 as below:

Significant past weather for the preceding bi-weekly of Imphal - West (19th to 21st January, 2016)

During last three days the sky was mainly clear to partly cloudy. 0.2 mm rainfall was recorded on 21st Jan. 2016, with maximum temperature range observed was 22.0 – 24.6°C, range of minimum temperature observed was 2.9 –9.0 °C, mean relative humidity observed in morning and afternoon were 93.67% and 51.67% respectively. The average wind speed range was 2.6 – 3.4 km/hr.

Weather forecast for next five days of Chandel (23rd to 27th January, 2016)

There may be no rain on Thursday. The Sky will remain mainly clear to partly foggy during morning hours, with maximum temp. of 19.0 – 20.0 °C & minimum temp. 4.0 – 5.0 °C. Relative humidity may reach upto 95 % (max RH) & 32 % (min RH). The average wind speed may be 4-5 km/ hr during next five days. The Wind direction may be west westerly to north west westerly during next five days.

Weather Forecast for next 5 days *upto 0830 hours IST of 27th January

Moderate to dense fog very likely at isolated places over Manipur during next 2 days

Weather Outlook for subsequent 2 days from 27th to 29th January 2016

Rain/thundershowers is likely at isolated places over northeast India

Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Field Crops			
Rabi Maize	Early vegetative		- Light irrigation may be given to keep soil moist if the soil is dry. - Thinning may be done in vegetative stage for mustard or pea intercropped with maize.
Rapeseed/ Mustard	Vegetative to flowering		- Apply foliar application of urea {5 pumps/ sangam @ 10g/ pump} or top dress with ½ N (@ 66kg Urea/ ha) just after irrigation. - If flowering has started, then four bee boxes may be kept for one hectare crop area. Avoid spray of insecticides in such area.
		Aphid	- Spray Monocrotophos @ 1.5 ml/lit of water to the infected patches only or use yellow plates filled with water.
Field Pea/ Lentil	Vegetative	Powdery Mildew	- Spray Bayleton (Triaclemefon 25% WP) @ 7.5 g/ pump or Dinocap (Karathane 46% EC) @ 7.5ml/ pump.
		Aphid	- Spray Cypermethrin @2ml/lit of water or Nuvan @ 0.5ml/ lit of water against aphid. - For organic control against aphid, sprays that consist of Azadirachtin/ neem oil or pyrethrin may be used.
Storage (seed)			- Clean and store the seeds in RC-seed bins after proper drying of the desiccant charcoal. Check for moisture content or weight of seeds. For grains, store in well ventilated godowns with regular check for moisture.
Horticulture Crops			
Turmeric	Harvesting		-Already mature plants whose leaves are more than 70 % dried may be harvested. -Harvested rhizome should be spread in the shed for at least 4- 5 days and screen for seed purposes. -Curing process should be started within one week of harvest.
French bean			-land preparation for sowing of French bean may be started.
Potato	Vegetative		- Spray Mencozeb 2gm/litre of water at least twice at an interval of 7-10 days to prevent spreading of early blight as the temperature rises. - If aphid is observed in small patches, wash off it with plain water but if it is severe, spray Cypermethrin 20 E.C. @ 1ml/ litre of water or Nuvan @ 0.5ml/litre of water. - Give light irrigation to attain optimum soil moisture.
Cucurbits	Nursery		- Nurseries of bitter gourd, cucumber, sponge gourd, bottle gourd, pumpkin, etc. may be done under polyhouse.
Tomato	Transplanting		- Proper field preparation for transplanting of 4 leaf stage should be done. -Due consideration for removal of potentially infected material such as



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Gramin Krishi Mausam Seva”

			vines and stubbles for early blight should be done while ploughing.
	Flowering		- prophylactic measures against tomato fruit borer should be done.
Animal Sciences			
Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
Poultry (Chicken/ Turkey/ Duck)	All age groups	Care & management	- To protect birds of all age groups from inclement weather e.g. frost, snowfall etc., there should be adequate arrangements for artificial lighting and heating. - Needed to increase energy giving ingredients of feed (i.e. Maize, Wheat, Bajra etc.) and can be added 5% more.
		Coccidiosis	- After routine preventive and prophylaxis medication for the said disease, if still there is any defaecation of reddish colour faeces, somnolence, drooping of wings, offed etc., Amprolium agent can be used in feed and water. Add vitamin k, vit. A and electrolytes etc. also add mineral mixture in the feeds and water.
Sheep/ Goat	All age groups excepts the young ones	PPR (Peste Des Petits Ruminants)	- If the animals have not yet been vaccinated, give vaccine now for PPR (Peste Des Petits Ruminants).
		Enterotoxemia	- If the Lambs and Kids in particular should be given the vaccination to prevent Enterotoxaemia.
		Ectoparasitism	- Keep the sheds clean. Hang Bouquets of Basil (Ocimum sactum) or Lemon Grass (Cymbopogon citrates) in the animals sheds to keep away the ecto-parasites. Neem oil based disinfectants can be sprayed. Butox solution (2 C.C. in 1 litre of water) can also be used.
Pig	All age groups excepts the young ones	Care & management for newly born one	-Artificial lighting and heating to the piglets. -Provide adequate bedding materials.
		Deworming	-Give Morantel citrate tablet @ 15 mg/kg body weight.
		Ectoparasitism	- Keep the sheds clean. Hang Bouquets of Basil (Ocimum sactum) or Lemon Grass (Cymbopogon citrates) in the animals sheds to keep away the ecto-parasites. Neem oil based disinfectants can be sprayed. Butox solution (2 C.C. in 1 litre of water) can also be used.
Cattle	All age groups excepts the young ones	B.Q./ H.S/FMD	-If the animals have not yet been vaccinated, give vaccine, now for Black Quarter, Haemorrhagic septicaemia, Foot & mouth Disease.
		Ectoparasitism	- Keep the sheds clean. Hang Bouquets of Basil (Ocimum sactum) or Lemon Grass (Cymbopogon citrates) in the animals sheds to keep away the ecto-parasites. Neem oil based disinfectants can be sprayed. Butox solution (2 C.C. in 1 litre of water) can also be used.
		Fodder cultivation	-Irrigate the crops like Berseem Clover every 20-30 days. -At every 20-22 days for Oat.
Fishery:			
Type of Farming	Stage	Management	Agro-metrological advisories
Carp culture	Fries/ fingerlings/ brooders	Pond and water	Remove aquatic weeds. Maintain minimum depth of 1.5 – 2.0 m. Apply Agri-Lime in brooder pond at small quantity at an interval of 30 days regularly.
		Feed	Provide feed 1-2 times @ 2 % body weight per day.
		Fish	- Male and female brooders should be stock in separate ponds for early breeding programs. - harvesting of fish may be done in area where water availability is low.

(I. Meghachandra Singh)
Scientist i/c, IAAS



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Gramin Krishi Mausam Seva”

MANIPUR BULLETIN

22nd January, 2016

Churachandpur

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 23rd to 27th January, 2016 as below:

<p>Significant past weather for the preceding bi-weekly of Imphal - West (19th to 21st January, 2016)</p> <p>During last three days the sky was mainly clear to partly cloudy. 0.2 mm rainfall was recorded on 21st Jan. 2016, with maximum temperature range observed was 22.0 – 24.6°C, range of minimum temperature observed was 2.9 –9.0 °C, mean relative humidity observed in morning and afternoon were 93.67% and 51.67% respectively. The average wind speed range was 2.6 – 3.4 km/hr.</p>	<p>Weather forecast for next five days of Churachandpur (23rd to 27th January, 2016)</p> <p>There may be no rain on Thursday. The Sky will remain mainly clear to partly foggy during morning hours, with maximum temp. of 20.0 – 21.0 °C & minimum temp. 4.0 – 5.0 °C. Relative humidity may reach upto 98 % (max RH) & 28 % (min RH). The average wind speed may be 4-5 km/ hr during next five days. The Wind direction may be northerly to south south easterly during next five days.</p>
--	---

Weather Forecast for next 5 days *upto 0830 hours IST of 27th January

Moderate to dense fog very likely at isolated places over Manipur during next 2 days

Weather Outlook for subsequent 2 days from 27th to 29th January 2016

Rain/thundershowers is likely at isolated places over northeast India

Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Field Crops			
Rabi Maize	Early vegetative		- Light irrigation may be given to keep soil moist if the soil is dry. - Thinning may be done in vegetative stage for mustard or pea intercropped with maize.
Rapeseed/ Mustard	Vegetative to flowering		- Apply foliar application of urea {5 pumps/ sangam @ 10g/ pump} or top dress with ½ N (@ 66kg Urea/ ha) just after irrigation. - If flowering has started, then four bee boxes may be kept for one hectare crop area. Avoid spray of insecticides in such area.
		Aphid	- Spray Monocrotophos @ 1.5 ml/lit of water to the infected patches only or use yellow plates filled with water.
Field Pea/ Lentil	Vegetative	Powdery Mildew	- Spray Bayleton (Triaclemefon 25% WP) @ 7.5 g/ pump or Dinocap (Karathane 46% EC) @ 7.5ml/ pump.
		Aphid	- Spray Cypermethrin @2ml/lit of water or Nuvan @ 0.5ml/ lit of water against aphid. - For organic control against aphid, sprays that consist of Azadirachtin/ neem oil or pyrethrin may be used.
Storage (seed)			- Clean and store the seeds in RC-seed bins after proper drying of the desiccant charcoal. Check for moisture content or weight of seeds. For grains, store in well ventilated godowns with regular check for moisture.
Horticulture Crops			
Turmeric	Harvesting		-Already mature plants whose leaves are more than 70 % dried may be harvested. -Harvested rhizome should be spread in the shed for at least 4- 5 days and screen for seed purposes. -Curing process should be started within one week of harvest.
French bean			-land preparation for sowing of French bean may be started.
Potato	Vegetative		- Spray Mencozeb 2gm/litre of water at least twice at an interval of 7-10 days to prevent spreading of early blight as the temperature rises. - If aphid is observed in small patches, wash off it with plain water but if it is severe, spray Cypermethrin 20 E.C. @ 1ml/ litre of water or Nuvan @ 0.5ml/litre of water. - Give light irrigation to attain optimum soil moisture.
Cucurbits	Nursery		- Nurseries of bitter gourd, cucumber, sponge gourd, bottle gourd, pumpkin, etc. may be done under polyhouse.
Tomato	Transplanting		- Proper field preparation for transplanting of 4 leaf stage should be done. -Due consideration for removal of potentially infected material such as



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Gramin Krishi Mausam Seva”

			vines and stubbles for early blight should be done while ploughing.
	Flowering		- prophylactic measures against tomato fruit borer should be done.
Animal Sciences			
Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
Poultry (Chicken/ Turkey/ Duck)	All age groups	Care & management	- To protect birds of all age groups from inclement weather e.g. frost, snowfall etc., there should be adequate arrangements for artificial lighting and heating. - Needed to increase energy giving ingredients of feed (i.e. Maize, Wheat, Bajra etc.) and can be added 5% more.
		Coccidiosis	- After routine preventive and prophylaxis medication for the said disease, if still there is any defaecation of reddish colour faeces, somnolence, drooping of wings, offed etc., Amprolium agent can be used in feed and water. Add vitamin k, vit. A and electrolytes etc. also add mineral mixture in the feeds and water.
Sheep/ Goat	All age groups excepts the young ones	PPR (Peste Des Petits Ruminants)	- If the animals have not yet been vaccinated, give vaccine now for PPR (Peste Des Petits Ruminants).
		Enterotoxemia	- If the Lambs and Kids in particular should be given the vaccination to prevent Enterotoxaemia.
		Ectoparasitism	- Keep the sheds clean. Hang Bouquets of Basil (Ocimum sanctum) or Lemon Grass (Cymbopogon citrates) in the animals sheds to keep away the ecto-parasites. Neem oil based disinfectants can be sprayed. Butox solution (2 C.C. in 1 litre of water) can also be used.
Pig	All age groups excepts the young ones	Care & management for newly born one	-Artificial lighting and heating to the piglets. -Provide adequate bedding materials.
		Deworming	-Give Morantel citrate tablet @ 15 mg/kg body weight.
		Ectoparasitism	- Keep the sheds clean. Hang Bouquets of Basil (Ocimum sanctum) or Lemon Grass (Cymbopogon citrates) in the animals sheds to keep away the ecto-parasites. Neem oil based disinfectants can be sprayed. Butox solution (2 C.C. in 1 litre of water) can also be used.
Cattle	All age groups excepts the young ones	B.Q./ H.S/FMD	-If the animals have not yet been vaccinated, give vaccine, now for Black Quarter, Haemorrhagic septicaemia, Foot & mouth Disease.
		Ectoparasitism	- Keep the sheds clean. Hang Bouquets of Basil (Ocimum sanctum) or Lemon Grass (Cymbopogon citrates) in the animals sheds to keep away the ecto-parasites. Neem oil based disinfectants can be sprayed. Butox solution (2 C.C. in 1 litre of water) can also be used.
		Fodder cultivation	-Irrigate the crops like Berseem Clover every 20-30 days. -At every 20-22 days for Oat.
Fishery:			
Type of Farming	Stage	Management	Agro-metrological advisories
Carp culture	Fries/ fingerlings/ brooders	Pond and water	Remove aquatic weeds. Maintain minimum depth of 1.5 – 2.0 m. Apply Agri-Lime in brooder pond at small quantity at an interval of 30 days regularly.
		Feed	Provide feed 1-2 times @ 2 % body weight per day.
		Fish	- Male and female brooders should be stock in separate ponds for early breeding programs. - harvesting of fish may be done in area where water availability is low.

(I. Meghachandra Singh)
Scientist i/c, IAAS



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Gramin Krishi Mausam Seva”

MANIPUR BULLETIN

22nd January, 2016

Imphal-East

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 23rd to 27th January, 2016 as below:

<p>Significant past weather for the preceding bi-weekly of Imphal - West (19th to 21st January, 2016)</p> <p>During last three days the sky was mainly clear to partly cloudy. 0.2 mm rainfall was recorded on 21st Jan. 2016, with maximum temperature range observed was 22.0 – 24.6°C, range of minimum temperature observed was 2.9 –9.0 °C, mean relative humidity observed in morning and afternoon were 93.67% and 51.67% respectively. The average wind speed range was 2.6 – 3.4 km/hr.</p>	<p>Weather forecast for next five days of Imphal-East (23rd to 27th January, 2016)</p> <p>There may be no rain for next five days. The Sky will remain mainly clear to partly foggy during morning hours, with maximum temp. of 21.0 – 22.0 °C & minimum temp. 4.0 – 5.0 °C. Relative humidity may reach upto 99 % (max RH) & 24 % (min RH). The average wind speed may be 2-4 km/ hr during next five days. The Wind direction may be north easterly to north west westerly during next five days.</p>
--	--

Weather Forecast for next 5 days * upto 0830 hours IST of 27th January

Moderate to dense fog very likely at isolated places over Manipur during next 2 days

Weather Outlook for subsequent 2 days from 27th to 29th January 2016

Rain/thundershowers is likely at isolated places over northeast India

Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Field Crops			
Rabi Maize	Early vegetative		- Light irrigation may be given to keep soil moist if the soil is dry. - Thinning may be done in vegetative stage for mustard or pea intercropped with maize.
Rapeseed/ Mustard	Vegetative to flowering		- Apply foliar application of urea {5 pumps/ sangam @ 10g/ pump} or top dress with ½ N (@ 66kg Urea/ ha) just after irrigation. - If flowering has started, then four bee boxes may be kept for one hectare crop area. Avoid spray of insecticides in such area.
		Aphid	- Spray Monocrotophos @ 1.5 ml/lit of water to the infected patches only or use yellow plates filled with water.
Field Pea/ Lentil	Vegetative	Powdery Mildew	- Spray Bayleton (Triaclemefon 25% WP) @ 7.5 g/ pump or Dinocap (Karathane 46% EC) @ 7.5ml/ pump.
		Aphid	- Spray Cypermethrin @2ml/lit of water or Nuvan @ 0.5ml/ lit of water against aphid. - For organic control against aphid, sprays that consist of Azadirachtin/ neem oil or pyrethrin may be used.
Storage (seed)			- Clean and store the seeds in RC-seed bins after proper drying of the desiccant charcoal. Check for moisture content or weight of seeds. For grains, store in well ventilated godowns with regular check for moisture.
Horticulture Crops			
Turmeric	Harvesting		-Already mature plants whose leaves are more than 70 % dried may be harvested. -Harvested rhizome should be spread in the shed for at least 4- 5 days and screen for seed purposes. -Curing process should be started within one week of harvest.
French bean			-land preparation for sowing of French bean may be started.
Potato	Vegetative		- Spray Mencozeb 2gm/litre of water at least twice at an interval of 7-10 days to prevent spreading of early blight as the temperature rises. - If aphid is observed in small patches, wash off it with plain water but if it is severe, spray Cypermethrin 20 E.C. @ 1ml/ litre of water or Nuvan @ 0.5ml/litre of water. - Give light irrigation to attain optimum soil moisture.
Cucurbits	Nursery		- Nurseries of bitter gourd, cucumber, sponge gourd, bottle gourd, pumpkin, etc. may be done under polyhouse.
Tomato	Transplanting		- Proper field preparation for transplanting of 4 leaf stage should be done. -Due consideration for removal of potentially infected material such as



**ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004**



“Gramin Krishi Mausam Seva”

			vines and stubbles for early blight should be done while ploughing.
	Flowering		- prophylactic measures against tomato fruit borer should be done.
Animal Sciences			
Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
Poultry (Chicken/ Turkey/ Duck)	All age groups	Care & management	- To protect birds of all age groups from inclement weather e.g. frost, snowfall etc., there should be adequate arrangements for artificial lighting and heating. - Needed to increase energy giving ingredients of feed (i.e. Maize, Wheat, Bajra etc.) and can be added 5% more.
		Coccidiosis	- After routine preventive and prophylaxis medication for the said disease, if still there is any defaecation of reddish colour faeces, somnolence, drooping of wings, offed etc., Amprolium agent can be used in feed and water. Add vitamin k, vit. A and electrolytes etc. also add mineral mixture in the feeds and water.
Sheep/ Goat	All age groups excepts the young ones	PPR (Peste Des Petits Ruminants)	- If the animals have not yet been vaccinated, give vaccine now for PPR (Peste Des Petits Ruminants).
		Enterotoxemia	- If the Lambs and Kids in particular should be given the vaccination to prevent Enterotoxaemia.
		Ectoparasitism	- Keep the sheds clean. Hang Bouquets of Basil (Ocimum sanctum) or Lemon Grass (Cymbopogon citrates) in the animals sheds to keep away the ecto-parasites. Neem oil based disinfectants can be sprayed. Butox solution (2 C.C. in 1 litre of water) can also be used.
Pig	All age groups excepts the young ones	Care & management for newly born one	-Artificial lighting and heating to the piglets. -Provide adequate bedding materials.
		Deworming	-Give Morantel citrate tablet @ 15 mg/kg body weight.
		Ectoparasitism	- Keep the sheds clean. Hang Bouquets of Basil (Ocimum sanctum) or Lemon Grass (Cymbopogon citrates) in the animals sheds to keep away the ecto-parasites. Neem oil based disinfectants can be sprayed. Butox solution (2 C.C. in 1 litre of water) can also be used.
Cattle	All age groups excepts the young ones	B.Q./ H.S/FMD	-If the animals have not yet been vaccinated, give vaccine, now for Black Quarter, Haemorrhagic septicaemia, Foot & mouth Disease.
		Ectoparasitism	- Keep the sheds clean. Hang Bouquets of Basil (Ocimum sanctum) or Lemon Grass (Cymbopogon citrates) in the animals sheds to keep away the ecto-parasites. Neem oil based disinfectants can be sprayed. Butox solution (2 C.C. in 1 litre of water) can also be used.
		Fodder cultivation	-Irrigate the crops like Berseem Clover every 20-30 days. -At every 20-22 days for Oat.
Fishery:			
Type of Farming	Stage	Management	Agro-metrological advisories
Carp culture	Fries/ fingerlings/ brooders	Pond and water	Remove aquatic weeds. Maintain minimum depth of 1.5 – 2.0 m. Apply Agri-Lime in brooder pond at small quantity at an interval of 30 days regularly.
		Feed	Provide feed 1-2 times @ 2 % body weight per day.
		Fish	- Male and female brooders should be stock in separate ponds for early breeding programs. - harvesting of fish may be done in area where water availability is low.

(I. Meghachandra Singh)
Scientist i/c, IAAS