



Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 14-02-2020

Weather Forecast of KANGPOKPI(Manipur) Issued On : 2020-02-14(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-15	0.0	24.0	8.0	71	32	5.4	255	1
2020-02-16	0.0	24.0	9.0	74	33	5.6	18	0
2020-02-17	0.0	24.0	10.0	72	26	5.5	22	0
2020-02-18	0.0	25.0	12.0	66	20	5.2	342	0
2020-02-19	0.0	25.0	12.0	46	18	5.8	22	0

Weather Summary/Alert:

Likely to have no rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 26-29°C & mini. temp. 12-14 °C. Relative humidity may reach upto 80 % (max) & 26% (min). The average wind speed may be 5.1-5.5 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 26-29°C amadi ingbagina 12-14 °C oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 76% amadi nungthildana 26% oiraba yai. Nungsit sitpigi changna pung amada 5.1-5.5 km oiraba yai.

General Advisory:

Monitor regularly insect pest infestation in crops. Mulching may be done in the crop field using straw/ dried leaves which is not infected by insect-pest to preserve soil moisture for the crop. Keep birds/ animals under your jurisdiction and avoid contact with migratory or wild birds/ animals.

Loufamda laina tilna taba ningthina toina yengsinbiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Sha uchek yariba makhei sennajaba mafamda thadokpiyu amadi lamlakki sha-uchek/ mapandagi paisinlakpa uchekka samnadanaba shinbiyu.

SMS Advisory:

Insect pest sprays may be given before rains. As weather is changing proper care of sheds, food and drinking water must be taken & administer vaccines in time.

Nong laktingaida laina tilnagi hidak kaphthokpiyu. Aying asa onrakpada sha uchek naraktanaba makol, ishing amadi machinjak afaba pibiduna matam chana hidak thabiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
TOMATO	Tomato nurseries should be raised under low cost polyhouse as the temperature is very low in open field. - Nursery bed of 0.75m x 4m may be prepared and mix Carbendazim @ 5g/ 3m ² area, keep the bed covered with gunny bag for at least 24 hrs. - Sow seeds @ 10g/ 3m ² area keep the bed covered with gunny bag. Seeds will germinate in 4-5 days.

Crop(Varieties)	Crop Specific Advisory
	- Ingba matamda polyhouse makhada khamen ashinbi chaar hunbiyu. - 0.75m x 4m pak-chaoba nursery bed samba loisanlakpada 3m2 khudingda 5g gi changda Carbendazim happaga bora na kupsinlaga pung 24 thambiyu. - 3m2 pak-chaoba nursery bed ta 10g gi changda maru hidak totlaga hunbibagi matungda bora kupsinlaga thambiyu. Numit 4-5ni gi manungda maton thokpaga chang yengbiraga bora louthokpiba yarani.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
POTATO	- For seed: • Strict supervision on removal of unwanted plant type should be maintained. • Maturity days should be strictly maintained as per variety. - Harvest the crop after 10-15 days after haulm cutting - Maru oinaba lokpige hairabadi: • Masak matou manadaba pambi makhal khandokpiyu.. • Alu na chaoba mapung faba matam variety makhal gi leijaba numit shurakpaga matungda lokpiyu. - Alu loktoubu numit 10-15ni watlingaeida makhok khara khakaga pambi kakaga thambiyu.
FRENCH BEAN	- Sowing of French bean may be done in prepared land. Seed rate for dwarf varieties like Pant Anupam & Arka Komal is 65kg/ha (spacing is 30x45 cm from seed to seed and row to row) and 25-30kg/ha for pole-type (spacing is 60x60cm with 2-3 seeds per hill). Treat seeds with Carbendazim @2g/kg seeds before sowing. - Give light irrigation after 2-3 days of sowing. - Koli hawai thanaba loufam semjinbiba yarani. Nemna panbi variety Pant Anupam amadi Arka Komal pari amada thanaba 65 kg maru lousinbiraga 30x45 cm lapna thabiyu. Pambi awangbagina 25-30 kg changani amadi 60x60cm gi changda lapna makhul amada 2-3 maru thabiyu. - Thadringeida maru kg amada 2g gi changda Carbendazim totpiyu amadi thabagi matung leibak leinil marang kaina leinaba nong churaktagadi ishing khara chaibiyu.



Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 14-02-2020

Weather Forecast of NONEY (Manipur) Issued On : 2020-02-14 (Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-15	0.0	24.0	8.0	80	34	3.8	252	0
2020-02-16	0.0	24.0	9.0	76	31	4.2	68	0
2020-02-17	0.0	24.0	10.0	64	28	4.4	68	0
2020-02-18	0.0	25.0	12.0	75	22	4.1	45	0
2020-02-19	0.0	25.0	12.0	52	21	4.7	63	0

Weather Summary/Alert:

Likely to have no rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 25-29 oC & mini. temp. 12-14 oC. Relative humidity may reach upto 84% (max) & 23% (min). The average wind speed may be 3.9-4.5 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 25-29 oC amadi ingbagina 12-14 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 84 % amadi nungthildana 23 % oiraba yai. Nungsit sitpaga changna pung amada 3.9-4.5 km oiraba yai.

General Advisory:

Monitor regularly insect pest infestation in crops. Mulching may be done in the crop field using straw/ dried leaves which is not infected by insect-pest to preserve soil moisture for the crop. Keep birds/ animals under your jurisdiction and avoid contact with migratory or wild birds/ animals.

Loufamda laina tilna taba ningthina toina yengsinbiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Sha uchek yariba makhei sennajaba mafamda thadokpiyu amadi lamlakki sha-uchek/ mapandagi paisinlakpa ucheikka samnadanaba shinbiyu.

SMS Advisory:

Insect pest sprays may be given before rains. As weather is changing proper care of sheds, food and drinking water must be taken & administer vaccines in time.

Nong laktingaida laina tilnagi hidak kaphthokpiyu. Aying asa onrakpada sha uchek naraktanaba makol, ishing amadi machinjak afaba pibiduna matam chana hidak thabiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
TOMATO	Tomato nurseries should be raised under low cost polyhouse as the temperature is very low in open field. - Nursery bed of 0.75m x 4m may be prepared and mix Carbendazim @ 5g/ 3m ² area, keep the bed covered with gunny bag for at least 24 hrs. - Sow seeds @ 10g/ 3m ² area keep the bed covered with gunny bag. Seeds will germinate in 4-5 days.

Crop(Varieties)	Crop Specific Advisory
	- Ingba matamda polyhouse makhada khamen ashinbi chaar hunbiyu. - 0.75m x 4m pak-chaoba nursery bed samba loisanlakpada 3m2 khudingda 5g gi changda Carbendazim happaga bora na kupsinlaga pung 24 thambiyu. - 3m2 pak-chaoba nursery bed ta 10g gi changda maru hidak totlaga hunbibagi matungda bora kupsinlaga thambiyu. Numit 4-5ni gi manungda maton thokpaga chang yengbiraga bora louthokpiba yarani.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
POTATO	- For seed: • Strict supervision on removal of unwanted plant type should be maintained. • Maturity days should be strictly maintained as per variety. - Harvest the crop after 10-15 days after haulm cutting - Maru oinaba lokpige hairabadi: • Masak matou manadaba pambi makhal khandokpiyu.. • Alu na chaoba mapung faba matam variety makhal gi leijaba numit shurakpaga matungda lokpiyu. - Alu loktoubu numit 10-15ni watlingaeida makhok khara khakaga pambi kakaga thambiyu.
FRENCH BEAN	- Sowing of French bean may be done in prepared land. Seed rate for dwarf varieties like Pant Anupam & Arka Komal is 65kg/ha (spacing is 30x45 cm from seed to seed and row to row) and 25-30kg/ha for pole-type (spacing is 60x60cm with 2-3 seeds per hill). Treat seeds with Carbendazim @2g/kg seeds before sowing. - Give light irrigation after 2-3 days of sowing. - Koli hawai thanaba loufam semjinbiba yarani. Nemna panbi variety Pant Anupam amadi Arka Komal pari amada thanaba 65 kg maru lousinbiraga 30x45 cm lapna thabiyu. Pambi awangbagina 25-30 kg changani amadi 60x60cm gi changda lapna makhul amada 2-3 maru thabiyu. - Thadringeida maru kg amada 2g gi changda Carbendazim totpiyu amadi thabagi matung leibak leinil marang kaina leinaba nong churaktagadi ishing khara chaibiyu.



Gramin Krishi Mausam Sewa

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 14-02-2020

Weather Forecast of PATSOI Block in IMPHAL-WEST(Manipur) Issued On :2020-02-14(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-15	0.0	25.9	12.2	78	43	8.0	283	1
2020-02-16	0.0	26.1	13.0	86	37	7.0	360	0
2020-02-17	0.0	29.2	13.6	67	27	6.0	23	0
2020-02-18	0.0	29.9	12.7	79	23	6.0	334	0
2020-02-19	0.0	28.8	12.0	64	25	7.0	252	0

Weather Summary/Alert:

Likely to have no rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 25.9-29.9 oC & mini. temp. 12-13.6 oC. Relative humidity may reach upto 86 % (max) & 43 % (min). The average wind speed may be 6-8 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 25.9-29.9oC amadi ingbagina 12-13.6 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 86 % amadi nungthildana 43 % oiraba yai. Nungsit sitpagi changna pung amada 6-8 km oiraba yai.

General Advisory:

Monitor regularly insect pest infestation in crops. Mulching may be done in the crop field using straw/ dried leaves which is not infected by insect-pest to preserve soil moisture for the crop. Keep birds/ animals under your jurisdiction and avoid contact with migratory or wild birds/ animals.

Loufamda laina tilna taba ningthina toina yengsinbiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Sha uchek yariba makhei sennajaba mafamda thadokpiyu amadi lamlakki sha-uchek/ mapandagi paisinlakpa uchekka samnadanaba shinbiyu.

SMS Advisory:

Insect pest sprays may be given before rains. As weather is changing proper care of sheds, food and drinking water must be taken & administer vaccines in time.

Nong laktingaida laina tilnagi hidak kaphthokpiyu. Aying asa onrakpada sha uchek naraktanaba makol, ishing amadi machinjak afaba pibiduna matam chana hidak thabiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
TOMATO	Tomato nurseries should be raised under low cost polyhouse as the temperature is very low in open field. - Nursery bed of 0.75m x 4m may be prepared and mix Carbendazim @ 5g/ 3m2 area, keep the bed covered with gunny bag for at least 24 hrs. - Sow seeds @ 10g/ 3m2 area keep the bed covered with gunny bag. Seeds will germinate in 4-5 days.
	- Ingba matamda polyhouse makhada khamen ashinbi chaar hunbiyu. - 0.75m x 4m pak-chaoba nursery bed samba loisanlakpada 3m2 khudingda 5g gi changda Carbendazim happaga bora na kupsinlaga pung 24 thambiyu. - 3m2 pak-chaoba nursery bed ta 10g gi changda maru hidak totlaga hunbibagi matungda bora kupsinlaga thambiyu. Numit 4-5ni gi manungda maton thokpagi chang yengbiraga bora louthokpiba yarani.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
POTATO	- For seed: • Strict supervision on removal of unwanted plant type should be maintained. • Maturity days should be strictly maintained as per variety. - Harvest the crop after 10-15 days after haulm cutting - Maru oinaba lokpige hairabadi: • Masak matou manadaba pambi makhal khandokpiyu.. • Alu na chaoba mapung faba matam variety makhal gi leijaba numit shurakpagi matungda lokpiyu. - Alu loktoubu numit 10-15ni watlingaeida makhok khara khakaga pambi kakaga thambiyu.
FRENCH BEAN	- Sowing of French bean may be done in prepared land. Seed rate for dwarf varieties like Pant Anupam & Arka Komal is 65kg/ha (spacing is 30x45 cm from seed to seed and row to row) and 25-30kg/ha for pole-type (spacing is 60x60cm with 2-3 seeds per hill). Treat seeds with Carbendazim @2g/kg seeds before sowing. - Give light irrigation after 2-3 days of sowing. - Koli hawai thanaba loufam semjinbiba yarani. Nemna panbi variety Pant Anupam amadi Arka Komal pari amada thanaba 65 kg maru lousinbiraga 30x45 cm lapna thabiyu. Pambi awangbagina 25-30 kg changani amadi 60x60cm gi changda lapna makhul amada 2-3 maru thabiyu. - Thadringeida maru kg amada 2g gi changda Carbendazim totpiyu amadi thabagi matung leibak leinil marang kaina leinaba nong churaktagadi ishing khara chaibiyu.



Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 14-02-2020

Weather Forecast of PHERZAWL(Manipur) Issued On : 2020-02-14(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-15	0.0	24.0	8.0	78	36	4.6	112	0
2020-02-16	0.0	24.0	9.0	74	30	4.6	110	0
2020-02-17	0.0	24.0	10.0	72	31	4.4	110	0
2020-02-18	0.0	25.0	12.0	72	28	4.3	112	0
2020-02-19	0.0	25.0	12.0	59	22	5.0	109	0

Weather Summary/Alert:

Likely to have no rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 25-30 oC & mini. temp. 12-14 oC. Relative humidity may reach upto 78% (max) & 38 % (min). The average wind speed may be 4.3-4.8 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 26-30 oC amadi ingbagina 12-14 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 78 % amadi nungthildana 38 % oiraba yai. Nungsit sitpigi changna pung amada 4.3-4.8km oiraba yai.

General Advisory:

Monitor regularly insect pest infestation in crops. Mulching may be done in the crop field using straw/ dried leaves which is not infected by insect-pest to preserve soil moisture for the crop. Keep birds/ animals under your jurisdiction and avoid contact with migratory or wild birds/ animals.

Loufamda laina tilna taba ningthina toina yengsinbiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Sha uchek yariba makhei sennajaba mafamda thadokpiyu amadi lamlakki sha-uchek/ mapandagi paisinlakpa uchekka samnadanaba shinbiyu.

SMS Advisory:

Insect pest sprays may be given before rains. As weather is changing proper care of sheds, food and drinking water must be taken & administer vaccines in time.

Nong laktingaida laina tilnagi hidak kaphthokpiyu. Aying asa onrakpada sha uchek naraktanaba makol, ishing amadi machinjak afaba pibiduna matam chana hidak thabiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
TOMATO	Tomato nurseries should be raised under low cost polyhouse as the temperature is very low in open field. - Nursery bed of 0.75m x 4m may be prepared and mix Carbendazim @ 5g/ 3m2 area, keep the bed covered with gunny bag for at least 24 hrs. - Sow seeds @ 10g/ 3m2 area keep the bed covered with gunny bag. Seeds will germinate in 4-5 days.

Crop(Varieties)	Crop Specific Advisory
	- Ingba matamda polyhouse makhada khamen ashinbi chaar hunbiyu. - 0.75m x 4m pak-chaoba nursery bed samba loisanlakpada 3m2 khudingda 5g gi changda Carbendazim happaga bora na kupsinlaga pung 24 thambiyu. - 3m2 pak-chaoba nursery bed ta 10g gi changda maru hidak totlaga hunbibagi matungda bora kupsinlaga thambiyu. Numit 4-5ni gi manungda maton thokpagi chang yengbiraga bora louthokpiba yarani.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
POTATO	- For seed: • Strict supervision on removal of unwanted plant type should be maintained. • Maturity days should be strictly maintained as per variety. - Harvest the crop after 10-15 days after haulm cutting. - Maru oinaba lokpige hairabadi: • Masak matou manadaba pambi makhal khandokpiyu.. • Alu na chaoba mapung faba matam variety makhal gi leijaba numit shurakpagi matungda lokpiyu. - Alu loktoubu numit 10-15ni watlingaeida makhok khara khakaga pambi kakaga thambiyu.
FRENCH BEAN	- Sowing of French bean may be done in prepared land. Seed rate for dwarf varieties like Pant Anupam & Arka Komal is 65kg/ha (spacing is 30x45 cm from seed to seed and row to row) and 25-30kg/ha for pole-type (spacing is 60x60cm with 2-3 seeds per hill). Treat seeds with Carbendazim @2g/kg seeds before sowing. - Give light irrigation after 2-3 days of sowing. - Koli hawai thanaba loufam semjinbiba yarani. Nemna panbi variety Pant Anupam amadi Arka Komal pari amada thanaba 65 kg maru lousinbiraga 30x45 cm lapna thabiyu. Pambi awangbagina 25-30 kg changani amadi 60x60cm gi changda lapna makhul amada 2-3 maru thabiyu. - Thadringeida maru kg amada 2g gi changda Carbendazim totpiyu amadi thabagi matung leibak leinil marang kaina leinaba nong churaktagadi ishing khara chaibiyu.



Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 14-02-2020

Weather Forecast of SENAPATI(Manipur) Issued On : 2020-02-14(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-15	0.0	21.0	5.0	73	33	5.2	255	1
2020-02-16	0.0	21.0	6.0	78	34	5.1	18	0
2020-02-17	0.0	21.0	8.0	80	29	5.2	22	0
2020-02-18	0.0	22.0	10.0	69	22	4.7	338	0
2020-02-19	0.0	22.0	10.0	52	20	5.1	22	0

Weather Summary/Alert:

Likely to have no rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 25-29°C & mini. temp. 11-13 °C. Relative humidity may reach upto 85 % (max) & 40 % (min). The average wind speed may be 4.6-5.0 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 25-29°C amadi ingbagina 11-13 °C oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 85 % amadi nungthildana 40 % oiraba yai. Nungsit sitpagi changna pung amada 4.6-5.0 km oiraba yai.

General Advisory:

Monitor regularly insect pest infestation in crops. Mulching may be done in the crop field using straw/ dried leaves which is not infected by insect-pest to preserve soil moisture for the crop. Keep birds/ animals under your jurisdiction and avoid contact with migratory or wild birds/ animals.

Loufamda laina tilna taba ningthina toina yengsinbiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Sha uchek yariba makhei sennajaba mafamda thadokpiyu amadi lamlakki sha-uchek/ mapandagi paisinlakpa uchekka samnadanaba shinbiyu.

SMS Advisory:

Insect pest sprays may be given before rains. As weather is changing proper care of sheds, food and drinking water must be taken & administer vaccines in time.

Nong laktingaida laina tilnagi hidak kaphthokpiyu. Aying asa onrakpada sha uchek naraktanaba makol, ishing amadi machinjak afaba pibiduna matam chana hidak thabiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
TOMATO	Tomato nurseries should be raised under low cost polyhouse as the temperature is very low in open field. - Nursery bed of 0.75m x 4m may be prepared and mix Carbendazim @ 5g/ 3m ² area, keep the bed covered with gunny bag for at least 24 hrs. - Sow seeds @ 10g/ 3m ² area keep the bed covered with gunny bag. Seeds will germinate in 4-5 days.

Crop(Varieties)	Crop Specific Advisory
	- Ingba matamda polyhouse makhada khamen ashinbi chaar hunbiyu. - 0.75m x 4m pak-chaoba nursery bed samba loisanlakpada 3m2 khudingda 5g gi changda Carbendazim happaga bora na kupsinlaga pung 24 thambiyu. - 3m2 pak-chaoba nursery bed ta 10g gi changda maru hidak totlaga hunbibagi matungda bora kupsinlaga thambiyu. Numit 4-5ni gi manungda maton thokpaga chang yengbiraga bora louthokpiba yarani.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
POTATO	- For seed: • Strict supervision on removal of unwanted plant type should be maintained. • Maturity days should be strictly maintained as per variety. - Harvest the crop after 10-15 days after haulm cutting - Maru oinaba lokpige hairabadi: • Masak matou manadaba pambi makhal khandokpiyu.. • Alu na chaoba mapung faba matam variety makhal gi leijaba numit shurakpaga matungda lokpiyu. - Alu loktoubu numit 10-15ni watlingaeida makhok khara khakaga pambi kakaga thambiyu.
FRENCH BEAN	- Sowing of French bean may be done in prepared land. Seed rate for dwarf varieties like Pant Anupam & Arka Komal is 65kg/ha (spacing is 30x45 cm from seed to seed and row to row) and 25-30kg/ha for pole-type (spacing is 60x60cm with 2-3 seeds per hill). Treat seeds with Carbendazim @2g/kg seeds before sowing. - Give light irrigation after 2-3 days of sowing. - Koli hawai thanaba loufam semjinbiba yarani. Nemna panbi variety Pant Anupam amadi Arka Komal pari amada thanaba 65 kg maru lousinbiraga 30x45 cm lapna thabiyu. Pambi awangbagina 25-30 kg changani amadi 60x60cm gi changda lapna makhul amada 2-3 maru thabiyu. - Thadringeida maru kg amada 2g gi changda Carbendazim totpiyu amadi thabagi matung leibak leinil marang kaina leinaba nong churaktagadi ishing khara chaibiyu.



Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 14-02-2020

Weather Forecast of TAMENGLONG(Manipur) Issued On : 2020-02-14(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-15	0.0	22.0	7.0	74	33	4.6	68	1
2020-02-16	0.0	22.0	8.0	73	32	4.9	71	0
2020-02-17	0.0	22.0	9.0	60	28	4.9	68	0
2020-02-18	0.0	23.0	10.0	69	21	4.9	68	0
2020-02-19	0.0	23.0	10.0	49	22	5.3	68	0

Weather Summary/Alert:

Likely to have no rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 25-29 oC & mini. temp. 12-14 oC. Relative humidity may reach upto 79% (max) & 43 % (min). The average wind speed may be 4.8-5.1 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 25-29 oC amadi ingbagina 12-14 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 79% amadi nungthildana 43 % oiraba yai. Nungsit sitpagi changna pung amada 4.8-5.1 km oiraba yai.

General Advisory:

Monitor regularly insect pest infestation in crops. Mulching may be done in the crop field using straw/ dried leaves which is not infected by insect-pest to preserve soil moisture for the crop. Keep birds/ animals under your jurisdiction and avoid contact with migratory or wild birds/ animals.

Loufamda laina tilna taba ningthina toina yengsinbiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Sha uchek yariba makhei sennajaba mafamda thadokpiyu amadi lamlakki sha-uchek/ mapandagi paisinlakpa ucheikka samnadanaba shinbiyu.

SMS Advisory:

Insect pest sprays may be given before rains. As weather is changing proper care of sheds, food and drinking water must be taken & administer vaccines in time.

Nong laktingaida laina tilnagi hidak kaphthokpiyu. Aying asa onrakpada sha uchek naraktanaba makol, ishing amadi machinjak afaba pibiduna matam chana hidak thabiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
TOMATO	Tomato nurseries should be raised under low cost polyhouse as the temperature is very low in open field. - Nursery bed of 0.75m x 4m may be prepared and mix Carbendazim @ 5g/ 3m2 area, keep the bed covered with gunny bag for at least 24 hrs. - Sow seeds @ 10g/ 3m2 area keep the bed covered with gunny bag. Seeds will germinate in 4-5 days.

Crop(Varieties)	Crop Specific Advisory
	- Ingba matamda polyhouse makhada khamen ashinbi chaar hunbiyu. - 0.75m x 4m pak-chaoba nursery bed samba loisanlakpada 3m2 khudingda 5g gi changda Carbendazim happaga bora na kupsinlaga pung 24 thambiyu. - 3m2 pak-chaoba nursery bed ta 10g gi changda maru hidak totlaga hunbibagi matungda bora kupsinlaga thambiyu. Numit 4-5ni gi manungda maton thokpagi chang yengbiraga bora louthokpiba yarani.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
POTATO	- For seed: • Strict supervision on removal of unwanted plant type should be maintained. • Maturity days should be strictly maintained as per variety. - Harvest the crop after 10-15 days after haulm cutting - Maru oinaba lokpige hairabadi: • Masak matou manadaba pambi makhal khandokpiyu.. • Alu na chaoba mapung faba matam variety makhal gi leijaba numit shurakpagi matungda lokpiyu. - Alu loktoubu numit 10-15ni watlingaeida makhok khara khakaga pambi kakaga thambiyu.
FRENCH BEAN	- Sowing of French bean may be done in prepared land. Seed rate for dwarf varieties like Pant Anupam & Arka Komal is 65kg/ha (spacing is 30x45 cm from seed to seed and row to row) and 25-30kg/ha for pole-type (spacing is 60x60cm with 2-3 seeds per hill). Treat seeds with Carbendazim @2g/kg seeds before sowing. - Give light irrigation after 2-3 days of sowing. - Koli hawai thanaba loufam semjinbiba yarani. Nemna panbi variety Pant Anupam amadi Arka Komal pari amada thanaba 65 kg maru lousinbiraga 30x45 cm lapna thabiyu. Pambi awangbagina 25-30 kg changani amadi 60x60cm gi changda lapna makhul amada 2-3 maru thabiyu. - Thadringeida maru kg amada 2g gi changda Carbendazim totpiyu amadi thabagi matung leibak leinil marang kaina leinaba nong churaktagadi ishing khara chaibiyu.



Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 14-02-2020

Weather Forecast of TENGNOUPAL(Manipur) Issued On : 2020-02-14(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-15	0.0	24.0	8.0	78	39	6.1	290	1
2020-02-16	0.0	24.0	9.0	72	30	6.6	292	0
2020-02-17	0.0	24.0	10.0	74	33	4.2	289	0
2020-02-18	0.0	25.0	12.0	74	31	5.7	290	0
2020-02-19	0.0	25.0	12.0	64	20	8.2	292	0

Weather Summary/Alert:

Likely to have no rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 25-30 oC & mini. temp. 13-14 oC. Relative humidity may reach upto 82 % (max) & 44 % (min). The average wind speed may be 4.1-7.5 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 25-30 oC amadi ingbagina 13-14 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 82 % amadi nungthildana 44 % oiraba yai. Nungsit sitpagi changna pung amada 4.4-7.5 km oiraba yai.

General Advisory:

Monitor regularly insect pest infestation in crops. Mulching may be done in the crop field using straw/ dried leaves which is not infected by insect-pest to preserve soil moisture for the crop. Keep birds/ animals under your jurisdiction and avoid contact with migratory or wild birds/ animals.

Loufamda laina tilna taba ningthina toina yengsinbiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Sha uchek yariba makhei sennajaba mafamda thadokpiyu amadi lamlakki sha-uchek/ mapandagi paisinlakpa uchekka samnadanaba shinbiyu.

SMS Advisory:

Insect pest sprays may be given before rains. As weather is changing proper care of sheds, food and drinking water must be taken & administer vaccines in time.

Nong laktingaida laina tilnagi hidak kaphthokpiyu. Aying asa onrakpada sha uchek naraktanaba makol, ishing amadi machinjak afaba pibiduna matam chana hidak thabiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
TOMATO	Tomato nurseries should be raised under low cost polyhouse as the temperature is very low in open field. - Nursery bed of 0.75m x 4m may be prepared and mix Carbendazim @ 5g/ 3m2 area, keep the bed covered with gunny bag for at least 24 hrs. - Sow seeds @ 10g/ 3m2 area keep the bed covered with gunny bag. Seeds will germinate in 4-5 days.

Crop(Varieties)	Crop Specific Advisory
	- Ingba matamda polyhouse makhada khamen ashinbi chaar hunbiyu. - 0.75m x 4m pak-chaoba nursery bed samba loisanlakpada 3m2 khudingda 5g gi changda Carbendazim happaga bora na kupsinlaga pung 24 thambiyu. - 3m2 pak-chaoba nursery bed ta 10g gi changda maru hidak totlaga hunbibagi matungda bora kupsinlaga thambiyu. Numit 4-5ni gi manungda maton thokpagi chang yengbiraga bora louthokpiba yarani.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
POTATO	- For seed: • Strict supervision on removal of unwanted plant type should be maintained. • Maturity days should be strictly maintained as per variety. - Harvest the crop after 10-15 days after haulm cutting - Maru oinaba lokpige hairabadi: • Masak matou manadaba pambi makhal khandokpiyu.. • Alu na chaoba mapung faba matam variety makhal gi leijaba numit shurakpagi matungda lokpiyu. - Alu loktoubu numit 10-15ni watlingaeida makhok khara khakaga pambi kakaga thambiyu.
FRENCH BEAN	- Sowing of French bean may be done in prepared land. Seed rate for dwarf varieties like Pant Anupam & Arka Komal is 65kg/ha (spacing is 30x45 cm from seed to seed and row to row) and 25-30kg/ha for pole-type (spacing is 60x60cm with 2-3 seeds per hill). Treat seeds with Carbendazim @2g/kg seeds before sowing. - Give light irrigation after 2-3 days of sowing. - Koli hawai thanaba loufam semjinbiba yarani. Nemna panbi variety Pant Anupam amadi Arka Komal pari amada thanaba 65 kg maru lousinbiraga 30x45 cm lapna thabiyu. Pambi awangbagina 25-30 kg changani amadi 60x60cm gi changda lapna makhul amada 2-3 maru thabiyu. - Thadringeida maru kg amada 2g gi changda Carbendazim totpiyu amadi thabagi matung leibak leinil marang kaina leinaba nong churaktagadi ishing khara chaibiyu.



Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 14-02-2020

Weather Forecast of THOUBAL(Manipur) Issued On : 2020-02-14(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-15	0.0	25.0	10.0	79	40	7.5	279	1
2020-02-16	0.0	25.0	10.0	76	35	7.0	22	0
2020-02-17	0.0	25.0	12.0	73	29	3.9	338	0
2020-02-18	0.0	26.0	14.0	81	25	5.5	290	0
2020-02-19	0.0	26.0	14.0	63	24	8.5	304	0

Weather Summary/Alert:

Likely to have no rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 26-30 oC & mini. temp. 12-13 oC. Relative humidity may reach upto 85 % (max) & 45 % (min). The average wind speed may be 4.0-7.9 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 26-30 oC amadi ingbagina 12-13 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 85 % amadi nungthildana 43 % oiraba yai. Nungsit sitpigi changna pung amada 4.0-7.9 km oiraba yai.

General Advisory:

Monitor regularly insect pest infestation in crops. Mulching may be done in the crop field using straw/ dried leaves which is not infected by insect-pest to preserve soil moisture for the crop. Keep birds/ animals under your jurisdiction and avoid contact with migratory or wild birds/ animals.

Loufamda laina tilna taba ningthina toina yengsinbiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Sha uchek yariba makhei sennajaba mafamda thadokpiyu amadi lamlakki sha-uchek/ mapandagi paisinlakpa uchekka samnadanaba shinbiyu.

SMS Advisory:

Insect pest sprays may be given before rains. As weather is changing proper care of sheds, food and drinking water must be taken & administer vaccines in time.

Nong laktingaida laina tilnagi hidak kaphthokpiyu. Aying asa onrakpada sha uchek naraktanaba makol, ishing amadi machinjak afaba pibiduna matam chana hidak thabiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
TOMATO	Tomato nurseries should be raised under low cost polyhouse as the temperature is very low in open field. - Nursery bed of 0.75m x 4m may be prepared and mix Carbendazim @ 5g/ 3m2 area, keep the bed covered with gunny bag for at least 24 hrs. - Sow seeds @ 10g/ 3m2 area keep the bed covered with gunny bag. Seeds will germinate in 4-5 days.

Crop(Varieties)	Crop Specific Advisory
	- Ingba matamda polyhouse makhada khamen ashinbi chaar hunbiyu. - 0.75m x 4m pak-chaoba nursery bed samba loisanlakpada 3m2 khudingda 5g gi changda Carbendazim happaga bora na kupsinlaga pung 24 thambiyu. - 3m2 pak-chaoba nursery bed ta 10g gi changda maru hidak totlaga hunbibagi matungda bora kupsinlaga thambiyu. Numit 4-5ni gi manungda maton thokpaga chang yengbiraga bora louthokpiba yarani.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
POTATO	- For seed: • Strict supervision on removal of unwanted plant type should be maintained. • Maturity days should be strictly maintained as per variety. - Harvest the crop after 10-15 days after haulm cutting. - Maru oinaba lokpige hairabadi: • Masak matou manadaba pambi makhal khandokpiyu.. • Alu na chaoba mapung faba matam variety makhal gi leijaba numit shurakpaga matungda lokpiyu. - Alu loktoubu numit 10-15ni watlingaeida makhok khara khakaga pambi kakaga thambiyu.
FRENCH BEAN	- Sowing of French bean may be done in prepared land. Seed rate for dwarf varieties like Pant Anupam & Arka Komal is 65kg/ha (spacing is 30x45 cm from seed to seed and row to row) and 25-30kg/ha for pole-type (spacing is 60x60cm with 2-3 seeds per hill). Treat seeds with Carbendazim @2g/kg seeds before sowing. - Give light irrigation after 2-3 days of sowing. - Koli hawai thanaba loufam semjinbiba yarani. Nemna panbi variety Pant Anupam amadi Arka Komal pari amada thanaba 65 kg maru lousinbiraga 30x45 cm lapna thabiyu. Pambi awangbagina 25-30 kg changani amadi 60x60cm gi changda lapna makhul amada 2-3 maru thabiyu. - Thadringeida maru kg amada 2g gi changda Carbendazim totpiyu amadi thabagi matung leibak leinil marang kaina leinaba nong churaktagadi ishing khara chaibiyu.



Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 14-02-2020

Weather Forecast of UKHRUL(Manipur) Issued On : 2020-02-14(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-15	0.0	22.0	7.0	81	38	5.1	207	2
2020-02-16	0.0	22.0	8.0	67	34	5.7	22	1
2020-02-17	0.0	22.0	9.0	77	30	5.6	22	0
2020-02-18	0.0	23.0	10.0	69	27	5.5	117	0
2020-02-19	0.0	23.0	10.0	43	17	7.0	22	0

Weather Summary/Alert:

Likely to have no rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 25-30 oC & mini. temp. 12-14 oC. Relative humidity may reach upto 88 % (max) & 45 % (min). The average wind speed may be 5.0-6.6 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 25-30 oC amadi ingbagina 12-14 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 88 % amadi nungthildana 45 % oiraba yai. Nungsit sitpagi changna pung amada 5.0-6.6 km oiraba yai.

General Advisory:

Monitor regularly insect pest infestation in crops. Mulching may be done in the crop field using straw/ dried leaves which is not infected by insect-pest to preserve soil moisture for the crop. Keep birds/ animals under your jurisdiction and avoid contact with migratory or wild birds/ animals.

Loufamda laina tilna taba ningthina toina yengsinbiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Sha uchek yariba makhei sennajaba mafamda thadokpiyu amadi lamlakki sha-uchek/ mapandagi paisinlakpa ucheikka samnadanaba shinbiyu.

SMS Advisory:

Insect pest sprays may be given before rains. As weather is changing proper care of sheds, food and drinking water must be taken & administer vaccines in time.

Nong laktingaida laina tilnagi hidak kaphthokpiyu. Aying asa onrakpada sha uchek naraktanaba makol, ishing amadi machinjak afaba pibiduna matam chana hidak thabiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
TOMATO	Tomato nurseries should be raised under low cost polyhouse as the temperature is very low in open field. - Nursery bed of 0.75m x 4m may be prepared and mix Carbendazim @ 5g/ 3m2 area, keep the bed covered with gunny bag for at least 24 hrs. - Sow seeds @ 10g/ 3m2 area keep the bed covered with gunny bag. Seeds will germinate in 4-5 days.

Crop(Varieties)	Crop Specific Advisory
	- Ingba matamda polyhouse makhada khamen ashinbi chaar hunbiyu. - 0.75m x 4m pak-chaoba nursery bed samba loisanlakpada 3m2 khudingda 5g gi changda Carbendazim happaga bora na kupsinlaga pung 24 thambiyu. - 3m2 pak-chaoba nursery bed ta 10g gi changda maru hidak totlaga hunbibagi matungda bora kupsinlaga thambiyu. Numit 4-5ni gi manungda maton thokpaga chang yengbiraga bora louthokpiba yarani.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
POTATO	- For seed: • Strict supervision on removal of unwanted plant type should be maintained. • Maturity days should be strictly maintained as per variety. - Harvest the crop after 10-15 days after haulm cutting. - Maru oinaba lokpige hairabadi: • Masak matou manadaba pambi makhal khandokpiyu.. • Alu na chaoba mapung faba matam variety makhal gi leijaba numit shurakpaga matungda lokpiyu. - Alu loktoubu numit 10-15ni watlingaeida makhok khara khakaga pambi kakaga thambiyu.
FRENCH BEAN	- Sowing of French bean may be done in prepared land. Seed rate for dwarf varieties like Pant Anupam & Arka Komal is 65kg/ha (spacing is 30x45 cm from seed to seed and row to row) and 25-30kg/ha for pole-type (spacing is 60x60cm with 2-3 seeds per hill). Treat seeds with Carbendazim @2g/kg seeds before sowing. - Give light irrigation after 2-3 days of sowing. - Koli hawai thanaba loufam semjinbiba yarani. Nemna panbi variety Pant Anupam amadi Arka Komal pari amada thanaba 65 kg maru lousinbiraga 30x45 cm lapna thabiyu. Pambi awangbagina 25-30 kg changani amadi 60x60cm gi changda lapna makhul amada 2-3 maru thabiyu. - Thadringeida maru kg amada 2g gi changda Carbendazim totpiyu amadi thabagi matung leibak leinil marang kaina leinaba nong churaktagadi ishing khara chaibiyu.



Gramin Krishi Mausam Sewa

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 14-02-2020

Weather Forecast of WANGOI Block in IMPHAL-WEST(Manipur) Issued On :2020-02-14(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-15	0.0	25.5	12.4	78	44	9.0	270	1
2020-02-16	0.0	25.9	13.0	85	37	6.0	345	0
2020-02-17	0.0	29.5	13.5	70	27	4.0	360	0
2020-02-18	0.0	30.0	12.7	81	24	5.0	287	0
2020-02-19	0.0	28.6	11.9	67	26	6.0	293	0

Weather Summary/Alert:

Likely to have no rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 25.5-30 oC & mini. temp. 11.9-13.5 oC. Relative humidity may reach upto 85 % (max) & 44 % (min). The average wind speed may be 4-9 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 25.5-30 oC amadi ingbagina 11.9-13.5 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 85 % amadi nungthildana 44 % oiraba yai. Nungsit sitpagi changna pung amada 4-9 km oiraba yai.

General Advisory:

Monitor regularly insect pest infestation in crops. Mulching may be done in the crop field using straw/ dried leaves which is not infected by insect-pest to preserve soil moisture for the crop. Keep birds/ animals under your jurisdiction and avoid contact with migratory or wild birds/ animals.

Loufamda laina tilna taba ningthina toina yengsinbiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Sha uchek yariba makhei sennajaba mafamda thadokpiyu amadi lamlakki sha-uchek/ mapandagi paisinlakpa uchekka samnadanaba shinbiyu.

SMS Advisory:

Insect pest sprays may be given before rains. As weather is changing proper care of sheds, food and drinking water must be taken & administer vaccines in time.

Nong laktingaida laina tilnagi hidak kaphthokpiyu. Aying asa onrakpada sha uchek naraktanaba makol, ishing amadi machinjak afaba pibiduna matam chana hidak thabiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
TOMATO	Tomato nurseries should be raised under low cost polyhouse as the temperature is very low in open field. - Nursery bed of 0.75m x 4m may be prepared and mix Carbendazim @ 5g/ 3m2 area, keep the bed covered with gunny bag for at least 24 hrs. - Sow seeds @ 10g/ 3m2 area keep the bed covered with gunny bag. Seeds will germinate in 4-5 days.
	- Ingba matamda polyhouse makhada khamen ashinbi chaar hunbiyu. - 0.75m x 4m pak-chaoba nursery bed samba loisanlakpada 3m2 khudingda 5g gi changda Carbendazim happaga bora na kupsinlaga pung 24 thambiyu. - 3m2 pak-chaoba nursery bed ta 10g gi changda maru hidak totlaga hunbibagi matungda bora kupsinlaga thambiyu. Numit 4-5ni gi manungda maton thokpagi chang yengbiraga bora louthokpiba yarani.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
POTATO	- For seed: • Strict supervision on removal of unwanted plant type should be maintained. • Maturity days should be strictly maintained as per variety. - Harvest the crop after 10-15 days after haulm cutting
	- Maru oinaba lokpige hairabadi: • Masak matou manadaba pambi makhal khandokpiyu.. • Alu na chaoba mapung faba matam variety makhal gi leijaba numit shurakpagi matungda lokpiyu. - Alu loktoubu numit 10-15ni watlingaeida makhok khara khakaga pambi kakaga thambiyu.
FRENCH BEAN	- Sowing of French bean may be done in prepared land. Seed rate for dwarf varieties like Pant Anupam & Arka Komal is 65kg/ha (spacing is 30x45 cm from seed to seed and row to row) and 25-30kg/ha for pole-type (spacing is 60x60cm with 2-3 seeds per hill). Treat seeds with Carbendazim @2g/kg seeds before sowing. - Give light irrigation after 2-3 days of sowing.
	- Koli hawai thanaba loufam semjinbiba yarani. Nemna panbi variety Pant Anupam amadi Arka Komal pari amada thanaba 65 kg maru lousinbiraga 30x45 cm lapna thabiyu. Pambi awangbagina 25-30 kg changani amadi 60x60cm gi changda lapna makhul amada 2-3 maru thabiyu. - Thadringaeida maru kg amada 2g gi changda Carbendazim totpiyu amadi thabagi matung leibak leinil marang kaina leinaba nong churaktagadi ishing khara chaibiyu.



Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 14-02-2020

Weather Forecast of BISHNUPUR(Manipur) Issued On : 2020-02-14(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-15	0.0	25.0	10.0	84	38	4.3	261	0
2020-02-16	0.0	25.0	10.0	80	34	3.6	68	0
2020-02-17	0.0	25.0	12.0	75	28	2.6	68	0
2020-02-18	0.0	26.0	14.0	82	23	2.8	270	0
2020-02-19	0.0	26.0	14.0	65	23	4.0	292	0

Weather Summary/Alert:

Likely to have no rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 24-31 oC & mini. temp. 12-14 oC. Relative humidity may reach upto 89% (max) & 24% (min). The average wind speed may be 2.7-4.9 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 24-31 oC amadi ingbagina 12-14 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 89% amadi nungthildana 24% oiraba yai. Nungsit sitpigi changna pung amada 2.7-4.9 km oiraba yai.

General Advisory:

Monitor regularly insect pest infestation in crops. Mulching may be done in the crop field using straw/ dried leaves which is not infected by insect-pest to preserve soil moisture for the crop. Keep birds/ animals under your jurisdiction and avoid contact with migratory or wild birds/ animals.

Loufamda laina tilna taba ningthina toina yengsinbiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Sha uchek yariba makhei sennajaba mafamda thadokpiyu amadi lamlakki sha-uchek/ mapandagi paisinlakpa uchekka samnadanaba shinbiyu.

SMS Advisory:

Insect pest sprays may be given before rains. As weather is changing proper care of sheds, food and drinking water must be taken & administer vaccines in time.

Nong laktingaida laina tilnagi hidak kaphthokpiyu. Aying asa onrakpada sha uchek naraktanaba makol, ishing amadi machinjak afaba pibiduna matam chana hidak thabiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
TOMATO	Tomato nurseries should be raised under low cost polyhouse as the temperature is very low in open field. - Nursery bed of 0.75m x 4m may be prepared and mix Carbendazim @ 5g/ 3m ² area, keep the bed covered with gunny bag for at least 24 hrs. - Sow seeds @ 10g/ 3m ² area keep the bed covered with gunny bag. Seeds will germinate in 4-5 days.

Crop(Varieties)	Crop Specific Advisory
	- Ingba matamda polyhouse makhada khamen ashinbi chaar hunbiyu. - 0.75m x 4m pak-chaoba nursery bed samba loisanlakpada 3m2 khudingda 5g gi changda Carbendazim happaga bora na kupsinlaga pung 24 thambiyu. - 3m2 pak-chaoba nursery bed ta 10g gi changda maru hidak totlaga hunbibagi matungda bora kupsinlaga thambiyu. Numit 4-5ni gi manungda maton thokpaga chang yengbiraga bora louthokpiba yarani.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
POTATO	- For seed: • Strict supervision on removal of unwanted plant type should be maintained. • Maturity days should be strictly maintained as per variety. - Harvest the crop after 10-15 days after haulm cutting - Maru oinaba lokpige hairabadi: • Masak matou manadaba pambi makhal khandokpiyu.. • Alu na chaoba mapung faba matam variety makhal gi leijaba numit shurakpaga matungda lokpiyu. - Alu loktoubu numit 10-15ni watlingaeida makhok khara khakaga pambi kakaga thambiyu.
FRENCH BEAN	- Sowing of French bean may be done in prepared land. Seed rate for dwarf varieties like Pant Anupam & Arka Komal is 65kg/ha (spacing is 30x45 cm from seed to seed and row to row) and 25-30kg/ha for pole-type (spacing is 60x60cm with 2-3 seeds per hill). Treat seeds with Carbendazim @2g/kg seeds before sowing. - Give light irrigation after 2-3 days of sowing. - Koli hawai thanaba loufam semjinbiba yarani. Nemna panbi variety Pant Anupam amadi Arka Komal pari amada thanaba 65 kg maru lousinbiraga 30x45 cm lapna thabiyu. Pambi awangbagina 25-30 kg changani amadi 60x60cm gi changda lapna makhul amada 2-3 maru thabiyu. - Thadringeida maru kg amada 2g gi changda Carbendazim totpiyu amadi thabagi matung leibak leinil marang kaina leinaba nong churaktagadi ishing khara chaibiyu.



Gramin Krishi Mausam Sewa
India Meteorological Department
ICAR Complex Branch
Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 14-02-2020

Weather Forecast of CHANDEL(Manipur) Issued On : 2020-02-14(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-15	0.0	24.0	8.0	77	36	5.8	292	0
2020-02-16	0.0	24.0	9.0	76	31	5.8	292	0
2020-02-17	0.0	24.0	10.0	76	29	3.3	259	0
2020-02-18	0.0	25.0	12.0	75	24	4.8	285	0
2020-02-19	0.0	25.0	12.0	65	20	7.6	292	0

Weather Summary/Alert:

Likely to have no rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 25-30 oC & mini. temp. 13 oC. Relative humidity may reach upto 85% (max) & 21% (min). The average wind speed may be 3.1-7.4 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 25-30 oC amadi ingbagina 13 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 85% amadi nunghildana 21% oiraba yai. Nungsit sitpigi changna pung amada 3.1-7.4 km oiraba yai.

General Advisory:

Monitor regularly insect pest infestation in crops. Mulching may be done in the crop field using straw/ dried leaves which is not infected by insect-pest to preserve soil moisture for the crop. Keep birds/ animals under your jurisdiction and avoid contact with migratory or wild birds/ animals.

Loufamda laina tilna taba ningthina toina yengsinbiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Sha uchek yariba makhei sennajaba mafamda thadokpiyu amadi lamlakki sha-uchek/ mapandagi paisinlakpa uchekka samnadanaba shinbiyu.

SMS Advisory:

Insect pest sprays may be given before rains. As weather is changing proper care of sheds, food and drinking water must be taken & administer vaccines in time.

Nong laktingaida laina tilnagi hidak kaphthokpiyu. Aying asa onrakpada sha uchek naraktanaba makol, ishing amadi machinjak afaba pibiduna matam chana hidak thabiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
TOMATO	Tomato nurseries should be raised under low cost polyhouse as the temperature is very low in open field. - Nursery bed of 0.75m x 4m may be prepared and mix Carbendazim @ 5g/ 3m ² area, keep the bed covered with gunny bag for at least 24 hrs. - Sow seeds @ 10g/ 3m ² area keep the bed covered with gunny bag. Seeds will germinate in 4-5 days.

Crop(Varieties)	Crop Specific Advisory
	- Ingba matamda polyhouse makhada khamen ashinbi chaar hunbiyu. - 0.75m x 4m pak-chaoba nursery bed samba loisanlakpada 3m2 khudingda 5g gi changda Carbendazim happaga bora na kupsinlaga pung 24 thambiyu. - 3m2 pak-chaoba nursery bed ta 10g gi changda maru hidak totlaga hunbibagi matungda bora kupsinlaga thambiyu. Numit 4-5ni gi manungda maton thokpagi chang yengbiraga bora louthokpiba yarani.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
POTATO	- For seed: • Strict supervision on removal of unwanted plant type should be maintained. • Maturity days should be strictly maintained as per variety. - Harvest the crop after 10-15 days after haulm cutting - Maru oinaba lokpige hairabadi: • Masak matou manadaba pambi makhal khandokpiyu.. • Alu na chaoba mapung faba matam variety makhal gi leijaba numit shurakpagi matungda lokpiyu. - Alu loktoubu numit 10-15ni watlingaeida makhok khara khakaga pambi kakaga thambiyu.
FRENCH BEAN	- Sowing of French bean may be done in prepared land. Seed rate for dwarf varieties like Pant Anupam & Arka Komal is 65kg/ha (spacing is 30x45 cm from seed to seed and row to row) and 25-30kg/ha for pole-type (spacing is 60x60cm with 2-3 seeds per hill). Treat seeds with Carbendazim @2g/kg seeds before sowing. - Give light irrigation after 2-3 days of sowing. - Koli hawai thanaba loufam semjinbiba yarani. Nemna panbi variety Pant Anupam amadi Arka Komal pari amada thanaba 65 kg maru lousinbiraga 30x45 cm lapna thabiyu. Pambi awangbagina 25-30 kg changani amadi 60x60cm gi changda lapna makhul amada 2-3 maru thabiyu. - Thadringeida maru kg amada 2g gi changda Carbendazim totpiyu amadi thabagi matung leibak leinil marang kaina leinaba nong churaktagadi ishing khara chaibiyu.



Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 14-02-2020

Weather Forecast of CHURACHANDPUR(Manipur) Issued On : 2020-02-14(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-15	0.0	24.0	8.0	86	38	4.9	248	0
2020-02-16	0.0	24.0	9.0	83	35	4.4	153	0
2020-02-17	0.0	24.0	10.0	83	28	3.3	158	0
2020-02-18	0.0	25.0	12.0	79	22	4.2	207	0
2020-02-19	0.0	25.0	12.0	66	23	5.6	202	0

Weather Summary/Alert:

Likely to have no rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 24-31 oC & mini. temp. 12-13 oC. Relative humidity may reach upto 90 % (max) & 24% (min). The average wind speed may be 3.4-5.8 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 24-31 oC amadi ingbagina 12-13 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 90 % amadi nungthildana 24% oiraba yai. Nungsit sitpagi changna pung amada 3.4-5.8 km oiraba yai.

General Advisory:

Monitor regularly insect pest infestation in crops. Mulching may be done in the crop field using straw/ dried leaves which is not infected by insect-pest to preserve soil moisture for the crop. Keep birds/ animals under your jurisdiction and avoid contact with migratory or wild birds/ animals.

Loufamda laina tilna taba ningthina toina yengsinbiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Sha uchek yariba makhei sennajaba mafamda thadokpiyu amadi lamlakki sha-uchek/ mapandagi paisinlakpa uchekka samnadanaba shinbiyu.

SMS Advisory:

Insect pest sprays may be given before rains. As weather is changing proper care of sheds, food and drinking water must be taken & administer vaccines in time.

Nong laktingaida laina tilnagi hidak kaphokpiyu. Aying asa onrakpada sha uchek naraktanaba makol, ishing amadi machinjak afaba pibiduna matam chana hidak thabiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
TOMATO	Tomato nurseries should be raised under low cost polyhouse as the temperature is very low in open field. - Nursery bed of 0.75m x 4m may be prepared and mix Carbendazim @ 5g/ 3m ² area, keep the bed covered with gunny bag for at least 24 hrs. - Sow seeds @ 10g/ 3m ² area keep the bed covered with gunny bag. Seeds will germinate in 4-5 days.

Crop(Varieties)	Crop Specific Advisory
	- Ingba matamda polyhouse makhada khamen ashinbi chaar hunbiyu. - 0.75m x 4m pak-chaoba nursery bed samba loisanlakpada 3m2 khudingda 5g gi changda Carbendazim happaga bora na kupsinlaga pung 24 thambiyu. - 3m2 pak-chaoba nursery bed ta 10g gi changda maru hidak totlaga hunbibagi matungda bora kupsinlaga thambiyu. Numit 4-5ni gi manungda maton thokpaga chang yengbiraga bora louthokpiba yarani.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
POTATO	- For seed: • Strict supervision on removal of unwanted plant type should be maintained. • Maturity days should be strictly maintained as per variety. - Harvest the crop after 10-15 days after haulm cutting - Maru oinaba lokpige hairabadi: • Masak matou manadaba pambi makhal khandokpiyu.. • Alu na chaoba mapung faba matam variety makhal gi leijaba numit shurakpaga matungda lokpiyu. - Alu loktoubu numit 10-15ni watlingaeida makhok khara khakaga pambi kakaga thambiyu.
FRENCH BEAN	- Sowing of French bean may be done in prepared land. Seed rate for dwarf varieties like Pant Anupam & Arka Komal is 65kg/ha (spacing is 30x45 cm from seed to seed and row to row) and 25-30kg/ha for pole-type (spacing is 60x60cm with 2-3 seeds per hill). Treat seeds with Carbendazim @2g/kg seeds before sowing. - Give light irrigation after 2-3 days of sowing. - Koli hawai thanaba loufam semjinbiba yarani. Nemna panbi variety Pant Anupam amadi Arka Komal pari amada thanaba 65 kg maru lousinbiraga 30x45 cm lapna thabiyu. Pambi awangbagina 25-30 kg changani amadi 60x60cm gi changda lapna makhul amada 2-3 maru thabiyu. - Thadringeida maru kg amada 2g gi changda Carbendazim totpiyu amadi thabagi matung leibak leinil marang kaina leinaba nong churaktagadi ishing khara chaibiyu.



Gramin Krishi Mausam Sewa

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 14-02-2020

Weather Forecast of HAORANGSABAL Block in IMPHAL-WEST(Manipur) Issued On :2020-02-14(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-15	0.0	25.8	12.1	78	43	8.0	281	1
2020-02-16	0.0	26.0	13.0	86	37	6.0	360	0
2020-02-17	0.0	29.1	13.7	67	27	6.0	23	0
2020-02-18	0.0	29.8	12.7	79	23	6.0	334	0
2020-02-19	0.0	28.7	12.1	64	25	6.0	338	0

Weather Summary/Alert:

Likely to have no rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 25.8-29.8 oC & mini. temp. 12.1-13.7 oC. Relative humidity may reach upto 86 % (max) & 43 % (min). The average wind speed may be 6-8 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 25.8-29.8 oC amadi ingbagina 12.1-13.7 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 86 % amadi nungthildana 43 % oiraba yai. Nungsit sitpagi changna pung amada 6-8 km oiraba yai.

General Advisory:

Monitor regularly insect pest infestation in crops. Mulching may be done in the crop field using straw/ dried leaves which is not infected by insect-pest to preserve soil moisture for the crop. Keep birds/ animals under your jurisdiction and avoid contact with migratory or wild birds/ animals.

Loufamda laina tilna taba ningthina toina yengsinbiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Sha uchek yariba makhei sennajaba mafamda thadokpiyu amadi lamlakki sha-uchek/ mapandagi paisinlakpa ucheikka samnadanaba shinbiyu.

SMS Advisory:

Insect pest sprays may be given before rains. As weather is changing proper care of sheds, food and drinking water must be taken & administer vaccines in time.

Nong laktingaida laina tilnagi hidak kaphthokpiyu. Aying asa onrakpada sha uchek naraktanaba makol, ishing amadi machinjak afaba pibiduna matam chana hidak thabiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
TOMATO	Tomato nurseries should be raised under low cost polyhouse as the temperature is very low in open field. - Nursery bed of 0.75m x 4m may be prepared and mix Carbendazim @ 5g/ 3m2 area, keep the bed covered with gunny bag for at least 24 hrs. - Sow seeds @ 10g/ 3m2 area keep the bed covered with gunny bag. Seeds will germinate in 4-5 days.
	- Ingba matamda polyhouse makhada khamen ashinbi chaar hunbiyu. - 0.75m x 4m pak-chaoba nursery bed samba loisanlakpada 3m2 khudingda 5g gi changda Carbendazim happaga bora na kupsinlaga pung 24 thambiyu. - 3m2 pak-chaoba nursery bed ta 10g gi changda maru hidak totlaga hunbibagi matungda bora kupsinlaga thambiyu. Numit 4-5ni gi manungda maton thokpagi chang yengbiraga bora louthokpiba yarani.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
POTATO	- For seed: • Strict supervision on removal of unwanted plant type should be maintained. • Maturity days should be strictly maintained as per variety. - Harvest the crop after 10-15 days after haulm cutting - Maru oinaba lokpige hairabadi: • Masak matou manadaba pambi makhal khandokpiyu.. • Alu na chaoba mapung faba matam variety makhal gi leijaba numit shurakpagi matungda lokpiyu. - Alu loktoubu numit 10-15ni watlingaeida makhok khara khakaga pambi kakaga thambiyu.
FRENCH BEAN	- Sowing of French bean may be done in prepared land. Seed rate for dwarf varieties like Pant Anupam & Arka Komal is 65kg/ha (spacing is 30x45 cm from seed to seed and row to row) and 25-30kg/ha for pole-type (spacing is 60x60cm with 2-3 seeds per hill). Treat seeds with Carbendazim @2g/kg seeds before sowing. - Give light irrigation after 2-3 days of sowing. - Koli hawai thanaba loufam semjinbiba yarani. Nemna panbi variety Pant Anupam amadi Arka Komal pari amada thanaba 65 kg maru lousinbiraga 30x45 cm lapna thabiyu. Pambi awangbagina 25-30 kg changani amadi 60x60cm gi changda lapna makhul amada 2-3 maru thabiyu. - Thadringeida maru kg amada 2g gi changda Carbendazim totpiyu amadi thabagi matung leibak leinil marang kaina leinaba nong churaktagadi ishing khara chaibiyu.



Gramin Krishi Mausam Sewa

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 14-02-2020

Weather Forecast of HIYANGTHANG Block in IMPHAL-WEST(Manipur) Issued On :2020-02-14(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-15	0.0	25.9	12.3	78	43	8.0	283	1
2020-02-16	0.0	26.1	13.0	86	37	7.0	345	0
2020-02-17	0.0	29.3	13.6	68	27	5.0	26	0
2020-02-18	0.0	30.0	12.7	80	23	5.0	315	0
2020-02-19	0.0	28.8	11.9	65	25	6.0	252	0

Weather Summary/Alert:

Likely to have no rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 25.9-30 oC & mini. temp. 11.9-13.6 oC. Relative humidity may reach upto 86 % (max) & 43 % (min). The average wind speed may be 5-8 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 25.9-30 oC amadi ingbagina 11.9-13.6 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 86 % amadi nungthildana 43 % oiraba yai. Nungsit sitpagi changna pung amada 5-8 km oiraba yai.

General Advisory:

Monitor regularly insect pest infestation in crops. Mulching may be done in the crop field using straw/ dried leaves which is not infected by insect-pest to preserve soil moisture for the crop. Keep birds/ animals under your jurisdiction and avoid contact with migratory or wild birds/ animals.

Loufamda laina tilna taba ningthina toina yengsinbiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Sha uchek yariba makhei sennajaba mafamda thadokpiyu amadi lamlakki sha-uchek/ mapandagi paisinlakpa ucheikka samnadanaba shinbiyu.

SMS Advisory:

Insect pest sprays may be given before rains. As weather is changing proper care of sheds, food and drinking water must be taken & administer vaccines in time.

Nong laktingaida laina tilnagi hidak kaphthokpiyu. Aying asa onrakpada sha uchek naraktanaba makol, ishing amadi machinjak afaba pibiduna matam chana hidak thabiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
TOMATO	Tomato nurseries should be raised under low cost polyhouse as the temperature is very low in open field. - Nursery bed of 0.75m x 4m may be prepared and mix Carbendazim @ 5g/ 3m2 area, keep the bed covered with gunny bag for at least 24 hrs. - Sow seeds @ 10g/ 3m2 area keep the bed covered with gunny bag. Seeds will germinate in 4-5 days.
	- Ingba matamda polyhouse makhada khamen ashinbi chaar hunbiyu. - 0.75m x 4m pak-chaoba nursery bed samba loisanlakpada 3m2 khudingda 5g gi changda Carbendazim happaga bora na kupsinlaga pung 24 thambiyu. - 3m2 pak-chaoba nursery bed ta 10g gi changda maru hidak totlaga hunbibagi matungda bora kupsinlaga thambiyu. Numit 4-5ni gi manungda maton thokpagi chang yengbiraga bora louthokpiba yarani.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
POTATO	- For seed: • Strict supervision on removal of unwanted plant type should be maintained. • Maturity days should be strictly maintained as per variety. - Harvest the crop after 10-15 days after haulm cutting
	- Maru oinaba lokpige hairabadi: • Masak matou manadaba pambi makhal khandokpiyu.. • Alu na chaoba mapung faba matam variety makhal gi leijaba numit shurakpagi matungda lokpiyu. - Alu loktoubu numit 10-15ni watlingaeida makhok khara khakaga pambi kakaga thambiyu.
FRENCH BEAN	- Sowing of French bean may be done in prepared land. Seed rate for dwarf varieties like Pant Anupam & Arka Komal is 65kg/ha (spacing is 30x45 cm from seed to seed and row to row) and 25-30kg/ha for pole-type (spacing is 60x60cm with 2-3 seeds per hill). Treat seeds with Carbendazim @2g/kg seeds before sowing. - Give light irrigation after 2-3 days of sowing.
	- Koli hawai thanaba loufam semjinbiba yarani. Nemna panbi variety Pant Anupam amadi Arka Komal pari amada thanaba 65 kg maru lousinbiraga 30x45 cm lapna thabiyu. Pambi awangbagina 25-30 kg changani amadi 60x60cm gi changda lapna makhul amada 2-3 maru thabiyu. - Thadringeida maru kg amada 2g gi changda Carbendazim totpiyu amadi thabagi matung leibak leinil marang kaina leinaba nong churaktagadi ishing khara chaibiyu.



Gramin Krishi Mausam Sewa
India Meteorological Department
ICAR Complex Branch
Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 14-02-2020

Weather Forecast of IMPHAL-EAST(Manipur) Issued On : 2020-02-14(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-15	0.0	24.0	8.0	78	37	7.0	287	1
2020-02-16	0.0	24.0	9.0	72	33	7.1	26	0
2020-02-17	0.0	24.0	10.0	68	28	4.9	18	0
2020-02-18	0.0	25.0	12.0	76	24	5.3	326	0
2020-02-19	0.0	25.0	12.0	55	22	7.5	345	0

Weather Summary/Alert:

Likely to have no rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 26-30 oC & mini. temp. 12-14 oC. Relative humidity may reach upto 82 % (max) & 24% (min). The average wind speed may be 5.1-8.6 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 26-30 oC amadi ingbagina 12-14 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 82 % amadi nungthildana 24% oiraba yai. Nungsit sitpagi changna pung amada 5.1-8.6 km oiraba yai.

General Advisory:

Monitor regularly insect pest infestation in crops. Mulching may be done in the crop field using straw/ dried leaves which is not infected by insect-pest to preserve soil moisture for the crop. Keep birds/ animals under your jurisdiction and avoid contact with migratory or wild birds/ animals.

Loufamda laina tilna taba ningthina toina yengsinbiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Sha uchek yariba makhei sennajaba mafamda thadokpiyu amadi lamlakki sha-uchek/ mapandagi paisinlakpa uchekka samnadanaba shinbiyu.

SMS Advisory:

Insect pest sprays may be given before rains. As weather is changing proper care of sheds, food and drinking water must be taken & administer vaccines in time.

Nong laktingaida laina tilnagi hidak kaphthokpiyu. Aying asa onrakpada sha uchek naraktanaba makol, ishing amadi machinjak afaba pibiduna matam chana hidak thabiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
TOMATO	Tomato nurseries should be raised under low cost polyhouse as the temperature is very low in open field. - Nursery bed of 0.75m x 4m may be prepared and mix Carbendazim @ 5g/ 3m ² area, keep the bed covered with gunny bag for at least 24 hrs. - Sow seeds @ 10g/ 3m ² area keep the bed covered with gunny bag. Seeds will germinate in 4-5 days.

Crop(Varieties)	Crop Specific Advisory
	- Ingba matamda polyhouse makhada khamen ashinbi chaar hunbiyu. - 0.75m x 4m pak-chaoba nursery bed samba loisanlakpada 3m2 khudingda 5g gi changda Carbendazim happaga bora na kupsinlaga pung 24 thambiyu. - 3m2 pak-chaoba nursery bed ta 10g gi changda maru hidak totlaga hunbibagi matungda bora kupsinlaga thambiyu. Numit 4-5ni gi manungda maton thokpagi chang yengbiraga bora louthokpiba yarani.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
POTATO	- For seed: • Strict supervision on removal of unwanted plant type should be maintained. • Maturity days should be strictly maintained as per variety. - Harvest the crop after 10-15 days after haulm cutting - Maru oinaba lokpige hairabadi: • Masak matou manadaba pambi makhal khandokpiyu.. • Alu na chaoba mapung faba matam variety makhal gi leijaba numit shurakpagi matungda lokpiyu. - Alu loktoubu numit 10-15ni watlingaeida makhok khara khakaga pambi kakaga thambiyu.
FRENCH BEAN	- Sowing of French bean may be done in prepared land. Seed rate for dwarf varieties like Pant Anupam & Arka Komal is 65kg/ha (spacing is 30x45 cm from seed to seed and row to row) and 25-30kg/ha for pole-type (spacing is 60x60cm with 2-3 seeds per hill). Treat seeds with Carbendazim @2g/kg seeds before sowing. - Give light irrigation after 2-3 days of sowing. - Koli hawai thanaba loufam semjinbiba yarani. Nemna panbi variety Pant Anupam amadi Arka Komal pari amada thanaba 65 kg maru lousinbiraga 30x45 cm lapna thabiyu. Pambi awangbagina 25-30 kg changani amadi 60x60cm gi changda lapna makhul amada 2-3 maru thabiyu. - Thadringeida maru kg amada 2g gi changda Carbendazim totpiyu amadi thabagi matung leibak leinil marang kaina leinaba nong churaktagadi ishing khara chaibiyu.



Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 14-02-2020

Weather Forecast of IMPHAL-WEST(Manipur) Issued On : 2020-02-14(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-15	0.0	24.0	8.0	79	37	7.2	281	1
2020-02-16	0.0	24.0	9.0	74	34	6.9	22	0
2020-02-17	0.0	24.0	10.0	67	27	5.6	22	0
2020-02-18	0.0	25.0	12.0	75	22	5.8	338	0
2020-02-19	0.0	25.0	12.0	54	22	7.1	338	0

Weather Summary/Alert:

Likely to have no rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 26-30 oC & mini. temp. 12-14 oC. Relative humidity may reach upto 80 % (max) & 23 % (min). The average wind speed may be 5.8-8.1 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 26-30 oC amadi ingbagina 12-14 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 80 % amadi nungthildana 23 % oiraba yai. Nungsit sitpigi changna pung amada 5.8-8.1 km oiraba yai.

General Advisory:

Monitor regularly insect pest infestation in crops. Mulching may be done in the crop field using straw/ dried leaves which is not infected by insect-pest to preserve soil moisture for the crop. Keep birds/ animals under your jurisdiction and avoid contact with migratory or wild birds/ animals.

Loufamda laina tilna taba ningthina toina yengsinbiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Sha uchek yariba makhei sennajaba mafamda thadokpiyu amadi lamlakki sha-uchek/ mapandagi paisinlakpa uchekka samnadanaba shinbiyu.

SMS Advisory:

Insect pest sprays may be given before rains. As weather is changing proper care of sheds, food and drinking water must be taken & administer vaccines in time.

Nong laktingaida laina tilnagi hidak kaphthokpiyu. Aying asa onrakpada sha uchek naraktanaba makol, ishing amadi machinjak afaba pibiduna matam chana hidak thabiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
TOMATO	Tomato nurseries should be raised under low cost polyhouse as the temperature is very low in open field. - Nursery bed of 0.75m x 4m may be prepared and mix Carbendazim @ 5g/ 3m ² area, keep the bed covered with gunny bag for at least 24 hrs. - Sow seeds @ 10g/ 3m ² area keep the bed covered with gunny bag. Seeds will germinate in 4-5 days.

Crop(Varieties)	Crop Specific Advisory
	- Ingba matamda polyhouse makhada khamen ashinbi chaar hunbiyu. - 0.75m x 4m pak-chaoba nursery bed samba loisanlakpada 3m2 khudingda 5g gi changda Carbendazim happaga bora na kupsinlaga pung 24 thambiyu. - 3m2 pak-chaoba nursery bed ta 10g gi changda maru hidak totlaga hunbibagi matungda bora kupsinlaga thambiyu. Numit 4-5ni gi manungda maton thokpaga chang yengbiraga bora louthokpiba yarani.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
POTATO	- For seed: • Strict supervision on removal of unwanted plant type should be maintained. • Maturity days should be strictly maintained as per variety. - Harvest the crop after 10-15 days after haulm cutting - Maru oinaba lokpige hairabadi: • Masak matou manadaba pambi makhal khandokpiyu.. • Alu na chaoba mapung faba matam variety makhal gi leijaba numit shurakpaga matungda lokpiyu. - Alu loktoubu numit 10-15ni watlingaeida makhok khara khakaga pambi kakaga thambiyu.
FRENCH BEAN	- Sowing of French bean may be done in prepared land. Seed rate for dwarf varieties like Pant Anupam & Arka Komal is 65kg/ha (spacing is 30x45 cm from seed to seed and row to row) and 25-30kg/ha for pole-type (spacing is 60x60cm with 2-3 seeds per hill). Treat seeds with Carbendazim @2g/kg seeds before sowing. - Give light irrigation after 2-3 days of sowing. - Koli hawai thanaba loufam semjinbiba yarani. Nemna panbi variety Pant Anupam amadi Arka Komal pari amada thanaba 65 kg maru lousinbiraga 30x45 cm lapna thabiyu. Pambi awangbagina 25-30 kg changani amadi 60x60cm gi changda lapna makhul amada 2-3 maru thabiyu. - Thadringeida maru kg amada 2g gi changda Carbendazim totpiyu amadi thabagi matung leibak leinil marang kaina leinaba nong churaktagadi ishing khara chaibiyu.



Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 14-02-2020

Weather Forecast of JIRIBAM(Manipur) Issued On : 2020-02-14(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-15	0.0	24.0	8.0	76	35	4.8	68	0
2020-02-16	0.0	24.0	9.0	72	32	4.8	68	0
2020-02-17	0.0	24.0	10.0	68	30	4.3	68	0
2020-02-18	0.0	25.0	12.0	69	27	4.3	65	0
2020-02-19	0.0	25.0	12.0	58	23	5.3	65	0

Weather Summary/Alert:

Likely to have no rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 25-29 oC & mini. temp. 12-13 oC. Relative humidity may reach upto 75% (max) & 26 % (min). The average wind speed may be 4.3-4.8 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 25-29 oC amadi ingbagina 12-13 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 75% amadi nungthildana 26 % oiraba yai. Nungsit sitpagi changna pung amada 4.3-4.8 km oiraba yai.

General Advisory:

Monitor regularly insect pest infestation in crops. Mulching may be done in the crop field using straw/ dried leaves which is not infected by insect-pest to preserve soil moisture for the crop. Keep birds/ animals under your jurisdiction and avoid contact with migratory or wild birds/ animals.

Loufamda laina tilna taba ningthina toina yengsinbiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Sha uchek yariba makhei sennajaba mafamda thadokpiyu amadi lamlakki sha-uchek/ mapandagi paisinlakpa uchekka samnadanaba shinbiyu.

SMS Advisory:

Insect pest sprays may be given before rains. As weather is changing proper care of sheds, food and drinking water must be taken & administer vaccines in time.

Nong laktingaida laina tilnagi hidak kaphthokpiyu. Aying asa onrakpada sha uchek naraktanaba makol, ishing amadi machinjak afaba pibiduna matam chana hidak thabiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
TOMATO	Tomato nurseries should be raised under low cost polyhouse as the temperature is very low in open field. - Nursery bed of 0.75m x 4m may be prepared and mix Carbendazim @ 5g/ 3m2 area, keep the bed covered with gunny bag for at least 24 hrs. - Sow seeds @ 10g/ 3m2 area keep the bed covered with gunny bag. Seeds will germinate in 4-5 days.

Crop(Varieties)	Crop Specific Advisory
	- Ingba matamda polyhouse makhada khamen ashinbi chaar hunbiyu. - 0.75m x 4m pak-chaoba nursery bed samba loisanlakpada 3m2 khudingda 5g gi changda Carbendazim happaga bora na kupsinlaga pung 24 thambiyu. - 3m2 pak-chaoba nursery bed ta 10g gi changda maru hidak totlaga hunbibagi matungda bora kupsinlaga thambiyu. Numit 4-5ni gi manungda maton thokpaga chang yengbiraga bora louthokpiba yarani.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
POTATO	- For seed: • Strict supervision on removal of unwanted plant type should be maintained. • Maturity days should be strictly maintained as per variety. - Harvest the crop after 10-15 days after haulm cutting - Maru oinaba lokpige hairabadi: • Masak matou manadaba pambi makhal khandokpiyu.. • Alu na chaoba mapung faba matam variety makhal gi leijaba numit shurakpaga matungda lokpiyu. - Alu loktoubu numit 10-15ni watlingaeida makhok khara khakaga pambi kakaga thambiyu.
FRENCH BEAN	- Sowing of French bean may be done in prepared land. Seed rate for dwarf varieties like Pant Anupam & Arka Komal is 65kg/ha (spacing is 30x45 cm from seed to seed and row to row) and 25-30kg/ha for pole-type (spacing is 60x60cm with 2-3 seeds per hill). Treat seeds with Carbendazim @2g/kg seeds before sowing. - Give light irrigation after 2-3 days of sowing. - Koli hawai thanaba loufam semjinbiba yarani. Nemna panbi variety Pant Anupam amadi Arka Komal pari amada thanaba 65 kg maru lousinbiraga 30x45 cm lapna thabiyu. Pambi awangbagina 25-30 kg changani amadi 60x60cm gi changda lapna makhul amada 2-3 maru thabiyu. - Thadringeida maru kg amada 2g gi changda Carbendazim totpiyu amadi thabagi matung leibak leinil marang kaina leinaba nong churaktagadi ishing khara chaibiyu.



Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 14-02-2020

Weather Forecast of KAKCHING(Manipur) Issued On : 2020-02-14(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-15	0.0	24.0	8.0	78	40	7.0	290	0
2020-02-16	0.0	24.0	9.0	78	34	6.5	298	0
2020-02-17	0.0	24.0	10.0	75	30	3.4	289	0
2020-02-18	0.0	25.0	12.0	80	25	5.5	285	0
2020-02-19	0.0	25.0	12.0	67	23	8.8	292	0

Weather Summary/Alert:

Likely to have no rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 25-30 oC & mini. temp. 12-13 oC. Relative humidity may reach upto 86% (max) & 25% (min). The average wind speed may be 3.3-9.3 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 25-30 oC amadi ingbagina 12-13 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 86 % amadi nungthildana 25% oiraba yai. Nungsit sitpagi changna pung amada 3.3-9.3 km oiraba yai.

General Advisory:

Monitor regularly insect pest infestation in crops. Mulching may be done in the crop field using straw/ dried leaves which is not infected by insect-pest to preserve soil moisture for the crop. Keep birds/ animals under your jurisdiction and avoid contact with migratory or wild birds/ animals.

Loufamda laina tilna taba ningthina toina yengsinbiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Sha uchek yariba makhei sennajaba mafamda thadokpiyu amadi lamlakki sha-uchek/ mapandagi paisinlakpa uchekka samnadanaba shinbiyu.

SMS Advisory:

Insect pest sprays may be given before rains. As weather is changing proper care of sheds, food and drinking water must be taken & administer vaccines in time.

Nong laktingaida laina tilnagi hidak kaphthokpiyu. Aying asa onrakpada sha uchek naraktanaba makol, ishing amadi machinjak afaba pibiduna matam chana hidak thabiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
TOMATO	Tomato nurseries should be raised under low cost polyhouse as the temperature is very low in open field. - Nursery bed of 0.75m x 4m may be prepared and mix Carbendazim @ 5g/ 3m2 area, keep the bed covered with gunny bag for at least 24 hrs. - Sow seeds @ 10g/ 3m2 area keep the bed covered with gunny bag. Seeds will germinate in 4-5 days.

Crop(Varieties)	Crop Specific Advisory
	- Ingba matamda polyhouse makhada khamen ashinbi chaar hunbiyu. - 0.75m x 4m pak-chaoba nursery bed samba loisanlakpada 3m2 khudingda 5g gi changda Carbendazim happaga bora na kupsinlaga pung 24 thambiyu. - 3m2 pak-chaoba nursery bed ta 10g gi changda maru hidak totlaga hunbibagi matungda bora kupsinlaga thambiyu. Numit 4-5ni gi manungda maton thokpaga chang yengbiraga bora louthokpiba yarani.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
POTATO	- For seed: • Strict supervision on removal of unwanted plant type should be maintained. • Maturity days should be strictly maintained as per variety. - Harvest the crop after 10-15 days after haulm cutting - Maru oinaba lokpige hairabadi: • Masak matou manadaba pambi makhal khandokpiyu.. • Alu na chaoba mapung faba matam variety makhal gi leijaba numit shurakpaga matungda lokpiyu. - Alu loktoubu numit 10-15ni watlingaeida makhok khara khakaga pambi kakaga thambiyu.
FRENCH BEAN	- Sowing of French bean may be done in prepared land. Seed rate for dwarf varieties like Pant Anupam & Arka Komal is 65kg/ha (spacing is 30x45 cm from seed to seed and row to row) and 25-30kg/ha for pole-type (spacing is 60x60cm with 2-3 seeds per hill). Treat seeds with Carbendazim @2g/kg seeds before sowing. - Give light irrigation after 2-3 days of sowing. - Koli hawai thanaba loufam semjinbiba yarani. Nemna panbi variety Pant Anupam amadi Arka Komal pari amada thanaba 65 kg maru lousinbiraga 30x45 cm lapna thabiyu. Pambi awangbagina 25-30 kg changani amadi 60x60cm gi changda lapna makhul amada 2-3 maru thabiyu. - Thadringeida maru kg amada 2g gi changda Carbendazim totpiyu amadi thabagi matung leibak leinil marang kaina leinaba nong churaktagadi ishing khara chaibiyu.



Gramin Krishi Mausam Sewa
India Meteorological Department
ICAR Complex Branch
Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 14-02-2020

Weather Forecast of KAMJONG(Manipur) Issued On : 2020-02-14(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-15	0.0	24.0	8.0	72	40	6.0	330	1
2020-02-16	0.0	24.0	9.0	60	31	5.5	334	0
2020-02-17	0.0	24.0	10.0	69	34	5.2	341	0
2020-02-18	0.0	25.0	12.0	68	30	5.5	338	0
2020-02-19	0.0	25.0	12.0	32	16	7.8	334	0

Weather Summary/Alert:

Likely to have no rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 26-30 oC & mini. temp. 12-14 oC. Relative humidity may reach upto 76% (max) & 18% (min). The average wind speed may be 5.1-6.7 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 26-30 oC amadi ingbagina 12-14 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 76% amadi nungthildana 18% oiraba yai. Nungsit sitpigi changna pung amada 5.1-6.7 km oiraba yai.

General Advisory:

Monitor regularly insect pest infestation in crops. Mulching may be done in the crop field using straw/ dried leaves which is not infected by insect-pest to preserve soil moisture for the crop. Keep birds/ animals under your jurisdiction and avoid contact with migratory or wild birds/ animals.

Loufamda laina tilna taba ningthina toina yengsinbiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Sha uchek yariba makhei sennajaba mafamda thadokpiyu amadi lamlakki sha-uchek/ mapandagi paisinlakpa uchekka samnadanaba shinbiyu.

SMS Advisory:

Insect pest sprays may be given before rains. As weather is changing proper care of sheds, food and drinking water must be taken & administer vaccines in time.

Nong laktingaida laina tilnagi hidak kaphthokpiyu. Aying asa onrakpada sha uchek naraktanaba makol, ishing amadi machinjak afaba pibiduna matam chana hidak thabiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
TOMATO	Tomato nurseries should be raised under low cost polyhouse as the temperature is very low in open field. - Nursery bed of 0.75m x 4m may be prepared and mix Carbendazim @ 5g/ 3m ² area, keep the bed covered with gunny bag for at least 24 hrs. - Sow seeds @ 10g/ 3m ² area keep the bed covered with gunny bag. Seeds will germinate in 4-5 days.

Crop(Varieties)	Crop Specific Advisory
	- Ingba matamda polyhouse makhada khamen ashinbi chaar hunbiyu. - 0.75m x 4m pak-chaoba nursery bed samba loisanlakpada 3m2 khudingda 5g gi changda Carbendazim happaga bora na kupsinlaga pung 24 thambiyu. - 3m2 pak-chaoba nursery bed ta 10g gi changda maru hidak totlaga hunbibagi matungda bora kupsinlaga thambiyu. Numit 4-5ni gi manungda maton thokpagi chang yengbiraga bora louthokpiba yarani.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
POTATO	- For seed: • Strict supervision on removal of unwanted plant type should be maintained. • Maturity days should be strictly maintained as per variety. - Harvest the crop after 10-15 days after haulm cutting - Maru oinaba lokpige hairabadi: • Masak matou manadaba pambi makhal khandokpiyu.. • Alu na chaoba mapung faba matam variety makhal gi leijaba numit shurakpagi matungda lokpiyu. - Alu loktoubu numit 10-15ni watlingaeida makhok khara khakaga pambi kakaga thambiyu.
FRENCH BEAN	- Sowing of French bean may be done in prepared land. Seed rate for dwarf varieties like Pant Anupam & Arka Komal is 65kg/ha (spacing is 30x45 cm from seed to seed and row to row) and 25-30kg/ha for pole-type (spacing is 60x60cm with 2-3 seeds per hill). Treat seeds with Carbendazim @2g/kg seeds before sowing. - Give light irrigation after 2-3 days of sowing. - Koli hawai thanaba loufam semjinbiba yarani. Nemna panbi variety Pant Anupam amadi Arka Komal pari amada thanaba 65 kg maru lousinbiraga 30x45 cm lapna thabiyu. Pambi awangbagina 25-30 kg changani amadi 60x60cm gi changda lapna makhul amada 2-3 maru thabiyu. - Thadringeida maru kg amada 2g gi changda Carbendazim totpiyu amadi thabagi matung leibak leinil marang kaina leinaba nong churaktagadi ishing khara chaibiyu.