



## Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



# Agromet Advisory Bulletin

Date : 18-02-2020

**Weather Forecast of KAKCHING(Manipur) Issued On : 2020-02-18(Valid Till 08:30 IST of the next 5 days)**

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-19	0.0	24.0	7.0	78	29	8.7	294	0
2020-02-20	0.0	24.0	7.0	70	25	5.5	290	0
2020-02-21	0.0	26.0	8.0	53	21	4.8	292	1
2020-02-22	0.0	26.0	9.0	68	25	4.1	109	4
2020-02-23	0.0	27.0	9.0	55	20	4.7	279	1

### **Weather Summary/Alert:**

Likely to have light rains on Saturday. The sky will remain mainly clear to partly cloudy with max. temp. 24-27°C & mini. temp. 7-9 °C. Relative humidity may reach upto 78 % (max) & 29 % (min). The average wind speed may be 4.1-8.7 km/hr.

Lakliba Thangjada nong khara churaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 24-27 °C amadi ingbagina 7-9 °C oiraba yai. Nungsitta ishingga manilgi changna yammabada ayukta 78 % amadi nunghildana 29 % oiraba yai. Nungsit sitpagi changna pung amada 4.1-8.7 km oiraba yai.

### **General Advisory:**

Sprays against insect pest may be given before rains. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free to preserve soil moisture for the crop. Keep sheds and surroundings of birds/ animals clean to avoid mosquitoes/ flies population.

Loufamda laina tilna ngakthoknaba hidak nong laktringeida kappiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Kang-hayinggi yamma indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nann thambiyu.

### **SMS Advisory:**

Spray Hexaconazole @ 2ml/L water against late blight in potato. Prepare land & mix well decomposed FYM @ 20-25 t/ha 15 days before planting ginger/ turmeric.

Alu da late blight ngakthoknaba ishing 1L da 2ml Hexaconazole happaga kappiyu. Shing/ Yaingang thadouba numit 15ni watlingeida pari 1da 20-25t patharaba FYM happyiu.

### **Crop Specific Advisory:**

Crop(Varieties)	Crop Specific Advisory
LENTIL	- Spray Bayleton @ 7g/ 10 litres of water or spray Neem Seed Kernel Extract at 5% concentration for powdery mildew and Nimbicidin (0.3%) to control rust disease. - Spray Imidachloprid 17.8 SL @ 7-10ml/ 15 litre of water against aphid.

Crop(Varieties)	Crop Specific Advisory
	- Ishing litre 10da 7g Bayleton happaga kappiyu natraga 5% concentration ta Neem Seed Kernel Extract at happaga chaibibada Kokfai kaba laina ngakthokpa yani. Kolhing kaba manbi laina ngakthoknaba Nimbicidin (0.3%) sijinabiyu. - Wahik ngakthoknaba ishing litre 15da 7-10 ml Imidachloprid 17.8 SL happaga kappiyu.
GARLIC	<p>- To prevent thrips in garlic crop install blue sticky trap in the field or spray Neem Seed Kernel extract @2ml/litre of water. Put a layer of mulch on the ground near the garlic plant to conserve soil moisture.</p> <p>Chanam mana maton ngangsinlakpa natraga wahik ngakthoknaba blue sticky trap sijinabiyu/ ishing litre amada 2ml Neem Seed Kernel Extract happaga kappiyu. Leibak leinil kannaba pambi mahongda charu/ una akangba thambiyu.</p>

### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
CHILLI	<p>- Raise chilies and capsicum nursery. Spray Carbendazim + Mancozeb (Sixer) @ 2g/litre of water to control the fungal attack.</p> <p>Morok Chaar thabiba houba yare. Kokfai kabi laina ngakthoknaba ishing litre amada 2g Carbendazim + Mancozeb (Sixer) happaga kappiyu.</p>

### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
BIRD	<p>- Amprolium powder may be added in low dose in both feed (@25 g per 1000kg of mash) and drinking water (30g/ 25 lit of water) of poultry above 3 weeks of age.</p> <p>- Chayol 3 shuraba yendagina mapi faraba yen gi chinzak 1000kg da 25g amadi thaknaba ishing litre 25da 30g Amprolium powder happeniraga amangbada ee yaorakpagi lainadagi ngakthokpiyu.</p>



## Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



# Agromet Advisory Bulletin

Date : 18-02-2020

**Weather Forecast of KAMJONG(Manipur) Issued On : 2020-02-18(Valid Till 08:30 IST of the next 5 days)**

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-19	0.0	28.0	13.0	56	20	7.2	334	0
2020-02-20	0.0	27.0	13.0	55	23	5.5	334	0
2020-02-21	0.0	28.0	14.0	47	21	5.4	338	1
2020-02-22	0.0	28.0	16.0	58	25	5.3	341	4
2020-02-23	0.0	29.0	16.0	52	26	5.2	341	1

### **Weather Summary/Alert:**

Likely to have light rains on Saturday. The sky will remain mainly clear to partly cloudy with max. temp. 27- 29 oC & mini. temp. 13-16 oC. Relative humidity may reach upto 58 % (max) & 23 % (min). The average wind speed may be 5.2-7.2 km/hr.

Lakliba Thangjada nong khara churaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 27- 29 oC amadi ingbagina 13-16 oC oiraba yai. Nungsitta ishingga manilgi changna yammabada ayukta 58 % amadi nunghildana 23 % oiraba yai. Nungsit sitpagi changna pung amada 5.2-7.2 km oiraba yai.

### **General Advisory:**

Sprays against insect pest may be given before rains. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free to preserve soil moisture for the crop. Keep sheds and surroundings of birds/ animals clean to avoid mosquitoes/ flies population.

Loufamda laina tilna ngakthoknaba hidak nong laktringeida kappiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Kang-hayinggi yamma indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nann thambiyu.

### **SMS Advisory:**

Spray Hexaconazole @ 2ml/L water against late blight in potato. Prepare land & mix well decomposed FYM @ 20-25 t/ha 15 days before planting ginger/ turmeric.

Alu da late blight ngakthoknaba ishing 1L da 2ml Hexaconazole happaga kappiyu. Shing/ Yaingang thadouba numit 15ni watlingeida pari 1da 20-25t patharaba FYM happyiu.

### **Crop Specific Advisory:**

Crop(Varieties)	Crop Specific Advisory
LENTIL	- Spray Bayleton @ 7g/ 10 litres of water or spray Neem Seed Kernel Extract at 5% concentration for powdery mildew and Nimbicidin (0.3%) to control rust disease. - Spray Imidachloprid 17.8 SL @ 7-10ml/ 15 litre of water against aphid.

Crop(Varieties)	Crop Specific Advisory
	- Ishing litre 10da 7g Bayleton happaga kappiyu natraga 5% concentration ta Neem Seed Kernel Extract at happaga chaibibada Kokfai kaba laina ngakthokpa yani. Kolhing kaba manbi laina ngakthoknaba Nimbicidin (0.3%) sijinabiyu. - Wahik ngakthoknaba ishing litre 15da 7-10 ml Imidachloprid 17.8 SL happaga kappiyu.
GARLIC	- To prevent thrips in garlic crop install blue sticky trap in the field or spray Neem Seed Kernel extract @2ml/litre of water. Put a layer of mulch on the ground near the garlic plant to conserve soil moisture. Chanam mana maton ngangsinlakpa natraga wahik ngakthoknaba blue sticky trap sijinabiyu/ ishing litre amada 2ml Neem Seed Kernel Extract happaga kappiyu. Leibak leinil kannaba pambi mahongda charu/ una akangba thambiyu.

### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
CHILLI	- Raise chilies and capsicum nursery. Spray Carbendazim + Mancozeb (Sixer) @ 2g/litre of water to control the fungal attack. Morok Chaar thabiba houba yare. Kokfai kabi laina ngakthoknaba ishing litre amada 2g Carbendazim + Mancozeb (Sixer) happaga kappiyu.

### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
BIRD	- Amprolium powder may be added in low dose in both feed (@25 g per 1000kg of mash) and drinking water (30g/ 25 lit of water) of poultry above 3 weeks of age. - Chayol 3 shuraba yendagina mapi faraba yen gi chinzak 1000kg da 25g amadi thaknaba ishing litre 25da 30g Amprolium powder happeniraga amangbada ee yaorakpagi lainadagi ngakthokpiyu.



## Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



# Agromet Advisory Bulletin

Date : 18-02-2020

**Weather Forecast of KANGPOKPI(Manipur) Issued On : 2020-02-18(Valid Till 08:30 IST of the next 5 days)**

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-19	0.0	27.0	13.0	74	21	5.5	18	0
2020-02-20	0.0	26.0	13.0	65	18	5.6	0	0
2020-02-21	0.0	27.0	13.0	62	19	5.4	18	1
2020-02-22	0.0	28.0	16.0	55	14	5.4	19	3
2020-02-23	0.0	29.0	16.0	52	19	5.4	18	1

### **Weather Summary/Alert:**

Likely to have light rains on Saturday. The sky will remain mainly clear to partly cloudy with max. temp. 26-29 oC & mini. temp. 13-16 oC. Relative humidity may reach upto 74 % (max) & 21 % (min). The average wind speed may be 5.4-5.6 km/hr.

Lakliba Thangjada nong khara churaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 26-29 oC amadi ingbagina 13-16 oC oiraba yai. Nungsitta ishingga manilgi changna yammabada ayukta 74 % amadi nunghildana 21 % oiraba yai. Nungsit sitpagi changna pung amada 5.4-5.6 km oiraba yai.

### **General Advisory:**

Sprays against insect pest may be given before rains. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free to preserve soil moisture for the crop. Keep sheds and surroundings of birds/ animals clean to avoid mosquitoes/ flies population.

Loufamda laina tilna ngakthoknaba hidak nong laktringeida kappiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Kang-hayinggi yamma indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nann thambiyu.

### **SMS Advisory:**

Spray Hexaconazole @ 2ml/L water against late blight in potato. Prepare land & mix well decomposed FYM @ 20-25 t/ha 15 days before planting ginger/ turmeric.

Alu da late blight ngakthoknaba ishing 1L da 2ml Hexaconazole happaga kappiyu. Shing/ Yaingang thadouba numit 15ni watlingeida pari 1da 20-25t patharaba FYM happyiu.

### **Crop Specific Advisory:**

Crop(Varieties)	Crop Specific Advisory
LENTIL	- Spray Bayleton @ 7g/ 10 litres of water or spray Neem Seed Kernel Extract at 5% concentration for powdery mildew and Nimbicidin (0.3%) to control rust disease. - Spray Imidachloprid 17.8 SL @ 7-10ml/ 15 litre of water against aphid.

Crop(Varieties)	Crop Specific Advisory
	- Ishing litre 10da 7g Bayleton happaga kappiyu natraga 5% concentration ta Neem Seed Kernel Extract at happaga chaibibada Kokfai kaba laina ngakthokpa yani. Kolhing kaba manbi laina ngakthoknaba Nimbicidin (0.3%) sijinabiyu. - Wahik ngakthoknaba ishing litre 15da 7-10 ml Imidachloprid 17.8 SL happaga kappiyu.
GARLIC	<p>- To prevent thrips in garlic crop install blue sticky trap in the field or spray Neem Seed Kernel extract @2ml/litre of water. Put a layer of mulch on the ground near the garlic plant to conserve soil moisture.</p> <p>Chanam mana maton ngangsinlakpa natraga wahik ngakthoknaba blue sticky trap sijinabiyu/ ishing litre amada 2ml Neem Seed Kernel Extract happaga kappiyu. Leibak leinil kannaba pambi mahongda charu/ una akangba thambiyu.</p>

### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
CHILLI	<p>- Raise chilies and capsicum nursery. Spray Carbendazim + Mancozeb (Sixer) @ 2g/litre of water to control the fungal attack.</p> <p>Morok Chaar thabiba houba yare. Kokfai kabi laina ngakthoknaba ishing litre amada 2g Carbendazim + Mancozeb (Sixer) happaga kappiyu.</p>

### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
BIRD	<p>- Amprolium powder may be added in low dose in both feed (@25 g per 1000kg of mash) and drinking water (30g/ 25 lit of water) of poultry above 3 weeks of age.</p> <p>- Chayol 3 shuraba yendagina mapi faraba yen gi chinzak 1000kg da 25g amadi thaknaba ishing litre 25da 30g Amprolium powder happeniraga amangbada ee yaorakpagi lainadagi ngakthokpiyu.</p>



## Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



# Agromet Advisory Bulletin

Date : 18-02-2020

**Weather Forecast of NONEY(Manipur) Issued On : 2020-02-18(Valid Till 08:30 IST of the next 5 days)**

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-19	0.0	27.0	13.0	64	26	4.3	63	0
2020-02-20	0.0	27.0	13.0	62	22	4.3	63	0
2020-02-21	0.0	27.0	13.0	51	19	4.6	68	3
2020-02-22	0.0	29.0	15.0	55	15	4.4	71	3
2020-02-23	0.0	30.0	16.0	51	18	4.6	63	1

### **Weather Summary/Alert:**

Likely to have light rains on Saturday. The sky will remain mainly clear to partly cloudy with max. temp. 27- 29 oC & mini. temp. 13-16 oC. Relative humidity may reach upto 64 % (max) & 26 % (min). The average wind speed may be 4.3-4.6km/hr.

Lakliba Thangjada nong khara churaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 27-29 oC amadi ingbagina 13-16 oC oiraba yai. Nungsitta ishingga manilgi changna yammabada ayukta 64 % amadi nunghildana 26 % oiraba yai. Nungsit sitpagi changna pung amada 4.3-4.6 km oiraba yai.

### **General Advisory:**

Sprays against insect pest may be given before rains. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free to preserve soil moisture for the crop. Keep sheds and surroundings of birds/ animals clean to avoid mosquitoes/ flies population.

Loufamda laina tilna ngakthoknaba hidak nong laktringeida kappiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Kang-hayinggi yamma indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nann thambiyu.

### **SMS Advisory:**

Spray Hexaconazole @ 2ml/L water against late blight in potato. Prepare land & mix well decomposed FYM @ 20-25 t/ha 15 days before planting ginger/ turmeric.

Alu da late blight ngakthoknaba ishing 1L da 2ml Hexaconazole happaga kappiyu. Shing/ Yaingang thadouba numit 15ni watlingeida pari 1da 20-25t patharaba FYM happyiu.

### **Crop Specific Advisory:**

Crop(Varieties)	Crop Specific Advisory
LENTIL	- Spray Bayleton @ 7g/ 10 litres of water or spray Neem Seed Kernel Extract at 5% concentration for powdery mildew and Nimbicidin (0.3%) to control rust disease. - Spray Imidachloprid 17.8 SL @ 7-10ml/ 15 litre of water against aphid.

Crop(Varieties)	Crop Specific Advisory
	- Ishing litre 10da 7g Bayleton happaga kappiyu natraga 5% concentration ta Neem Seed Kernel Extract at happaga chaibibada Kokfai kaba laina ngakthokpa yani. Kolhing kaba manbi laina ngakthoknaba Nimbicidin (0.3%) sijinabiyu. - Wahik ngakthoknaba ishing litre 15da 7-10 ml Imidachloprid 17.8 SL happaga kappiyu.
GARLIC	<p>- To prevent thrips in garlic crop install blue sticky trap in the field or spray Neem Seed Kernel extract @2ml/litre of water. Put a layer of mulch on the ground near the garlic plant to conserve soil moisture.</p> <p>Chanam mana maton ngangsinlakpa natraga wahik ngakthoknaba blue sticky trap sijinabiyu/ ishing litre amada 2ml Neem Seed Kernel Extract happaga kappiyu. Leibak leinil kannaba pambi mahongda charu/ una akangba thambiyu.</p>

### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
CHILLI	<p>- Raise chilies and capsicum nursery. Spray Carbendazim + Mancozeb (Sixer) @ 2g/litre of water to control the fungal attack.</p> <p>Morok Chaar thabiba houba yare. Kokfai kabi laina ngakthoknaba ishing litre amada 2g Carbendazim + Mancozeb (Sixer) happaga kappiyu.</p>

### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
BIRD	<p>- Amprolium powder may be added in low dose in both feed (@25 g per 1000kg of mash) and drinking water (30g/ 25 lit of water) of poultry above 3 weeks of age.</p> <p>- Chayol 3 shuraba yendagina mapi faraba yen gi chinzak 1000kg da 25g amadi thaknaba ishing litre 25da 30g Amprolium powder happeniraga amangbada ee yaorakpagi lainadagi ngakthokpiyu.</p>



**Gramin Krishi Mausam Sewa**  
 Experimental Block Level Agromet Advisory Bulletin  
 (A Joint Initiative of IMD & ICAR)



## Agromet Advisory Bulletin

Date : 18-02-2020

Weather Forecast of PATSOI Block in IMPHAL-WEST(Manipur) Issued On :2020-02-18(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-19	0.0	26.8	12.5	88	33	7.0	330	0
2020-02-20	0.0	26.9	12.9	71	25	5.0	360	0
2020-02-21	0.0	27.8	13.3	51	21	6.0	338	1
2020-02-22	0.0	28.2	15.3	60	19	5.0	27	5
2020-02-23	0.0	29.6	15.3	59	20	5.0	27	1

### Weather Summary/Alert:

Likely to have light rains on Saturday. The sky will remain mainly clear to partly cloudy with max. temp. 26.8-29.6°C & mini. temp. 12.5-15.3 °C. Relative humidity may reach upto 88% (max) & 33% (min). The average wind speed may be 5-7km/hr.

Lakliba Thangjada nong khara churaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 26.8-29.6 °C amadi ingbagina 12.5-15.3 °C oiraba yai. Nungsitta ishingga manilgi changna yammabada ayukta 88% amadi nungthildana 33 % oiraba yai. Nungsit sitpagi changna pung amada 5-7 km oiraba yai.

### General Advisory:

Sprays against insect pest may be given before rains. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free to preserve soil moisture for the crop. Keep sheds and surroundings of birds/ animals clean to avoid mosquitoes/ flies population.

Loufamda laina tilna ngakthoknaba hidak nong laktringeida kappiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Kang-hayinggi yamma indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nann thambiyu.

### SMS Advisory:

Spray Hexaconazole @ 2ml/L water against late blight in potato. Prepare land & mix well decomposed FYM @ 20-25 t/ha 15 days before planting ginger/ turmeric.

Alu da late blight ngakthoknaba ishing 1L da 2ml Hexaconazole happaga kappiyu. Shing/ Yaingang thadouba numit 15ni watlingeida pari 1da 20-25t patharaba FYM happyiu.

### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
LENTIL	- Spray Bayleton @ 7g/ 10 litres of water or spray Neem Seed Kernel Extract at 5% concentration for powdery mildew and Nimbicidin (0.3%) to control rust disease. - Spray Imidachloprid 17.8 SL @ 7-10ml/ 15 litre of water against aphid.
	- Isching litre 10da 7g Bayleton happaga kappiyu natraga 5% concentration ta Neem Seed Kernel Extract at happaga chaibibada Kokfai kaba laina ngakthokpa yani. Kolhing kaba manbi laina ngakthoknaba Nimbicidin (0.3%) sijinabiyu. - Wahik ngakthoknaba ishing litre 15da 7-10 ml Imidachloprid 17.8 SL happaga kappiyu.
GARLIC	- To prevent thrips in garlic crop install blue sticky trap in the field or spray Neem Seed Kernel extract @2ml/litre of water. Put a layer of mulch on the ground near the garlic plant to conserve soil moisture.
	Chanam mana maton ngangsinlakpa natraga wahik ngakthoknaba blue sticky trap sijinabiyu/ ishing litre amada 2ml Neem Seed Kernel Extract happaga kappiyu. Leibak leinil kannaba pambi mahongda charu/ una akangba thambiyu.

**Horticulture Specific Advisory:**

<b>Horticulture (Varieties)</b>	<b>Horticulture Specific Advisory</b>
CHILLI	<ul style="list-style-type: none"> <li>- Raise chilies and capsicum nursery. Spray Carbendazim + Mancozeb (Sixer) @ 2g/litre of water to control the fungal attack.</li> <li>Morok Chaar thabiba houba yare. Kokfai kabi laina ngakthoknaba ishing litre amada 2g Carbendazim + Mancozeb (Sixer) happaga kappiyu.</li> </ul>

**Poultry Specific Advisory:**

<b>Poultry(Varieties)</b>	<b>Poultry Specific Advisory</b>
BIRD	<ul style="list-style-type: none"> <li>- Amprolium powder may be added in low dose in both feed (@25 g per 1000kg of mash) and drinking water (30g/ 25 lit of water) of poultry above 3 weeks of age.</li> <li>- Chayol 3 shuraba yendagina mapi faraba yen gi chinzak 1000kg da 25g amadi thaknaba ishing litre25da 30g Amprolium powder happiraga amangbada ee yaorakpagi lainadagi ngakthokpiyu.</li> </ul>



## Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



# Agromet Advisory Bulletin

Date : 18-02-2020

**Weather Forecast of PHERZAWL(Manipur) Issued On : 2020-02-18(Valid Till 08:30 IST of the next 5 days)**

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-19	0.0	28.0	14.0	68	29	4.7	109	0
2020-02-20	0.0	28.0	14.0	68	26	4.4	109	0
2020-02-21	0.0	28.0	13.0	53	20	4.4	109	2
2020-02-22	0.0	30.0	14.0	61	18	4.0	112	3
2020-02-23	0.0	31.0	16.0	54	19	4.2	105	1

### **Weather Summary/Alert:**

Likely to have light rains on Saturday. The sky will remain mainly clear to partly cloudy with max. temp. 28-31 oC & mini. temp. 13-16 oC. Relative humidity may reach upto 68 % (max) & 29 % (min). The average wind speed may be 4.0-4.7km/hr.

Lakliba Thangjada nong khara churaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 28-31 oC amadi ingbagina 13-16 oC oiraba yai. Nungsitta ishingga manilgi changna yammabada ayukta 68 % amadi nunghildana 29 % oiraba yai. Nungsit sitpagi changna pung amada 4.0-4.7km oiraba yai.

### **General Advisory:**

Sprays against insect pest may be given before rains. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free to preserve soil moisture for the crop. Keep sheds and surroundings of birds/ animals clean to avoid mosquitoes/ flies population.

Loufamda laina tilna ngakthoknaba hidak nong laktringeida kappiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Kang-hayinggi yamma indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nann thambiyu.

### **SMS Advisory:**

Spray Hexaconazole @ 2ml/L water against late blight in potato. Prepare land & mix well decomposed FYM @ 20-25 t/ha 15 days before planting ginger/ turmeric.

Alu da late blight ngakthoknaba ishing 1L da 2ml Hexaconazole happaga kappiyu. Shing/ Yaingang thadouba numit 15ni watlingeida pari 1da 20-25t patharaba FYM happyiu.

### **Crop Specific Advisory:**

Crop(Varieties)	Crop Specific Advisory
LENTIL	- Spray Bayleton @ 7g/ 10 litres of water or spray Neem Seed Kernel Extract at 5% concentration for powdery mildew and Nimbicidin (0.3%) to control rust disease. - Spray Imidachloprid 17.8 SL @ 7-10ml/ 15 litre of water against aphid.

Crop(Varieties)	Crop Specific Advisory
	- Ishing litre 10da 7g Bayleton happaga kappiyu natraga 5% concentration ta Neem Seed Kernel Extract at happaga chaibibada Kokfai kaba laina ngakthokpa yani. Kolhing kaba manbi laina ngakthoknaba Nimbicidin (0.3%) sijinabiyu. - Wahik ngakthoknaba ishing litre 15da 7-10 ml Imidachloprid 17.8 SL happaga kappiyu.
GARLIC	<p>- To prevent thrips in garlic crop install blue sticky trap in the field or spray Neem Seed Kernel extract @2ml/litre of water. Put a layer of mulch on the ground near the garlic plant to conserve soil moisture.</p> <p>Chanam mana maton ngangsinlakpa natraga wahik ngakthoknaba blue sticky trap sijinabiyu/ ishing litre amada 2ml Neem Seed Kernel Extract happaga kappiyu. Leibak leinil kannaba pambi mahongda charu/ una akangba thambiyu.</p>

### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
CHILLI	<p>- Raise chilies and capsicum nursery. Spray Carbendazim + Mancozeb (Sixer) @ 2g/litre of water to control the fungal attack.</p> <p>Morok Chaar thabiba houba yare. Kokfai kabi laina ngakthoknaba ishing litre amada 2g Carbendazim + Mancozeb (Sixer) happaga kappiyu.</p>

### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
BIRD	<p>- Amprolium powder may be added in low dose in both feed (@25 g per 1000kg of mash) and drinking water (30g/ 25 lit of water) of poultry above 3 weeks of age.</p> <p>- Chayol 3 shuraba yendagina mapi faraba yen gi chinzak 1000kg da 25g amadi thaknaba ishing litre 25da 30g Amprolium powder happeniraga amangbada ee yaorakpagi lainadagi ngakthokpiyu.</p>



## Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



# Agromet Advisory Bulletin

Date : 18-02-2020

**Weather Forecast of SENAPATI(Manipur) Issued On : 2020-02-18(Valid Till 08:30 IST of the next 5 days)**

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-19	0.0	23.0	6.0	83	24	4.8	18	0
2020-02-20	0.0	23.0	6.0	74	21	5.1	0	0
2020-02-21	0.0	24.0	7.0	71	23	4.9	18	2
2020-02-22	0.0	25.0	8.0	59	16	5.1	22	4
2020-02-23	0.0	25.0	8.0	58	21	4.7	18	2

### **Weather Summary/Alert:**

Likely to have light rains on Saturday. The sky will remain mainly clear to partly cloudy with max. temp. 23-25°C & mini. temp. 6-8 °C. Relative humidity may reach upto 83 % (max) & 24 % (min). The average wind speed may be 4.7-5.1 km/hr.

Lakliba Thangjada nong khara churaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 23-25 °C amadi ingbagina 6-8 °C oiraba yai. Nungsitta ishingga manilgi changna yammabada ayukta 83 % amadi nunghildana 24 % oiraba yai. Nungsit sitpagi changna pung amada 4.7-51 km oiraba yai.

### **General Advisory:**

Sprays against insect pest may be given before rains. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free to preserve soil moisture for the crop. Keep sheds and surroundings of birds/ animals clean to avoid mosquitoes/ flies population.

Loufamda laina tilna ngakthoknaba hidak nong laktringeida kappiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Kang-hayinggi yamma indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nann thambiyu.

### **SMS Advisory:**

Spray Hexaconazole @ 2ml/L water against late blight in potato. Prepare land & mix well decomposed FYM @ 20-25 t/ha 15 days before planting ginger/ turmeric.

Alu da late blight ngakthoknaba ishing 1L da 2ml Hexaconazole happaga kappiyu. Shing/ Yaingang thadouba numit 15ni watlingeida pari 1da 20-25t patharaba FYM happyiu.

### **Crop Specific Advisory:**

Crop(Varieties)	Crop Specific Advisory
LENTIL	- Spray Bayleton @ 7g/ 10 litres of water or spray Neem Seed Kernel Extract at 5% concentration for powdery mildew and Nimbicidin (0.3%) to control rust disease. - Spray Imidachloprid 17.8 SL @ 7-10ml/ 15 litre of water against aphid.

Crop(Varieties)	Crop Specific Advisory
	- Ishing litre 10da 7g Bayleton happaga kappiyu natraga 5% concentration ta Neem Seed Kernel Extract at happaga chaibibada Kokfai kaba laina ngakthokpa yani. Kolhing kaba manbi laina ngakthoknaba Nimbicidin (0.3%) sijinabiyu. - Wahik ngakthoknaba ishing litre 15da 7-10 ml Imidachloprid 17.8 SL happaga kappiyu.
GARLIC	- To prevent thrips in garlic crop install blue sticky trap in the field or spray Neem Seed Kernel extract @2ml/litre of water. Put a layer of mulch on the ground near the garlic plant to conserve soil moisture. Chanam mana maton ngangsinlakpa natraga wahik ngakthoknaba blue sticky trap sijinabiyu/ ishing litre amada 2ml Neem Seed Kernel Extract happaga kappiyu. Leibak leinil kannaba pambi mahongda charu/ una akangba thambiyu.

### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
CHILLI	- Raise chilies and capsicum nursery. Spray Carbendazim + Mancozeb (Sixer) @ 2g/litre of water to control the fungal attack. Morok Chaar thabiba houba yare. Kokfai kabi laina ngakthoknaba ishing litre amada 2g Carbendazim + Mancozeb (Sixer) happaga kappiyu.

### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
BIRD	- Amprolium powder may be added in low dose in both feed (@25 g per 1000kg of mash) and drinking water (30g/ 25 lit of water) of poultry above 3 weeks of age. - Chayol 3 shuraba yendagina mapi faraba yen gi chinzak 1000kg da 25g amadi thaknaba ishing litre 25da 30g Amprolium powder happeniraga amangbada ee yaorakpagi lainadagi ngakthokpiyu.



## Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



# Agromet Advisory Bulletin

Date : 18-02-2020

**Weather Forecast of TAMENGLONG(Manipur) Issued On : 2020-02-18(Valid Till 08:30 IST of the next 5 days)**

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-19	0.0	24.0	7.0	60	27	5.0	68	0
2020-02-20	0.0	24.0	7.0	56	23	4.9	68	0
2020-02-21	0.0	26.0	8.0	53	20	5.0	68	3
2020-02-22	0.0	26.0	9.0	55	17	4.6	68	3
2020-02-23	0.0	27.0	9.0	47	18	4.6	68	2

### **Weather Summary/Alert:**

Likely to have light rains on Saturday. The sky will remain mainly clear to partly cloudy with max. temp. 24-27°C & mini. temp. 7-9 °C. Relative humidity may reach upto 60 % (max) & 27% (min). The average wind speed may be 4.6-5.0 km/hr.

Lakliba Thangjada nong khara churaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 24-27 °C amadi ingbagina 7-9 °C oiraba yai. Nungsitta ishingga manilgi changna yammabada ayukta 60% amadi nunghildana 27% oiraba yai. Nungsit sitpagi changna pung amada 4.6-5.0 km oiraba yai.

### **General Advisory:**

Sprays against insect pest may be given before rains. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free to preserve soil moisture for the crop. Keep sheds and surroundings of birds/ animals clean to avoid mosquitoes/ flies population.

Loufamda laina tilna ngakthoknaba hidak nong laktringeida kappiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Kang-hayinggi yamma indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nann thambiyu.

### **SMS Advisory:**

Spray Hexaconazole @ 2ml/L water against late blight in potato. Prepare land & mix well decomposed FYM @ 20-25 t/ha 15 days before planting ginger/ turmeric.

Alu da late blight ngakthoknaba ishing 1L da 2ml Hexaconazole happaga kappiyu. Shing/ Yaingang thadouba numit 15ni watlingeida pari 1da 20-25t patharaba FYM happyiu.

### **Crop Specific Advisory:**

Crop(Varieties)	Crop Specific Advisory
LENTIL	- Spray Bayleton @ 7g/ 10 litres of water or spray Neem Seed Kernel Extract at 5% concentration for powdery mildew and Nimbicidin (0.3%) to control rust disease. - Spray Imidachloprid 17.8 SL @ 7-10ml/ 15 litre of water against aphid.

Crop(Varieties)	Crop Specific Advisory
	- Ishing litre 10da 7g Bayleton happaga kappiyu natraga 5% concentration ta Neem Seed Kernel Extract at happaga chaibibada Kokfai kaba laina ngakthokpa yani. Kolhing kaba manbi laina ngakthoknaba Nimbicidin (0.3%) sijinabiyu. - Wahik ngakthoknaba ishing litre 15da 7-10 ml Imidachloprid 17.8 SL happaga kappiyu.
GARLIC	- To prevent thrips in garlic crop install blue sticky trap in the field or spray Neem Seed Kernel extract @2ml/litre of water. Put a layer of mulch on the ground near the garlic plant to conserve soil moisture. Chanam mana maton ngangsinlakpa natraga wahik ngakthoknaba blue sticky trap sijinabiyu/ ishing litre amada 2ml Neem Seed Kernel Extract happaga kappiyu. Leibak leinil kannaba pambi mahongda charu/ una akangba thambiyu.

### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
CHILLI	- Raise chilies and capsicum nursery. Spray Carbendazim + Mancozeb (Sixer) @ 2g/litre of water to control the fungal attack. Morok Chaar thabiba houba yare. Kokfai kabi laina ngakthoknaba ishing litre amada 2g Carbendazim + Mancozeb (Sixer) happaga kappiyu.

### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
BIRD	- Amprolium powder may be added in low dose in both feed (@25 g per 1000kg of mash) and drinking water (30g/ 25 lit of water) of poultry above 3 weeks of age. - Chayol 3 shuraba yendagina mapi faraba yen gi chinzak 1000kg da 25g amadi thaknaba ishing litre 25da 30g Amprolium powder happeniraga amangbada ee yaorakpagi lainadagi ngakthokpiyu.



## Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



# Agromet Advisory Bulletin

Date : 18-02-2020

**Weather Forecast of TENGNOUPAL(Manipur) Issued On : 2020-02-18(Valid Till 08:30 IST of the next 5 days)**

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-19	0.0	24.0	7.0	73	25	7.9	292	0
2020-02-20	0.0	24.0	7.0	66	25	4.9	292	0
2020-02-21	0.0	25.0	8.0	50	22	4.1	290	1
2020-02-22	0.0	26.0	9.0	68	28	4.4	112	5
2020-02-23	0.0	27.0	9.0	52	23	4.5	290	2

### **Weather Summary/Alert:**

Likely to have light rains on Saturday. The sky will remain mainly clear to partly cloudy with max. temp. 24-27°C & mini. temp. 7-9 °C. Relative humidity may reach upto 73 % (max) & 28 % (min). The average wind speed may be 4.1-7.9km/hr.

Lakliba Thangjada nong khara churaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 24-27 °C amadi ingbagina 7-9 °C oiraba yai. Nungsitta ishingga manilgi changna yammabada ayukta 73 % amadi nunghildana 28 % oiraba yai. Nungsit sitpagi changna pung amada 4.1-7.9km oiraba yai.

### **General Advisory:**

Sprays against insect pest may be given before rains. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free to preserve soil moisture for the crop. Keep sheds and surroundings of birds/ animals clean to avoid mosquitoes/ flies population.

Loufamda laina tilna ngakthoknaba hidak nong laktringeida kappiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Kang-hayinggi yamma indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nann thambiyu.

### **SMS Advisory:**

Spray Hexaconazole @ 2ml/L water against late blight in potato. Prepare land & mix well decomposed FYM @ 20-25 t/ha 15 days before planting ginger/ turmeric.

Alu da late blight ngakthoknaba ishing 1L da 2ml Hexaconazole happaga kappiyu. Shing/ Yaingang thadouba numit 15ni watlingeida pari 1da 20-25t patharaba FYM happyiu.

### **Crop Specific Advisory:**

Crop(Varieties)	Crop Specific Advisory
LENTIL	- Spray Bayleton @ 7g/ 10 litres of water or spray Neem Seed Kernel Extract at 5% concentration for powdery mildew and Nimbicidin (0.3%) to control rust disease. - Spray Imidachloprid 17.8 SL @ 7-10ml/ 15 litre of water against aphid.

Crop(Varieties)	Crop Specific Advisory
	- Ishing litre 10da 7g Bayleton happaga kappiyu natraga 5% concentration ta Neem Seed Kernel Extract at happaga chaibibada Kokfai kaba laina ngakthokpa yani. Kolhing kaba manbi laina ngakthoknaba Nimbicidin (0.3%) sijinabiyu. - Wahik ngakthoknaba ishing litre 15da 7-10 ml Imidachloprid 17.8 SL happaga kappiyu.
GARLIC	<p>- To prevent thrips in garlic crop install blue sticky trap in the field or spray Neem Seed Kernel extract @2ml/litre of water. Put a layer of mulch on the ground near the garlic plant to conserve soil moisture.</p> <p>Chanam mana maton ngangsinlakpa natraga wahik ngakthoknaba blue sticky trap sijinabiyu/ ishing litre amada 2ml Neem Seed Kernel Extract happaga kappiyu. Leibak leinil kannaba pambi mahongda charu/ una akangba thambiyu.</p>

### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
CHILLI	<p>- Raise chilies and capsicum nursery. Spray Carbendazim + Mancozeb (Sixer) @ 2g/litre of water to control the fungal attack.</p> <p>Morok Chaar thabiba houba yare. Kokfai kabi laina ngakthoknaba ishing litre amada 2g Carbendazim + Mancozeb (Sixer) happaga kappiyu.</p>

### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
BIRD	<p>- Amprolium powder may be added in low dose in both feed (@25 g per 1000kg of mash) and drinking water (30g/ 25 lit of water) of poultry above 3 weeks of age.</p> <p>- Chayol 3 shuraba yendagina mapi faraba yen gi chinzak 1000kg da 25g amadi thaknaba ishing litre 25da 30g Amprolium powder happeniraga amangbada ee yaorakpagi lainadagi ngakthokpiyu.</p>



## Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



# Agromet Advisory Bulletin

Date : 18-02-2020

**Weather Forecast of THOUBAL(Manipur) Issued On : 2020-02-18(Valid Till 08:30 IST of the next 5 days)**

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-19	0.0	24.0	7.0	76	30	8.5	292	0
2020-02-20	0.0	24.0	7.0	70	26	5.7	292	0
2020-02-21	0.0	26.0	8.0	53	21	5.6	300	1
2020-02-22	0.0	26.0	9.0	65	24	5.1	27	4
2020-02-23	0.0	27.0	9.0	55	20	4.9	248	1

### **Weather Summary/Alert:**

Likely to have light rains on Saturday. The sky will remain mainly clear to partly cloudy with max. temp. 24-27°C & mini. temp. 7-9 °C. Relative humidity may reach upto 76 % (max) & 30 % (min). The average wind speed may be 4.9-8.5 km/hr.

Lakliba Thangjada nong khara churaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 24-27 °C amadi ingbagina 7-9 °C oiraba yai. Nungsitta ishingga manilgi changna yammabada ayukta 76 % amadi nunghildana 30 % oiraba yai. Nungsit sitpagi changna pung amada 4.9-8.5 km oiraba yai.

### **General Advisory:**

Sprays against insect pest may be given before rains. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free to preserve soil moisture for the crop. Keep sheds and surroundings of birds/ animals clean to avoid mosquitoes/ flies population.

Loufamda laina tilna ngakthoknaba hidak nong laktringeida kappiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Kang-hayinggi yamma indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nann thambiyu.

### **SMS Advisory:**

Spray Hexaconazole @ 2ml/L water against late blight in potato. Prepare land & mix well decomposed FYM @ 20-25 t/ha 15 days before planting ginger/ turmeric.

Alu da late blight ngakthoknaba ishing 1L da 2ml Hexaconazole happaga kappiyu. Shing/ Yaingang thadouba numit 15ni watlingeida pari 1da 20-25t patharaba FYM happyiu.

### **Crop Specific Advisory:**

Crop(Varieties)	Crop Specific Advisory
LENTIL	- Spray Bayleton @ 7g/ 10 litres of water or spray Neem Seed Kernel Extract at 5% concentration for powdery mildew and Nimbicidin (0.3%) to control rust disease. - Spray Imidachloprid 17.8 SL @ 7-10ml/ 15 litre of water against aphid.

Crop(Varieties)	Crop Specific Advisory
	- Ishing litre 10da 7g Bayleton happaga kappiyu natraga 5% concentration ta Neem Seed Kernel Extract at happaga chaibibada Kokfai kaba laina ngakthokpa yani. Kolhing kaba manbi laina ngakthoknaba Nimbicidin (0.3%) sijinabiyu. - Wahik ngakthoknaba ishing litre 15da 7-10 ml Imidachloprid 17.8 SL happaga kappiyu.
GARLIC	<p>- To prevent thrips in garlic crop install blue sticky trap in the field or spray Neem Seed Kernel extract @2ml/litre of water. Put a layer of mulch on the ground near the garlic plant to conserve soil moisture.</p> <p>Chanam mana maton ngangsinlakpa natraga wahik ngakthoknaba blue sticky trap sijinabiyu/ ishing litre amada 2ml Neem Seed Kernel Extract happaga kappiyu. Leibak leinil kannaba pambi mahongda charu/ una akangba thambiyu.</p>

### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
CHILLI	<p>- Raise chilies and capsicum nursery. Spray Carbendazim + Mancozeb (Sixer) @ 2g/litre of water to control the fungal attack.</p> <p>Morok Chaar thabiba houba yare. Kokfai kabi laina ngakthoknaba ishing litre amada 2g Carbendazim + Mancozeb (Sixer) happaga kappiyu.</p>

### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
BIRD	<p>- Amprolium powder may be added in low dose in both feed (@25 g per 1000kg of mash) and drinking water (30g/ 25 lit of water) of poultry above 3 weeks of age.</p> <p>- Chayol 3 shuraba yendagina mapi faraba yen gi chinzak 1000kg da 25g amadi thaknaba ishing litre 25da 30g Amprolium powder happeniraga amangbada ee yaorakpagi lainadagi ngakthokpiyu.</p>



## Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



# Agromet Advisory Bulletin

Date : 18-02-2020

**Weather Forecast of UKHRUL(Manipur) Issued On : 2020-02-18(Valid Till 08:30 IST of the next 5 days)**

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-19	0.0	25.0	8.0	74	19	6.7	22	0
2020-02-20	0.0	25.0	8.0	69	19	5.8	22	0
2020-02-21	0.0	26.0	8.0	59	20	5.6	22	1
2020-02-22	0.0	26.0	9.0	62	19	5.8	22	4
2020-02-23	0.0	27.0	9.0	55	23	5.7	22	2

### **Weather Summary/Alert:**

Likely to have light rains on Saturday. The sky will remain mainly clear to partly cloudy with max. temp. 25-27°C & mini. temp. 8-9 °C. Relative humidity may reach upto 74 % (max) & 23 % (min). The average wind speed may be 5.6-6.7km/hr.

Lakliba Thangjada nong khara churaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 25-27 °C amadi ingbagina 8-9 °C oiraba yai. Nungsitta ishingga manilgi changna yammabada ayukta 74 % amadi nunghildana 23 % oiraba yai. Nungsit sitpagi changna pung amada 5.6-6.7 km oiraba yai.

### **General Advisory:**

Sprays against insect pest may be given before rains. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free to preserve soil moisture for the crop. Keep sheds and surroundings of birds/ animals clean to avoid mosquitoes/ flies population.

Loufamda laina tilna ngakthoknaba hidak nong laktringeida kappiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Kang-hayinggi yamma indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nann thambiyu.

### **SMS Advisory:**

Spray Hexaconazole @ 2ml/L water against late blight in potato. Prepare land & mix well decomposed FYM @ 20-25 t/ha 15 days before planting ginger/ turmeric.

Alu da late blight ngakthoknaba ishing 1L da 2ml Hexaconazole happaga kappiyu. Shing/ Yaingang thadouba numit 15ni watlingeida pari 1da 20-25t patharaba FYM happyiu.

### **Crop Specific Advisory:**

Crop(Varieties)	Crop Specific Advisory
LENTIL	- Spray Bayleton @ 7g/ 10 litres of water or spray Neem Seed Kernel Extract at 5% concentration for powdery mildew and Nimbicidin (0.3%) to control rust disease. - Spray Imidachloprid 17.8 SL @ 7-10ml/ 15 litre of water against aphid.

Crop(Varieties)	Crop Specific Advisory
	- Ishing litre 10da 7g Bayleton happaga kappiyu natraga 5% concentration ta Neem Seed Kernel Extract at happaga chaibibada Kokfai kaba laina ngakthokpa yani. Kolhing kaba manbi laina ngakthoknaba Nimbicidin (0.3%) sijinabiyu. - Wahik ngakthoknaba ishing litre 15da 7-10 ml Imidachloprid 17.8 SL happaga kappiyu.
GARLIC	- To prevent thrips in garlic crop install blue sticky trap in the field or spray Neem Seed Kernel extract @2ml/litre of water. Put a layer of mulch on the ground near the garlic plant to conserve soil moisture. Chanam mana maton ngangsinlakpa natraga wahik ngakthoknaba blue sticky trap sijinabiyu/ ishing litre amada 2ml Neem Seed Kernel Extract happaga kappiyu. Leibak leinil kannaba pambi mahongda charu/ una akangba thambiyu.

### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
CHILLI	- Raise chilies and capsicum nursery. Spray Carbendazim + Mancozeb (Sixer) @ 2g/litre of water to control the fungal attack. Morok Chaar thabiba houba yare. Kokfai kabi laina ngakthoknaba ishing litre amada 2g Carbendazim + Mancozeb (Sixer) happaga kappiyu.

### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
BIRD	- Amprolium powder may be added in low dose in both feed (@25 g per 1000kg of mash) and drinking water (30g/ 25 lit of water) of poultry above 3 weeks of age. - Chayol 3 shuraba yendagina mapi faraba yen gi chinzak 1000kg da 25g amadi thaknaba ishing litre 25da 30g Amprolium powder happeniraga amangbada ee yaorakpagi lainadagi ngakthokpiyu.



**Gramin Krishi Mausam Sewa**  
 Experimental Block Level Agromet Advisory Bulletin  
 (A Joint Initiative of IMD & ICAR)



## Agromet Advisory Bulletin

Date : 18-02-2020

**Weather Forecast of WANGOI Block in IMPHAL-WEST(Manipur) Issued On :2020-02-18(Valid Till 08:30 IST of the next 5 days)**

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-19	0.0	26.9	12.5	88	34	7.0	296	0
2020-02-20	0.0	27.1	13.0	73	25	5.0	297	0
2020-02-21	0.0	27.8	13.2	52	22	5.0	270	1
2020-02-22	0.0	28.2	15.4	63	21	4.0	27	5
2020-02-23	0.0	29.8	15.2	59	20	4.0	162	1

### **Weather Summary/Alert:**

Likely to have light rains on Saturday. The sky will remain mainly clear to partly cloudy with max. temp. 26.9-29.8°C & mini. temp. 12.5-15.4 °C. Relative humidity may reach upto 88% (max) & 34% (min). The average wind speed may be 4-7km/hr.

Lakliba Thangjada nong khara churaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 26.9-29.8 °C amadi ingbagina 12.5-15.4 °C oiraba yai. Nungsitta ishingga manilgi changna yammabada ayukta 88% amadi nungthildana 34 % oiraba yai. Nungsit sitpagi changna pung amada 4-7 km oiraba yai.

### **General Advisory:**

Sprays against insect pest may be given before rains. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free to preserve soil moisture for the crop. Keep sheds and surroundings of birds/ animals clean to avoid mosquitoes/ flies population.

Loufamda laina tilna ngakthoknaba hidak nong laktringeida kappiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Kang-hayinggi yamma indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nann thambiyu.

### **SMS Advisory:**

Spray Hexaconazole @ 2ml/L water against late blight in potato. Prepare land & mix well decomposed FYM @ 20-25 t/ha 15 days before planting ginger/ turmeric.

Alu da late blight ngakthoknaba ishing 1L da 2ml Hexaconazole happaga kappiyu. Shing/ Yaingang thadouba numit 15ni watlingeida pari 1da 20-25t patharaba FYM happyiu.

### **Crop Specific Advisory:**

Crop(Varieties)	Crop Specific Advisory
LENTIL	- Spray Bayleton @ 7g/ 10 litres of water or spray Neem Seed Kernel Extract at 5% concentration for powdery mildew and Nimbicidin (0.3%) to control rust disease. - Spray Imidachloprid 17.8 SL @ 7-10ml/ 15 litre of water against aphid.
	- Isching litre 10da 7g Bayleton happaga kappiyu natraga 5% concentration ta Neem Seed Kernel Extract at happaga chaibibada Kokfai kaba laina ngakthokpa yani. Kolhing kaba manbi laina ngakthoknaba Nimbicidin (0.3%) sijinabiyu. - Wahik ngakthoknaba ishing litre 15da 7-10 ml Imidachloprid 17.8 SL happaga kappiyu.
GARLIC	- To prevent thrips in garlic crop install blue sticky trap in the field or spray Neem Seed Kernel extract @2ml/litre of water. Put a layer of mulch on the ground near the garlic plant to conserve soil moisture.
	Chanam mana maton ngangsinlakpa natraga wahik ngakthoknaba blue sticky trap sijinabiyu/ ishing litre amada 2ml Neem Seed Kernel Extract happaga kappiyu. Leibak leinil kannaba pambi mahongda charu/ una akangba thambiyu.

**Horticulture Specific Advisory:**

<b>Horticulture (Varieties)</b>	<b>Horticulture Specific Advisory</b>
CHILLI	<ul style="list-style-type: none"> <li>- Raise chilies and capsicum nursery. Spray Carbendazim + Mancozeb (Sixer) @ 2g/litre of water to control the fungal attack.</li> <li>Morok Chaar thabiba houba yare. Kokfai kabi laina ngakthoknaba ishing litre amada 2g Carbendazim + Mancozeb (Sixer) happaga kappiyu.</li> </ul>

**Poultry Specific Advisory:**

<b>Poultry(Varieties)</b>	<b>Poultry Specific Advisory</b>
BIRD	<ul style="list-style-type: none"> <li>- Amprolium powder may be added in low dose in both feed (@25 g per 1000kg of mash) and drinking water (30g/ 25 lit of water) of poultry above 3 weeks of age.</li> <li>- Chayol 3 shuraba yendagina mapi faraba yen gi chinzak 1000kg da 25g amadi thaknaba ishing litre25da 30g Amprolium powder happiraga amangbada ee yaorakpagi lainadagi ngakthokpiyu.</li> </ul>



## Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



# Agromet Advisory Bulletin

Date : 18-02-2020

**Weather Forecast of BISHNUPUR(Manipur) Issued On : 2020-02-18(Valid Till 08:30 IST of the next 5 days)**

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-19	0.0	24.0	7.0	75	30	3.5	292	0
2020-02-20	0.0	24.0	7.0	73	25	3.3	292	0
2020-02-21	0.0	26.0	8.0	55	20	3.5	288	1
2020-02-22	0.0	26.0	9.0	65	19	2.4	112	3
2020-02-23	0.0	27.0	9.0	58	19	3.1	112	1

### **Weather Summary/Alert:**

Likely to have light rains on Saturday. The sky will remain mainly clear to partly cloudy with max. temp. 24-27°C & mini. temp. 7-9 °C. Relative humidity may reach upto 75 % (max) & 30 % (min). The average wind speed may be 2.4-3.5 km/hr.

Lakliba Thangjada nong khara churaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 24-27 °C amadi ingbagina 7-9 °C oiraba yai. Nungsitta ishingga manilgi changna yammabada ayukta 75 % amadi nunghildana 30 % oiraba yai. Nungsit sitpagi changna pung amada 2.4-3.5 km oiraba yai.

### **General Advisory:**

Sprays against insect pest may be given before rains. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free to preserve soil moisture for the crop. Keep sheds and surroundings of birds/ animals clean to avoid mosquitoes/ flies population.

Loufamda laina tilna ngakthoknaba hidak nong laktringeida kappiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Kang-hayinggi yamma indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nann thambiyu.

### **SMS Advisory:**

Spray Hexaconazole @ 2ml/L water against late blight in potato. Prepare land & mix well decomposed FYM @ 20-25 t/ha 15 days before planting ginger/ turmeric.

Alu da late blight ngakthoknaba ishing 1L da 2ml Hexaconazole happaga kappiyu. Shing/ Yaingang thadouba numit 15ni watlingeida pari 1da 20-25t patharaba FYM happyiu.

### **Crop Specific Advisory:**

Crop(Varieties)	Crop Specific Advisory
LENTIL	- Spray Bayleton @ 7g/ 10 litres of water or spray Neem Seed Kernel Extract at 5% concentration for powdery mildew and Nimbicidin (0.3%) to control rust disease. - Spray Imidachloprid 17.8 SL @ 7-10ml/ 15 litre of water against aphid.

Crop(Varieties)	Crop Specific Advisory
	- Ishing litre 10da 7g Bayleton happaga kappiyu natraga 5% concentration ta Neem Seed Kernel Extract at happaga chaibibada Kokfai kaba laina ngakthokpa yani. Kolhing kaba manbi laina ngakthoknaba Nimbicidin (0.3%) sijinabiyu. - Wahik ngakthoknaba ishing litre 15da 7-10 ml Imidachloprid 17.8 SL happaga kappiyu.
GARLIC	<p>- To prevent thrips in garlic crop install blue sticky trap in the field or spray Neem Seed Kernel extract @2ml/litre of water. Put a layer of mulch on the ground near the garlic plant to conserve soil moisture.</p> <p>Chanam mana maton ngangsinlakpa natraga wahik ngakthoknaba blue sticky trap sijinabiyu/ ishing litre amada 2ml Neem Seed Kernel Extract happaga kappiyu. Leibak leinil kannaba pambi mahongda charu/ una akangba thambiyu.</p>

### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
CHILLI	<p>- Raise chilies and capsicum nursery. Spray Carbendazim + Mancozeb (Sixer) @ 2g/ litre of water to control the fungal attack.</p> <p>Morok Chaar thabiba houba yare. Kokfai kabi laina ngakthoknaba ishing litre amada 2g Carbendazim + Mancozeb (Sixer) happaga kappiyu. Morok Chaar thabiba houba yare. Kokfai kabi laina ngakthoknaba ishing litre amada 2g Carbendazim + Mancozeb (Sixer) happaga kappiyu.</p>

### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
BIRD	<p>- Amprolium powder may be added in low dose in both feed (@25 g per 1000kg of mash) and drinking water (30g/ 25 lit of water) of poultry above 3 weeks of age.</p> <p>- Chayol 3 shuraba yendagina mapi faraba yen gi chinzak 1000kg da 25g amadi thaknaba ishing litre 25da 30g Amprolium powder happiraga amangbada ee yaorakpagi lainadagi ngakthokpiyu.</p>



## Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



# Agromet Advisory Bulletin

Date : 18-02-2020

**Weather Forecast of CHANDEL(Manipur) Issued On : 2020-02-18(Valid Till 08:30 IST of the next 5 days)**

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-19	0.0	24.0	7.0	77	26	7.4	294	0
2020-02-20	0.0	24.0	7.0	69	23	4.6	287	0
2020-02-21	0.0	26.0	8.0	53	20	3.9	287	0
2020-02-22	0.0	26.0	9.0	69	24	3.7	112	5
2020-02-23	0.0	27.0	9.0	53	20	4.1	279	2

### **Weather Summary/Alert:**

Likely to have light rains on Saturday. The sky will remain mainly clear to partly cloudy with max. temp. 24-27 oC & mini. temp. 7-9 oC. Relative humidity may reach upto 77 % (max) & 26% (min). The average wind speed may be 3.7-7.4 km/hr.

Lakliba Thangjada nong khara churaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 24-27 oC amadi ingbagina 7-9 oC oiraba yai. Nungsitta ishingga manilgi changna yammabada ayukta 77% amadi nunghildana 26% oiraba yai. Nungsit sitpagi changna pung amada 3.7-7.4 km oiraba yai.

### **General Advisory:**

Sprays against insect pest may be given before rains. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free to preserve soil moisture for the crop. Keep sheds and surroundings of birds/ animals clean to avoid mosquitoes/ flies population.

Loufamda laina tilna ngakthoknaba hidak nong laktringeida kappiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Kang-hayinggi yamma indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nann thambiyu.

### **SMS Advisory:**

Spray Hexaconazole @ 2ml/L water against late blight in potato. Prepare land & mix well decomposed FYM @ 20-25 t/ha 15 days before planting ginger/ turmeric.

Alu da late blight ngakthoknaba ishing 1L da 2ml Hexaconazole happaga kappiyu. Shing/ Yaingang thadouba numit 15ni watlingeida pari 1da 20-25t patharaba FYM happyiu.

### **Crop Specific Advisory:**

Crop(Varieties)	Crop Specific Advisory
LENTIL	- Spray Bayleton @ 7g/ 10 litres of water or spray Neem Seed Kernel Extract at 5% concentration for powdery mildew and Nimbicidin (0.3%) to control rust disease. - Spray Imidachloprid 17.8 SL @ 7-10ml/ 15 litre of water against aphid.

Crop(Varieties)	Crop Specific Advisory
	- Ishing litre 10da 7g Bayleton happaga kappiyu natraga 5% concentration ta Neem Seed Kernel Extract at happaga chaibibada Kokfai kaba laina ngakthokpa yani. Kolhing kaba manbi laina ngakthoknaba Nimbicidin (0.3%) sijinabiyu. - Wahik ngakthoknaba ishing litre 15da 7-10 ml Imidachloprid 17.8 SL happaga kappiyu.
GARLIC	<p>- To prevent thrips in garlic crop install blue sticky trap in the field or spray Neem Seed Kernel extract @2ml/litre of water. Put a layer of mulch on the ground near the garlic plant to conserve soil moisture.</p> <p>Chanam mana maton ngangsinlakpa natraga wahik ngakthoknaba blue sticky trap sijinabiyu/ ishing litre amada 2ml Neem Seed Kernel Extract happaga kappiyu. Leibak leinil kannaba pambi mahongda charu/ una akangba thambiyu.</p>

### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
CHILLI	<p>- Raise chilies and capsicum nursery. Spray Carbendazim + Mancozeb (Sixer) @ 2g/ litre of water to control the fungal attack.</p> <p>Morok Chaar thabiba houba yare. Kokfai kabi laina ngakthoknaba ishing litre amada 2g Carbendazim + Mancozeb (Sixer) happaga kappiyu. Morok Chaar thabiba houba yare. Kokfai kabi laina ngakthoknaba ishing litre amada 2g Carbendazim + Mancozeb (Sixer) happaga kappiyu.</p>

### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
BIRD	<p>- Amprolium powder may be added in low dose in both feed (@25 g per 1000kg of mash) and drinking water (30g/ 25 lit of water) of poultry above 3 weeks of age.</p> <p>- Chayol 3 shuraba yendagina mapi faraba yen gi chinzak 1000kg da 25g amadi thaknaba ishing litre 25da 30g Amprolium powder happiraga amangbada ee yaorakpagi lainadagi ngakthokpiyu.</p>

**Gramin Krishi Mausam Sewa**

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya

**Agromet Advisory Bulletin**

Date : 18-02-2020

Weather Forecast of CHURACHANDPUR(Manipur) Issued On : 2020-02-18(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-19	0.0	24.0	7.0	81	31	5.1	202	0
2020-02-20	0.0	24.0	7.0	73	25	4.6	288	0
2020-02-21	0.0	26.0	8.0	59	21	4.0	146	1
2020-02-22	0.0	26.0	9.0	69	21	2.8	124	4
2020-02-23	0.0	27.0	9.0	57	19	4.6	158	2

**Weather Summary/Alert:**

Likely to have light rains on Saturday. The sky will remain mainly clear to partly cloudy with max. temp. 24-27 oC & mini. temp. 7-9 oC. Relative humidity may reach upto 81% (max) & 31 % (min). The average wind speed may be 2.8-5.1 km/hr.

Lakliba Thangjada nong khara churaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 24-27 oC amadi ingbagina 7-9 oC oiraba yai. Nungsitta ishingga manilgi changna yammabada ayukta 81% amadi nunghildana 31 % oiraba yai. Nungsit sitpagi changna pung amada 2.8-5.1 km oiraba yai.

**General Advisory:**

Sprays against insect pest may be given before rains. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free to preserve soil moisture for the crop. Keep sheds and surroundings of birds/ animals clean to avoid mosquitoes/ flies population.

Loufamda laina tilna ngakthoknaba hidak nong laktringeida kappiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Kang-hayinggi yamma indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nann thambiyu.

**SMS Advisory:**

Spray Hexaconazole @ 2ml/L water against late blight in potato. Prepare land & mix well decomposed FYM @ 20-25 t/ha 15 days before planting ginger/ turmeric.

Alu da late blight ngakthoknaba ishing 1L da 2ml Hexaconazole happaga kappiyu. Shing/ Yaingang thadouba numit 15ni watlingeida pari 1da 20-25t patharaba FYM happyiu.

**Crop Specific Advisory:**

Crop(Varieties)	Crop Specific Advisory
LENTIL	- Spray Bayleton @ 7g/ 10 litres of water or spray Neem Seed Kernel Extract at 5% concentration for powdery mildew and Nimbicidin (0.3%) to control rust disease. - Spray Imidachloprid 17.8 SL @ 7-10ml/ 15 litre of water against aphid.

Crop(Varieties)	Crop Specific Advisory
	- Ishing litre 10da 7g Bayleton happaga kappiyu natraga 5% concentration ta Neem Seed Kernel Extract at happaga chaibibada Kokfai kaba laina ngakthokpa yani. Kolhing kaba manbi laina ngakthoknaba Nimbicidin (0.3%) sijinabiyu. - Wahik ngakthoknaba ishing litre 15da 7-10 ml Imidachloprid 17.8 SL happaga kappiyu.
GARLIC	<p>- To prevent thrips in garlic crop install blue sticky trap in the field or spray Neem Seed Kernel extract @2ml/litre of water. Put a layer of mulch on the ground near the garlic plant to conserve soil moisture.</p> <p>Chanam mana maton ngangsinlakpa natraga wahik ngakthoknaba blue sticky trap sijinabiyu/ ishing litre amada 2ml Neem Seed Kernel Extract happaga kappiyu. Leibak leinil kannaba pambi mahongda charu/ una akangba thambiyu.</p>

### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
CHILLI	<p>- Raise chilies and capsicum nursery. Spray Carbendazim + Mancozeb (Sixer) @ 2g/ litre of water to control the fungal attack.</p> <p>Morok Chaar thabiba houba yare. Kokfai kabi laina ngakthoknaba ishing litre amada 2g Carbendazim + Mancozeb (Sixer) happaga kappiyu. Morok Chaar thabiba houba yare. Kokfai kabi laina ngakthoknaba ishing litre amada 2g Carbendazim + Mancozeb (Sixer) happaga kappiyu.</p>

### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
BIRD	<p>- Amprolium powder may be added in low dose in both feed (@25 g per 1000kg of mash) and drinking water (30g/ 25 lit of water) of poultry above 3 weeks of age.</p> <p>- Chayol 3 shuraba yendagina mapi faraba yen gi chinzak 1000kg da 25g amadi thaknaba ishing litre 25da 30g Amprolium powder happiraga amangbada ee yaorakpagi lainadagi ngakthokpiyu.</p>



**Gramin Krishi Mausam Sewa**  
 Experimental Block Level Agromet Advisory Bulletin  
 (A Joint Initiative of IMD & ICAR)



## Agromet Advisory Bulletin

Date : 18-02-2020

Weather Forecast of HAORANGSABAL Block in IMPHAL-WEST(Manipur) Issued On :2020-02-18(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-19	0.0	26.7	12.6	88	33	7.0	338	0
2020-02-20	0.0	26.9	12.9	71	25	6.0	360	0
2020-02-21	0.0	27.8	13.3	52	20	6.0	360	2
2020-02-22	0.0	28.2	15.2	59	18	5.0	27	5
2020-02-23	0.0	29.6	15.4	60	20	6.0	27	1

### **Weather Summary/Alert:**

Likely to have light rains on Saturday. The sky will remain mainly clear to partly cloudy with max. temp. 26.7-29.6°C & mini. temp. 12.6-15.4 °C. Relative humidity may reach upto 88% (max) & 33% (min). The average wind speed may be 5-7km/hr.

Lakliba Thangjada nong khara churaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 26.7-29.6 °C amadi ingbagina 12.6-15.4 °C oiraba yai. Nungsitta ishingga manilgi changna yammabada ayukta 88% amadi nunghildana 33 % oiraba yai. Nungsit sitpagi changna pung amada 5-7 km oiraba yai.

### **General Advisory:**

Sprays against insect pest may be given before rains. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free to preserve soil moisture for the crop. Keep sheds and surroundings of birds/ animals clean to avoid mosquitoes/ flies population.

Loufamda laina tilna ngakthoknaba hidak nong laktringeida kappiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Kang-hayinggi yamma indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nann thambiyu.

### **SMS Advisory:**

Spray Hexaconazole @ 2ml/L water against late blight in potato. Prepare land & mix well decomposed FYM @ 20-25 t/ha 15 days before planting ginger/ turmeric.

Alu da late blight ngakthoknaba ishing 1L da 2ml Hexaconazole happaga kappiyu. Shing/ Yaingang thadouba numit 15ni watlingeida pari 1da 20-25t patharaba FYM happyiu.

### **Crop Specific Advisory:**

Crop(Varieties)	Crop Specific Advisory
LENTIL	- Spray Bayleton @ 7g/ 10 litres of water or spray Neem Seed Kernel Extract at 5% concentration for powdery mildew and Nimbicidin (0.3%) to control rust disease. - Spray Imidachloprid 17.8 SL @ 7-10ml/ 15 litre of water against aphid.
	- Isching litre 10da 7g Bayleton happaga kappiyu natraga 5% concentration ta Neem Seed Kernel Extract at happaga chaibibada Kokfai kaba laina ngakthokpa yani. Kolhing kaba manbi laina ngakthoknaba Nimbicidin (0.3%) sijinabiyu. - Wahik ngakthoknaba ishing litre 15da 7-10 ml Imidachloprid 17.8 SL happaga kappiyu.
GARLIC	- To prevent thrips in garlic crop install blue sticky trap in the field or spray Neem Seed Kernel extract @2ml/litre of water. Put a layer of mulch on the ground near the garlic plant to conserve soil moisture.
	Chanam mana maton ngangsinlakpa natraga wahik ngakthoknaba blue sticky trap sijinabiyu/ ishing litre amada 2ml Neem Seed Kernel Extract happaga kappiyu. Leibak leinil kannaba pambi mahongda charu/ una akangba thambiyu.

**Horticulture Specific Advisory:**

<b>Horticulture (Varieties)</b>	<b>Horticulture Specific Advisory</b>
CHILLI	<ul style="list-style-type: none"> <li>- Raise chilies and capsicum nursery. Spray Carbendazim + Mancozeb (Sixer) @ 2g/litre of water to control the fungal attack.</li> <li>Morok Chaar thabiba houba yare. Kokfai kabi laina ngakthoknaba ishing litre amada 2g Carbendazim + Mancozeb (Sixer) happaga kappiyu.</li> </ul>

**Poultry Specific Advisory:**

<b>Poultry(Varieties)</b>	<b>Poultry Specific Advisory</b>
BIRD	<ul style="list-style-type: none"> <li>- Amprolium powder may be added in low dose in both feed (@25 g per 1000kg of mash) and drinking water (30g/ 25 lit of water) of poultry above 3 weeks of age.</li> <li>- Chayol 3 shuraba yendagina mapi faraba yen gi chinzak 1000kg da 25g amadi thaknaba ishing litre25da 30g Amprolium powder happiraga amangbada ee yaorakpagi lainadagi ngakthokpiyu.</li> </ul>



**Gramin Krishi Mausam Sewa**  
 Experimental Block Level Agromet Advisory Bulletin  
 (A Joint Initiative of IMD & ICAR)



## Agromet Advisory Bulletin

Date : 18-02-2020

**Weather Forecast of HIYANGTHANG Block in IMPHAL-WEST(Manipur) Issued On :2020-02-18(Valid Till 08:30 IST of the next 5 days)**

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-19	0.0	26.9	12.5	88	34	7.0	330	0
2020-02-20	0.0	26.9	13.0	72	25	5.0	289	0
2020-02-21	0.0	27.9	13.3	51	21	6.0	270	1
2020-02-22	0.0	28.2	15.4	60	19	5.0	27	5
2020-02-23	0.0	29.6	15.3	59	20	5.0	23	1

### **Weather Summary/Alert:**

Likely to have light rains on Saturday. The sky will remain mainly clear to partly cloudy with max. temp. 26.9-29.6°C & mini. temp. 12.5-15.4 °C. Relative humidity may reach upto 88% (max) & 34% (min). The average wind speed may be 5-7km/hr.

Lakliba Thangjada nong khara churaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 26.9-29.6 °C amadi ingbagina 12.5-15.4 °C oiraba yai. Nungsitta ishingga manilgi changna yammabada ayukta 88% amadi nungthildana 34 % oiraba yai. Nungsit sitpagi changna pung amada 5-7 km oiraba yai.

### **General Advisory:**

Sprays against insect pest may be given before rains. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free to preserve soil moisture for the crop. Keep sheds and surroundings of birds/ animals clean to avoid mosquitoes/ flies population.

Loufamda laina tilna ngakthoknaba hidak nong laktringeida kappiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Kang-hayinggi yamma indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nann thambiyu.

### **SMS Advisory:**

Spray Hexaconazole @ 2ml/L water against late blight in potato. Prepare land & mix well decomposed FYM @ 20-25 t/ha 15 days before planting ginger/ turmeric.

Alu da late blight ngakthoknaba ishing 1L da 2ml Hexaconazole happaga kappiyu. Shing/ Yaingang thadouba numit 15ni watlingeida pari 1da 20-25t patharaba FYM happyiu.

### **Crop Specific Advisory:**

Crop(Varieties)	Crop Specific Advisory
LENTIL	- Spray Bayleton @ 7g/ 10 litres of water or spray Neem Seed Kernel Extract at 5% concentration for powdery mildew and Nimbicidin (0.3%) to control rust disease. - Spray Imidachloprid 17.8 SL @ 7-10ml/ 15 litre of water against aphid.
	- Isching litre 10da 7g Bayleton happaga kappiyu natraga 5% concentration ta Neem Seed Kernel Extract at happaga chaibibada Kokfai kaba laina ngakthokpa yani. Kolhing kaba manbi laina ngakthoknaba Nimbicidin (0.3%) sijinabiyu. - Wahik ngakthoknaba ishing litre 15da 7-10 ml Imidachloprid 17.8 SL happaga kappiyu.
GARLIC	- To prevent thrips in garlic crop install blue sticky trap in the field or spray Neem Seed Kernel extract @2ml/litre of water. Put a layer of mulch on the ground near the garlic plant to conserve soil moisture.
	Chanam mana maton ngangsinlakpa natraga wahik ngakthoknaba blue sticky trap sijinabiyu/ ishing litre amada 2ml Neem Seed Kernel Extract happaga kappiyu. Leibak leinil kannaba pambi mahongda charu/ una akangba thambiyu.

**Horticulture Specific Advisory:**

<b>Horticulture (Varieties)</b>	<b>Horticulture Specific Advisory</b>
CHILLI	<ul style="list-style-type: none"> <li>- Raise chilies and capsicum nursery. Spray Carbendazim + Mancozeb (Sixer) @ 2g/litre of water to control the fungal attack.</li> <li>Morok Chaar thabiba houba yare. Kokfai kabi laina ngakthoknaba ishing litre amada 2g Carbendazim + Mancozeb (Sixer) happaga kappiyu.</li> </ul>

**Poultry Specific Advisory:**

<b>Poultry(Varieties)</b>	<b>Poultry Specific Advisory</b>
BIRD	<ul style="list-style-type: none"> <li>- Amprolium powder may be added in low dose in both feed (@25 g per 1000kg of mash) and drinking water (30g/ 25 lit of water) of poultry above 3 weeks of age.</li> <li>- Chayol 3 shuraba yendagina mapi faraba yen gi chinzak 1000kg da 25g amadi thaknaba ishing litre25da 30g Amprolium powder happiraga amangbada ee yaorakpagi lainadagi ngakthokpiyu.</li> </ul>



## Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



# Agromet Advisory Bulletin

Date : 18-02-2020

**Weather Forecast of IMPHAL-EAST(Manipur) Issued On : 2020-02-18(Valid Till 08:30 IST of the next 5 days)**

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-19	0.0	24.0	7.0	71	27	7.5	330	0
2020-02-20	0.0	24.0	7.0	67	24	5.7	330	0
2020-02-21	0.0	26.0	8.0	51	19	6.2	0	1
2020-02-22	0.0	26.0	9.0	60	21	5.9	22	3
2020-02-23	0.0	27.0	9.0	52	19	5.3	248	0

### **Weather Summary/Alert:**

Likely to have light rains on Saturday. The sky will remain mainly clear to partly cloudy with max. temp. 24-27°C & mini. temp. 7-9 °C. Relative humidity may reach upto 71 % (max) & 27 % (min). The average wind speed may be 5.3-7.5 km/hr.

Lakliba Thangjada nong khara churaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 24-27 °C amadi ingbagina 7-9 °C oiraba yai. Nungsitta ishingga manilgi changna yammabada ayukta 71 % amadi nunghildana 27 % oiraba yai. Nungsit sitpagi changna pung amada 5.3-7.5km oiraba yai.

### **General Advisory:**

Sprays against insect pest may be given before rains. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free to preserve soil moisture for the crop. Keep sheds and surroundings of birds/ animals clean to avoid mosquitoes/ flies population.

Loufamda laina tilna ngakthoknaba hidak nong laktringeida kappiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Kang-hayinggi yamma indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nann thambiyu.

### **SMS Advisory:**

Spray Hexaconazole @ 2ml/L water against late blight in potato. Prepare land & mix well decomposed FYM @ 20-25 t/ha 15 days before planting ginger/ turmeric.

Alu da late blight ngakthoknaba ishing 1L da 2ml Hexaconazole happaga kappiyu. Shing/ Yaingang thadouba numit 15ni watlingeida pari 1da 20-25t patharaba FYM happyiu.

### **Crop Specific Advisory:**

Crop(Varieties)	Crop Specific Advisory
LENTIL	- Spray Bayleton @ 7g/ 10 litres of water or spray Neem Seed Kernel Extract at 5% concentration for powdery mildew and Nimbicidin (0.3%) to control rust disease. - Spray Imidachloprid 17.8 SL @ 7-10ml/ 15 litre of water against aphid.

Crop(Varieties)	Crop Specific Advisory
	- Ishing litre 10da 7g Bayleton happaga kappiyu natraga 5% concentration ta Neem Seed Kernel Extract at happaga chaibibada Kokfai kaba laina ngakthokpa yani. Kolhing kaba manbi laina ngakthoknaba Nimbicidin (0.3%) sijinabiyu. - Wahik ngakthoknaba ishing litre 15da 7-10 ml Imidachloprid 17.8 SL happaga kappiyu.
GARLIC	<p>- To prevent thrips in garlic crop install blue sticky trap in the field or spray Neem Seed Kernel extract @2ml/litre of water. Put a layer of mulch on the ground near the garlic plant to conserve soil moisture.</p> <p>Chanam mana maton ngangsinlakpa natraga wahik ngakthoknaba blue sticky trap sijinabiyu/ ishing litre amada 2ml Neem Seed Kernel Extract happaga kappiyu. Leibak leinil kannaba pambi mahongda charu/ una akangba thambiyu.</p>

### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
CHILLI	<p>- Raise chilies and capsicum nursery. Spray Carbendazim + Mancozeb (Sixer) @ 2g/ litre of water to control the fungal attack.</p> <p>Morok Chaar thabiba houba yare. Kokfai kabi laina ngakthoknaba ishing litre amada 2g Carbendazim + Mancozeb (Sixer) happaga kappiyu. Morok Chaar thabiba houba yare. Kokfai kabi laina ngakthoknaba ishing litre amada 2g Carbendazim + Mancozeb (Sixer) happaga kappiyu.</p>

### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
BIRD	<p>- Amprolium powder may be added in low dose in both feed (@25 g per 1000kg of mash) and drinking water (30g/ 25 lit of water) of poultry above 3 weeks of age.</p> <p>- Chayol 3 shuraba yendagina mapi faraba yen gi chinzak 1000kg da 25g amadi thaknaba ishing litre 25da 30g Amprolium powder happiraga amangbada ee yaorakpagi lainadagi ngakthokpiyu.</p>



## Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



# Agromet Advisory Bulletin

Date : 18-02-2020

**Weather Forecast of IMPHAL-WEST(Manipur) Issued On : 2020-02-18(Valid Till 08:30 IST of the next 5 days)**

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-19	0.0	25.0	8.0	70	27	7.1	252	0
2020-02-20	0.0	25.0	8.0	66	23	6.0	315	0
2020-02-21	0.0	26.0	8.0	53	19	6.6	15	2
2020-02-22	0.0	26.0	9.0	58	19	6.0	22	3
2020-02-23	0.0	27.0	9.0	52	19	5.8	243	1

### **Weather Summary/Alert:**

Likely to have light rains on Saturday. The sky will remain mainly clear to partly cloudy with max. temp. 25-27°C & mini. temp. 8-9 °C. Relative humidity may reach upto 70 % (max) & 27 % (min). The average wind speed may be 5.8-7.1km/hr.

Lakliba Thangjada nong khara churaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 24-27 °C amadi ingbagina 7-9 °C oiraba yai. Nungsitta ishingga manilgi changna yammabada ayukta 70 % amadi nunghildana 27 % oiraba yai. Nungsit sitpagi changna pung amada 5.8-7.1km oiraba yai.

### **General Advisory:**

Sprays against insect pest may be given before rains. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free to preserve soil moisture for the crop. Keep sheds and surroundings of birds/ animals clean to avoid mosquitoes/ flies population.

Loufamda laina tilna ngakthoknaba hidak nong laktringeida kappiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Kang-hayinggi yamma indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nann thambiyu.

### **SMS Advisory:**

Spray Hexaconazole @ 2ml/L water against late blight in potato. Prepare land & mix well decomposed FYM @ 20-25 t/ha 15 days before planting ginger/ turmeric.

Alu da late blight ngakthoknaba ishing 1L da 2ml Hexaconazole happaga kappiyu. Shing/ Yaingang thadouba numit 15ni watlingeida pari 1da 20-25t patharaba FYM happyiu.

### **Crop Specific Advisory:**

Crop(Varieties)	Crop Specific Advisory
LENTIL	- Spray Bayleton @ 7g/ 10 litres of water or spray Neem Seed Kernel Extract at 5% concentration for powdery mildew and Nimbicidin (0.3%) to control rust disease. - Spray Imidachloprid 17.8 SL @ 7-10ml/ 15 litre of water against aphid.

Crop(Varieties)	Crop Specific Advisory
	- Ishing litre 10da 7g Bayleton happaga kappiyu natraga 5% concentration ta Neem Seed Kernel Extract at happaga chaibibada Kokfai kaba laina ngakthokpa yani. Kolhing kaba manbi laina ngakthoknaba Nimbicidin (0.3%) sijinabiyu. - Wahik ngakthoknaba ishing litre 15da 7-10 ml Imidachloprid 17.8 SL happaga kappiyu.
GARLIC	<p>- To prevent thrips in garlic crop install blue sticky trap in the field or spray Neem Seed Kernel extract @2ml/litre of water. Put a layer of mulch on the ground near the garlic plant to conserve soil moisture.</p> <p>Chanam mana maton ngangsinlakpa natraga wahik ngakthoknaba blue sticky trap sijinabiyu/ ishing litre amada 2ml Neem Seed Kernel Extract happaga kappiyu. Leibak leinil kannaba pambi mahongda charu/ una akangba thambiyu.</p>

### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
CHILLI	<p>- Raise chilies and capsicum nursery. Spray Carbendazim + Mancozeb (Sixer) @ 2g/ litre of water to control the fungal attack.</p> <p>Morok Chaar thabiba houba yare. Kokfai kabi laina ngakthoknaba ishing litre amada 2g Carbendazim + Mancozeb (Sixer) happaga kappiyu. Morok Chaar thabiba houba yare. Kokfai kabi laina ngakthoknaba ishing litre amada 2g Carbendazim + Mancozeb (Sixer) happaga kappiyu.</p>

### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
BIRD	<p>- Amprolium powder may be added in low dose in both feed (@25 g per 1000kg of mash) and drinking water (30g/ 25 lit of water) of poultry above 3 weeks of age.</p> <p>- Chayol 3 shuraba yendagina mapi faraba yen gi chinzak 1000kg da 25g amadi thaknaba ishing litre 25da 30g Amprolium powder happiraga amangbada ee yaorakpagi lainadagi ngakthokpiyu.</p>

**Gramin Krishi Mausam Sewa**

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya

**Agromet Advisory Bulletin**

Date : 18-02-2020

Weather Forecast of JIRIBAM(Manipur) Issued On : 2020-02-18(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-19	0.0	27.0	13.0	67	30	4.8	65	0
2020-02-20	0.0	28.0	13.0	65	26	4.5	65	0
2020-02-21	0.0	29.0	13.0	55	21	4.6	65	3
2020-02-22	0.0	30.0	14.0	63	17	4.1	65	3
2020-02-23	0.0	31.0	16.0	51	18	4.3	68	1

**Weather Summary/Alert:**

Likely to have light rains on Saturday. The sky will remain mainly clear to partly cloudy with max. temp. 27-31 oC & mini. temp. 13-16 oC. Relative humidity may reach upto 67 % (max) & 30 % (min). The average wind speed may be 4.1-4.8 km/hr.

Lakliba Thangjada nong khara churaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 27-31 oC amadi ingbagina 13-16 oC oiraba yai. Nungsitta ishingga manilgi changna yammabada ayukta 67 % amadi nunghildana 30 % oiraba yai. Nungsit sitpagi changna pung amada 4.1-4.8 km oiraba yai.

**General Advisory:**

Sprays against insect pest may be given before rains. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free to preserve soil moisture for the crop. Keep sheds and surroundings of birds/ animals clean to avoid mosquitoes/ flies population.

Loufamda laina tilna ngakthoknaba hidak nong laktringeida kappiyu. Nong churakpada leibak leinil leihounaba pambi makhongda laina tilna paktaba charu usa-wasa na kupsinbiyu. Kang-hayinggi yamma indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nann thambiyu.

**SMS Advisory:**

Spray Hexaconazole @ 2ml/L water against late blight in potato. Prepare land & mix well decomposed FYM @ 20-25 t/ha 15 days before planting ginger/ turmeric.

Alu da late blight ngakthoknaba ishing 1L da 2ml Hexaconazole happaga kappiyu. Shing/ Yaingang thadouba numit 15ni watlingeida pari 1da 20-25t patharaba FYM happyiu.

**Crop Specific Advisory:**

Crop(Varieties)	Crop Specific Advisory
GARLIC	- To prevent thrips in garlic crop install blue sticky trap in the field or spray Neem Seed Kernel extract @2ml/litre of water. Put a layer of mulch on the ground near the garlic plant to conserve soil moisture.
	Chanam mana maton ngangsinlakpa natraga wahik ngakthoknaba blue sticky trap sijinabiyu/ ishing litre amada 2ml Neem Seed Kernel Extract happaga kappiyu. Leibak leinil kannaba pambi mahongda charu/ una akangba thambiyu.

Crop(Varieties)	Crop Specific Advisory
LENTIL	- Spray Bayleton @ 7g/ 10 litres of water or spray Neem Seed Kernel Extract at 5% concentration for powdery mildew and Nimbicidin (0.3%) to control rust disease. - Spray Imidachloprid 17.8 SL @ 7-10ml/ 15 litre of water against aphid.
	- Ishing litre 10da 7g Bayleton happaga kappiyu natraga 5% concentration ta Neem Seed Kernel Extract at happaga chaibibada Kokfai kaba laina ngakthokpa yani. Kolhing kaba manbi laina ngakthoknaba Nimbicidin (0.3%) sijinabiyu. - Wahik ngakthoknaba ishing litre 15da 7-10 ml Imidachloprid 17.8 SL happaga kappiyu.

### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
CHILLI	- Raise chilies and capsicum nursery. Spray Carbendazim + Mancozeb (Sixer) @ 2g/litre of water to control the fungal attack.
	Morok Chaar thabiba houba yare. Kokfai kabi laina ngakthoknaba ishing litre amada 2g Carbendazim + Mancozeb (Sixer) happaga kappiyu.

### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
BIRD	<ul style="list-style-type: none"> <li>- Amprolium powder may be added in low dose in both feed (@25 g per 1000kg of mash) and drinking water (30g/ 25 lit of water) of poultry above 3 weeks of age.</li> <li>- Chayol 3 shuraba yendagina mapi faraba yen gi chinzak 1000kg da 25g amadi thaknaba ishing litre 25da 30g Amprolium powder happeniraga amangbada ee yaorakpagi lainadagi ngakthokpiyu.</li> </ul>