



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 13-03-2020

Weather Forecast of KANGPOKPI(Manipur) Issued On : 2020-03-13(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-03-14	0.0	25.0	10.0	61	20	5.2	19	2
2020-03-15	0.0	25.0	10.0	66	23	4.7	19	6
2020-03-16	1.0	25.0	9.0	66	25	4.8	11	5
2020-03-17	0.0	25.0	10.0	57	19	4.9	248	6
2020-03-18	0.0	26.0	10.0	61	30	4.6	19	3

Weather Summary/Alert:

Likely to have no light rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 25-26 oC & mini. temp. 9-10 oC. Relative humidity may reach upto 66% (max) & 19% (min). The average wind speed may be 4.6-5.2 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 25-26 oC amadi ingbagina 9-10oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 66% amadi nunghildana 19% oiraba yai. Nungsit sitpangi changna pung amada 4.6-5.2 km oiraba yai.

General Advisory:

Disinfect the farm premises of all livestock/ poultry (surrounding area of 3ft width) using the repellent agent/ insecticides spray. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free or polythene sheets to preserve soil moisture for the crop. Monitor the soil moisture status regularly and give life-saving irrigation, especially at growth stages when water requirement is higher.

Kang-hayinggi yamna indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nanna thambiyu. Leibak leinil kuina leinaba pambi makhongda laina tilna paktaba charu usa-wasa natraga polythene na kupsinbiyu. Loufamda pambi houbada ishinggi awatpa leitanaba toina sennabiraga darkar leibada ishing thajinbiyu.

SMS Advisory:

Harvest the mature crops like rapeseed/ pea/ lentil/ broad bean/ potato; select pure healthy insect pest free seeds & sundry properly before storing as seed.

Hangam, hawai tharak/ moobi, masoori amadi alu lokpiraga laina paktaba mapung faba mahei khangatpiraga nungshada ningthina kanghanlaga maru oinaba thamjinbiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
-----------------	------------------------

Crop(Varieties)	Crop Specific Advisory
SOYABEAN	Procure insect pest free healthy seeds of soybean/ groundnut. Get the soil testing done for proper fertilizer application. If soil is acidic i.e., PH level is below 6.5 then liming in furrows is required @ 500kg/ ha for having better yield.
	Nung-hawai thanaba afaba mahal laina tilna paktaba maru khanlaga lousinbiba yare. Leibak leihao amadi PH gi changyeng toubiraga PH 6.5 gi makhada leirabadi pari amada 500kg shunu paring naina churi sembiraga happiradi pothook hengathanba ngamgani

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
ONION	Spray Sixer @ 2 gm/lit of water as prophylactic measure against Purple Bloch and bulb rot. Spray is to be given twice at an interval of 7 to 10 days.
	Tilhou manada angouba mapak paklakpa amadi kuisinlakpada khamen machu onsinlakpa natraga tilhou maru patharakpa laina ngakthoknaba numit 7-10ni kannaraga ishing litre amada 2g gi chaangda Sixer happaga kappiyu.

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	- Any type of influenza may be given symptomatic treatment with antibiotics. - Vaccinate against swine fever after the age of 3 months. - Give Ivermectin (Indectin tab.) @ 200mcg/ kg body weight or Ivermec 1% injection @ 1ml/50 kg body weight once only to avoid skin infection. - Vaccinate against F.M.D.
	- Lok thunglakpaga mawong matou ubiradi checksin thourang loubiraga matam chana hidak thabiyu. - Tha 3 shuraba oakta Swine Fever naraktanaba hidak kappiyu. - Unshada tourakpa ana-ayek ngakthoknaba masagi arumba kg amada 200mcg gi chaangda Ivermectin (Indectin tab.) pibiyu natraga masagi arumba kg 50 khudingda 1ml gi chaangda Ivermec 1% injection kappiyu. - F.M.D. laina ngakthoknaba hidak kappiyu.



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 13-03-2020

Weather Forecast of NONEY (Manipur) Issued On : 2020-03-13 (Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-03-14	0.0	25.0	10.0	55	18	4.9	68	2
2020-03-15	0.0	25.0	10.0	64	24	4.6	292	7
2020-03-16	0.0	25.0	10.0	56	25	4.5	63	6
2020-03-17	0.0	25.0	11.0	47	16	4.8	248	5
2020-03-18	0.0	26.0	11.0	58	24	6.1	63	3

Weather Summary/Alert:

Likely to have no light rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 25-26 oC & mini. temp. 10-11 oC. Relative humidity may reach upto 64% (max) & 16% (min). The average wind speed may be 4.5-6.1 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 25-26 oC amadi ingbagina 10-11 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 64% amadi nunghildana 16% oiraba yai. Nungsit sitpangi changna pung amada 4.5-6.1 km oiraba yai.

General Advisory:

Disinfect the farm premises of all livestock/ poultry (surrounding area of 3ft width) using the repellent agent/ insecticides spray. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free or polythene sheets to preserve soil moisture for the crop. Monitor the soil moisture status regularly and give life-saving irrigation, especially at growth stages when water requirement is higher.

Kang-hayinggi yamna indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nanna thambiyu. Leibak leinil kuina leinaba pambi makhongda laina tilna paktaba charu usa-wasa natraga polythene na kupsinbiyu. Loufamda pambi houbada ishinggi awatpa leitanaba toina sennabiraga darkar leibada ishing thajinbiyu.

SMS Advisory:

Harvest the mature crops like rapeseed/ pea/ lentil/ broad bean/ potato; select pure healthy insect pest free seeds & sundry properly before storing as seed.

Hangam, hawai tharak/ moobi, masoori amadi alu lokpiraga laina paktaba mapung faba mahei khangatpiraga nunghshada ningthina kanghanlaga maru oinaba thamjinbiyu.

Crop Specific Advisory:

Crop (Varieties)	Crop Specific Advisory
------------------	------------------------

Crop(Varieties)	Crop Specific Advisory
SOYABEAN	Procure insect pest free healthy seeds of soybean/ groundnut. Get the soil testing done for proper fertilizer application. If soil is acidic i.e., PH level is below 6.5 then liming in furrows is required @ 500kg/ ha for having better yield.
	Nung-hawai thanaba afaba mahal laina tilna paktaba maru khanlaga lousinbiba yare. Leibak leihao amadi PH gi changyeng toubiraga PH 6.5 gi makhada leirabadi pari amada 500kg shunu paring naina churi sembiraga happiradi pothook hengathanba ngamgani

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
ONION	Spray Sixer @ 2 gm/lit of water as prophylactic measure against Purple Bloch and bulb rot. Spray is to be given twice at an interval of 7 to 10 days.
	Tilhou manada angouba mapak paklakpa amadi kuisinlakpada khamen machu onsinlakpa natraga tilhou maru patharakpa laina ngakthoknaba numit 7-10ni kannaraga ishing litre amada 2g gi chaangda Sixer happaga kappiyu.

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	- Any type of influenza may be given symptomatic treatment with antibiotics. - Vaccinate against swine fever after the age of 3 months. - Give Ivermectin (Indectin tab.) @ 200mcg/ kg body weight or Ivermec 1% injection @ 1ml/50 kg body weight once only to avoid skin infection. - Vaccinate against F.M.D.
	- Lok thunglakpaga mawong matou ubiradi checksin thourang loubiraga matam chana hidak thabiyu. - Tha 3 shuraba oakta Swine Fever naraktanaba hidak kappiyu. - Unshada tourakpa ana-ayek ngakthoknaba masagi arumba kg amada 200mcg gi chaangda Ivermectin (Indectin tab.) pibiyu natraga masagi arumba kg 50 khudingda 1ml gi chaangda Ivermec 1% injection kappiyu. - F.M.D. laina ngakthoknaba hidak kappiyu.



Gramin Krishi Mausam Sewa
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 13-03-2020

Weather Forecast of PATSOI Block in IMPHAL-WEST(Manipur) Issued On :2020-03-13(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-03-14	0.0	33.4	19.6	63	22	6.0	360	3
2020-03-15	0.1	34.7	19.7	73	22	6.0	248	3
2020-03-16	0.0	33.0	17.3	77	27	9.0	283	2
2020-03-17	0.0	29.9	16.0	70	20	7.0	248	6
2020-03-18	0.0	31.6	15.6	58	24	7.0	248	4

Weather Summary/Alert:

Likely to have no light rains upto Wednesday. The sky will remain mainly clear to partly cloudy but mainly cloudy on Tuesday with max. temp. 29.9-33.4 oC & mini. temp. 15.6-19.7 oC. Relative humidity may reach upto 77% (max) & 27 % (min). The average wind speed may be 6-9 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai adubu Leibakpokpadadi khara yamna khanlaba yai. Khwaidagi sabagi changna 29.9-33.4 oC amadi ingbagina 15.6-19.7 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 77% amadi nungthildana 27 % oiraba yai. Nungsit sitpagi changna pung amada 6-9 km oiraba yai.

General Advisory:

Disinfect the farm premises of all livestock/ poultry (surrounding area of 3ft width) using the repellent agent/ insecticides spray. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free or polythene sheets to preserve soil moisture for the crop. Monitor the soil moisture status regularly and give life-saving irrigation, especially at growth stages when water requirement is higher.

Kang-hayinggi yamna indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nanna thambiyeu. Leibak leinil kuina leinaba pambi makhongda laina tilna paktaba charu usa-wasa natraga polythene na kupsinbiyu. Loufamda pambi houbada ishinggi awatpa leitanaba toina sennabiraga darkar leibada ishing thajinbiyu.

SMS Advisory:

Harvest the mature crops like rapeseed/ pea/ lentil/ broad bean/ potato; select pure healthy insect pest free seeds & sundry properly before storing as seed.

Hangam, hawai tharak/ moobi, masoori amadi alu lokpiraga laina paktaba mapung faba mahei khangatpiraga nungshada ningthina kanghanlaga maru oinaba thamjinbiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
SOYABEAN	Procure insect pest free healthy seeds of soybean/ groundnut. Get the soil testing done for proper fertilizer application. If soil is acidic i.e., PH level is below 6.5 then liming in furrows is required @ 500kg/ ha for having better yield.
	Nung-hawai thanaba afaba mahal laina tilna paktaba maru khanlaga lousinbiba yare. Leibak leihao amadi PH gi changyeng toubiraga PH 6.5 gi makhada leirabadi pari amada 500kg shunu paring naina churi sembiraga happiradi pothook hengathanba ngamgani.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
ONION	<p>Spray Sixer @ 2 gm/lit of water as prophylactic measure against Purple Bloch and bulb rot. Spray is to be given twice at an interval of 7 to 10 days.</p> <p>Tilhou manada angouba mapak paklakpa amadi kuisinlakpada khamen machu onsinlakpa natraga tilhou maru patharakpa laina ngakthoknaba numit 7-10ni kannaraga ishing litre amada 2g gi chaangda Sixer happaga kappiyu.</p>

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	<p>- Any type of influenza may be given symptomatic treatment with antibiotics. - Vaccinate against swine fever after the age of 3 months. - Give Ivermectin (Indectin tab.) @ 200mcg/ kg body weight or Ivermec 1% injection @ 1ml/50 kg body weight once only to avoid skin infection. - Vaccinate against F.M.D.</p> <p>- Lok thunglakpagi mawong matou ubiradi checksin thourang loubiraga matam chana hidak thabiyu. - Tha 3 shuraba oakta Swine Fever naraktanaba hidak kappiyu. - Unshada tourakpa ana-ayek ngakthoknaba masagi arumba kg amada 200mcg gi chaangda Ivermectin (Indectin tab.) pibiyu natraga masagi arumba kg 50 khudingda 1ml gi chaangda Ivermec 1% injection kappiyu. - F.M.D. laina ngakthoknaba hidak kappiyu.</p>



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 13-03-2020

Weather Forecast of PHERZAWL(Manipur) Issued On : 2020-03-13(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-03-14	0.0	25.0	10.0	52	18	3.9	109	2
2020-03-15	0.0	25.0	10.0	65	25	4.6	112	7
2020-03-16	0.0	25.0	10.0	57	23	3.9	112	5
2020-03-17	0.0	25.0	11.0	44	16	4.4	116	5
2020-03-18	0.0	26.0	11.0	54	21	5.5	112	4

Weather Summary/Alert:

Likely to have no light rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 25-26 oC & mini. temp. 10-11 oC. Relative humidity may reach upto 65% (max) & 16% (min). The average wind speed may be 3.9-5.5 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 25-26 oC amadi ingbagina 10-11 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 65% amadi nunghildana 16% oiraba yai. Nungsit sitpangi changna pung amada 3.9-5.5 km oiraba yai.

General Advisory:

Disinfect the farm premises of all livestock/ poultry (surrounding area of 3ft width) using the repellent agent/ insecticides spray. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free or polythene sheets to preserve soil moisture for the crop. Monitor the soil moisture status regularly and give life-saving irrigation, especially at growth stages when water requirement is higher.

Kang-hayinggi yamna indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nanna thambiyu. Leibak leinil kuina leinaba pambi makhongda laina tilna paktaba charu usa-wasa natraga polythene na kupsinbiyu. Loufamda pambi houbada ishinggi awatpa leitanaba toina sennabiraga darkar leibada ishing thajinbiyu.

SMS Advisory:

Harvest the mature crops like rapeseed/ pea/ lentil/ broad bean/ potato; select pure healthy insect pest free seeds & sundry properly before storing as seed.

Hangam, hawai tharak/ moobi, masoori amadi alu lokpiraga laina paktaba mapung faba mahei khangatpiraga nunghshada ningthina kanghanlaga maru oinaba thamjinbiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
-----------------	------------------------

Crop(Varieties)	Crop Specific Advisory
SOYABEAN	Procure insect pest free healthy seeds of soybean/ groundnut. Get the soil testing done for proper fertilizer application. If soil is acidic i.e., PH level is below 6.5 then liming in furrows is required @ 500kg/ ha for having better yield.
	Nung-hawai thanaba afaba mahal laina tilna paktaba maru khanlaga lousinbiba yare. Leibak leihao amadi PH gi changyeng toubiraga PH 6.5 gi makhada leirabadi pari amada 500kg shunu paring naina churi sembiraga happiradi pothook hengathanba ngamgani

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
ONION	Spray Sixer @ 2 gm/lit of water as prophylactic measure against Purple Bloch and bulb rot. Spray is to be given twice at an interval of 7 to 10 days.
	Tilhou manada angouba mapak paklakpa amadi kuisinlakpada khamen machu onsinlakpa natraga tilhou maru patharakpa laina ngakthoknaba numit 7-10ni kannaraga ishing litre amada 2g gi chaangda Sixer happaga kappiyu.

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	- Any type of influenza may be given symptomatic treatment with antibiotics. - Vaccinate against swine fever after the age of 3 months. - Give Ivermectin (Indectin tab.) @ 200mcg/ kg body weight or Ivermec 1% injection @ 1ml/50 kg body weight once only to avoid skin infection. - Vaccinate against F.M.D.
	- Lok thunglakpaga mawong matou ubiradi checksin thourang loubiraga matam chana hidak thabiyu. - Tha 3 shuraba oakta Swine Fever naraktanaba hidak kappiyu. - Unshada tourakpa ana-ayek ngakthoknaba masagi arumba kg amada 200mcg gi chaangda Ivermectin (Indectin tab.) pibiyu natraga masagi arumba kg 50 khudingda 1ml gi chaangda Ivermec 1% injection kappiyu. - F.M.D. laina ngakthoknaba hidak kappiyu.



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 13-03-2020

Weather Forecast of SENAPATI(Manipur) Issued On : 2020-03-13(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-03-14	0.0	25.0	9.0	68	24	4.8	20	2
2020-03-15	2.0	24.0	8.0	74	27	4.5	15	7
2020-03-16	3.0	24.0	8.0	73	30	3.9	19	7
2020-03-17	0.0	25.0	9.0	63	24	4.3	252	6
2020-03-18	0.0	26.0	10.0	66	36	3.7	19	4

Weather Summary/Alert:

Likely to have no light rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 24-26 oC & mini. temp. 8-10 oC. Relative humidity may reach upto 74% (max) & 24% (min). The average wind speed may be 3.7-4.8 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 28-30 oC amadi ingbagina 12-13 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 62 % amadi nungthildana 25 % oiraba yai. Nungsit sitpagi changna pung amada 6-8 km oiraba yai.

General Advisory:

Disinfect the farm premises of all livestock/ poultry (surrounding area of 3ft width) using the repellent agent/ insecticides spray. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free or polythene sheets to preserve soil moisture for the crop. Monitor the soil moisture status regularly and give life-saving irrigation, especially at growth stages when water requirement is higher.

Kang-hayinggi yamna indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nanna thambiyu. Leibak leinil kuina leinaba pambi makhongda laina tilna paktaba charu usa-wasa natraga polythene na kupsinbiyu. Loufamda pambi houbada ishinggi awatpa leitanaba toina sennabiraga darkar leibada ishing thajinbiyu.

SMS Advisory:

Harvest the mature crops like rapeseed/ pea/ lentil/ broad bean/ potato; select pure healthy insect pest free seeds & sundry properly before storing as seed.

Hangam, hawai tharak/ moobi, masoori amadi alu lokpiraga laina paktaba mapung faba mahei khangatpiraga nungshada ningthina kanghanlaga maru oinaba thamjinbiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
-----------------	------------------------

Crop(Varieties)	Crop Specific Advisory
SOYABEAN	Procure insect pest free healthy seeds of soybean/ groundnut. Get the soil testing done for proper fertilizer application. If soil is acidic i.e., PH level is below 6.5 then liming in furrows is required @ 500kg/ ha for having better yield.
	Nung-hawai thanaba afaba mahal laina tilna paktaba maru khanlaga lousinbiba yare. Leibak leihao amadi PH gi changyeng toubiraga PH 6.5 gi makhada leirabadi pari amada 500kg shunu paring naina churi sembiraga happiradi pothook hengathanba ngamgani

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
ONION	Spray Sixer @ 2 gm/lit of water as prophylactic measure against Purple Bloch and bulb rot. Spray is to be given twice at an interval of 7 to 10 days.
	Tilhou manada angouba mapak paklakpa amadi kuisinlakpada khamen machu onsinlakpa natraga tilhou maru patharakpa laina ngakthoknaba numit 7-10ni kannaraga ishing litre amada 2g gi chaangda Sixer happaga kappiyu.

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	- Any type of influenza may be given symptomatic treatment with antibiotics. - Vaccinate against swine fever after the age of 3 months. - Give Ivermectin (Indectin tab.) @ 200mcg/ kg body weight or Ivermec 1% injection @ 1ml/50 kg body weight once only to avoid skin infection. - Vaccinate against F.M.D.
	- Lok thunglakpaga mawong matou ubiradi checksin thourang loubiraga matam chana hidak thabiyu. - Tha 3 shuraba oakta Swine Fever naraktanaba hidak kappiyu. - Unshada tourakpa ana-ayek ngakthoknaba masagi arumba kg amada 200mcg gi chaangda Ivermectin (Indectin tab.) pibiyu natraga masagi arumba kg 50 khudingda 1ml gi chaangda Ivermec 1% injection kappiyu. - F.M.D. laina ngakthoknaba hidak kappiyu.



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 13-03-2020

Weather Forecast of TAMENGLONG(Manipur) Issued On : 2020-03-13(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-03-14	0.0	25.0	10.0	58	19	5.4	63	3
2020-03-15	1.0	25.0	10.0	66	26	5.1	22	7
2020-03-16	2.0	25.0	9.0	56	24	4.6	68	7
2020-03-17	0.0	25.0	10.0	46	18	5.3	68	6
2020-03-18	0.0	26.0	10.0	56	26	5.6	68	4

Weather Summary/Alert:

Likely to have no light rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 25-26 oC & mini. temp. 9-10 oC. Relative humidity may reach upto 66% (max) & 18% (min). The average wind speed may be 4.6-5.6 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 25-26 oC amadi ingbagina 9-10 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 66% amadi nunghildana 18% oiraba yai. Nungsit sitpangi changna pung amada 4.6-5.6 km oiraba yai.

General Advisory:

Disinfect the farm premises of all livestock/ poultry (surrounding area of 3ft width) using the repellent agent/ insecticides spray. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free or polythene sheets to preserve soil moisture for the crop. Monitor the soil moisture status regularly and give life-saving irrigation, especially at growth stages when water requirement is higher.

Kang-hayinggi yamna indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nanna thambiyu. Leibak leinil kuina leinaba pambi makhongda laina tilna paktaba charu usa-wasa natraga polythene na kupsinbiyu. Loufamda pambi houbada ishinggi awatpa leitanaba toina sennabiraga darkar leibada ishing thajinbiyu.

SMS Advisory:

Harvest the mature crops like rapeseed/ pea/ lentil/ broad bean/ potato; select pure healthy insect pest free seeds & sundry properly before storing as seed.

Hangam, hawai tharak/ moobi, masoori amadi alu lokpiraga laina paktaba mapung faba mahei khangatpiraga nunghshada ningthina kanghanlaga maru oinaba thamjinbiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
-----------------	------------------------

Crop(Varieties)	Crop Specific Advisory
SOYABEAN	Procure insect pest free healthy seeds of soybean/ groundnut. Get the soil testing done for proper fertilizer application. If soil is acidic i.e., PH level is below 6.5 then liming in furrows is required @ 500kg/ ha for having better yield.
	Nung-hawai thanaba afaba mahal laina tilna paktaba maru khanlaga lousinbiba yare. Leibak leihao amadi PH gi changyeng toubiraga PH 6.5 gi makhada leirabadi pari amada 500kg shunu paring naina churi sembiraga happiradi pothook hengathanba ngamgani

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
ONION	Spray Sixer @ 2 gm/lit of water as prophylactic measure against Purple Bloch and bulb rot. Spray is to be given twice at an interval of 7 to 10 days.
	Tilhou manada angouba mapak paklakpa amadi kuisinlakpada khamen machu onsinlakpa natraga tilhou maru patharakpa laina ngakthoknaba numit 7-10ni kannaraga ishing litre amada 2g gi chaangda Sixer happaga kappiyu.

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	- Any type of influenza may be given symptomatic treatment with antibiotics. - Vaccinate against swine fever after the age of 3 months. - Give Ivermectin (Indectin tab.) @ 200mcg/ kg body weight or Ivermec 1% injection @ 1ml/50 kg body weight once only to avoid skin infection. - Vaccinate against F.M.D.
	- Lok thunglakpaga mawong matou ubiradi checksin thourang loubiraga matam chana hidak thabiyu. - Tha 3 shuraba oakta Swine Fever naraktanaba hidak kappiyu. - Unshada tourakpa ana-ayek ngakthoknaba masagi arumba kg amada 200mcg gi chaangda Ivermectin (Indectin tab.) pibiyu natraga masagi arumba kg 50 khudingda 1ml gi chaangda Ivermec 1% injection kappiyu. - F.M.D. laina ngakthoknaba hidak kappiyu.



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 13-03-2020

Weather Forecast of TENGNOUPAL(Manipur) Issued On : 2020-03-13(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-03-14	0.0	27.0	11.0	52	26	3.5	292	1
2020-03-15	0.0	28.0	11.0	62	25	3.7	292	7
2020-03-16	1.0	28.0	11.0	72	27	5.0	292	5
2020-03-17	0.0	28.0	12.0	51	21	7.7	292	6
2020-03-18	0.0	29.0	12.0	58	26	9.2	292	4

Weather Summary/Alert:

Likely to have no light rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 27-30 oC & mini. temp. 11-12 oC. Relative humidity may reach upto 72% (max) & 21% (min). The average wind speed may be 3.5-9.2 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 27-30 oC amadi ingbagina 11-12 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 72% amadi nunghildana 21% oiraba yai. Nungsit sitpangi changna pung amada 3.5-9.2 km oiraba yai.

General Advisory:

Disinfect the farm premises of all livestock/ poultry (surrounding area of 3ft width) using the repellent agent/ insecticides spray. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free or polythene sheets to preserve soil moisture for the crop. Monitor the soil moisture status regularly and give life-saving irrigation, especially at growth stages when water requirement is higher.

Kang-hayinggi yamna indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nanna thambiyu. Leibak leinil kuina leinaba pambi makhongda laina tilna paktaba charu usa-wasa natraga polythene na kupsinbiyu. Loufamda pambi houbada ishinggi awatpa leitanaba toina sennabiraga darkar leibada ishing thajinbiyu.

SMS Advisory:

Harvest the mature crops like rapeseed/ pea/ lentil/ broad bean/ potato; select pure healthy insect pest free seeds & sundry properly before storing as seed.

Hangam, hawai tharak/ moobi, masoori amadi alu lokpiraga laina paktaba mapung faba mahei khangatpiraga nunghshada ningthina kanghanlaga maru oinaba thamjinbiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
-----------------	------------------------

Crop(Varieties)	Crop Specific Advisory
SOYABEAN	Procure insect pest free healthy seeds of soybean/ groundnut. Get the soil testing done for proper fertilizer application. If soil is acidic i.e., PH level is below 6.5 then liming in furrows is required @ 500kg/ ha for having better yield.
	Nung-hawai thanaba afaba mahal laina tilna paktaba maru khanlaga lousinbiba yare. Leibak leihao amadi PH gi changyeng toubiraga PH 6.5 gi makhada leirabadi pari amada 500kg shunu paring naina churi sembiraga happiradi pothook hengathanba ngamgani

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
ONION	Spray Sixer @ 2 gm/lit of water as prophylactic measure against Purple Bloch and bulb rot. Spray is to be given twice at an interval of 7 to 10 days.
	Tilhou manada angouba mapak paklakpa amadi kuisinlakpada khamen machu onsinlakpa natraga tilhou maru patharakpa laina ngakthoknaba numit 7-10ni kannaraga ishing litre amada 2g gi chaangda Sixer happaga kappiyu.

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	- Any type of influenza may be given symptomatic treatment with antibiotics. - Vaccinate against swine fever after the age of 3 months. - Give Ivermectin (Indectin tab.) @ 200mcg/ kg body weight or Ivermec 1% injection @ 1ml/50 kg body weight once only to avoid skin infection. - Vaccinate against F.M.D.
	- Lok thunglakpaga mawong matou ubiradi checksin thourang loubiraga matam chana hidak thabiyu. - Tha 3 shuraba oakta Swine Fever naraktanaba hidak kappiyu. - Unshada tourakpa ana-ayek ngakthoknaba masagi arumba kg amada 200mcg gi chaangda Ivermectin (Indectin tab.) pibiyu natraga masagi arumba kg 50 khudingda 1ml gi chaangda Ivermec 1% injection kappiyu. - F.M.D. laina ngakthoknaba hidak kappiyu.



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 13-03-2020

Weather Forecast of THOUBAL(Manipur) Issued On : 2020-03-13(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-03-14	0.0	29.0	13.0	54	23	4.1	243	2
2020-03-15	0.0	29.0	13.0	63	24	5.6	304	6
2020-03-16	0.0	29.0	13.0	59	24	7.4	288	4
2020-03-17	0.0	30.0	14.0	53	18	9.0	288	6
2020-03-18	0.0	30.0	14.0	60	25	10.0	292	4

Weather Summary/Alert:

Likely to have no light rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 29-30 oC & mini. temp. 13-14 oC. Relative humidity may reach upto 63% (max) & 18% (min). The average wind speed may be 4.1-10.0 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 29-30 oC amadi ingbagina 13-14 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 63% amadi nunghildana 18% oiraba yai. Nungsit sitpangi changna pung amada 4.1-10km oiraba yai.

General Advisory:

Disinfect the farm premises of all livestock/ poultry (surrounding area of 3ft width) using the repellent agent/ insecticides spray. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free or polythene sheets to preserve soil moisture for the crop. Monitor the soil moisture status regularly and give life-saving irrigation, especially at growth stages when water requirement is higher.

Kang-hayinggi yamna indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nanna thambiyu. Leibak leinil kuina leinaba pambi makhongda laina tilna paktaba charu usa-wasa natraga polythene na kupsinbiyu. Loufamda pambi houbada ishinggi awatpa leitanaba toina sennabiraga darkar leibada ishing thajinbiyu.

SMS Advisory:

Harvest the mature crops like rapeseed/ pea/ lentil/ broad bean/ potato; select pure healthy insect pest free seeds & sundry properly before storing as seed.

Hangam, hawai tharak/ moobi, masoori amadi alu lokpiraga laina paktaba mapung faba mahei khangatpiraga nungshada ningthina kanghanlaga maru oinaba thamjinbiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
-----------------	------------------------

Crop(Varieties)	Crop Specific Advisory
SOYABEAN	Procure insect pest free healthy seeds of soybean/ groundnut. Get the soil testing done for proper fertilizer application. If soil is acidic i.e., PH level is below 6.5 then liming in furrows is required @ 500kg/ ha for having better yield.
	Nung-hawai thanaba afaba mahal laina tilna paktaba maru khanlaga lousinbiba yare. Leibak leihao amadi PH gi changyeng toubiraga PH 6.5 gi makhada leirabadi pari amada 500kg shunu paring naina churi sembiraga happiradi pothook hengathanba ngamgani

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
ONION	Spray Sixer @ 2 gm/lit of water as prophylactic measure against Purple Bloch and bulb rot. Spray is to be given twice at an interval of 7 to 10 days.
	Tilhou manada angouba mapak paklakpa amadi kuisinlakpada khamen machu onsinlakpa natraga tilhou maru patharakpa laina ngakthoknaba numit 7-10ni kannaraga ishing litre amada 2g gi chaangda Sixer happaga kappiyu.

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	- Any type of influenza may be given symptomatic treatment with antibiotics. - Vaccinate against swine fever after the age of 3 months. - Give Ivermectin (Indectin tab.) @ 200mcg/ kg body weight or Ivermec 1% injection @ 1ml/50 kg body weight once only to avoid skin infection. - Vaccinate against F.M.D.
	- Lok thunglakpaga mawong matou ubiradi checksin thourang loubiraga matam chana hidak thabiyu. - Tha 3 shuraba oakta Swine Fever naraktanaba hidak kappiyu. - Unshada tourakpa ana-ayek ngakthoknaba masagi arumba kg amada 200mcg gi chaangda Ivermectin (Indectin tab.) pibiyu natraga masagi arumba kg 50 khudingda 1ml gi chaangda Ivermec 1% injection kappiyu. - F.M.D. laina ngakthoknaba hidak kappiyu.



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 13-03-2020

Weather Forecast of UKHRUL(Manipur) Issued On : 2020-03-13(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-03-14	0.0	25.0	10.0	63	25	5.1	22	2
2020-03-15	1.0	25.0	10.0	69	27	5.1	22	7
2020-03-16	2.0	25.0	9.0	70	27	5.9	19	6
2020-03-17	0.0	25.0	10.0	57	18	6.4	112	6
2020-03-18	0.0	26.0	10.0	66	32	6.2	112	5

Weather Summary/Alert:

Likely to have no light rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 25-26 oC & mini. temp. 9-10 oC. Relative humidity may reach upto 70% (max) & 18% (min). The average wind speed may be 5.1-6.4 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 25-26 oC amadi ingbagina 9-10 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 70% amadi nunghildana 18% oiraba yai. Nungsit sitpangi changna pung amada 5.1-6.4 km oiraba yai.

General Advisory:

Disinfect the farm premises of all livestock/ poultry (surrounding area of 3ft width) using the repellent agent/ insecticides spray. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free or polythene sheets to preserve soil moisture for the crop. Monitor the soil moisture status regularly and give life-saving irrigation, especially at growth stages when water requirement is higher.

Kang-hayinggi yamna indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nanna thambiyu. Leibak leinil kuina leinaba pambi makhongda laina tilna paktaba charu usa-wasa natraga polythene na kupsinbiyu. Loufamda pambi houbada ishinggi awatpa leitanaba toina sennabiraga darkar leibada ishing thajinbiyu.

SMS Advisory:

Harvest the mature crops like rapeseed/ pea/ lentil/ broad bean/ potato; select pure healthy insect pest free seeds & sundry properly before storing as seed.

Hangam, hawai tharak/ moobi, masoori amadi alu lokpiraga laina paktaba mapung faba mahei khangatpiraga nungshada ningthina kanghanlaga maru oinaba thamjinbiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
-----------------	------------------------

Crop(Varieties)	Crop Specific Advisory
SOYABEAN	Procure insect pest free healthy seeds of soybean/ groundnut. Get the soil testing done for proper fertilizer application. If soil is acidic i.e., PH level is below 6.5 then liming in furrows is required @ 500kg/ ha for having better yield.
	Nung-hawai thanaba afaba mahal laina tilna paktaba maru khanlaga lousinbiba yare. Leibak leihao amadi PH gi changyeng toubiraga PH 6.5 gi makhada leirabadi pari amada 500kg shunu paring naina churi sembiraga happiradi pothook hengathanba ngamgani

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
ONION	Spray Sixer @ 2 gm/lit of water as prophylactic measure against Purple Bloch and bulb rot. Spray is to be given twice at an interval of 7 to 10 days.
	Tilhou manada angouba mapak paklakpa amadi kuisinlakpada khamen machu onsinlakpa natraga tilhou maru patharakpa laina ngakthoknaba numit 7-10ni kannaraga ishing litre amada 2g gi chaangda Sixer happaga kappiyu.

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	- Any type of influenza may be given symptomatic treatment with antibiotics. - Vaccinate against swine fever after the age of 3 months. - Give Ivermectin (Indectin tab.) @ 200mcg/ kg body weight or Ivermec 1% injection @ 1ml/50 kg body weight once only to avoid skin infection. - Vaccinate against F.M.D.
	- Lok thunglakpaga mawong matou ubiradi checksin thourang loubiraga matam chana hidak thabiyu. - Tha 3 shuraba oakta Swine Fever naraktanaba hidak kappiyu. - Unshada tourakpa ana-ayek ngakthoknaba masagi arumba kg amada 200mcg gi chaangda Ivermectin (Indectin tab.) pibiyu natraga masagi arumba kg 50 khudingda 1ml gi chaangda Ivermec 1% injection kappiyu. - F.M.D. laina ngakthoknaba hidak kappiyu.



Gramin Krishi Mausam Sewa
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 13-03-2020

Weather Forecast of WANGOI Block in IMPHAL-WEST(Manipur) Issued On :2020-03-13(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-03-14	0.0	33.5	19.2	61	23	5.0	330	2
2020-03-15	0.1	34.9	19.6	73	22	6.0	288	3
2020-03-16	0.0	33.0	17.5	77	28	9.0	270	2
2020-03-17	0.0	30.0	15.6	71	22	7.0	259	6
2020-03-18	0.0	31.7	15.7	59	24	8.0	270	5

Weather Summary/Alert:

Likely to have no light rains upto Wednesday. The sky will remain mainly clear to partly cloudy but mainly cloudy on Tuesday with max. temp. 30-33.5 oC & mini. temp. 15.6-19.6 oC. Relative humidity may reach upto 77% (max) & 28 % (min). The average wind speed may be 5-8 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai adubu Leibakpokpadadi khara yamna khanlaba yai. Khwaidagi sabagi changna 30-33.5 oC amadi ingbagina 15.6-19.6 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 77% amadi nungthildana 28 % oiraba yai. Nungsit sitpagi changna pung amada 5-8 km oiraba yai.

General Advisory:

Disinfect the farm premises of all livestock/ poultry (surrounding area of 3ft width) using the repellent agent/ insecticides spray. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free or polythene sheets to preserve soil moisture for the crop. Monitor the soil moisture status regularly and give life-saving irrigation, especially at growth stages when water requirement is higher.

Kang-hayinggi yamna indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nanna thambiye. Leibak leinil kuina leinaba pambi makhongda laina tilna paktaba charu usa-wasa natraga polythene na kupsinbiyu. Loufamda pambi houbada ishinggi awatpa leitanaba toina sennabiraga darkar leibada ishing thajinbiyu.

SMS Advisory:

Harvest the mature crops like rapeseed/ pea/ lentil/ broad bean/ potato; select pure healthy insect pest free seeds & sundry properly before storing as seed.

Hangam, hawai tharak/ moobi, masoori amadi alu lokpiraga laina paktaba mapung faba mahei khangatpiraga nungshada ningthina kanghanlaga maru oinaba thamjinbiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
SOYABEAN	Procure insect pest free healthy seeds of soybean/ groundnut. Get the soil testing done for proper fertilizer application. If soil is acidic i.e., PH level is below 6.5 then liming in furrows is required @ 500kg/ ha for having better yield.
	Nung-hawai thanaba afaba mahal laina tilna paktaba maru khanlaga lousinbiba yare. Leibak leihao amadi PH gi changyeng toubiraga PH 6.5 gi makhada leirabadi pari amada 500kg shunu paring naina churi sembiraga happiradi pothook hengathanba ngamgani.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
ONION	<p>Spray Sixer @ 2 gm/lit of water as prophylactic measure against Purple Bloch and bulb rot. Spray is to be given twice at an interval of 7 to 10 days.</p> <p>Tilhou manada angouba mapak paklakpa amadi kuisinlakpada khamen machu onsinlakpa natraga tilhou maru patharakpa laina ngakthoknaba numit 7-10ni kannaraga ishing litre amada 2g gi chaangda Sixer happaga kappiyu.</p>

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	<p>- Any type of influenza may be given symptomatic treatment with antibiotics. - Vaccinate against swine fever after the age of 3 months. - Give Ivermectin (Indectin tab.) @ 200mcg/ kg body weight or Ivermec 1% injection @ 1ml/50 kg body weight once only to avoid skin infection. - Vaccinate against F.M.D.</p> <p>- Lok thunglakpangi mawong matou ubiradi checksin thourang loubiraga matam chana hidak thabiyu. - Tha 3 shuraba oakta Swine Fever naraktanaba hidak kappiyu. - Unshada tourakpa ana-ayek ngakthoknaba masagi arumba kg amada 200mcg gi chaangda Ivermectin (Indectin tab.) pibiyu natraga masagi arumba kg 50 khudingda 1ml gi chaangda Ivermec 1% injection kappiyu. - F.M.D. laina ngakthoknaba hidak kappiyu.</p>



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 13-03-2020

Weather Forecast of BISHNUPUR(Manipur) Issued On : 2020-03-13(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-03-14	0.0	29.0	13.0	54	21	3.3	60	2
2020-03-15	0.0	29.0	13.0	65	25	3.3	292	5
2020-03-16	0.0	29.0	13.0	61	25	3.8	292	4
2020-03-17	0.0	30.0	14.0	54	18	4.7	281	5
2020-03-18	0.0	30.0	14.0	62	25	4.9	120	4

Weather Summary/Alert:

Likely to have no light rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 29-30 oC & mini. temp. 13-14 oC. Relative humidity may reach upto 65% (max) & 18% (min). The average wind speed may be 3.3-4.9 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 29-30 oC amadi ingbagina 13-14 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 65% amadi nunghildana 18% oiraba yai. Nungsit sitpangi changna pung amada 3.3-4.9 km oiraba yai.

General Advisory:

Disinfect the farm premises of all livestock/ poultry (surrounding area of 3ft width) using the repellent agent/ insecticides spray. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free or polythene sheets to preserve soil moisture for the crop. Monitor the soil moisture status regularly and give life-saving irrigation, especially at growth stages when water requirement is higher.

Kang-hayinggi yamna indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nanna thambiyu. Leibak leinil kuina leinaba pambi makhongda laina tilna paktaba charu usa-wasa natraga polythene na kupsinbiyu. Loufamda pambi houbada ishinggi awatpa leitanaba toina sennabiraga darkar leibada ishing thajinbiyu.

SMS Advisory:

Harvest the mature crops like rapeseed/ pea/ lentil/ broad bean/ potato; select pure healthy insect pest free seeds & sundry properly before storing as seed.

Hangam, hawai tharak/ moobi, masoori amadi alu lokpiraga laina paktaba mapung faba mahei khangatpiraga nungshada ningthina kanghanlaga maru oinaba thamjinbiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
-----------------	------------------------

Crop(Varieties)	Crop Specific Advisory
SOYABEAN	Procure insect pest free healthy seeds of soybean/ groundnut. Get the soil testing done for proper fertilizer application. If soil is acidic i.e., PH level is below 6.5 then liming in furrows is required @ 500kg/ ha for having better yield.
	Nung-hawai thanaba afaba mahal laina tilna paktaba maru khanlaga lousinbiba yare. Leibak leihao amadi PH gi changyeng toubiraga PH 6.5 gi makhada leirabadi pari amada 500kg shunu paring naina churi sembiraga happiradi pothook hengathanba ngamgani

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
ONION	Spray Sixer @ 2 gm/lit of water as prophylactic measure against Purple Bloch and bulb rot. Spray is to be given twice at an interval of 7 to 10 days.
	Tilhou manada angouba mapak paklakpa amadi kuisinlakpada khamen machu onsinlakpa natraga tilhou maru patharakpa laina ngakthoknaba numit 7-10ni kannaraga ishing litre amada 2g gi chaangda Sixer happaga kappiyu.

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	- Any type of influenza may be given symptomatic treatment with antibiotics. - Vaccinate against swine fever after the age of 3 months. - Give Ivermectin (Indectin tab.) @ 200mcg/ kg body weight or Ivermec 1% injection @ 1ml/50 kg body weight once only to avoid skin infection. - Vaccinate against F.M.D.
	- Lok thunglakpagi mawong matou ubiradi checksin thourang loubiraga matam chana hidak thabiyu. - Tha 3 shuraba oakta Swine Fever naraktanaba hidak kappiyu. - Unshada tourakpa ana-ayek ngakthoknaba masagi arumba kg amada 200mcg gi chaangda Ivermectin (Indectin tab.) pibiyu natraga masagi arumba kg 50 khudingda 1ml gi chaangda Ivermec 1% injection kappiyu. - F.M.D. laina ngakthoknaba hidak kappiyu.



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 13-03-2020

Weather Forecast of CHANDEL(Manipur) Issued On : 2020-03-13(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-03-14	0.0	27.0	11.0	52	24	2.6	281	1
2020-03-15	0.0	28.0	11.0	61	24	3.6	292	6
2020-03-16	0.0	28.0	11.0	63	25	4.9	288	5
2020-03-17	0.0	28.0	12.0	54	21	7.6	292	5
2020-03-18	0.0	29.0	12.0	58	27	8.7	292	5

Weather Summary/Alert:

Likely to have no light rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 27-29 oC & mini. temp. 11-12 oC. Relative humidity may reach upto 61% (max) & 21% (min). The average wind speed may be 2.6-8.7 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 27-29 oC amadi ingbagina 11-12 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 61% amadi nunghildana 21% oiraba yai. Nungsit sitpangi changna pung amada 2.6-8.7 km oiraba yai.

General Advisory:

Disinfect the farm premises of all livestock/ poultry (surrounding area of 3ft width) using the repellent agent/ insecticides spray. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free or polythene sheets to preserve soil moisture for the crop. Monitor the soil moisture status regularly and give life-saving irrigation, especially at growth stages when water requirement is higher.

Kang-hayinggi yamna indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nanna thambiyu. Leibak leinil kuina leinaba pambi makhongda laina tilna paktaba charu usa-wasa natraga polythene na kupsinbiyu. Loufamda pambi houbada ishinggi awatpa leitanaba toina sennabiraga darkar leibada ishing thajinbiyu.

SMS Advisory:

Harvest the mature crops like rapeseed/ pea/ lentil/ broad bean/ potato; select pure healthy insect pest free seeds & sundry properly before storing as seed.

Hangam, hawai tharak/ moobi, masoori amadi alu lokpiraga laina paktaba mapung faba mahei khangatpiraga nunghshada ningthina kanghanlaga maru oinaba thamjinbiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
-----------------	------------------------

Crop(Varieties)	Crop Specific Advisory
SOYABEAN	Procure insect pest free healthy seeds of soybean/ groundnut. Get the soil testing done for proper fertilizer application. If soil is acidic i.e., PH level is below 6.5 then liming in furrows is required @ 500kg/ ha for having better yield.
	Nung-hawai thanaba afaba mahal laina tilna paktaba maru khanlaga lousinbiba yare. Leibak leihao amadi PH gi changyeng toubiraga PH 6.5 gi makhada leirabadi pari amada 500kg shunu paring naina churi sembiraga happiradi pothook hengathanba ngamgani

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
ONION	Spray Sixer @ 2 gm/lit of water as prophylactic measure against Purple Bloch and bulb rot. Spray is to be given twice at an interval of 7 to 10 days.
	Tilhou manada angouba mapak paklakpa amadi kuisinlakpada khamen machu onsinlakpa natraga tilhou maru patharakpa laina ngakthoknaba numit 7-10ni kannaraga ishing litre amada 2g gi chaangda Sixer happaga kappiyu.

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	- Any type of influenza may be given symptomatic treatment with antibiotics. - Vaccinate against swine fever after the age of 3 months. - Give Ivermectin (Indectin tab.) @ 200mcg/ kg body weight or Ivermec 1% injection @ 1ml/50 kg body weight once only to avoid skin infection. - Vaccinate against F.M.D.
	- Lok thunglakpaga mawong matou ubiradi checksin thourang loubiraga matam chana hidak thabiyu. - Tha 3 shuraba oakta Swine Fever naraktanaba hidak kappiyu. - Unshada tourakpa ana-ayek ngakthoknaba masagi arumba kg amada 200mcg gi chaangda Ivermectin (Indectin tab.) pibiyu natraga masagi arumba kg 50 khudingda 1ml gi chaangda Ivermec 1% injection kappiyu. - F.M.D. laina ngakthoknaba hidak kappiyu.



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 13-03-2020

Weather Forecast of CHURACHANDPUR(Manipur) Issued On : 2020-03-13(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-03-14	0.0	28.0	12.0	54	24	3.6	158	1
2020-03-15	0.0	29.0	12.0	63	25	4.1	191	6
2020-03-16	0.0	29.0	12.0	65	26	4.6	292	5
2020-03-17	0.0	29.0	12.0	56	21	6.4	288	4
2020-03-18	0.0	30.0	13.0	62	28	6.3	240	4

Weather Summary/Alert:

Likely to have no light rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 28-30 oC & mini. temp. 12-13 oC. Relative humidity may reach upto 65% (max) & 21% (min). The average wind speed may be 3.6-6.3 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 28-30 oC amadi ingbagina 12-13 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 65% amadi nunghildana 21% oiraba yai. Nungsit sitpangi changna pung amada 3.6-6.3 km oiraba yai.

General Advisory:

Disinfect the farm premises of all livestock/ poultry (surrounding area of 3ft width) using the repellent agent/ insecticides spray. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free or polythene sheets to preserve soil moisture for the crop. Monitor the soil moisture status regularly and give life-saving irrigation, especially at growth stages when water requirement is higher.

Kang-hayinggi yamna indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nanna thambiyu. Leibak leinil kuina leinaba pambi makhongda laina tilna paktaba charu usa-wasa natraga polythene na kupsinbiyu. Loufamda pambi houbada ishinggi awatpa leitanaba toina sennabiraga darkar leibada ishing thajinbiyu.

SMS Advisory:

Harvest the mature crops like rapeseed/ pea/ lentil/ broad bean/ potato; select pure healthy insect pest free seeds & sundry properly before storing as seed.

Hangam, hawai tharak/ moobi, masoori amadi alu lokpiraga laina paktaba mapung faba mahei khangatpiraga nunghshada ningthina kanghanlaga maru oinaba thamjinbiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
-----------------	------------------------

Crop(Varieties)	Crop Specific Advisory
SOYABEAN	Procure insect pest free healthy seeds of soybean/ groundnut. Get the soil testing done for proper fertilizer application. If soil is acidic i.e., PH level is below 6.5 then liming in furrows is required @ 500kg/ ha for having better yield.
	Nung-hawai thanaba afaba mahal laina tilna paktaba maru khanlaga lousinbiba yare. Leibak leihao amadi PH gi changyeng toubiraga PH 6.5 gi makhada leirabadi pari amada 500kg shunu paring naina churi sembiraga happiradi pothook hengathanba ngamgani

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
ONION	Spray Sixer @ 2 gm/lit of water as prophylactic measure against Purple Bloch and bulb rot. Spray is to be given twice at an interval of 7 to 10 days.
	Tilhou manada angouba mapak paklakpa amadi kuisinlakpada khamen machu onsinlakpa natraga tilhou maru patharakpa laina ngakthoknaba numit 7-10ni kannaraga ishing litre amada 2g gi chaangda Sixer happaga kappiyu.

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	- Any type of influenza may be given symptomatic treatment with antibiotics. - Vaccinate against swine fever after the age of 3 months. - Give Ivermectin (Indectin tab.) @ 200mcg/ kg body weight or Ivermec 1% injection @ 1ml/50 kg body weight once only to avoid skin infection. - Vaccinate against F.M.D.
	- Lok thunglakpaga mawong matou ubiradi checksin thourang loubiraga matam chana hidak thabiyu. - Tha 3 shuraba oakta Swine Fever naraktanaba hidak kappiyu. - Unshada tourakpa ana-ayek ngakthoknaba masagi arumba kg amada 200mcg gi chaangda Ivermectin (Indectin tab.) pibiyu natraga masagi arumba kg 50 khudingda 1ml gi chaangda Ivermec 1% injection kappiyu. - F.M.D. laina ngakthoknaba hidak kappiyu.



Gramin Krishi Mausam Sewa
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 13-03-2020

Weather Forecast of HAORANGSABAL Block in IMPHAL-WEST(Manipur) Issued On :2020-03-13(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-03-14	0.0	33.5	19.7	64	22	6.0	23	3
2020-03-15	0.1	34.7	19.7	74	22	6.0	248	3
2020-03-16	0.0	32.9	17.3	77	27	8.0	283	2
2020-03-17	0.0	29.8	16.1	70	20	7.0	248	7
2020-03-18	0.0	31.4	15.6	58	24	7.0	248	4

Weather Summary/Alert:

Likely to have no light rains upto Wednesday. The sky will remain mainly clear to partly cloudy but mainly cloudy on Tuesday with max. temp. 29.8-33.5 oC & mini. temp. 15.6-19.7 oC. Relative humidity may reach upto 77% (max) & 27 % (min). The average wind speed may be 6-8 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai adubu Leibakpokpadadi khara yamna khanlaba yai. Khwaidagi sabagi changna 29.8-33.5 oC amadi ingbagina 15.6-19.7 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 77% amadi nungthildana 27 % oiraba yai. Nungsit sitpagi changna pung amada 6-8 km oiraba yai.

General Advisory:

Disinfect the farm premises of all livestock/ poultry (surrounding area of 3ft width) using the repellent agent/ insecticides spray. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free or polythene sheets to preserve soil moisture for the crop. Monitor the soil moisture status regularly and give life-saving irrigation, especially at growth stages when water requirement is higher.

Kang-hayinggi yamna indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nanna thambiyeu. Leibak leinil kuina leinaba pambi makhongda laina tilna paktaba charu usa-wasa natraga polythene na kupsinbiyu. Loufamda pambi houbada ishinggi awatpa leitanaba toina sennabiraga darkar leibada ishing thajinbiyu.

SMS Advisory:

Harvest the mature crops like rapeseed/ pea/ lentil/ broad bean/ potato; select pure healthy insect pest free seeds & sundry properly before storing as seed.

Hangam, hawai tharak/ moobi, masoori amadi alu lokpiraga laina paktaba mapung faba mahei khangatpiraga nungshada ningthina kanghanlaga maru oinaba thamjinbiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
SOYABEAN	Procure insect pest free healthy seeds of soybean/ groundnut. Get the soil testing done for proper fertilizer application. If soil is acidic i.e., PH level is below 6.5 then liming in furrows is required @ 500kg/ ha for having better yield.
	Nung-hawai thanaba afaba mahal laina tilna paktaba maru khanlaga lousinbiba yare. Leibak leihao amadi PH gi changyeng toubiraga PH 6.5 gi makhada leirabadi pari amada 500kg shunu paring naina churi sembiraga happiradi pothook hengathanba ngamgani.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
ONION	<p>Spray Sixer @ 2 gm/lit of water as prophylactic measure against Purple Bloch and bulb rot. Spray is to be given twice at an interval of 7 to 10 days.</p> <p>Tilhou manada angouba mapak paklakpa amadi kuisinlakpada khamen machu onsinlakpa natraga tilhou maru patharakpa laina ngakthoknaba numit 7-10ni kannaraga ishing litre amada 2g gi chaangda Sixer happaga kappiyu.</p>

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	<p>- Any type of influenza may be given symptomatic treatment with antibiotics. - Vaccinate against swine fever after the age of 3 months. - Give Ivermectin (Indectin tab.) @ 200mcg/ kg body weight or Ivermec 1% injection @ 1ml/50 kg body weight once only to avoid skin infection. - Vaccinate against F.M.D.</p> <p>- Lok thunglakpagi mawong matou ubiradi checksin thourang loubiraga matam chana hidak thabiyu. - Tha 3 shuraba oakta Swine Fever naraktanaba hidak kappiyu. - Unshada tourakpa ana-ayek ngakthoknaba masagi arumba kg amada 200mcg gi chaangda Ivermectin (Indectin tab.) pibiyu natraga masagi arumba kg 50 khudingda 1ml gi chaangda Ivermec 1% injection kappiyu. - F.M.D. laina ngakthoknaba hidak kappiyu.</p>



Gramin Krishi Mausam Sewa
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 13-03-2020

Weather Forecast of HIYANGTHANG Block in IMPHAL-WEST(Manipur) Issued On :2020-03-13(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-03-14	0.0	33.4	19.5	62	22	6.0	360	3
2020-03-15	0.0	34.7	19.7	73	22	6.0	293	3
2020-03-16	0.0	33.0	17.4	77	27	9.0	270	2
2020-03-17	0.0	30.0	15.9	70	20	7.0	248	6
2020-03-18	0.0	31.8	15.6	58	24	7.0	248	4

Weather Summary/Alert:

Likely to have no light rains upto Wednesday. The sky will remain mainly clear to partly cloudy but mainly cloudy on Tuesday with max. temp. 30-33.4 oC & mini. temp. 15.6-19.7 oC. Relative humidity may reach upto 77% (max) & 27 % (min). The average wind speed may be 6-9 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai adubu Leibakpokpadadi khara yamna khanlaba yai. Khwaidagi sabagi changna 30-33.4 oC amadi ingbagina 15.6-19.7 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 77% amadi nungthildana 27% oiraba yai. Nungsit sitpangi changna pung amada 6-9 km oiraba yai.

General Advisory:

Disinfect the farm premises of all livestock/ poultry (surrounding area of 3ft width) using the repellent agent/ insecticides spray. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free or polythene sheets to preserve soil moisture for the crop. Monitor the soil moisture status regularly and give life-saving irrigation, especially at growth stages when water requirement is higher.

Kang-hayinggi yamna indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nanna thambiyyu. Leibak leinil kuina leinaba pambi makhongda laina tilna paktaba charu usa-wasa natraga polythene na kupsinbiyyu. Loufamda pambi houbada ishinggi awatpa leitanaba toina sennabiraga darkar leibada ishing thajinbiyyu.

SMS Advisory:

Harvest the mature crops like rapeseed/ pea/ lentil/ broad bean/ potato; select pure healthy insect pest free seeds & sundry properly before storing as seed.

Hangam, hawai tharak/ moobi, masoori amadi alu lokpiraga laina paktaba mapung faba mahei khangatpiraga nungshada ningthina kanghanlaga maru oinaba thamjinbiyyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
SOYABEAN	Procure insect pest free healthy seeds of soybean/ groundnut. Get the soil testing done for proper fertilizer application. If soil is acidic i.e., PH level is below 6.5 then liming in furrows is required @ 500kg/ ha for having better yield.
	Nung-hawai thanaba afaba mahal laina tilna paktaba maru khanlaga lousinbiba yare. Leibak leihao amadi PH gi changyeng toubiraga PH 6.5 gi makhada leirabadi pari amada 500kg shunu paring naina churi sembiraga happiradi pothook hengathanba ngamgani.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
ONION	<p>Spray Sixer @ 2 gm/lit of water as prophylactic measure against Purple Bloch and bulb rot. Spray is to be given twice at an interval of 7 to 10 days.</p> <p>Tilhou manada angouba mapak paklakpa amadi kuisinlakpada khamen machu onsinlakpa natraga tilhou maru patharakpa laina ngakthoknaba numit 7-10ni kannaraga ishing litre amada 2g gi chaangda Sixer happaga kappiyu.</p>

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	<p>- Any type of influenza may be given symptomatic treatment with antibiotics. - Vaccinate against swine fever after the age of 3 months. - Give Ivermectin (Indectin tab.) @ 200mcg/ kg body weight or Ivermec 1% injection @ 1ml/50 kg body weight once only to avoid skin infection. - Vaccinate against F.M.D.</p> <p>- Lok thunglakpagi mawong matou ubiradi checksin thourang loubiraga matam chana hidak thabiyu. - Tha 3 shuraba oakta Swine Fever naraktanaba hidak kappiyu. - Unshada tourakpa ana-ayek ngakthoknaba masagi arumba kg amada 200mcg gi chaangda Ivermectin (Indectin tab.) pibiyu natraga masagi arumba kg 50 khudingda 1ml gi chaangda Ivermec 1% injection kappiyu. - F.M.D. laina ngakthoknaba hidak kappiyu.</p>



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 13-03-2020

Weather Forecast of IMPHAL-EAST(Manipur) Issued On : 2020-03-13(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-03-14	0.0	28.0	12.0	54	20	5.3	22	2
2020-03-15	0.0	29.0	12.0	62	22	5.7	252	6
2020-03-16	0.0	29.0	12.0	57	23	7.1	315	5
2020-03-17	0.0	29.0	12.0	52	17	7.4	281	6
2020-03-18	0.0	30.0	13.0	59	24	8.9	330	3

Weather Summary/Alert:

Likely to have no light rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 28-30 oC & mini. temp. 12-13 oC. Relative humidity may reach upto 62 % (max) & 17% (min). The average wind speed may be 5.3-8.9 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 28-30 oC amadi ingbagina 12-13 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 62 % amadi nungthildana 17% oiraba yai. Nungsit sitpagi changna pung amada 5.3-8.9 km oiraba yai.

General Advisory:

Disinfect the farm premises of all livestock/ poultry (surrounding area of 3ft width) using the repellent agent/ insecticides spray. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free or polythene sheets to preserve soil moisture for the crop. Monitor the soil moisture status regularly and give life-saving irrigation, especially at growth stages when water requirement is higher.

Kang-hayinggi yamna indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nanna thambiyu. Leibak leinil kuina leinaba pambi makhongda laina tilna paktaba charu usa-wasa natraga polythene na kupsinbiyu. Loufamda pambi houbada ishinggi awatpa leitanaba toina sennabiraga darkar leibada ishing thajinbiyu.

SMS Advisory:

Harvest the mature crops like rapeseed/ pea/ lentil/ broad bean/ potato; select pure healthy insect pest free seeds & sundry properly before storing as seed.

Hangam, hawai tharak/ moobi, masoori amadi alu lokpiraga laina paktaba mapung faba mahei khangatpiraga nungshada ningthina kanghanlaga maru oinaba thamjinbiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
-----------------	------------------------

Crop(Varieties)	Crop Specific Advisory
SOYABEAN	Procure insect pest free healthy seeds of soybean/ groundnut. Get the soil testing done for proper fertilizer application. If soil is acidic i.e., PH level is below 6.5 then liming in furrows is required @ 500kg/ ha for having better yield.
	Nung-hawai thanaba afaba mahal laina tilna paktaba maru khanlaga lousinbiba yare. Leibak leihao amadi PH gi changyeng toubiraga PH 6.5 gi makhada leirabadi pari amada 500kg shunu paring naina churi sembiraga happiradi pothook hengathanba ngamgani

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
ONION	Spray Sixer @ 2 gm/lit of water as prophylactic measure against Purple Bloch and bulb rot. Spray is to be given twice at an interval of 7 to 10 days.
	Tilhou manada angouba mapak paklakpa amadi kuisinlakpada khamen machu onsinlakpa natraga tilhou maru patharakpa laina ngakthoknaba numit 7-10ni kannaraga ishing litre amada 2g gi chaangda Sixer happaga kappiyu.

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	- Any type of influenza may be given symptomatic treatment with antibiotics. - Vaccinate against swine fever after the age of 3 months. - Give Ivermectin (Indectin tab.) @ 200mcg/ kg body weight or Ivermec 1% injection @ 1ml/50 kg body weight once only to avoid skin infection. - Vaccinate against F.M.D.
	- Lok thunglakpaga mawong matou ubiradi checksin thourang loubiraga matam chana hidak thabiyu. - Tha 3 shuraba oakta Swine Fever naraktanaba hidak kappiyu. - Unshada tourakpa ana-ayek ngakthoknaba masagi arumba kg amada 200mcg gi chaangda Ivermectin (Indectin tab.) pibiyu natraga masagi arumba kg 50 khudingda 1ml gi chaangda Ivermec 1% injection kappiyu. - F.M.D. laina ngakthoknaba hidak kappiyu.



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 13-03-2020

Weather Forecast of IMPHAL-WEST(Manipur) Issued On : 2020-03-13(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-03-14	0.0	28.0	12.0	55	20	6.0	22	2
2020-03-15	0.0	29.0	12.0	62	23	5.8	281	6
2020-03-16	0.0	29.0	12.0	58	24	6.8	292	5
2020-03-17	0.0	29.0	12.0	52	17	6.6	248	6
2020-03-18	0.0	30.0	13.0	60	25	8.0	68	3

Weather Summary/Alert:

Likely to have no light rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 28-30 oC & mini. temp. 12-13 oC. Relative humidity may reach upto 62 % (max) & 17% (min). The average wind speed may be 5.8-8.0 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 28-30 oC amadi ingbagina 12-13 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 62 % amadi nungthildana 17% oiraba yai. Nungsit sitpangi changna pung amada 5.8-8.0 km oiraba yai.

General Advisory:

Disinfect the farm premises of all livestock/ poultry (surrounding area of 3ft width) using the repellent agent/ insecticides spray. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free or polythene sheets to preserve soil moisture for the crop. Monitor the soil moisture status regularly and give life-saving irrigation, especially at growth stages when water requirement is higher.

Kang-hayinggi yamna indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nanna thambiyu. Leibak leinil kuina leinaba pambi makhongda laina tilna paktaba charu usa-wasa natraga polythene na kupsinbiyu. Loufamda pambi houbada ishinggi awatpa leitanaba toina sennabiraga darkar leibada ishing thajinbiyu.

SMS Advisory:

Harvest the mature crops like rapeseed/ pea/ lentil/ broad bean/ potato; select pure healthy insect pest free seeds & sundry properly before storing as seed.

Hangam, hawai tharak/ moobi, masoori amadi alu lokpiraga laina paktaba mapung faba mahei khangatpiraga nungshada ningthina kanghanlaga maru oinaba thamjinbiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
-----------------	------------------------

Crop(Varieties)	Crop Specific Advisory
SOYABEAN	Procure insect pest free healthy seeds of soybean/ groundnut. Get the soil testing done for proper fertilizer application. If soil is acidic i.e., PH level is below 6.5 then liming in furrows is required @ 500kg/ ha for having better yield.
	Nung-hawai thanaba afaba mahal laina tilna paktaba maru khanlaga lousinbiba yare. Leibak leihao amadi PH gi changyeng toubiraga PH 6.5 gi makhada leirabadi pari amada 500kg shunu paring naina churi sembiraga happiradi pothook hengathanba ngamgani

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
ONION	Spray Sixer @ 2 gm/lit of water as prophylactic measure against Purple Bloch and bulb rot. Spray is to be given twice at an interval of 7 to 10 days.
	Tilhou manada angouba mapak paklakpa amadi kuisinlakpada khamen machu onsinlakpa natraga tilhou maru patharakpa laina ngakthoknaba numit 7-10ni kannaraga ishing litre amada 2g gi chaangda Sixer happaga kappiyu.

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	- Any type of influenza may be given symptomatic treatment with antibiotics. - Vaccinate against swine fever after the age of 3 months. - Give Ivermectin (Indectin tab.) @ 200mcg/ kg body weight or Ivermec 1% injection @ 1ml/50 kg body weight once only to avoid skin infection. - Vaccinate against F.M.D.
	- Lok thunglakpaga mawong matou ubiradi checksin thourang loubiraga matam chana hidak thabiyu. - Tha 3 shuraba oakta Swine Fever naraktanaba hidak kappiyu. - Unshada tourakpa ana-ayek ngakthoknaba masagi arumba kg amada 200mcg gi chaangda Ivermectin (Indectin tab.) pibiyu natraga masagi arumba kg 50 khudingda 1ml gi chaangda Ivermec 1% injection kappiyu. - F.M.D. laina ngakthoknaba hidak kappiyu.



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 13-03-2020

Weather Forecast of JIRIBAM(Manipur) Issued On : 2020-03-13(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-03-14	0.0	25.0	10.0	55	17	4.1	64	2
2020-03-15	0.0	25.0	10.0	67	25	5.2	65	7
2020-03-16	0.0	25.0	9.0	55	23	4.0	71	6
2020-03-17	0.0	25.0	10.0	43	16	4.6	71	5
2020-03-18	0.0	26.0	10.0	53	22	5.9	68	3

Weather Summary/Alert:

Likely to have no light rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 25-26 oC & mini. temp. 9-10 oC. Relative humidity may reach upto 67% (max) & 16% (min). The average wind speed may be 4.1-5.9 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 25-26 oC amadi ingbagina 9-10 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 67% amadi nunghildana 16% oiraba yai. Nungsit sitpangi changna pung amada 4.1-5.9 km oiraba yai.

General Advisory:

Disinfect the farm premises of all livestock/ poultry (surrounding area of 3ft width) using the repellent agent/ insecticides spray. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free or polythene sheets to preserve soil moisture for the crop. Monitor the soil moisture status regularly and give life-saving irrigation, especially at growth stages when water requirement is higher.

Kang-hayinggi yamna indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nanna thambiyu. Leibak leinil kuina leinaba pambi makhongda laina tilna paktaba charu usa-wasa natraga polythene na kupsinbiyu. Loufamda pambi houbada ishinggi awatpa leitanaba toina sennabiraga darkar leibada ishing thajinbiyu.

SMS Advisory:

Harvest the mature crops like rapeseed/ pea/ lentil/ broad bean/ potato; select pure healthy insect pest free seeds & sundry properly before storing as seed.

Hangam, hawai tharak/ moobi, masoori amadi alu lokpiraga laina paktaba mapung faba mahei khangatpiraga nunghshada ningthina kanghanlaga maru oinaba thamjinbiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
-----------------	------------------------

Crop(Varieties)	Crop Specific Advisory
SOYABEAN	Procure insect pest free healthy seeds of soybean/ groundnut. Get the soil testing done for proper fertilizer application. If soil is acidic i.e., PH level is below 6.5 then liming in furrows is required @ 500kg/ ha for having better yield.
	Nung-hawai thanaba afaba mahal laina tilna paktaba maru khanlaga lousinbiba yare. Leibak leihao amadi PH gi changyeng toubiraga PH 6.5 gi makhada leirabadi pari amada 500kg shunu paring naina churi sembiraga happiradi pothook hengathanba ngamgani

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
ONION	Spray Sixer @ 2 gm/lit of water as prophylactic measure against Purple Bloch and bulb rot. Spray is to be given twice at an interval of 7 to 10 days.
	Tilhou manada angouba mapak paklakpa amadi kuisinlakpada khamen machu onsinlakpa natraga tilhou maru patharakpa laina ngakthoknaba numit 7-10ni kannaraga ishing litre amada 2g gi chaangda Sixer happaga kappiyu.

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	- Any type of influenza may be given symptomatic treatment with antibiotics. - Vaccinate against swine fever after the age of 3 months. - Give Ivermectin (Indectin tab.) @ 200mcg/ kg body weight or Ivermec 1% injection @ 1ml/50 kg body weight once only to avoid skin infection. - Vaccinate against F.M.D.
	- Lok thunglakpaga mawong matou ubiradi checksin thourang loubiraga matam chana hidak thabiyu. - Tha 3 shuraba oakta Swine Fever naraktanaba hidak kappiyu. - Unshada tourakpa ana-ayek ngakthoknaba masagi arumba kg amada 200mcg gi chaangda Ivermectin (Indectin tab.) pibiyu natraga masagi arumba kg 50 khudingda 1ml gi chaangda Ivermec 1% injection kappiyu. - F.M.D. laina ngakthoknaba hidak kappiyu.



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 13-03-2020

Weather Forecast of KAKCHING(Manipur) Issued On : 2020-03-13(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-03-14	0.0	27.0	11.0	53	24	3.2	338	1
2020-03-15	0.0	28.0	11.0	63	25	4.9	289	5
2020-03-16	0.0	28.0	11.0	60	25	6.7	288	4
2020-03-17	0.0	28.0	12.0	53	19	9.2	270	5
2020-03-18	0.0	29.0	12.0	59	26	10.1	292	4

Weather Summary/Alert:

Likely to have no light rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 27-29oC & mini. temp. 11-12 oC. Relative humidity may reach upto 63% (max) & 19% (min). The average wind speed may be 3.2-10.1 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 27-29oC amadi ingbagina 11-12 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 63% amadi nunghildana 19% oiraba yai. Nungsit sitpagi changna pung amada 3.2-10.1 km oiraba yai.

General Advisory:

Disinfect the farm premises of all livestock/ poultry (surrounding area of 3ft width) using the repellent agent/ insecticides spray. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free or polythene sheets to preserve soil moisture for the crop. Monitor the soil moisture status regularly and give life-saving irrigation, especially at growth stages when water requirement is higher.

Kang-hayinggi yamna indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nanna thambiyu. Leibak leinil kuina leinaba pambi makhongda laina tilna paktaba charu usa-wasa natraga polythene na kupsinbiyu. Loufamda pambi houbada ishinggi awatpa leitanaba toina sennabiraga darkar leibada ishing thajinbiyu.

SMS Advisory:

Harvest the mature crops like rapeseed/ pea/ lentil/ broad bean/ potato; select pure healthy insect pest free seeds & sundry properly before storing as seed.

Hangam, hawai tharak/ moobi, masoori amadi alu lokpiraga laina paktaba mapung faba mahei khangatpiraga nunghshada ningthina kanghanlaga maru oinaba thamjinbiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
-----------------	------------------------

Crop(Varieties)	Crop Specific Advisory
SOYABEAN	Procure insect pest free healthy seeds of soybean/ groundnut. Get the soil testing done for proper fertilizer application. If soil is acidic i.e., PH level is below 6.5 then liming in furrows is required @ 500kg/ ha for having better yield.
	Nung-hawai thanaba afaba mahal laina tilna paktaba maru khanlaga lousinbiba yare. Leibak leihao amadi PH gi changyeng toubiraga PH 6.5 gi makhada leirabadi pari amada 500kg shunu paring naina churi sembiraga happiradi pothook hengathanba ngamgani

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
ONION	Spray Sixer @ 2 gm/lit of water as prophylactic measure against Purple Bloch and bulb rot. Spray is to be given twice at an interval of 7 to 10 days.
	Tilhou manada angouba mapak paklakpa amadi kuisinlakpada khamen machu onsinlakpa natraga tilhou maru patharakpa laina ngakthoknaba numit 7-10ni kannaraga ishing litre amada 2g gi chaangda Sixer happaga kappiyu.

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	- Any type of influenza may be given symptomatic treatment with antibiotics. - Vaccinate against swine fever after the age of 3 months. - Give Ivermectin (Indectin tab.) @ 200mcg/ kg body weight or Ivermec 1% injection @ 1ml/50 kg body weight once only to avoid skin infection. - Vaccinate against F.M.D.
	- Lok thunglakpaga mawong matou ubiradi checksin thourang loubiraga matam chana hidak thabiyu. - Tha 3 shuraba oakta Swine Fever naraktanaba hidak kappiyu. - Unshada tourakpa ana-ayek ngakthoknaba masagi arumba kg amada 200mcg gi chaangda Ivermectin (Indectin tab.) pibiyu natraga masagi arumba kg 50 khudingda 1ml gi chaangda Ivermec 1% injection kappiyu. - F.M.D. laina ngakthoknaba hidak kappiyu.



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 13-03-2020

Weather Forecast of KAMJONG(Manipur) Issued On : 2020-03-13(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-03-14	0.0	25.0	10.0	55	26	4.9	338	1
2020-03-15	0.0	25.0	10.0	59	26	4.3	341	7
2020-03-16	1.0	25.0	9.0	57	24	5.5	338	5
2020-03-17	0.0	25.0	10.0	46	15	6.6	334	6
2020-03-18	0.0	26.0	10.0	56	24	6.8	324	4

Weather Summary/Alert:

Likely to have no light rains upto Wednesday. The sky will remain mainly clear to partly cloudy with max. temp. 25-26 oC & mini. temp. 9-10 oC. Relative humidity may reach upto 59% (max) & 15% (min). The average wind speed may be 4.3-6.8 km/hr.

Lakliba Yumjakeisa faoba nong chudaba yai. Atiyada leichil khara khanlaba yai. Khwaidagi sabagi changna 25-26 oC amadi ingbagina 9-10 oC oiraba yai. Nungsitta ishinggi manilgi changna yammabada ayukta 59% amadi nunghildana 15% oiraba yai. Nungsit sitpangi changna pung amada 4.3-6.8 km oiraba yai.

General Advisory:

Disinfect the farm premises of all livestock/ poultry (surrounding area of 3ft width) using the repellent agent/ insecticides spray. Mulching may be done in the crop field using straw/ dried leaves which are insect-pest free or polythene sheets to preserve soil moisture for the crop. Monitor the soil moisture status regularly and give life-saving irrigation, especially at growth stages when water requirement is higher.

Kang-hayinggi yamna indanaba sha uchekki makol manung amadi akoibada leiba mafam lu-nanna thambiyu. Leibak leinil kuina leinaba pambi makhongda laina tilna paktaba charu usa-wasa natraga polythene na kupsinbiyu. Loufamda pambi houbada ishinggi awatpa leitanaba toina sennabiraga darkar leibada ishing thajinbiyu.

SMS Advisory:

Harvest the mature crops like rapeseed/ pea/ lentil/ broad bean/ potato; select pure healthy insect pest free seeds & sundry properly before storing as seed.

Hangam, hawai tharak/ moobi, masoori amadi alu lokpiraga laina paktaba mapung faba mahei khangatpiraga nungshada ningthina kanghanlaga maru oinaba thamjinbiyu.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
-----------------	------------------------

Crop(Varieties)	Crop Specific Advisory
SOYABEAN	Procure insect pest free healthy seeds of soybean/ groundnut. Get the soil testing done for proper fertilizer application. If soil is acidic i.e., PH level is below 6.5 then liming in furrows is required @ 500kg/ ha for having better yield.
	Nung-hawai thanaba afaba mahal laina tilna paktaba maru khanlaga lousinbiba yare. Leibak leihao amadi PH gi changyeng toubiraga PH 6.5 gi makhada leirabadi pari amada 500kg shunu paring naina churi sembiraga happiradi pothook hengathanba ngamgani

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
ONION	Spray Sixer @ 2 gm/lit of water as prophylactic measure against Purple Bloch and bulb rot. Spray is to be given twice at an interval of 7 to 10 days.
	Tilhou manada angouba mapak paklakpa amadi kuisinlakpada khamen machu onsinlakpa natraga tilhou maru patharakpa laina ngakthoknaba numit 7-10ni kannaraga ishing litre amada 2g gi chaangda Sixer happaga kappiyu.

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	- Any type of influenza may be given symptomatic treatment with antibiotics. - Vaccinate against swine fever after the age of 3 months. - Give Ivermectin (Indectin tab.) @ 200mcg/ kg body weight or Ivermec 1% injection @ 1ml/50 kg body weight once only to avoid skin infection. - Vaccinate against F.M.D.
	- Lok thunglakpaga mawong matou ubiradi checksin thourang loubiraga matam chana hidak thabiyu. - Tha 3 shuraba oakta Swine Fever naraktanaba hidak kappiyu. - Unshada tourakpa ana-ayek ngakthoknaba masagi arumba kg amada 200mcg gi chaangda Ivermectin (Indectin tab.) pibiyu natraga masagi arumba kg 50 khudingda 1ml gi chaangda Ivermec 1% injection kappiyu. - F.M.D. laina ngakthoknaba hidak kappiyu.