



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
 Mizoram Centre, Kolasib- 796081, MIZORAM
(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



District: Lawngtlai

Period: 03 May – 07 May, 2017

Bulletin No: - 697/2017/ Bulletin/Mizo

Date of issue: 02nd May, 2017

Parameters	03.05.2017	04.05.2017	05.05.2017	06.05.2017	07.05.2017
Rainfall (mm)	0	0	0	0	0
Max Temp (°C)	32	33	35	34	33
Min Temp (°C)	20	22	21	21	21
Cloud Coverage	Mainly clear	Mainly clear	Mainly clear	Mainly clear	Mainly clear
Max RH (%)	97	81	88	83	87
Min RH (%)	35	32	32	30	36
Wind Speed (Kmph)	4	4	4	5	4
*Wind Direction	E	E	S-E	S-E	S-E

**Northerly- N, North-Easterly- N-E, Easterly- E, South-Easterly- S-E,
 Southerly- S, South-Westerly- S-W, Westerly- W, North-westerly- N-W.**

STATUS OF MONSOON- June 1-30, 2016 (Percent of deviation from normal in parenthesis)

Aizawl- 384.87mm (430.2mm)	Champhai- 105.48mm (359.89mm)	Saiha- 307.40 mm (507.7mm)	Kolasib- 236.00mm (428.1mm)
Lawngtlai-291.20mm (453.1mm)	Lunglei-326.00mm (465.14mm)	Mamit-204.87mm (442.80mm)	Serchhip-411.72mm (259.62mm)

Weather summary of the past three days

Maximum Tem. (°C):28-31°C
Minimum Tem. (°C):18-21°C
Maximum RH (%):91-98%
Minimum RH (%):55-82%
Wind Direction: southeasterly
Cloud cover: Mainly clear
Wind speed: 2-3 km/hr


Rainfall: 00.0 mm

**03rd May– 07th May, 2017 chhunga sik leh sa
 dinhmun tur tlangpui**

Tun ni 5 chhung lo awm turah hian ruahtui tla miahlo tura beisei a ni. Khua a lum lai berin 32-35°C a ni ang a. A vawh lai ber in 20-22°C ni tura beisei a ni. RH san lai berin 81-97% leh a hniam lai berin 30-36% ni tur a rin niin. Thli hi darkar khatah 4-5 km vela chakin chhaklam awi zawngin a tleh rin a ni. A tlangpuiin tun ni nga chung hian khawthiang tak hmuh beisei a ni.

Weekly cumulative rainfall: 00.0mm

NDVI for Mizoram



Moderately wet mildly dry/mildly wet conditions



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



Main Crop / Animal / Fisheries	Stage	Cultural practices/ Pest/ Diseases	Agricultural / Horticultural/ animal husbandry advisories
FRUITS CROPS			
KHASI MANDARIN AND ACID LIME	A kui atanga a seng hun		- Thlasik laia thlai bul khoro lutuk tur vennan chuan hnim hnah hring tlai bul velah dahkhawm tur ani. - Thlai naupang deuah chuan chawlh kar tin a tui pek thin tur ani. - Leia tha mamawh tawh a hmuh theihna turin a hmunhma a hnim awm te thlawhfai thin tur ani. - A seng hma kar 6 chhung chu tui tha taka pek hian a rah tla tur chelh nan leh a rah than that nan te leh a rah keh tur lakah t a veng thei ani.
BANANA			
STAR FRUIT			
PLUM AND PEACH			
		Gummosis, citrus canker, citrus greening and Dieback	Temperture hniam lutuk leh hnawng vang hian natna a a tam duh a . Soil bome natna laka vennan Bordeaux past hi thing zar leh a trangah te hnawih tur ani.
		Fruit fly	Huan zau takah chuan a par tan tirh leh a rah tan tirin chawlhkar hnih chhung chu heng te hian enkawl tur ani: carbaryl 0.2 percent emaw malathion 0.15 percent suspension containing sugar or jeggery at 10 g/l.
PLANTATION CROP			
COFFEE	All stages		Nursery stage - Thlai chi thlak hma in <i>Azospirillum</i> leh <i>Phosphobacterium</i> a enkawl tur ani. - A chi hi December – January ah hmun zawl/rualrem 1.5 - 2.5 cm a in hlatin tlar mumal tak siam in chin tur ani. - Chuan a chi chu lei tlem te a chhilh a buhpawla khuh tur ani. - Nitin tui pek tur ani a, a sat lutuka loh nan niin a chhun loh nan zar hlih tur ani. - Ni 45 hnu velah a tiak thin a,chu chu bag ah an sawn chhuak leh thin ani.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			Harvesting Stage - Coffe rah hmin hi thlasik lain an seng thin a ni. - A rah hmin tha lo ho chu nuai sawm hmain an thliar hrang leh vek thin ani.
		Coffee Berry borer	- A hun takah leh fimkhur taka seng tur ani. - Hmaih neih nuaih loh tur ani. - Sneg hmaih te lakkhawma paih vek tur ani. - Seng bang zawng zawng chu uluka taka paih fai vek tur ani. - A thlai vennis a rah thlai ho chu paih vek tur ani. - Hmun dam lutukah dah loh tur. - Boruak tha taka a hmuh theih nan leh a rawn chawr no theih nan thlai chu uluk tak a hlawi tur ani. - Chuan hei hian kah tur ani. Quinalphos 25 EC @ 340 ml/200 lit emaw lamda cyhalothrin 5 EC 120 – 160 ml / 200 lit. - In leh loa sawngbawl a a ro dan tur tawkah chuan: Arabica / robusta parchment 10 %, Arabica cherry 10.5 % leh robusta cherry 11.0 %.
		Coffee Rust	- Natna hrik in a khawih tawh a hnah leh a dang te paihpaih vek tur ani. - Bordeaux mixture 0.5% in February - March (Pre-bloom) a kah phawt a, Oxycarboxin 0.03% in May - June (Pre-monsoon) ah kah chhunzawm tur ani.
CEREALS AND PULSE CROPS			
Rabi Maize	A chin hun		- Vaimim chinna tur atan lei kan let phut darh anga. - Hei hian a rawn to chhuah na tur atan a pui dawn a ni. - A chi chu kan lei leh saah chuan kan dah ang. - A chi chu Thiram @4 g/kg seed hian ak sawngbawl hmasa anga. 20-25 kg/ha vel a chi thlak hi a tawk vel viau ani. - A chi tuh hma in lei leh FYM @5-10 t/ha leh 80:60:40 kg N, P₂O₅ leh



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			K ₂ O/ha pawlh chu hman phawt tur a ni. Nitrogen dose chanve chu a chi tuh hunlaia hman tur a ni a, tichuan a bang 25% chu thla khat hnu ah ani ang a adang leh 25% chu a par hunah hman tur a ni.
Soybean, pea, lentil toria, breen gram and black gram cultivation in rice fellow	All stage	Zero tillage	- A than a that theih nan nikhat danah tui pek thin tur ani. - Lei rih vur hian thlai kung te a veng ve ani. - Thlasik lai a lei khoro lutuk tur ven nan a chungah hnim leh thildanga khuh tur ani.
Potato	Sowing stage		- Muangchang loving alu chin na tur chu buatsaih vat tur ani. - Hei hian a than hun lain natna hrikin lakah a veng dawn ani. - Lei leh hmain a hmun hma chu fai taka thlawh hmasak tur ani. - A chi thlak hma in a chi chu en fiah hmasak tur ani. - A than a that theih nan nikhat danah tui pek thin tur ani.
VEGETABLE CROP			
Tomato	Bacterial Blight disease		- Tomato bikah chuan sik leh sa hi natna an kaina tlang lawn ber ani . - Hmun hnawng leh ni hmu lo lutuk hmunah chuan natna an kai hma bik ani. - Tomato hi a uai a, a thih mai loh nan Ridomil emaw Indofil emaw Mancozeb @ 2 gm hi tui liter 1 ah pawlh a kah tur ani .
Early Cole crop	Black spot disease		- A than a that theih nan nikhat danah tui pek thin tur ani. - Thlai bul vawn hnawn nana thlai bula hnim ring vawm khawm hi tui pek zawhah dah tur ani. - Zikhlum lam chi ah chuan sik leh sa vangin a hnah ah thil dum rawn



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			awm thin a , hei hi natna tlanglawn ber ani. Thlai hna lam chi leh zikhlum lam chi reng reng enkawl nan Mancozeb @ 2gm ah tui leter 1 pawlha kah tur ani.
Onion and capsicum	Nursery stage	Poly house	- A than a that theih nan nikhat danah tui pek thin tur ani. - Thlai bul vawn hnawn nana thlai bula hnim ring vawm khawm hi tui pek zawhah dah tur ani. - Thlai chhina hmun (nursery) hi hnim a to loh nan Pendimethalin @ 3.5ml hi tui liter 1 zelah pawlh a kah hi a tha hle ani.
		Phytophthora blight	- A chi ven that nan thiram 3g/kg seed emaw Trichoderma viride 4g+ metalaxyl 4g (Apron)/ kg seed hi a tha hle ani - Hneh taka 1% Bordeaux chawhpawlh emaw 2 g captan emaw 3 copper oxychloride a tui liter 1 hi 10-15 DAS a pek hi a tha hle ani.
French bean	Sowing stage		- Tui pek a hnihnah hringa khuh tur ani a. than a that theih nan tui pek hma in lei rin pan hmasak tur ani. - A than duna theih nan leh hnim to loh na turin a kung bulah lei vur chhoh zel tur ani.
Carrot and radish	Sowing stage		- A than a that theih nan nikhat danah tui pek thin tur ani. - Tui pek hnuah thlai bul vawn hnawn na tur siam tur ani. - Zikhlum lam chi ah chuan sik leh sa vangin a hnah ah thil dum a rawn awm thina, hei hi natna tlanglawn ber ani. - Thlai hna lam chi leh zikhlum lam chi reng reng enkawl nan Mancozeb @ 2gm ah tui leter 1 pawlha kah tur ani.






GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



ANIMAL HUSBANDRY			
Pig	All stages		- Khua a vawh hian vawk hian an mahni in tih lumna tur atan chakna an mamawhna a sang bik ani. - An hriselna that leh that loh enfiah renga, a chaw ei tur tlem tlema tih tam hret hret tur ani.. - Sangha tel ah hian omega-3 hi atam em a vangin an chakna muangchanga a in siam chhoh zel theih nan a tha hle ani.
			Porcine Reproductive Respiratory Syndrome (PRRS). 1. Vawknote emaw vawk lak hran.
	Adult stage		Swine fever. 2. SF vaccines hi thla 2 hnua pek tur ani a, chumi hnuah chuan kumtin thlaruk danah pek chhonzawm tur ani.
Cattle	All age group		Hun rei tak khua a ro avanga hnim hnah hring peh tur a awm loh laia bawngin an chaw ei in buk tawk tur leh an taksa tana mamawh tur atan buh kung urea molasses hmanga sawngbawl pek tur ani.
	All age group		Kar 16 hnuah FMD vaccine pek a, chuan thla tin thla 6 chhung chhonzawm tur ani.
	Young stage		- Black Quarter Vaccine (BQV). - Vaccinne hmasa ber hi thla 6 ah emaw a hnu lamah pek tur. - Chumi hnuah chuan Vaccine hi kum tin pek tur ani.
Poultry	Litter management		- Ar te hian hmun thawl nuam tawk, chaw tha an mamawh tawk leh tui thianghlim an mamawh tawk an hmu tur ani a. - An hriselna atan enkawltha tha tawk tak pek hian natna an kai mai theih tur lak atang a venna tha ber ani.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			- Tui an in tur chhawpna tur tha /lian tha tak leh tui thianghlim tak pek tur ani. - Chaw a hmuar/thing pek loh tur ani a, an chaw eitur thlak sak thut loh tur ani.
	Preventive measures	0-3rd week	Ranikhet Disease- an pian atanga ni 1-6 ah F1 vaccine pek tur ani a, chuan a puitlingh chuan R₂B vaccine pek tur ani. - B complex with antibodies
		4th weeks	Coccidiosis- Amprolium or coccidiostat
		4-5th Weeks	Calcium tonic fortified with B₁₂
FISHERY			
	Pond preparation (Dil buatsaih)	3rd-4th weeks	Dil a leitha hman hian sangha chaw kan tih mai planktons insiam nan a tanpui thin.
			- Bawngkek hring 10 tonnes/ha/year vel dil ah hman thin a ni a; bawngkek kumkhat a kan mamawh zat hmunthum a then a hmunkhat hi dil buatsaih nan hman tur ani. A bak zawng hi tui boruak a zirin semdarh a hman thin tur ani. - Single super phosphate hi dil hectare khat zel a zauah kg 250 ang a hman thin tur ani.
			- Leitha kan hman atang a karkhat hnuah tui rawng a tangin sangha chaw inseam that leh thatloh a hriat theih a. Tui rawng eng hring deuh nghalh ah hian sangha chaw planktons te an inseam tha ang a ngaih ani. - Tui nut zawng tehma transparency pawh 30-40 cm vel ani tur ani.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD,
Guwahati)



Expert committee members:

Dr. S.B. Singh	:	Joint Director	basantasinghsoibam@rediffmail.com
Dr. Saurav Saha	:	Scientist (Agril. Physics)	sauravs.saha@gmail.com
Dr. T. Boopathi	:	Scientist (Agril Entomology)	boopathiars@gmail.com
Dr. A. Ratankumar Singh	:	Scientist (Plant Pathology)	ratantplantpatho@gmail.com
Dr. Lungmuana	:	Scientist (Soil Fertility)	lmsingson@gmail.com
Mr. P.L. Lalrinsanga	:	Scientist (Aquaculture)	viensky2@gmail.com
Dr. Dr. V. Dayal	:	Scientist (Horticulture)	Vishambhai5009@gmail.com
Dr. Samuel Lalliansanga	:	Head & Sr. Scientist	samuelpachua10@gmail.com
Mr. Samik Chowdhury	:	Technical Officer	samikchowdhury33@gmail.com
Mr. Diktea chenkuai	:	Project Assistant	dikteachenkuaiboy@gmail.com

Collaborating Department:

Name of the KVK	Programme Coordinator Name and Designation	KVK Email Id	Phone no/ Mobile no
KVK Lunglei	Dr. Lalmuanzovi Head & Sr. Scientist	kvkhnahthial@gmail.com	9862803750 9436154614
KVK, Kolasib	Mr. Lalrosamga Khiangte Head & Sr. Scientist	kvkkolasib@gmail.com	9436152440
KVK, Serchhip	Mr. K. Laltlanmawia Head & Sr. Scientist	kvkserchhip@gmail.com	9436146115 9615389293
KVK, Champhai	Mrs. Lalrinawmi Renthlei Head & Sr. Scientist	kvkchawzawl@gmail.com	9436159788
KVK, Lawngtlai	Dr. Michel Lallawmkimi Head & Sr. Scientist	kvklawntlai@gmail.com	9436155858
KVK, Saiha	Dr. Vanlalhraia Hnampe Head & Sr. Scientist	kvksaiha@gmail.com	8974656509
KVK, Mamit	Dr. Samuel Lalliansanga Head & Sr. Scientist	kvkmamit@gmail.com	9436147625
KVK, Aizawl	Dr. K. P. Chaudhary Head & Sr. Scientist	Kpchy@rediffmail.com kvkaizawl@rediffmail.com	9436351669



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
 Mizoram Centre, Kolasib- 796081, MIZORAM
(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



District: Lunglei

Period: 03 May – 07 May, 2017

Bulletin No: - 697/2017/ Bulletin/English

Date of issue: 02nd May, 2017

Parameters	03.05.2017	04.05.2017	05.05.2017	06.05.2017	07.05.2017
Rainfall (mm)	0	0	0	0	0
Max Temp (°C)	32	32	32	33	33
Min Temp (°C)	22	22	22	22	22
Cloud Coverage	Mainly clear	Mainly clear	Mainly clear	Mainly clear	Mainly clear
Max RH (%)	99	87	83	85	93
Min RH (%)	37	33	31	29	33
Wind Speed (Kmph)	3	4	4	4	4
*Wind Direction	E	E	S-E	S-E	S

**Northerly- N, North-Easterly- N-E, Easterly- E, South-Easterly- S-E,
 Southerly- S, South-Westerly- S-W, Westerly- W, North-westerly- N-W.**

STATUS OF MONSOON- June 1-30, 2016 (Percent of deviation from normal in parenthesis)

Aizawl- 384.87mm (430.2mm)	Champhai- 105.48mm (359.89mm)	Saiha- 307.40 mm (507.7mm)	Kolasib- 236.00mm (428.1mm)
Lawngtlai-291.20mm (453.1mm)	Lunglei-326.00mm (465.14mm)	Mamit-204.87mm (442.80mm)	Serchhip-411.72mm (259.62mm)

Weather summary of the past three days

Maximum Tem. (°C):26-29°C
Minimum Tem. (°C):16-19°C
Maximum RH (%):95-98%
Minimum RH (%):44-78%
Wind Direction: Southwesterly
Cloud cover: Mainly clear
Wind Speed: 2-3 km/hr


Rainfall: 00.0 mm

Weather forecast valid from 03rd May, 2017 To 07thMay, 2017.

There are no chances of rainfall during the next 5 days. The maximum and minimum temperatures for the next 5 days may range for 32-33°C and 22°C. Maximum relative humidity is expected in the range of 83-99% and minimum may from 29-37%. Wind direction would be easterly to southeasterly and southerly with the wind speed of 3-4 km per hour. Mainly clear sky will prevail during the next five days.

Weekly cumulative rainfall: 00.0 mm

NDVI for Mizoram



Moderately wet mildly dry/mildly wet conditions



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



Main Crop / Animal / Fisheries	Stage	Cultural practices/ Pest/ Diseases	Agricultural / Horticultural/ animal husbandry advisories
FRUITS CROPS			
KHASI MANDARIN AND ACID LIME	Seedling transplanting stage		- Fruit plant should be planted in a sunny and wind-protected area. - In the citrus belt, trees can be planted at any time, however, spring is the best time for container grown plants. - Standard-size trees should be spaced 12 to 25 feet apart and dwarf trees should be set 6 to 10 feet apart. The exact distance depends on the variety. The bigger the fruit, the farther the distance. - If the soil is not well-drained, plant the trees on a slight mound to prevent water logging. - Mulching should be done with dry grasses near the tree base to conserve soil moisture during winter. - The young fruit plant must be irrigated at weekly interval for better establishment.
BANANA			
STAR FRUIT			
PLUM AND PEACH			
		Gummosis, citrus Canker, Citrus greening, Dieback, Lamon butterfly and leaf minor	Die back- Due to low temperature and humidity disease appearance will more. Use Bordeaux past in tree trunk, twigs and branches protect healthy plant from soil borne disease. Lamon butterfly- Spray monocrotophos @0.04% @1.2 ml/lt of water. Leaf minor- Spray confidor 0.05% (0.5 ml/lit of water) at each flush emergence. Citrus Canker- Apply bacterimycin @0.6 g/lt of water.
Passion Fruit	Transplanting stage		- High yielding mother vine with good quality fruits and free of virus diseases should be selected to provide cuttings. - A cutting should contain at least 3 buds and must be planted in sand beds.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			- Immediately after planting these should be kept inside a high humid chamber made out of bamboo and polythene. Grafting: - The root stock of yellow Passion fruit is planted in polythene sleeves and the section from Rahangala hybrid is grafted using wedge or approach method of grafting.
Pineapple	Flowering stage		- Apply flowering inducing chemical (Ethrel 10 PPM+2% urea+0.04% Sodium Carbonate) should be applied in the heart of the plant. In evening and only when plants have at least 32 leaves. - The flowering emergence will come out after 55-60 days after chemical spraying. - Apply split doses of fertilizer @ 60:50:60 g per plant. - Remove all unwanted leaves, branches and weed near to the plant.
PLANTATION CROP			
COFFEE	Blooming stage		- If day temperature and prolong dry spell occur it lead to Floral abnormalities like "Star Flower" in Arabica and "Pink Flower" in Robusta. - Irrigation of plants at alternate day's interval, 6 weeks before harvesting improves fruit retention and fruit development, and reduces fruit cracking and improves the fruit quality. - Mulching should be done with dry grasses near the tree base to conserve soil moisture during winter. - The young fruit plant must be irrigated at weekly interval for better establishment. - Foliar application of Mepiquat chloride @ 1000 PPM concentration or 0.75% SSP @ 1.5 g per 200 lt of water 15 days




GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



Rubber	Nursery stage		interval. ✚ Clearing operation may be done during the month of February to April. ✚ Make fire line to protect the young tree and seedlings. ✚ 10-12 kg of well rotten organic manure and 225 gm rock phosphate should be apply at time of planting to each pit as basal dose application.
CEREALS AND PULSE CROPS			
Pre Kharif Rice	Nursery stage		✚ Use only Well filled and healthy seeds. ✚ Put the seed in 2.5% salt solution i.e 250 g of common salt in 10 lts of water. ✚ Seed treated with Bavistin 50 WP @ 0.1% (2 g/ltr) solution. ✚ Mulching is requiring for better germination in nursery.
Jhum Rice	Sowing stage		✚ Sowing in ridge and furrow. ✚ Mulching with grass for better germination. ✚ Intercropping with greengram and bunchy type frenchbean.
Maize (Jhum)	Sowing stage		✚ Seed should be treated with Thiram @4 g/kg seed. ✚ Use optimum seed rate (20-25 kg/ha) for desire plant population. ✚ Apply well decomposed FYM/pig manure @ 5-10 t/ha along with 80:60:40 kg N, P₂O₅ and K₂O/ha incorporate with soil before sowing. Half nitrogen dose will use at the time of sowing and remaining 25% after one month and 25% at flowering stage.
Rabi Maize	vegetative stage		✚ Light irrigation on every week may be given for better establishment and smooth growth. ✚ Earthing up soil near to plant for better support. ✚ Maize rust disease will prevail due to high relative humidity with low temperature. Apply Mancozeb Kg/ha



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			for effective control. Remove the alternate host <i>Oxalis comiculata</i>.
VEGETABLE CROP			
Cowpea	Sowing stage	KOLASIB	- Plough the field properly, at least 2-3 times. - Mix fertilizer with FYM 50:60:60Kg/ha. - Sow 2-3 seed per whole. - Spacing should be 30 X 20 cm.
Okra	Sowing stage	Weeding and light irrigation in nursery bed. Provide irrigation in transplanted okra field.	- Plough the field with the help of spade. - Sow 2 seed 45 X 45 cm spacing. - Before sowing seed provide one or two irrigation. - Provide fertilizer @ 120: 60: 60 Kg/ha
Ginger and turmeric	Sowing stage	SERCHHIP	- Rhizome should be treated with Thiram @4 g/kg seed. - Use optimum seed rate (50-60 kg/ha) for desire plant population. - Apply well decomposed FYM/ pig manure @ 10-20 t/ha along with 120:80:60 kg N, P₂O₅ and K₂O/ha incorporate with soil before sowing. Half nitrogen dose will use at the time of sowing and remaining 25% after one month and 25% at flowering stage.
Colocasia	Sowing stage	LUNGLEI	- Planting is done well prepared land or pits filled up with FYM (12-15) t/ha - Sprouted corms or cormels are planted 5-7 deep at a spacing of 40-50 cm between and within rows in the pits. - Inorganic fertilizer like Urea, SSP and MOP @ 220: 375: 134 kg.
ANIMAL HUSBANDRY			
Pig	All stages	LAWNGTALA	- As the weather gets colder, your pigs' energy requirement will increase, as they need more energy to keep warm. - Regularly monitor their level of 'fitness'



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			and increase their feed intake to maintain. ✚ Fish oils are excellent for providing slow-release energy with the added advantage of a high level of omega-3.
		Porcine Reproductive Respiratory Syndrome (PRRS).	1. Culling of positive pigs or piglets.
	Adult stage	Swine fever.	2. Vaccination of pigs with SF vaccines at 2 months and yearly interval/6 month interval
Cattle	All age group		Due to prolong dry spell there is a shortage of green grass in the field. For balanced diet and nutrition to your cattle, provide urea molasses treated paddy straw.
	All age group	Foot and Mouth Disease (FMD)	FMD vaccine at 16 week and repeat every 6 month.
	Young stage	Black Quarter (BQ)	- Black Quarter Vaccine (BQV). ✦ Primary vaccination 6 month or above ✦ Revaccination annually
Poultry	Litter management		✚ Birds require adequate space, sufficient feed to meet their nutritional requirements and an adequate supply of good-quality water. ✚ Good management and sanitation are the best ways to avoid infectious disease in poultry. ✚ Provide ample quantity of clean drinking water. ✚ Avoid feeding of mouldy feed. Don't make sudden changes in feed
	Preventive measures	0-3rd week	✚ Ranikhet Disease- F1 vaccine at (1-6) days of birth and R₂B vaccine for adult birds. ✚ B complex with antibodies
		4th weeks	✚ Coccidiosis- Amprolium or coccidiostat



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



		4-5 th Weeks	✚ Calcium tonic fortified with B ₁₂
FISHERY			
	Pond preparation	3 rd -4 th weeks	✚ Application of fertilizers/manure helps in development of plankton which serve as natural feed for the fishes.
		KOLASIB	✚ Raw cowdung should be applied in the pond at the rate of 10 tonnes/ha/year. One third of the total dose should be applied initially and the rest may be applied in a spilt doses. ✚ Single super phosphate should also be applied at the rate of 250 kg/ha in the pond.
		AIZAWL	✚ After one week of application development of planktons could be observed in the pond depending on the colour of the water. Yellowish green colour is an indicator of the good plankton development. ✚ Transparency of the water needs to be maintained at 30-40 cm.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD,
Guwahati)



Expert committee members:

Dr. S.B. Singh	:	Joint Director	basantasinghsoibam@rediffmail.com
Dr. Saurav Saha	:	Scientist (Agril. Physics)	sauravs.saha@gmail.com
Dr. T. Boopathi	:	Scientist (Agril Entomology)	boopathiars@gmail.com
Dr. A. Ratankumar Singh	:	Scientist (Plant Pathology)	ratantplantpatho@gmail.com
Dr. Lungmuana	:	Scientist (Soil Fertility)	lmsingson@gmail.com
Mr. P.L. Lalrinsanga	:	Scientist (Aquaculture)	viensky2@gmail.com
Dr. Dr. V. Dayal	:	Scientist (Horticulture)	Vishambhai5009@gmail.com
Dr. Samuel Lalliansanga	:	Head & Sr. Scientist	samuelpachua10@gmail.com
Mr. Samik Chowdhury	:	Technical Officer	samikchowdhury33@gmail.com
Mr. Diktea chenkual	:	Project Assistant	dikteachenkualboy@gmail.com

Collaborating Department:

Name of the KVK	Programme Coordinator Name and Designation	KVK Email Id	Phone no/ Mobile no
KVK Lunglei	Dr. Lalmuanzovi Head & Sr. Scientist	kvkhnahthial@gmail.com	9862803750 9436154614
KVK, Kolasib	Mr. Lalrosamga Khiangte Head & Sr. Scientist	kvkkolasib@gmail.com	9436152440
KVK, Serchhip	Mr. K. Laltlanmawia Head & Sr. Scientist	kvkserchhip@gmail.com	9436146115 9615389293
KVK, Champhai	Mrs. Lalrinawmi Renthlei Head & Sr. Scientist	kvkchawzawl@gmail.com	9436159788
KVK, Lawngtlai	Dr. Michel Lallawmkimi Head & Sr. Scientist	kvklawntlai@gmail.com	9436155858
KVK, Saiha	Dr. Vanlalhruaia Hnampe Head & Sr. Scientist	kvksaiha@gmail.com	8974656509
KVK, Mamit	Dr. Samuel Lalliansanga Head & Sr. Scientist	kvkmamit@gmail.com	9436147625
KVK, Aizawl	Dr. K. P. Chaudhary Head & Sr. Scientist	Kpchy@rediffmail.com kvkaizawl@rediffmail.com	9436351669



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
 Mizoram Centre, Kolasib- 796081, MIZORAM
(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



District: Lunglei

Period: 03 May – 07 May, 2017

Bulletin No: - 697/2017/ Bulletin/Mizo

Date of issue: 02nd May, 2017

Parameters	03.05.2017	04.05.2017	05.05.2017	06.05.2017	07.05.2017
Rainfall (mm)	0	0	0	0	0
Max Temp (°C)	32	32	32	33	33
Min Temp (°C)	22	22	22	22	22
Cloud Coverage	Mainly clear	Mainly clear	Mainly clear	Mainly clear	Mainly clear
Max RH (%)	99	87	83	85	93
Min RH (%)	37	33	31	29	33
Wind Speed (Kmph)	3	4	4	4	4
*Wind Direction	E	E	S-E	S-E	S

**Northerly- N, North-Easterly- N-E, Easterly- E, South-Easterly- S-E,
 Southerly- S, South-Westerly- S-W, Westerly- W, North-westerly- N-W.**

STATUS OF MONSOON- June 1-30, 2016 (Percent of deviation from normal in parenthesis)

Aizawl- 384.87mm (430.2mm)	Champhai- 105.48mm (359.89mm)	Saiha- 307.40 mm (507.7mm)	Kolasib- 236.00mm (428.1mm)
Lawngtlai-291.20mm (453.1mm)	Lunglei-326.00mm (465.14mm)	Mamit-204.87mm (442.80mm)	Serchhip-411.72mm (259.62mm)

Weather summary of the past three days

Maximum Tem. (°C):26-29°C
Minimum Tem. (°C):16-19°C
Maximum RH (%):95-98%
Minimum RH (%):44-78%
Wind Direction: Southwesterly
Cloud cover: Mainly clear
Wind Speed: 2-3 km/hr


Rainfall: 00.0 mm

**03rd May– 07th May, 2017 chhunga sik leh sa
 dinhmun tur tlangpui**

Tun ni 5 chhung lo awm turah hian ruahtui tla miahlo tura beisei a ni. Khua a lum lai berin 32-33°C a ni ang a. A vawh lai ber in 22°C ni tura beisei a ni. RH san lai berin 83-99% leh a hniam lai berin 29-37% ni tur a rin niin. Thli hi darkar khatah 3-4 km vela chakin chhaklam awi zawngin a tleh rin a ni. A tlangpuiin tun ni nga chung hian khawthiang tak hmuh beisei a ni.

Weekly cumulative rainfall: 00.0mm

NDVI for Mizoram



Moderately wet mildly dry/mildly wet conditions



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



Main Crop / Animal / Fisheries	Stage	Cultural practices/ Pest/ Diseases	Agricultural / Horticultural/ animal husbandry advisories
FRUITS CROPS			
KHASI MANDARIN AND ACID LIME	A kui atanga a seng hun		- Thlasik laia thlai bul khoro lutuk tur vennan chuan hnim hnah hring tlai bul velah dahkhawm tur ani. - Thlai naupang deuah chuan chawlh kar tin a tui pek thin tur ani. - Leia tha mamawh tawh a hmuh theihna turin a hmunhma a hnim awm te thlawhfai thin tur ani. - A seng hma kar 6 chhung chu tui tha taka pek hian a rah tla tur chelh nan leh a rah than that nan te leh a rah keh tur lakah t a veng thei ani.
BANANA			
STAR FRUIT			
PLUM AND PEACH			
		Gummosis, citrus canker, citrus greening and Dieback	Temperture hniam lutuk leh hnawng vang hian natna a a tam duh a . Soil bome natna laka vennan Bordeaux past hi thing zar leh a trangah te hnawih tur ani.
		Fruit fly	Huan zau takah chuan a par tan tirh leh a rah tan tirin chawlhkar hnih chhung chu heng te hian enkawl tur ani: carbaryl 0.2 percent emaw malathion 0.15 percent suspension containing sugar or jeggery at 10 g/l.
PLANTATION CROP			
COFFEE	All stages		Nursery stage - Thlai chi thlak hma in <i>Azospirillum</i> leh <i>Phosphobacterium</i> a enkawl tur ani. - A chi hi December – January ah hmun zawl/rualrem 1.5 - 2.5 cm a in hlatin tlar mumal tak siam in chin tur ani. - Chuan a chi chu lei tlem te a chhilh a buhpawla khuh tur ani. - Nitin tui pek tur ani a, a sat lutuka loh nan niin a chhun loh nan zar hlih tur ani. - Ni 45 hnu velah a tiak thin a,chu chu bag ah an sawn chhuak leh thin ani.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			Harvesting Stage - Coffe rah hmin hi thlasik lain an seng thin a ni. - A rah hmin tha lo ho chu nuai sawm hmain an thliar hrang leh vek thin ani.
		Coffee Berry borer	- A hun takah leh fimkhur taka seng tur ani. - Hmaih neih nuaih loh tur ani. - Sneg hmaih te lakkhawma paih vek tur ani. - Seng bang zawng zawng chu uluka taka paih fai vek tur ani. - A thlai vennan a rah tlai ho chu paih vek tur ani. - Hmun dam lutukah dah loh tur. - Boruak tha taka a hmuh theih nan leh a rawn chawr no theih nan thlai chu uluk tak a hlawi tur ani. - Chuan hei hian kah tur ani. Quinalphos 25 EC @ 340 ml/200 lit emaw lamda cyhalothrin 5 EC 120 – 160 ml / 200 lit. - In leh loa sawngbawl a a ro dan tur tawkah chuan: Arabica / robusta parchment 10 %, Arabica cherry 10.5 % leh robusta cherry 11.0 %.
		Coffee Rust	- Natna hrik in a khawih tawh a hnah leh a dang te paihpaih vek tur ani. - Bordeaux mixture 0.5% in February - March (Pre-bloom) a kah phawt a, Oxycarboxin 0.03% in May - June (Pre-monsoon) ah kah chhunzawm tur ani.
CEREALS AND PULSE CROPS			
Rabi Maize	A chin hun		- Vaimim chinna tur atan lei kan let phut darh anga. - Hei hian a rawn to chhuah na tur atan a pui dawn a ni. - A chi chu kan lei leh saah chuan kan dah ang. - A chi chu Thiram @4 g/kg seed hian ak sawngbawl hmasa anga. 20-25 kg/ha vel a chi thlak hi a tawk vel viau ani. - A chi tuh hma in lei leh FYM @5-10 t/ha leh 80:60:40 kg N, P₂O₅ leh



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			K ₂ O/ha pawlh chu hman phawt tur a ni. Nitrogen dose chanve chu a chi tuh hunlaia hman tur a ni a, tichuan a bang 25% chu thla khat hnu ah ani ang a adang leh 25% chu a par hunah hman tur a ni.
Soybean, pea, lentil toria, breen gram and black gram cultivation in rice fellow	All stage	Zero tillage	- A than a that theih nan nikhat danah tui pek thin tur ani. - Lei rih vur hian thlai kung te a veng ve ani. - Thlasik lai a lei khoro lutuk tur ven nan a chungah hnim leh thildanga khuh tur ani.
Potato	Sowing stage		- Muangchang loving alu chin na tur chu buatsaih vat tur ani. - Hei hian a than hun lain natna hrikin lakah a veng dawn ani. - Lei leh hmain a hmun hma chu fai taka thlawh hmasak tur ani. - A chi thlak hma in a chi chu en fiah hmasak tur ani. - A than a that theih nan nikhat danah tui pek thin tur ani.
VEGETABLE CROP			
Tomato	Bacterial Blight disease		- Tomato bikah chuan sik leh sa hi natna an kaina tlang lawn ber ani . - Hmun hnawng leh ni hmu lo lutuk hmunah chuan natna an kai hma bik ani. - Tomato hi a uai a, a thih mai loh nan Ridomil emaw Indofil emaw Mancozeb @ 2 gm hi tui liter 1 ah pawlh a kah tur ani .
Early Cole crop	Black spot disease		- A than a that theih nan nikhat danah tui pek thin tur ani. - Thlai bul vawn hnawn nana thlai bula hnim ring vawm khawm hi tui pek zawhah dah tur ani. - Zikhlum lam chi ah chuan sik leh sa vangin a hnah ah thil dum rawn



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			awm thin a , hei hi natna tlanglawn ber ani. ✚ Thlai hna lam chi leh zikhlum lam chi reng reng enkawl nan Mancozeb @ 2gm ah tui leter 1 pawlha kah tur ani.
Onion and capsicum	Nursery stage	Poly house	✚ A than a that theih nan nikhat danah tui pek thin tur ani. ✚ Thlai bul vawn hnawn nana thlai bula hnim ring vawm khawm hi tui pek zawhah dah tur ani. ✚ Thlai chhina hmun (nursery) hi hnim a to loh nan Pendimethalin @ 3.5ml hi tui liter 1 zelah pawlh a kah hi a tha hle ani.
		Phytophthora blight	✚ A chi ven that nan thiram 3g/kg seed emaw Trichoderma viride 4g+ metalaxyl 4g (Apron)/ kg seed hi a tha hle ani ✚ Hneh taka 1% Bordeaux chawhpawlh emaw 2 g captan emaw 3 copper oxychloride a tui liter 1 hi 10-15 DAS a pek hi a tha hle ani.
French bean	Sowing stage		✚ Tui pek a hnihnah hringa khuh tur ani a. than a that theih nan tui pek hma in lei rin pan hmasak tur ani. ✚ A than duna theih nan leh hnim to loh na turin a kung bulah lei vur chhoh zel tur ani.
Carrot and radish	Sowing stage		✚ A than a that theih nan nikhat danah tui pek thin tur ani. ✚ Tui pek hnuah thlai bul vawn hnawn na tur siam tur ani. ✚ Zikhlum lam chi ah chuan sik leh sa vangin a hnah ah thil dum a rawn awm thina, hei hi natna tlanglawn ber ani. ✚ Thlai hna lam chi leh zikhlum lam chi reng reng enkawl nan Mancozeb @ 2gm ah tui leter 1 pawlha kah tur ani.






GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



ANIMAL HUSBANDRY			
Pig	All stages		- Khua a vawh hian vawk hian an mahni in tih lumna tur atan chakna an mamawhna a sang bik ani. - An hriselna that leh that loh enfiah renga, a chaw ei tur tlem tlema tih tam hret hret tur ani.. - Sangha tel ah hian omega-3 hi atam em a vanging an chakna muangchanga a in siam chhoh zel theih nan a tha hle ani.
			Porcine Reproductive Respiratory Syndrome (PRRS). 1. Vawknote emaw vawk lak hran.
	Adult stage		Swine fever. 2. SF vaccines hi thla 2 hnua pek tur ani a, chumi hnuah chuan kumtin thlaruk danah pek chhonzawm tur ani.
Cattle	All age group		Hun rei tak khua a ro avanga hnim hnah hring peh tur a awm loh laia bawngin an chaw ei in buk tawk tur leh an taksa tana mamawh tur atan buh kung urea molasses hmanga sawngbawl pek tur ani.
	All age group		Kar 16 hnuah FMD vaccine pek a, chuan thla tin thla 6 chhung chhonzawm tur ani.
	Young stage		- Black Quarter Vaccine (BQV). - Vaccinne hmasa ber hi thla 6 ah emaw a hnu lamah pek tur. - Chumi hnuah chuan Vaccine hi kum tin pek tur ani.
Poultry	Litter management		- Ar te hian hmun thawl nuam tawk, chaw tha an mamawh tawk leh tui thianghlim an mamawh tawk an hmu tur ani a. - An hriselna atan enkawltha tha tawk tak pek hian natna an kai mai theih tur lak atang a venna tha ber ani.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			- Tui an in tur chhawpna tur tha /lian tha tak leh tui thianghlim tak pek tur ani. - Chaw a hmuar/thing pek loh tur ani a, an chaw eitur thlak sak thut loh tur ani.
	Preventive measures	0-3rd week	Ranikhet Disease- an pian atanga ni 1-6 ah F1 vaccine pek tur ani a, chuan a puitlingh chuan R₂B vaccine pek tur ani. - B complex with antibodies
		4th weeks	Coccidiosis- Amprolium or coccidiostat
		4-5th Weeks	Calcium tonic fortified with B₁₂
FISHERY			
	Pond preparation (Dil buatsaih)	3rd-4th weeks	Dil a leitha hman hian sangha chaw kan tih mai planktons insiam nan a tanpui thin.
			- Bawngkek hring 10 tonnes/ha/year vel dil ah hman thin a ni a; bawngkek kumkhat a kan mamawh zat hmunthum a then a hmunkhat hi dil buatsaih nan hman tur ani. A bak zawng hi tui boruak a zirin semdarh a hman thin tur ani. - Single super phosphate hi dil hectare khat zel a zauah kg 250 ang a hman thin tur ani.
			- Leitha kan hman atang a karkhat hnuah tui rawng a tangin sangha chaw inseam that leh thatloh a hriat theih a. Tui rawng eng hring deuh nghalh ah hian sangha chaw planktons te an inseam tha ang a ngaih ani. - Tui nut zawng tehma transparency pawh 30-40 cm vel ani tur ani.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD,
Guwahati)



Expert committee members:

Dr. S.B. Singh	:	Joint Director	basantasinghsoibam@rediffmail.com
Dr. Saurav Saha	:	Scientist (Agril. Physics)	sauravs.saha@gmail.com
Dr. T. Boopathi	:	Scientist (Agril Entomology)	boopathiars@gmail.com
Dr. A. Ratankumar Singh	:	Scientist (Plant Pathology)	ratantplantpatho@gmail.com
Dr. Lungmuana	:	Scientist (Soil Fertility)	lmsingson@gmail.com
Mr. P.L. Lalrinsanga	:	Scientist (Aquaculture)	viensky2@gmail.com
Dr. Dr. V. Dayal	:	Scientist (Horticulture)	Vishambhai5009@gmail.com
Dr. Samuel Lalliansanga	:	Head & Sr. Scientist	samuelpachua10@gmail.com
Mr. Samik Chowdhury	:	Technical Officer	samikchowdhury33@gmail.com
Mr. Diktea chenkuai	:	Project Assistant	dikteachenkuaiboy@gmail.com

Collaborating Department:

Name of the KVK	Programme Coordinator Name and Designation	KVK Email Id	Phone no/ Mobile no
KVK Lunglei	Dr. Lalmuanzovi Head & Sr. Scientist	kvknhathial@gmail.com	9862803750 9436154614
KVK, Kolasib	Mr. Lalrosamga Khiangte Head & Sr. Scientist	kvkkolasib@gmail.com	9436152440
KVK, Serchhip	Mr. K. Laltlanmawia Head & Sr. Scientist	kvkserchhip@gmail.com	9436146115 9615389293
KVK, Champhai	Mrs. Lalrinawmi Renthlei Head & Sr. Scientist	kvkchawzawl@gmail.com	9436159788
KVK, Lawngtlai	Dr. Michel Lallawmkimi Head & Sr. Scientist	kvklawntlai@gmail.com	9436155858
KVK, Saiha	Dr. Vanlalhraia Hnampe Head & Sr. Scientist	kvksaiha@gmail.com	8974656509
KVK, Mamit	Dr. Samuel Lalliansanga Head & Sr. Scientist	kvkmamit@gmail.com	9436147625
KVK, Aizawl	Dr. K. P. Chaudhary Head & Sr. Scientist	Kpchy@rediffmail.com kvkaizawl@rediffmail.com	9436351669



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
 Mizoram Centre, Kolasib- 796081, MIZORAM
(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



District: Mamit

Period: 03 May – 07 May, 2017

Bulletin No: - 697/2017/ Bulletin/English

Date of issue: 02nd May, 2017

Parameters	03.05.2017	04.05.2017	05.05.2017	06.05.2017	07.05.2017
Rainfall (mm)	13	0	0	10	3
Max Temp (°C)	31	32	32	32	32
Min Temp (°C)	22	22	22	22	22
Cloud Coverage	Partially clear	Partially clear	Partially clear	Partially clear	Partially clear
Max RH (%)	98	92	89	92	96
Min RH (%)	45	45	42	36	42
Wind Speed (Kmph)	4	4	6	6	6
*Wind Direction	S-E	S-E	S-E	S-E	S

**Northerly- N, North-Easterly- N-E, Easterly- E, South-Easterly- S-E,
 Southerly- S, South-Westerly- S-W, Westerly- W, North-westerly- N-W.**

STATUS OF MONSOON- June 1-30, 2016 (Percent of deviation from normal in parenthesis)

Aizawl- 384.87mm (430.2mm)	Champhai- 105.48mm (359.89mm)	Saiha- 307.40 mm (507.7mm)	Kolasib- 236.00mm (428.1mm)
Lawngtlai-291.20mm (453.1mm)	Lunglei-326.00mm (465.14mm)	Mamit-204.87mm (442.80mm)	Serchhip-411.72mm (259.62mm)

Weather summary of the past three days

Maximum Tem. (°C):29-31°C
Minimum Tem. (°C):18-21°C
Maximum RH (%):97-99%
Minimum RH (%):52-85%
Wind Direction: southeasterly
Cloud cover: Mainly clear
Wind speed: 2-3 km/hr


Rainfall: 15.8 mm

Weather forecast valid from 03rd May, 2017 To 07thMay, 2017.

There are chances of moderate to light rainfall during the next 3 days. The maximum and minimum temperatures for the next 5 days may range for 31-32°C and 19-21°C. Maximum relative humidity is expected in the range of 89-98% and minimum may from 36-45%. Wind direction would be southeasterly to southerly with the wind speed of 4-6 km per hour. Partially clear will prevail during the next five days.

Weekly cumulative rainfall: 26.0 mm

NDVI for Mizoram



Moderately wet mildly dry/mildly wet conditions



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



Main Crop / Animal / Fisheries	Stage	Cultural practices/ Pest/ Diseases	Agricultural / Horticultural/ animal husbandry advisories
FRUITS CROPS			
KHASI MANDARIN AND ACID LIME	Seedling transplanting stage		- Fruit plant should be planted in a sunny and wind-protected area. - In the citrus belt, trees can be planted at any time, however, spring is the best time for container grown plants. - Standard-size trees should be spaced 12 to 25 feet apart and dwarf trees should be set 6 to 10 feet apart. The exact distance depends on the variety. The bigger the fruit, the farther the distance. - If the soil is not well-drained, plant the trees on a slight mound to prevent water logging. - Mulching should be done with dry grasses near the tree base to conserve soil moisture during winter. - The young fruit plant must be irrigated at weekly interval for better establishment.
BANANA			
STAR FRUIT			
PLUM AND PEACH			
		Gummosis, citrus Canker, Citrus greening, Dieback, Lamon butterfly and leaf minor	Die back- Due to low temperature and humidity disease appearance will more. Use Bordeaux past in tree trunk, twigs and branches protect healthy plant from soil borne disease. Lamon butterfly- Spray monocrotophos @0.04% @1.2 ml/lt of water. Leaf minor- Spray confidor 0.05% (0.5 ml/lit of water) at each flush emergence. Citrus Canker- Apply bacterimycin @0.6 g/lt of water.
Passion Fruit	Transplanting stage		- High yielding mother vine with good quality fruits and free of virus diseases should be selected to provide cuttings. - A cutting should contain at least 3 buds and must be planted in sand beds.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			- Immediately after planting these should be kept inside a high humid chamber made out of bamboo and polythene. Grafting: - The root stock of yellow Passion fruit is planted in polythene sleeves and the section from Rahangala hybrid is grafted using wedge or approach method of grafting.
Pineapple	Flowering stage		- Apply flowering inducing chemical (Ethrel 10 PPM+2% urea+0.04% Sodium Carbonate) should be applied in the heart of the plant. In evening and only when plants have at least 32 leaves. - The flowering emergence will come out after 55-60 days after chemical spraying. - Apply split doses of fertilizer @ 60:50:60 g per plant. - Remove all unwanted leaves, branches and weed near to the plant.
PLANTATION CROP			
COFFEE	Blooming stage		- If day temperature and prolong dry spell occur it lead to Floral abnormalities like "Star Flower" in Arabica and "Pink Flower" in Robusta. - Irrigation of plants at alternate day's interval, 6 weeks before harvesting improves fruit retention and fruit development, and reduces fruit cracking and improves the fruit quality. - Mulching should be done with dry grasses near the tree base to conserve soil moisture during winter. - The young fruit plant must be irrigated at weekly interval for better establishment. - Foliar application of Mepiquat chloride @ 1000 PPM concentration or 0.75% SSP @ 1.5 g per 200 lt of water 15 days




GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



Rubber	Nursery stage		interval. ✚ Clearing operation may be done during the month of February to April. ✚ Make fire line to protect the young tree and seedlings. ✚ 10-12 kg of well rotten organic manure and 225 gm rock phosphate should be apply at time of planting to each pit as basal dose application.
CEREALS AND PULSE CROPS			
Pre Kharif Rice	Nursery stage		✚ Use only Well filled and healthy seeds. ✚ Put the seed in 2.5% salt solution i.e 250 g of common salt in 10 lts of water. ✚ Seed treated with Bavistin 50 WP @ 0.1% (2 g/ltr) solution. ✚ Mulching is requiring for better germination in nursery.
Jhum Rice	Sowing stage		✚ Sowing in ridge and furrow. ✚ Mulching with grass for better germination. ✚ Intercropping with greengram and bunchy type frenchbean.
Maize (Jhum)	Sowing stage		✚ Seed should be treated with Thiram @4 g/kg seed. ✚ Use optimum seed rate (20-25 kg/ha) for desire plant population. ✚ Apply well decomposed FYM/pig manure @ 5-10 t/ha along with 80:60:40 kg N, P₂O₅ and K₂O/ha incorporate with soil before sowing. Half nitrogen dose will use at the time of sowing and remaining 25% after one month and 25% at flowering stage.
Rabi Maize	vegetative stage		✚ Light irrigation on every week may be given for better establishment and smooth growth. ✚ Earthing up soil near to plant for better support. ✚ Maize rust disease will prevail due to high relative humidity with low temperature. Apply Mancozeb Kg/ha



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			for effective control. Remove the alternate host <i>Oxalis comiculata</i>.
VEGETABLE CROP			
Cowpea	Sowing stage	KOLASIB	- Plough the field properly, at least 2-3 times. - Mix fertilizer with FYM 50:60:60Kg/ha. - Sow 2-3 seed per whole. - Spacing should be 30 X 20 cm.
Okra	Sowing stage	WEEDING AND light irrigation in nursery bed. Provide irrigation in transplanted okra field.	- Plough the field with the help of spade. - Sow 2 seed 45 X 45 cm spacing. - Before sowing seed provide one or two irrigation. - Provide fertilizer @ 120: 60: 60 Kg/ha
Ginger and turmeric	Sowing stage	SERCHHIP	- Rhizome should be treated with Thiram @4 g/kg seed. - Use optimum seed rate (50-60 kg/ha) for desire plant population. - Apply well decomposed FYM/ pig manure @ 10-20 t/ha along with 120:80:60 kg N, P₂O₅ and K₂O/ha incorporate with soil before sowing. Half nitrogen dose will use at the time of sowing and remaining 25% after one month and 25% at flowering stage.
Colocasia	Sowing stage	LUNGLEI	- Planting is done well prepared land or pits filled up with FYM (12-15) t/ha - Sprouted corms or cormels are planted 5-7 deep at a spacing of 40-50 cm between and within rows in the pits. - Inorganic fertilizer like Urea, SSP and MOP @ 220: 375: 134 kg.
ANIMAL HUSBANDRY			
Pig	All stages	LAWNGTIA SAIHA	- As the weather gets colder, your pigs' energy requirement will increase, as they need more energy to keep warm. - Regularly monitor their level of 'fitness'



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			and increase their feed intake to maintain. ✚ Fish oils are excellent for providing slow-release energy with the added advantage of a high level of omega-3.
		Porcine Reproductive Respiratory Syndrome (PRRS).	1. Culling of positive pigs or piglets.
	Adult stage	Swine fever.	2. Vaccination of pigs with SF vaccines at 2 months and yearly interval/6 month interval
Cattle	All age group		Due to prolong dry spell there is a shortage of green grass in the field. For balanced diet and nutrition to your cattle, provide urea molasses treated paddy straw.
	All age group	Foot and Mouth Disease (FMD)	FMD vaccine at 16 week and repeat every 6 month.
	Young stage	Black Quarter (BQ)	- Black Quarter Vaccine (BQV). ✦ Primary vaccination 6 month or above ✦ Revaccination annually
Poultry	Litter management		✚ Birds require adequate space, sufficient feed to meet their nutritional requirements and an adequate supply of good-quality water. ✚ Good management and sanitation are the best ways to avoid infectious disease in poultry. ✚ Provide ample quantity of clean drinking water. ✚ Avoid feeding of mouldy feed. Don't make sudden changes in feed
	Preventive measures	0-3rd week	✚ Ranikhet Disease- F1 vaccine at (1-6) days of birth and R₂B vaccine for adult birds. ✚ B complex with antibodies
		4th weeks	✚ Coccidiosis- Amprolium or coccidiostat



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



		4-5 th Weeks	✚ Calcium tonic fortified with B ₁₂
FISHERY			
	Pond preparation	3 rd -4 th weeks	✚ Application of fertilizers/manure helps in development of plankton which serve as natural feed for the fishes.
		KOLASIB	✚ Raw cowdung should be applied in the pond at the rate of 10 tonnes/ha/year. One third of the total dose should be applied initially and the rest may be applied in a spilt doses. ✚ Single super phosphate should also be applied at the rate of 250 kg/ha in the pond.
		AIZAWL	✚ After one week of application development of planktons could be observed in the pond depending on the colour of the water. Yellowish green colour is an indicator of the good plankton development. ✚ Transparency of the water needs to be maintained at 30-40 cm.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD,
Guwahati)



Expert committee members:

Dr. S.B. Singh	:	Joint Director	basantasinghsoibam@rediffmail.com
Dr. Saurav Saha	:	Scientist (Agril. Physics)	sauravs.saha@gmail.com
Dr. T. Boopathi	:	Scientist (Agril Entomology)	boopathiars@gmail.com
Dr. A. Ratankumar Singh	:	Scientist (Plant Pathology)	ratantplantpatho@gmail.com
Dr. Lungmuana	:	Scientist (Soil Fertility)	lmsingson@gmail.com
Mr. P.L. Lalrinsanga	:	Scientist (Aquaculture)	viensky2@gmail.com
Dr. Dr. V. Dayal	:	Scientist (Horticulture)	Vishambhai5009@gmail.com
Dr. Samuel Lalliansanga	:	Head & Sr. Scientist	samuelpachua10@gmail.com
Mr. Samik Chowdhury	:	Technical Officer	samikchowdhury33@gmail.com
Mr. Diktea chenkuai	:	Project Assistant	dikteachenkuaiboy@gmail.com

Collaborating Department:

Name of the KVK	Programme Coordinator Name and Designation	KVK Email Id	Phone no/ Mobile no
KVK Lunglei	Dr. Lalmuanzovi Head & Sr. Scientist	kvknhathial@gmail.com	9862803750 9436154614
KVK, Kolasib	Mr. Lalrosamga Khiangte Head & Sr. Scientist	kvkkolasib@gmail.com	9436152440
KVK, Serchhip	Mr. K. Laltlanmawia Head & Sr. Scientist	kvkserchhip@gmail.com	9436146115 9615389293
KVK, Champhai	Mrs. Lalrinawmi Renthlei Head & Sr. Scientist	kvkchawzawl@gmail.com	9436159788
KVK, Lawngtlai	Dr. Michel Lallawmkimi Head & Sr. Scientist	kvklawntlai@gmail.com	9436155858
KVK, Saiha	Dr. Vanlalhraia Hnampe Head & Sr. Scientist	kvksaiha@gmail.com	8974656509
KVK, Mamit	Dr. Samuel Lalliansanga Head & Sr. Scientist	kvkmamit@gmail.com	9436147625
KVK, Aizawl	Dr. K. P. Chaudhary Head & Sr. Scientist	Kpchy@rediffmail.com kvkaizawl@rediffmail.com	9436351669



GRAMIN KRISHI MAUSAM SEWA

ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM
(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



District: Mamit

Period: 03 May – 07 May, 2017

Bulletin No: - 697/2017/ Bulletin/Mizo

Date of issue: 02nd May, 2017

Parameters	03.05.2017	04.05.2017	05.05.2017	06.05.2017	07.05.2017
Rainfall (mm)	13	0	0	10	3
Max Temp (°C)	31	32	32	32	32
Min Temp (°C)	22	22	22	22	22
Cloud Coverage	Partially clear	Partially clear	Partially clear	Partially clear	Partially clear
Max RH (%)	98	92	89	92	96
Min RH (%)	45	45	42	36	42
Wind Speed (Kmph)	4	4	6	6	6
*Wind Direction	S-E	S-E	S-E	S-E	S

**Northerly- N, North-Easterly- N-E, Easterly- E, South-Easterly- S-E,
Southerly- S, South-Westerly- S-W, Westerly- W, North-westerly- N-W.**

STATUS OF MONSOON- June 1-30, 2016 (Percent of deviation from normal in parenthesis)

Aizawl- 384.87mm (430.2mm)	Champhai- 105.48mm (359.89mm)	Saiha- 307.40 mm (507.7mm)	Kolasib- 236.00mm (428.1mm)
Lawngtlai-291.20mm (453.1mm)	Lunglei-326.00mm (465.14mm)	Mamit-204.87mm (442.80mm)	Serchhip-411.72mm (259.62mm)

Weather summary of the past three days

Maximum Tem. (°C):29-31°C
Minimum Tem. (°C):18-21°C
Maximum RH (%):97-99%
Minimum RH (%):52-85%
Wind Direction: southeasterly
Cloud cover: Mainly clear
Wind speed: 2-3 km/hr


Rainfall: 15.8 mm

**03rd May– 07th May, 2017 chhunga sik leh sa
dinhmun tur tlangpui**

Tun ni 3 chhung lo awm turah hian ruahtui tla miahlo tura beisei a ni. Khua a lum lai berin 31-32°C a ni ang a. A vawh lai ber in 19-21°C ni tura beisei a ni. RH san lai berin 89-98% leh a hniam lai berin 36-45% ni tur a rin niin. Thli hi darkar khatah 4-6 km vela chakin chhaklam awi zawngin a tleh rin a ni. A tlangpuiin tun ni nga chung hian khawthiang tak hmuh beisei a ni.

Weekly cumulative rainfall: 26.0mm

NDVI for Mizoram



Moderately wet mildly dry/mildly wet conditions



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



Main Crop / Animal / Fisheries	Stage	Cultural practices/ Pest/ Diseases	Agricultural / Horticultural/ animal husbandry advisories
FRUITS CROPS			
KHASI MANDARIN AND ACID LIME	A kui atanga a seng hun		- Thlasik laia thlai bul khoro lutuk tur vennis chuan hnim hnah hring tlai bul velah dahkhawm tur ani. - Thlai naupang deuah chuan chawlh kar tin a tui pek thin tur ani. - Leia tha mamawh tawh a hmuh theihna turin a hmunhma a hnim awm te thlawhfai thin tur ani. - A seng hma kar 6 chhung chu tui tha taka pek hian a rah tla tur chelh nan leh a rah than that nan te leh a rah keh tur lakah t a veng thei ani.
BANANA			
STAR FRUIT			
PLUM AND PEACH			
		Gummosis, citrus canker, citrus greening and Dieback	Temperture hniam lutuk leh hnawng vang hian natna a a tam duh a . Soil bome natna laka vennis Bordeaux past hi thing zar leh a trangah te hnawih tur ani.
		Fruit fly	Huan zau takah chuan a par tan tirh leh a rah tan tirin chawlhkar hnih chhung chu heng te hian enkawl tur ani: carbaryl 0.2 percent emaw malathion 0.15 percent suspension containing sugar or jeggery at 10 g/l.
PLANTATION CROP			
COFFEE	All stages		Nursery stage - Thlai chi thlak hma in <i>Azospirillum</i> leh <i>Phosphobacterium</i> a enkawl tur ani. - A chi hi December – January ah hmun zawl/rualrem 1.5 - 2.5 cm a in hlatin tlar mumal tak siam in chin tur ani. - Chuan a chi chu lei tlem te a chhilh a buhpawla khuh tur ani. - Nitin tui pek tur ani a, a sat lutuka loh nan niin a chhun loh nan zar hlih tur ani. - Ni 45 hnu velah a tiak thin a,chu chu bag ah an sawn chhuak leh thin ani.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			Harvesting Stage - Coffe rah hmin hi thlasik lain an seng thin a ni. - A rah hmin tha lo ho chu nuai sawm hmain an thliar hrang leh vek thin ani.
		Coffee Berry borer	- A hun takah leh fimkhur taka seng tur ani. - Hmaih neih nuaih loh tur ani. - Sneg hmaih te lakkhawma paih vek tur ani. - Seng bang zawng zawng chu uluka taka paih fai vek tur ani. - A thlai vennis a rah tlai ho chu paih vek tur ani. - Hmun dam lutukah dah loh tur. - Boruak tha taka a hmuh theih nan leh a rawn chawr no theih nan thlai chu uluk tak a hlawi tur ani. - Chuan hei hian kah tur ani. Quinalphos 25 EC @ 340 ml/200 lit emaw lamda cyhalothrin 5 EC 120 – 160 ml / 200 lit. - In leh loa sawngbawl a a ro dan tur tawkah chuan: Arabica / robusta parchment 10 %, Arabica cherry 10.5 % leh robusta cherry 11.0 %.
		Coffee Rust	- Natna hrik in a khawih tawh a hnah leh a dang te paihpaih vek tur ani. - Bordeaux mixture 0.5% in February - March (Pre-bloom) a kah phawt a, Oxycarboxin 0.03% in May - June (Pre-monsoon) ah kah chhunzawm tur ani.
CEREALS AND PULSE CROPS			
Rabi Maize	A chin hun		- Vaimim chinna tur atan lei kan let phut darh anga. - Hei hian a rawn to chhuah na tur atan a pui dawn a ni. - A chi chu kan lei leh saah chuan kan dah ang. - A chi chu Thiram @4 g/kg seed hian ak sawngbawl hmasa anga. 20-25 kg/ha vel a chi thlak hi a tawk vel viau ani. - A chi tuh hma in lei leh FYM @5-10 t/ha leh 80:60:40 kg N, P₂O₅ leh



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			K ₂ O/ha pawlh chu hman phawt tur a ni. Nitrogen dose chanve chu a chi tuh hunlaia hman tur a ni a, tichuan a bang 25% chu thla khat hnu ah ani ang a adang leh 25% chu a par hunah hman tur a ni.
Soybean, pea, lentil toria, breen gram and black gram cultivation in rice fellow	All stage	Zero tillage	- A than a that theih nan nikhat danah tui pek thin tur ani. - Lei rih vur hian thlai kung te a veng ve ani. - Thlasik lai a lei khoro lutuk tur ven nan a chungah hnim leh thildanga khuh tur ani.
Potato	Sowing stage		- Muangchang loving alu chin na tur chu buatsaih vat tur ani. - Hei hian a than hun lain natna hrikin lakah a veng dawn ani. - Lei leh hmain a hmun hma chu fai taka thlawh hmasak tur ani. - A chi thlak hma in a chi chu en fiah hmasak tur ani. - A than a that theih nan nikhat danah tui pek thin tur ani.
VEGETABLE CROP			
Tomato	Bacterial Blight disease		- Tomato bikah chuan sik leh sa hi natna an kaina tlang lawn ber ani . - Hmun hnawng leh ni hmu lo lutuk hmunah chuan natna an kai hma bik ani. - Tomato hi a uai a, a thih mai loh nan Ridomil emaw Indofil emaw Mancozeb @ 2 gm hi tui liter 1 ah pawlh a kah tur ani .
Early Cole crop	Black spot disease		- A than a that theih nan nikhat danah tui pek thin tur ani. - Thlai bul vawn hnawn nana thlai bula hnim ring vawm khawm hi tui pek zawhah dah tur ani. - Zikhlum lam chi ah chuan sik leh sa vangin a hnah ah thil dum rawn



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			awm thin a , hei hi natna tlanglawn ber ani. Thlai hna lam chi leh zikhlum lam chi reng reng enkawl nan Mancozeb @ 2gm ah tui leter 1 pawlha kah tur ani.
Onion and capsicum	Nursery stage	Poly house	- A than a that theih nan nikhat danah tui pek thin tur ani. - Thlai bul vawn hnawn nana thlai bula hnim ring vawm khawm hi tui pek zawhah dah tur ani. - Thlai chhina hmun (nursery) hi hnim a to loh nan Pendimethalin @ 3.5ml hi tui liter 1 zelah pawlh a kah hi a tha hle ani.
		Phytophthora blight	- A chi ven that nan thiram 3g/kg seed emaw Trichoderma viride 4g+ metalaxyl 4g (Apron)/ kg seed hi a tha hle ani - Hneh taka 1% Bordeaux chawhpawlh emaw 2 g captan emaw 3 copper oxychloride a tui liter 1 hi 10-15 DAS a pek hi a tha hle ani.
French bean	Sowing stage		- Tui pek a hnihnah hringa khuh tur ani a. than a that theih nan tui pek hma in lei rin pan hmasak tur ani. - A than duna theih nan leh hnim to loh na turin a kung bulah lei vur chhoh zel tur ani.
Carrot and radish	Sowing stage		- A than a that theih nan nikhat danah tui pek thin tur ani. - Tui pek hnuah thlai bul vawn hnawn na tur siam tur ani. - Zikhlum lam chi ah chuan sik leh sa vangin a hnah ah thil dum a rawn awm thina, hei hi natna tlanglawn ber ani. - Thlai hna lam chi leh zikhlum lam chi reng reng enkawl nan Mancozeb @ 2gm ah tui leter 1 pawlha kah tur ani.






GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



ANIMAL HUSBANDRY			
Pig	All stages		- Khua a vawh hian vawk hian an mahni in tih lumna tur atan chakna an mamawhna a sang bik ani. - An hriselna that leh that loh enfiah renga, a chaw ei tur tlem tlema tih tam hret hret tur ani.. - Sangha tel ah hian omega-3 hi atam em a vangin an chakna muangchanga a in siam chhoh zel theih nan a tha hle ani.
			Porcine Reproductive Respiratory Syndrome (PRRS). 1. Vawknote emaw vawk lak hran.
	Adult stage		Swine fever. 2. SF vaccines hi thla 2 hnua pek tur ani a, chumi hnuah chuan kumtin thlaruk danah pek chhonzawm tur ani.
Cattle	All age group		Hun rei tak khua a ro avanga hnim hnah hring peh tur a awm loh laia bawngin an chaw ei in buk tawk tur leh an taksa tana mamawh tur atan buh kung urea molasses hmanga sawngbawl pek tur ani.
	All age group		Kar 16 hnuah FMD vaccine pek a, chuan thla tin thla 6 chhung chhonzawm tur ani.
	Young stage		- Black Quarter Vaccine (BQV). - Vaccinne hmasa ber hi thla 6 ah emaw a hnu lamah pek tur. - Chumi hnuah chuan Vaccine hi kum tin pek tur ani.
Poultry	Litter management		- Ar te hian hmun thawl nuam tawk, chaw tha an mamawh tawk leh tui thianghlim an mamawh tawk an hmu tur ani a. - An hriselna atan enkawltha tha tawk tak pek hian natna an kai mai theih tur lak atang a venna tha ber ani.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			- Tui an in tur chhawpna tur tha /lian tha tak leh tui thianghlim tak pek tur ani. - Chaw a hmuar/thing pek loh tur ani a, an chaw eitur thlak sak thut loh tur ani.
	Preventive measures	0-3rd week	Ranikhet Disease- an pian atanga ni 1-6 ah F1 vaccine pek tur ani a, chuan a puitlingh chuan R₂B vaccine pek tur ani. - B complex with antibodies
		4th weeks	Coccidiosis- Amprolium or coccidiostat
		4-5th Weeks	Calcium tonic fortified with B₁₂
FISHERY			
	Pond preparation (Dil buatsaih)	3rd-4th weeks	Dil a leitha hman hian sangha chaw kan tih mai planktons insiam nan a tanpui thin.
			- Bawngkek hring 10 tonnes/ha/year vel dil ah hman thin a ni a; bawngkek kumkhat a kan mamawh zat hmunthum a then a hmunkhat hi dil buatsaih nan hman tur ani. A bak zawng hi tui boruak a zirin semdarh a hman thin tur ani. - Single super phosphate hi dil hectare khat zel a zauah kg 250 ang a hman thin tur ani.
			- Leitha kan hman atang a karkhat hnuah tui rawng a tangin sangha chaw inseam that leh thatloh a hriat theih a. Tui rawng eng hring deuh nghalh ah hian sangha chaw planktons te an inseam tha ang a ngaih ani. - Tui nut zawng tehma transparency pawh 30-40 cm vel ani tur ani.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD,
Guwahati)



Expert committee members:

Dr. S.B. Singh	:	Joint Director	basantasinghsoibam@rediffmail.com
Dr. Saurav Saha	:	Scientist (Agril. Physics)	sauravs.saha@gmail.com
Dr. T. Boopathi	:	Scientist (Agril Entomology)	boopathiars@gmail.com
Dr. A. Ratankumar Singh	:	Scientist (Plant Pathology)	ratantplantpatho@gmail.com
Dr. Lungmuana	:	Scientist (Soil Fertility)	lmsingson@gmail.com
Mr. P.L. Lalrinsanga	:	Scientist (Aquaculture)	viensky2@gmail.com
Dr. Dr. V. Dayal	:	Scientist (Horticulture)	Vishambhai5009@gmail.com
Dr. Samuel Lalliansanga	:	Head & Sr. Scientist	samuelpachua10@gmail.com
Mr. Samik Chowdhury	:	Technical Officer	samikchowdhury33@gmail.com
Mr. Diktea chenkual	:	Project Assistant	dikteachenkualboy@gmail.com

Collaborating Department:

Name of the KVK	Programme Coordinator Name and Designation	KVK Email Id	Phone no/ Mobile no
KVK Lunglei	Dr. Lalmuanzovi Head & Sr. Scientist	kvknhathial@gmail.com	9862803750 9436154614
KVK, Kolasib	Mr. Lalrosamga Khiangte Head & Sr. Scientist	kvkkolasib@gmail.com	9436152440
KVK, Serchhip	Mr. K. Laltlanmawia Head & Sr. Scientist	kvkserchhip@gmail.com	9436146115 9615389293
KVK, Champhai	Mrs. Lalrinawmi Renthlei Head & Sr. Scientist	kvkchawzawl@gmail.com	9436159788
KVK, Lawngtlai	Dr. Michel Lallawmkimi Head & Sr. Scientist	kvklawntlai@gmail.com	9436155858
KVK, Saiha	Dr. Vanlalhruaia Hnampe Head & Sr. Scientist	kvksaiha@gmail.com	8974656509
KVK, Mamit	Dr. Samuel Lalliansanga Head & Sr. Scientist	kvkmamit@gmail.com	9436147625
KVK, Aizawl	Dr. K. P. Chaudhary Head & Sr. Scientist	Kpchy@rediffmail.com kvkaizawl@rediffmail.com	9436351669



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Mizoram Centre, Kolasib- 796081, MIZORAM
(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



District: Saiha

Period: 03 May – 07 May, 2017

Bulletin No: - 697/2017/ Bulletin/English

Date of issue: 02nd May, 2017

Parameters	03.05.2017	04.05.2017	05.05.2017	06.05.2017	07.05.2017
Rainfall (mm)	0	0	0	0	0
Max Temp (°C)	32	33	35	34	33
Min Temp (°C)	20	22	20	20	20
Cloud Coverage	Mainly clear	Mainly clear	Mainly clear	Mainly clear	Mainly clear
Max RH (%)	98	76	84	81	87
Min RH (%)	30	28	27	26	31
Wind Speed (Kmph)	4	4	2	3	2
*Wind Direction	N-E	E	E	S-E	E

**Northerly- N, North-Easterly- N-E, Easterly- E, South-Easterly- S-E,
Southerly- S, South-Westerly- S-W, Westerly- W, North-westerly- N-W.**

STATUS OF MONSOON- June 1-30, 2016 (Percent of deviation from normal in parenthesis)

Aizawl- 384.87mm (430.2mm)	Champhai- 105.48mm (359.89mm)	Saiha- 307.40 mm (507.7mm)	Kolasib- 236.00mm (428.1mm)
Lawngtlai-291.20mm (453.1mm)	Lunglei-326.00mm (465.14mm)	Mamit-204.87mm (442.80mm)	Serchhip-411.72mm (259.62mm)

Weather summary of the past three days

Maximum Tem. (°C):24-28°C
Minimum Tem. (°C):16-18°C
Maximum RH (%):92-98%
Minimum RH (%):48-89%
Wind Direction: Southeasterly
Cloud cover: Mainly clear
Wind Speed: 2-4 km/hr


Rainfall: 00.0 mm

Weather forecast valid from 03rd May, 2017 To 07thMay, 2017.

There are no chances of rainfall during the next 5 days. The maximum and minimum temperatures for the next 5 days may range for 32-35°C and 20-22°C. Maximum relative humidity is expected in the range of 76-98% and minimum may from 26-31%. Wind direction would northeaster to easterly to southeasterly and easterly with the wind speed of 2-4 km per hour. Mainly clear sky will prevail during the next five days.

Weekly cumulative rainfall: 00.0 mm

NDVI for Mizoram



Moderately wet mildly dry/mildly wet conditions



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



Main Crop / Animal / Fisheries	Stage	Cultural practices/ Pest/ Diseases	Agricultural / Horticultural/ animal husbandry advisories
FRUITS CROPS			
KHASI MANDARIN AND ACID LIME	Seedling transplanting stage		- Fruit plant should be planted in a sunny and wind-protected area. - In the citrus belt, trees can be planted at any time, however, spring is the best time for container grown plants. - Standard-size trees should be spaced 12 to 25 feet apart and dwarf trees should be set 6 to 10 feet apart. The exact distance depends on the variety. The bigger the fruit, the farther the distance. - If the soil is not well-drained, plant the trees on a slight mound to prevent water logging. - Mulching should be done with dry grasses near the tree base to conserve soil moisture during winter. - The young fruit plant must be irrigated at weekly interval for better establishment.
BANANA			
STAR FRUIT			
PLUM AND PEACH			
		Gummosis, citrus Canker, Citrus greening, Dieback, Lamon butterfly and leaf minor	Die back- Due to low temperature and humidity disease appearance will more. Use Bordeaux past in tree trunk, twigs and branches protect healthy plant from soil borne disease. Lamon butterfly- Spray monocrotophos @0.04% @1.2 ml/lt of water. Leaf minor- Spray confidor 0.05% (0.5 ml/lit of water) at each flush emergence. Citrus Canker- Apply bacterimycin @0.6 g/lt of water.
Passion Fruit	Transplanting stage		- High yielding mother vine with good quality fruits and free of virus diseases should be selected to provide cuttings. - A cutting should contain at least 3 buds and must be planted in sand beds.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			- Immediately after planting these should be kept inside a high humid chamber made out of bamboo and polythene. Grafting: - The root stock of yellow Passion fruit is planted in polythene sleeves and the section from Rahangala hybrid is grafted using wedge or approach method of grafting.
Pineapple	Flowering stage		- Apply flowering inducing chemical (Ethrel 10 PPM+2% urea+0.04% Sodium Carbonate) should be applied in the heart of the plant. In evening and only when plants have at least 32 leaves. - The flowering emergence will come out after 55-60 days after chemical spraying. - Apply split doses of fertilizer @ 60:50:60 g per plant. - Remove all unwanted leaves, branches and weed near to the plant.
PLANTATION CROP			
COFFEE	Blooming stage		- If day temperature and prolong dry spell occur it lead to Floral abnormalities like "Star Flower" in Arabica and "Pink Flower" in Robusta. - Irrigation of plants at alternate day's interval, 6 weeks before harvesting improves fruit retention and fruit development, and reduces fruit cracking and improves the fruit quality. - Mulching should be done with dry grasses near the tree base to conserve soil moisture during winter. - The young fruit plant must be irrigated at weekly interval for better establishment. - Foliar application of Mepiquat chloride @ 1000 PPM concentration or 0.75% SSP @ 1.5 g per 200 lt of water 15 days




GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



Rubber	Nursery stage		interval. ✚ Clearing operation may be done during the month of February to April. ✚ Make fire line to protect the young tree and seedlings. ✚ 10-12 kg of well rotten organic manure and 225 gm rock phosphate should be apply at time of planting to each pit as basal dose application.
CEREALS AND PULSE CROPS			
Pre Kharif Rice	Nursery stage		✚ Use only Well filled and healthy seeds. ✚ Put the seed in 2.5% salt solution i.e 250 g of common salt in 10 lts of water. ✚ Seed treated with Bavistin 50 WP @ 0.1% (2 g/ltr) solution. ✚ Mulching is requiring for better germination in nursery.
Jhum Rice	Sowing stage		✚ Sowing in ridge and furrow. ✚ Mulching with grass for better germination. ✚ Intercropping with greengram and bunchy type frenchbean.
Maize (Jhum)	Sowing stage		✚ Seed should be treated with Thiram @4 g/kg seed. ✚ Use optimum seed rate (20-25 kg/ha) for desire plant population. ✚ Apply well decomposed FYM/pig manure @ 5-10 t/ha along with 80:60:40 kg N, P₂O₅ and K₂O/ha incorporate with soil before sowing. Half nitrogen dose will use at the time of sowing and remaining 25% after one month and 25% at flowering stage.
Rabi Maize	vegetative stage		✚ Light irrigation on every week may be given for better establishment and smooth growth. ✚ Earthing up soil near to plant for better support. ✚ Maize rust disease will prevail due to high relative humidity with low temperature. Apply Mancozeb Kg/ha



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			for effective control. Remove the alternate host <i>Oxalis comiculata</i>.
VEGETABLE CROP			
Cowpea	Sowing stage	KOLASIB	- Plough the field properly, at least 2-3 times. - Mix fertilizer with FYM 50:60:60Kg/ha. - Sow 2-3 seed per whole. - Spacing should be 30 X 20 cm.
Okra	Sowing stage	WEEDING AND light irrigation in nursery bed. Provide irrigation in transplanted okra field.	- Plough the field with the help of spade. - Sow 2 seed 45 X 45 cm spacing. - Before sowing seed provide one or two irrigation. - Provide fertilizer @ 120: 60: 60 Kg/ha
Ginger and turmeric	Sowing stage	SERCHHIP	- Rhizome should be treated with Thiram @4 g/kg seed. - Use optimum seed rate (50-60 kg/ha) for desire plant population. - Apply well decomposed FYM/ pig manure @ 10-20 t/ha along with 120:80:60 kg N, P₂O₅ and K₂O/ha incorporate with soil before sowing. Half nitrogen dose will use at the time of sowing and remaining 25% after one month and 25% at flowering stage.
Colocasia	Sowing stage	LUNGLEI	- Planting is done well prepared land or pits filled up with FYM (12-15) t/ha - Sprouted corms or cormels are planted 5-7 deep at a spacing of 40-50 cm between and within rows in the pits. - Inorganic fertilizer like Urea, SSP and MOP @ 220: 375: 134 kg.
ANIMAL HUSBANDRY			
Pig	All stages	LAWNGTELA SAIHA	- As the weather gets colder, your pigs' energy requirement will increase, as they need more energy to keep warm. - Regularly monitor their level of 'fitness'



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			and increase their feed intake to maintain. ✚ Fish oils are excellent for providing slow-release energy with the added advantage of a high level of omega-3.
		Porcine Reproductive Respiratory Syndrome (PRRS).	1. Culling of positive pigs or piglets.
	Adult stage	Swine fever.	2. Vaccination of pigs with SF vaccines at 2 months and yearly interval/6 month interval
Cattle	All age group		Due to prolong dry spell there is a shortage of green grass in the field. For balanced diet and nutrition to your cattle, provide urea molasses treated paddy straw.
	All age group	Foot and Mouth Disease (FMD)	FMD vaccine at 16 week and repeat every 6 month.
	Young stage	Black Quarter (BQ)	- Black Quarter Vaccine (BQV). ✦ Primary vaccination 6 month or above ✦ Revaccination annually
Poultry	Litter management		✚ Birds require adequate space, sufficient feed to meet their nutritional requirements and an adequate supply of good-quality water. ✚ Good management and sanitation are the best ways to avoid infectious disease in poultry. ✚ Provide ample quantity of clean drinking water. ✚ Avoid feeding of mouldy feed. Don't make sudden changes in feed
	Preventive measures	0-3rd week	✚ Ranikhet Disease- F1 vaccine at (1-6) days of birth and R₂B vaccine for adult birds. ✚ B complex with antibodies
		4th weeks	✚ Coccidiosis- Amprolium or coccidiostat



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



		4-5 th Weeks	✚ Calcium tonic fortified with B ₁₂
FISHERY			
	Pond preparation	3 rd -4 th weeks	✚ Application of fertilizers/manure helps in development of plankton which serve as natural feed for the fishes.
		KOLASIB	✚ Raw cowdung should be applied in the pond at the rate of 10 tonnes/ha/year. One third of the total dose should be applied initially and the rest may be applied in a spilt doses. ✚ Single super phosphate should also be applied at the rate of 250 kg/ha in the pond.
		AIZAWL	✚ After one week of application development of planktons could be observed in the pond depending on the colour of the water. Yellowish green colour is an indicator of the good plankton development. ✚ Transparency of the water needs to be maintained at 30-40 cm.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD,
Guwahati)



Expert committee members:

Dr. S.B. Singh	:	Joint Director	basantasinghsoibam@rediffmail.com
Dr. Saurav Saha	:	Scientist (Agril. Physics)	sauravs.saha@gmail.com
Dr. T. Boopathi	:	Scientist (Agril Entomology)	boopathiars@gmail.com
Dr. A. Ratankumar Singh	:	Scientist (Plant Pathology)	ratantplantpatho@gmail.com
Dr. Lungmuana	:	Scientist (Soil Fertility)	lmsingson@gmail.com
Mr. P.L. Lalrinsanga	:	Scientist (Aquaculture)	viensky2@gmail.com
Dr. Dr. V. Dayal	:	Scientist (Horticulture)	Vishambhai5009@gmail.com
Dr. Samuel Lalliansanga	:	Head & Sr. Scientist	samuelpachua10@gmail.com
Mr. Samik Chowdhury	:	Technical Officer	samikchowdhury33@gmail.com
Mr. Diktea chenkuai	:	Project Assistant	dikteachenkuaiboy@gmail.com

Collaborating Department:

Name of the KVK	Programme Coordinator Name and Designation	KVK Email Id	Phone no/ Mobile no
KVK Lunglei	Dr. Lalmuanzovi Head & Sr. Scientist	kvknhathial@gmail.com	9862803750 9436154614
KVK, Kolasib	Mr. Lalrosamga Khiangte Head & Sr. Scientist	kvkkolasib@gmail.com	9436152440
KVK, Serchhip	Mr. K. Laltlanmawia Head & Sr. Scientist	kvkserchhip@gmail.com	9436146115 9615389293
KVK, Champhai	Mrs. Lalrinawmi Renthlei Head & Sr. Scientist	kvkchawzawl@gmail.com	9436159788
KVK, Lawngtlai	Dr. Michel Lallawmkimi Head & Sr. Scientist	kvklawntlai@gmail.com	9436155858
KVK, Saiha	Dr. Vanlalhraia Hnampe Head & Sr. Scientist	kvksaiha@gmail.com	8974656509
KVK, Mamit	Dr. Samuel Lalliansanga Head & Sr. Scientist	kvkmamit@gmail.com	9436147625
KVK, Aizawl	Dr. K. P. Chaudhary Head & Sr. Scientist	Kpchy@rediffmail.com kvkaizawl@rediffmail.com	9436351669



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
 Mizoram Centre, Kolasib- 796081, MIZORAM
(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



District: Saiha

Period: 03 May – 07 May, 2017

Bulletin No: - 697/2017/ Bulletin/Mizo

Date of issue: 02nd May, 2017

Parameters	03.05.2017	04.05.2017	05.05.2017	06.05.2017	07.05.2017
Rainfall (mm)	0	0	0	0	0
Max Temp (°C)	32	33	35	34	33
Min Temp (°C)	20	22	20	20	20
Cloud Coverage	Mainly clear	Mainly clear	Mainly clear	Mainly clear	Mainly clear
Max RH (%)	98	76	84	81	87
Min RH (%)	30	28	27	26	31
Wind Speed (Kmph)	4	4	2	3	2
*Wind Direction	N-E	E	E	S-E	E

**Northerly- N, North-Easterly- N-E, Easterly- E, South-Easterly- S-E,
 Southerly- S, South-Westerly- S-W, Westerly- W, North-westerly- N-W.**

STATUS OF MONSOON- June 1-30, 2016 (Percent of deviation from normal in parenthesis)

Aizawl- 384.87mm (430.2mm)	Champhai- 105.48mm (359.89mm)	Saiha- 307.40 mm (507.7mm)	Kolasib- 236.00mm (428.1mm)
Lawngtlai-291.20mm (453.1mm)	Lunglei-326.00mm (465.14mm)	Mamit-204.87mm (442.80mm)	Serchhip-411.72mm (259.62mm)

Weather summary of the past three days

Maximum Tem. (°C):24-28°C
Minimum Tem. (°C):16-18°C
Maximum RH (%):92-98%
Minimum RH (%):48-89%
Wind Direction: Southeasterly
Cloud cover: Mainly clear
Wind Speed: 2-4 km/hr


Rainfall: 00.0 mm

**03rd May– 07th May, 2017 chhunga sik leh sa
 dinhmun tur tlangpui**

Tun ni 5 chhung lo awm turah hian ruahtui tla miahlo tura beisei a ni. Khua a lum lai berin 32-35°C a ni ang a. A vawh lai ber in 20-22°C ni tura beisei a ni. RH san lai berin 76-98% leh a hniam lai berin 26-31% ni tur a rin niin. Thli hi darkar khatah 2-4 km vela chakin chhaklam awi zawngin a tleh rin a ni. A tlangpuiin tun ni nga chung hian khawthiang tak hmuh beisei a ni.

Weekly cumulative rainfall: 00.0mm

NDVI for Mizoram



Moderately wet mildly dry/mildly wet conditions



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



Main Crop / Animal / Fisheries	Stage	Cultural practices/ Pest/ Diseases	Agricultural / Horticultural/ animal husbandry advisories
FRUITS CROPS			
KHASI MANDARIN AND ACID LIME	A kui atanga a seng hun		- Thlasik laia thlai bul khoro lutuk tur vennan chuan hnim hnah hring tlai bul velah dahkhawm tur ani. - Thlai naupang deuah chuan chawlh kar tin a tui pek thin tur ani. - Leia tha mamawh tawh a hmuh theihna turin a hmunhma a hnim awm te thlawhfai thin tur ani. - A seng hma kar 6 chhung chu tui tha taka pek hian a rah tla tur chelh nan leh a rah than that nan te leh a rah keh tur lakah t a veng thei ani.
BANANA			
STAR FRUIT			
PLUM AND PEACH			
		Gummosis, citrus canker, citrus greening and Dieback	Temperture hniam lutuk leh hnawng vang hian natna a a tam duh a . Soil bome natna laka vennan Bordeaux past hi thing zar leh a trangah te hnawih tur ani.
		Fruit fly	Huan zau takah chuan a par tan tirh leh a rah tan tirin chawlhkar hnih chhung chu heng te hian enkawl tur ani: carbaryl 0.2 percent emaw malathion 0.15 percent suspension containing sugar or jeggery at 10 g/l.
PLANTATION CROP			
COFFEE	All stages		Nursery stage - Thlai chi thlak hma in <i>Azospirillum</i> leh <i>Phosphobacterium</i> a enkawl tur ani. - A chi hi December – January ah hmun zawl/rualrem 1.5 - 2.5 cm a in hlatin tlar mumal tak siam in chin tur ani. - Chuan a chi chu lei tlem te a chhilh a buhpawla khuh tur ani. - Nitin tui pek tur ani a, a sat lutuka loh nan niin a chhun loh nan zar hlih tur ani. - Ni 45 hnu velah a tiak thin a,chu chu bag ah an sawn chhuak leh thin ani.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			Harvesting Stage - Coffe rah hmin hi thlasik lain an seng thin a ni. - A rah hmin tha lo ho chu nuai sawm hmain an thliar hrang leh vek thin ani.
		Coffee Berry borer	- A hun takah leh fimkhur taka seng tur ani. - Hmaih neih nuaih loh tur ani. - Sneg hmaih te lakkhawma paih vek tur ani. - Seng bang zawng zawng chu uluka taka paih fai vek tur ani. - A thlai vennan a rah tlai ho chu paih vek tur ani. - Hmun dam lutukah dah loh tur. - Boruak tha taka a hmuh theih nan leh a rawn chawr no theih nan thlai chu uluk tak a hlawi tur ani. - Chuan hei hian kah tur ani. Quinalphos 25 EC @ 340 ml/200 lit emaw lamda cyhalothrin 5 EC 120 – 160 ml / 200 lit. - In leh loa sawngbawl a a ro dan tur tawkah chuan: Arabica / robusta parchment 10 %, Arabica cherry 10.5 % leh robusta cherry 11.0 %.
		Coffee Rust	- Natna hrik in a khawih tawh a hnah leh a dang te paihpaih vek tur ani. - Bordeaux mixture 0.5% in February - March (Pre-bloom) a kah phawt a, Oxycarboxin 0.03% in May - June (Pre-monsoon) ah kah chhunzawm tur ani.
CEREALS AND PULSE CROPS			
Rabi Maize	A chin hun		- Vaimim chinna tur atan lei kan let phut darh anga. - Hei hian a rawn to chhuah na tur atan a pui dawn a ni. - A chi chu kan lei leh saah chuan kan dah ang. - A chi chu Thiram @4 g/kg seed hian ak sawngbawl hmasa anga. 20-25 kg/ha vel a chi thlak hi a tawk vel viau ani. - A chi tuh hma in lei leh FYM @5-10 t/ha leh 80:60:40 kg N, P₂O₅ leh



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			K ₂ O/ha pawlh chu hman phawt tur a ni. Nitrogen dose chanve chu a chi tuh hunlaia hman tur a ni a, tichuan a bang 25% chu thla khat hnu ah ani ang a adang leh 25% chu a par hunah hman tur a ni.
Soybean, pea, lentil toria, breen gram and black gram cultivation in rice fellow	All stage	Zero tillage	- A than a that theih nan nikhat danah tui pek thin tur ani. - Lei rih vur hian thlai kung te a veng ve ani. - Thlasik lai a lei khoro lutuk tur ven nan a chungah hnim leh thildanga khuh tur ani.
Potato	Sowing stage		- Muangchang loving alu chin na tur chu buatsaih vat tur ani. - Hei hian a than hun lain natna hrikin lakah a veng dawn ani. - Lei leh hmain a hmun hma chu fai taka thlawh hmasak tur ani. - A chi thlak hma in a chi chu en fiah hmasak tur ani. - A than a that theih nan nikhat danah tui pek thin tur ani.
VEGETABLE CROP			
Tomato	Bacterial Blight disease		- Tomato bikah chuan sik leh sa hi natna an kaina tlang lawn ber ani . - Hmun hnawng leh ni hmu lo lutuk hmunah chuan natna an kai hma bik ani. - Tomato hi a uai a, a thih mai loh nan Ridomil emaw Indofil emaw Mancozeb @ 2 gm hi tui liter 1 ah pawlh a kah tur ani .
Early Cole crop	Black spot disease		- A than a that theih nan nikhat danah tui pek thin tur ani. - Thlai bul vawn hnawn nana thlai bula hnim ring vawm khawm hi tui pek zawhah dah tur ani. - Zikhlum lam chi ah chuan sik leh sa vangin a hnah ah thil dum rawn



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			awm thin a , hei hi natna tlanglawn ber ani. Thlai hna lam chi leh zikhlum lam chi reng reng enkawl nan Mancozeb @ 2gm ah tui leter 1 pawlha kah tur ani.
Onion and capsicum	Nursery stage	Poly house	- A than a that theih nan nikhat danah tui pek thin tur ani. - Thlai bul vawn hnawn nana thlai bula hnim ring vawm khawm hi tui pek zawhah dah tur ani. - Thlai chhina hmun (nursery) hi hnim a to loh nan Pendimethalin @ 3.5ml hi tui liter 1 zelah pawlh a kah hi a tha hle ani.
		Phytophthora blight	- A chi ven that nan thiram 3g/kg seed emaw Trichoderma viride 4g+ metalaxyl 4g (Apron)/ kg seed hi a tha hle ani - Hneh taka 1% Bordeaux chawhpawlh emaw 2 g captan emaw 3 copper oxychloride a tui liter 1 hi 10-15 DAS a pek hi a tha hle ani.
French bean	Sowing stage		- Tui pek a hnihnah hringa khuh tur ani a. than a that theih nan tui pek hma in lei rin pan hmasak tur ani. - A than duna theih nan leh hnim to loh na turin a kung bulah lei vur chhoh zel tur ani.
Carrot and radish	Sowing stage		- A than a that theih nan nikhat danah tui pek thin tur ani. - Tui pek hnuah thlai bul vawn hnawn na tur siam tur ani. - Zikhlum lam chi ah chuan sik leh sa vangin a hnah ah thil dum a rawn awm thina, hei hi natna tlanglawn ber ani. - Thlai hna lam chi leh zikhlum lam chi reng reng enkawl nan Mancozeb @ 2gm ah tui leter 1 pawlha kah tur ani.






GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



ANIMAL HUSBANDRY			
Pig	All stages		- Khua a vawh hian vawk hian an mahni in tih lumna tur atan chakna an mamawhna a sang bik ani. - An hriselna that leh that loh enfiah renga, a chaw ei tur tlem tlema tih tam hret hret tur ani.. - Sangha tel ah hian omega-3 hi atam em a vangin an chakna muangchanga a in siam chhoh zel theih nan a tha hle ani.
			Porcine Reproductive Respiratory Syndrome (PRRS). 1. Vawknote emaw vawk lak hran.
	Adult stage		Swine fever. 2. SF vaccines hi thla 2 hnua pek tur ani a, chumi hnuah chuan kumtin thlaruk danah pek chhonzawm tur ani.
Cattle	All age group		Hun rei tak khua a ro avanga hnim hnah hring peh tur a awm loh laia bawngin an chaw ei in buk tawk tur leh an taksa tana mamawh tur atan buh kung urea molasses hmanga sawngbawl pek tur ani.
	All age group		Kar 16 hnuah FMD vaccine pek a, chuan thla tin thla 6 chhung chhonzawm tur ani.
	Young stage		- Black Quarter Vaccine (BQV). - Vaccinne hmasa ber hi thla 6 ah emaw a hnu lamah pek tur. - Chumi hnuah chuan Vaccine hi kum tin pek tur ani.
Poultry	Litter management		- Ar te hian hmun thawl nuam tawk, chaw tha an mamawh tawk leh tui thianghlim an mamawh tawk an hmu tur ani a. - An hriselna atan enkawltha tha tawk tak pek hian natna an kai mai theih tur lak atang a venna tha ber ani.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			- Tui an in tur chhawpna tur tha /lian tha tak leh tui thianghlim tak pek tur ani. - Chaw a hmuar/thing pek loh tur ani a, an chaw eitur thlak sak thut loh tur ani.
	Preventive measures	0-3rd week	Ranikhet Disease- an pian atanga ni 1-6 ah F1 vaccine pek tur ani a, chuan a puitlingh chuan R₂B vaccine pek tur ani. - B complex with antibodies
		4th weeks	Coccidiosis- Amprolium or coccidiostat
		4-5th Weeks	Calcium tonic fortified with B₁₂
FISHERY			
	Pond preparation (Dil buatsaih)	3rd-4th weeks	Dil a leitha hman hian sangha chaw kan tih mai planktons insiam nan a tanpui thin.
			- Bawngkek hring 10 tonnes/ha/year vel dil ah hman thin a ni a; bawngkek kumkhat a kan mamawh zat hmunthum a then a hmunkhat hi dil buatsaih nan hman tur ani. A bak zawng hi tui boruak a zirin semdarh a hman thin tur ani. - Single super phosphate hi dil hectare khat zel a zauah kg 250 ang a hman thin tur ani.
			- Leitha kan hman atang a karkhat hnuah tui rawng a tangin sangha chaw inseam that leh thatloh a hriat theih a. Tui rawng eng hring deuh nghalh ah hian sangha chaw planktons te an inseam tha ang a ngaih ani. - Tui nut zawng tehma transparency pawh 30-40 cm vel ani tur ani.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD,
Guwahati)



Expert committee members:

Dr. S.B. Singh	:	Joint Director	basantasinghsoibam@rediffmail.com
Dr. Saurav Saha	:	Scientist (Agril. Physics)	sauravs.saha@gmail.com
Dr. T. Boopathi	:	Scientist (Agril Entomology)	boopathiars@gmail.com
Dr. A. Ratankumar Singh	:	Scientist (Plant Pathology)	ratantplantpatho@gmail.com
Dr. Lungmuana	:	Scientist (Soil Fertility)	lmsingson@gmail.com
Mr. P.L. Lalrinsanga	:	Scientist (Aquaculture)	viensky2@gmail.com
Dr. Dr. V. Dayal	:	Scientist (Horticulture)	Vishambhai5009@gmail.com
Dr. Samuel Lalliansanga	:	Head & Sr. Scientist	samuelpachua10@gmail.com
Mr. Samik Chowdhury	:	Technical Officer	samikchowdhury33@gmail.com
Mr. Diktea chenkuai	:	Project Assistant	dikteachenkuaiboy@gmail.com

Collaborating Department:

Name of the KVK	Programme Coordinator Name and Designation	KVK Email Id	Phone no/ Mobile no
KVK Lunglei	Dr. Lalmuanzovi Head & Sr. Scientist	kvkhnahthial@gmail.com	9862803750 9436154614
KVK, Kolasib	Mr. Lalrosamga Khiangte Head & Sr. Scientist	kvkkolasib@gmail.com	9436152440
KVK, Serchhip	Mr. K. Laltlanmawia Head & Sr. Scientist	kvkserchhip@gmail.com	9436146115 9615389293
KVK, Champhai	Mrs. Lalrinawmi Renthlei Head & Sr. Scientist	kvkchawzawl@gmail.com	9436159788
KVK, Lawngtlai	Dr. Michel Lallawmkimi Head & Sr. Scientist	kvklawntlai@gmail.com	9436155858
KVK, Saiha	Dr. Vanlalhraia Hnampe Head & Sr. Scientist	kvksaiha@gmail.com	8974656509
KVK, Mamit	Dr. Samuel Lalliansanga Head & Sr. Scientist	kvkmamit@gmail.com	9436147625
KVK, Aizawl	Dr. K. P. Chaudhary Head & Sr. Scientist	Kpchy@rediffmail.com kvkaizawl@rediffmail.com	9436351669



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
 Mizoram Centre, Kolasib- 796081, MIZORAM
(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



District: Serchhip

Period: 03 May – 07 May, 2017

Bulletin No: - 697/2017/ Bulletin/English

Date of issue: 02nd May, 2017

Parameters	03.05.2017	04.05.2017	05.05.2017	06.05.2017	07.05.2017
Rainfall (mm)	0	0	0	0	0
Max Temp (°C)	32	32	32	32	32
Min Temp (°C)	21	23	21	19	20
Cloud Coverage	Partially clear	Partially clear	Partially clear	Partially clear	Partially clear
Max RH (%)	99	85	77	84	94
Min RH (%)	37	34	32	29	31
Wind Speed (Kmph)	2	4	2	2	2
*Wind Direction	E	E	S-E	S-E	S

**Northerly- N, North-Easterly- N-E, Easterly- E, South-Easterly- S-E,
 Southerly- S, South-Westerly- S-W, Westerly- W, North-westerly- N-W.**

STATUS OF MONSOON- June 1-30, 2016 (Percent of deviation from normal in parenthesis)

Aizawl- 384.87mm (430.2mm)	Champhai- 105.48mm (359.89mm)	Saiha- 307.40 mm (507.7mm)	Kolasib- 236.00mm (428.1mm)
Lawngtlai-291.20mm (453.1mm)	Lunglei-326.00mm (465.14mm)	Mamit-204.87mm (442.80mm)	Serchhip-411.72mm (259.62mm)

Weather summary of the past three days

Maximum Tem. (°C):27-29°C
Minimum Tem. (°C):18-20°C
Maximum RH (%):97-99%
Minimum RH (%):47-84%
Wind Direction: Easterly
Cloud cover: Mainly clear
Wind speed: 2-4 km/hr


Rainfall: 08.3 mm

Weather forecast valid from 03rd May, 2017 To 07thMay, 2017.

There are no chances of rainfall during the next 5 days. The maximum and minimum temperatures for the next 5 days may range for 32°C and 19-23°C. Maximum relative humidity is expected in the range of 77-99% and minimum may from 29-37%. Wind direction would be easterly to southeasterly and southerly with the wind speed of 2-4 km per hour. Partially clear sky will prevail during the next five days.

Weekly cumulative rainfall: 00.0 mm

NDVI for Mizoram



Moderately wet mildly dry/mildly wet conditions



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



Main Crop / Animal / Fisheries	Stage	Cultural practices/ Pest/ Diseases	Agricultural / Horticultural/ animal husbandry advisories
FRUITS CROPS			
KHASI MANDARIN AND ACID LIME	Seedling transplanting stage		- Fruit palnt should be planted in a sunny and wind-protected area. - In the citrus belt, trees can be planted at any time, however, spring is the best time for container grown plants. - Standard-size trees should be spaced 12 to 25 feet apart and dwarf trees should be set 6 to 10 feet apart. The exact distance depends on the variety. The bigger the fruit, the farther the distance. - If the soil is not well-drained, plant the trees on a slight mound to prevent water logging. - Mulching should be done with dry grasses near the tree base to conserve soil moisture during winter. - The young fruit plant must be irrigated at weekly interval for better establishment.
BANANA			
STAR FRUIT			
PLUM AND PEACH			
		Gummosis, citrus Canker, Citrus greening, Dieback, Lamon butterfly and leaf minor	Die back- Due to low temperature and humidity disease appearance will more. Use Bordeaux past in tree trunk, twigs and branches protect healthy plant from soil borne disease. Lamon butterfly- Spray monocrotophos @0.04% @1.2 ml/lt of water. Leaf minor- Spray confidor 0.05% (0.5 ml/lit of water) at each flush emergence. Citrus Canker- Apply bacterimycin @0.6 g/lt of water.
Passion Fruit	Transplanting stage		- High yielding mother vine with good quality fruits and free of virus diseases should be selected to provide cuttings. - A cutting should contain at least 3 buds and must be planted in sand beds.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			- Immediately after planting these should be kept inside a high humid chamber made out of bamboo and polythene. Grafting: - The root stock of yellow Passion fruit is planted in polythene sleeves and the section from Rahangala hybrid is grafted using wedge or approach method of grafting.
Pineapple	Flowering stage		- Apply flowering inducing chemical (Ethrel 10 PPM+2% urea+0.04% Sodium Carbonate) should be applied in the heart of the plant. In evening and only when plants have at least 32 leaves. - The flowering emergence will come out after 55-60 days after chemical spraying. - Apply split doses of fertilizer @ 60:50:60 g per plant. - Remove all unwanted leaves, branches and weed near to the plant.
PLANTATION CROP			
COFFEE	Blooming stage		- If day temperature and prolong dry spell occur it lead to Floral abnormalities like "Star Flower" in Arabica and "Pink Flower" in Robusta. - Irrigation of plants at alternate day's interval, 6 weeks before harvesting improves fruit retention and fruit development, and reduces fruit cracking and improves the fruit quality. - Mulching should be done with dry grasses near the tree base to conserve soil moisture during winter. - The young fruit plant must be irrigated at weekly interval for better establishment. - Foliar application of Mepiquat chloride @ 1000 PPM concentration or 0.75% SSP @ 1.5 g per 200 lt of water 15 days




GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



Rubber	Nursery stage		interval. ✚ Clearing operation may be done during the month of February to April. ✚ Make fire line to protect the young tree and seedlings. ✚ 10-12 kg of well rotten organic manure and 225 gm rock phosphate should be apply at time of planting to each pit as basal dose application.
CEREALS AND PULSE CROPS			
Pre Kharif Rice	Nursery stage		✚ Use only Well filled and healthy seeds. ✚ Put the seed in 2.5% salt solution i.e 250 g of common salt in 10 lts of water. ✚ Seed treated with Bavistin 50 WP @ 0.1% (2 g/ltr) solution. ✚ Mulching is requiring for better germination in nursery.
Jhum Rice	Sowing stage		✚ Sowing in ridge and furrow. ✚ Mulching with grass for better germination. ✚ Intercropping with greengram and bunchy type frenchbean.
Maize (Jhum)	Sowing stage		✚ Seed should be treated with Thiram @4 g/kg seed. ✚ Use optimum seed rate (20-25 kg/ha) for desire plant population. ✚ Apply well decomposed FYM/pig manure @ 5-10 t/ha along with 80:60:40 kg N, P₂O₅ and K₂O/ha incorporate with soil before sowing. Half nitrogen dose will use at the time of sowing and remaining 25% after one month and 25% at flowering stage.
Rabi Maize	vegetative stage		✚ Light irrigation on every week may be given for better establishment and smooth growth. ✚ Earthing up soil near to plant for better support. ✚ Maize rust disease will prevail due to high relative humidity with low temperature. Apply Mancozeb Kg/ha



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			for effective control. Remove the alternate host <i>Oxalis comiculata</i>.
VEGETABLE CROP			
Cowpea	Sowing stage	KOLASIB	- Plough the field properly, at least 2-3 times. - Mix fertilizer with FYM 50:60:60Kg/ha. - Sow 2-3 seed per whole. - Spacing should be 30 X 20 cm.
Okra	Sowing stage	WEEDING AND light irrigation in nursery bed. Provide irrigation in transplanted okra field.	- Plough the field with the help of spade. - Sow 2 seed 45 X 45 cm spacing. - Before sowing seed provide one or two irrigation. - Provide fertilizer @ 120: 60: 60 Kg/ha
Ginger and turmeric	Sowing stage	SERCHHIP	- Rhizome should be treated with Thiram @4 g/kg seed. - Use optimum seed rate (50-60 kg/ha) for desire plant population. - Apply well decomposed FYM/ pig manure @ 10-20 t/ha along with 120:80:60 kg N, P₂O₅ and K₂O/ha incorporate with soil before sowing. Half nitrogen dose will use at the time of sowing and remaining 25% after one month and 25% at flowering stage.
Colocasia	Sowing stage	LUNGLEI	- Planting is done well prepared land or pits filled up with FYM (12-15) t/ha - Sprouted corms or cormels are planted 5-7 deep at a spacing of 40-50 cm between and within rows in the pits. - Inorganic fertilizer like Urea, SSP and MOP @ 220: 375: 134 kg.
ANIMAL HUSBANDRY			
Pig	All stages	LAWNGTELA SAIHA	- As the weather gets colder, your pigs' energy requirement will increase, as they need more energy to keep warm. - Regularly monitor their level of 'fitness'



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			and increase their feed intake to maintain. ✚ Fish oils are excellent for providing slow-release energy with the added advantage of a high level of omega-3.
		Porcine Reproductive Respiratory Syndrome (PRRS).	1. Culling of positive pigs or piglets.
	Adult stage	Swine fever.	2. Vaccination of pigs with SF vaccines at 2 months and yearly interval/6 month interval
Cattle	All age group		Due to prolong dry spell there is a shortage of green grass in the field. For balanced diet and nutrition to your cattle, provide urea molasses treated paddy straw.
	All age group	Foot and Mouth Disease (FMD)	FMD vaccine at 16 week and repeat every 6 month.
	Young stage	Black Quarter (BQ)	- Black Quarter Vaccine (BQV). ❖ Primary vaccination 6 month or above ❖ Revaccination annually
Poultry	Litter management		✚ Birds require adequate space, sufficient feed to meet their nutritional requirements and an adequate supply of good-quality water. ✚ Good management and sanitation are the best ways to avoid infectious disease in poultry. ✚ Provide ample quantity of clean drinking water. ✚ Avoid feeding of mouldy feed. Don't make sudden changes in feed
	Preventive measures	0-3rd week	✚ Ranikhet Disease- F1 vaccine at (1-6) days of birth and R₂B vaccine for adult birds. ✚ B complex with antibodies
		4th weeks	✚ Coccidiosis- Amprolium or coccidiostat



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



		4-5 th Weeks	✚ Calcium tonic fortified with B ₁₂
FISHERY			
	Pond preparation	3 rd -4 th weeks	✚ Application of fertilizers/manure helps in development of plankton which serve as natural feed for the fishes.
		KOLASIB	✚ Raw cowdung should be applied in the pond at the rate of 10 tonnes/ha/year. One third of the total dose should be applied initially and the rest may be applied in a spilt doses. ✚ Single super phosphate should also be applied at the rate of 250 kg/ha in the pond.
		AIZAWL	✚ After one week of application development of planktons could be observed in the pond depending on the colour of the water. Yellowish green colour is an indicator of the good plankton development. ✚ Transparency of the water needs to be maintained at 30-40 cm.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD,
Guwahati)



Expert committee members:

Dr. S.B. Singh	:	Joint Director	basantasinghsoibam@rediffmail.com
Dr. Saurav Saha	:	Scientist (Agril. Physics)	sauravs.saha@gmail.com
Dr. T. Boopathi	:	Scientist (Agril Entomology)	boopathiars@gmail.com
Dr. A. Ratankumar Singh	:	Scientist (Plant Pathology)	ratantplantpatho@gmail.com
Dr. Lungmuana	:	Scientist (Soil Fertility)	lmsingson@gmail.com
Mr. P.L. Lalrinsanga	:	Scientist (Aquaculture)	viensky2@gmail.com
Dr. Dr. V. Dayal	:	Scientist (Horticulture)	Vishambhai5009@gmail.com
Dr. Samuel Lalliansanga	:	Head & Sr. Scientist	samuelpachua10@gmail.com
Mr. Samik Chowdhury	:	Technical Officer	samikchowdhury33@gmail.com
Mr. Diktea chenkuai	:	Project Assistant	dikteachenkuai10@gmail.com

Collaborating Department:

Name of the KVK	Programme Coordinator Name and Designation	KVK Email Id	Phone no/ Mobile no
KVK Lunglei	Dr. Lalmuanzovi Head & Sr. Scientist	kvknhathial@gmail.com	9862803750 9436154614
KVK, Kolasib	Mr. Lalrosamga Khiangte Head & Sr. Scientist	kvkkolasib@gmail.com	9436152440
KVK, Serchhip	Mr. K. Laltlanmawia Head & Sr. Scientist	kvkserchhip@gmail.com	9436146115 9615389293
KVK, Champhai	Mrs. Lalrinawmi Renthlei Head & Sr. Scientist	kvkchawzawl@gmail.com	9436159788
KVK, Lawngtlai	Dr. Michel Lallawmkimi Head & Sr. Scientist	kvklawntlai@gmail.com	9436155858
KVK, Saiha	Dr. Vanlalhrui Hnampe Head & Sr. Scientist	kvksaiha@gmail.com	8974656509
KVK, Mamit	Dr. Samuel Lalliansanga Head & Sr. Scientist	kvkmamit@gmail.com	9436147625
KVK, Aizawl	Dr. K. P. Chaudhary Head & Sr. Scientist	Kpchy@rediffmail.com kvkaizawl@rediffmail.com	9436351669



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
 Mizoram Centre, Kolasib- 796081, MIZORAM
(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



District: Serchhip

Period: 03 May – 07 May, 2017

Bulletin No: - 697/2017/ Bulletin/Mizo

Date of issue: 02nd May, 2017

Parameters	03.05.2017	04.05.2017	05.05.2017	06.05.2017	07.05.2017
Rainfall (mm)	0	0	0	0	0
Max Temp (°C)	32	32	32	32	32
Min Temp (°C)	21	23	21	19	20
Cloud Coverage	Partially clear	Partially clear	Partially clear	Partially clear	Partially clear
Max RH (%)	99	85	77	84	94
Min RH (%)	37	34	32	29	31
Wind Speed (Kmph)	2	4	2	2	2
*Wind Direction	E	E	S-E	S-E	S

**Northerly- N, North-Easterly- N-E, Easterly- E, South-Easterly- S-E,
 Southerly- S, South-Westerly- S-W, Westerly- W, North-westerly- N-W.**

STATUS OF MONSOON- June 1-30, 2016 (Percent of deviation from normal in parenthesis)

Aizawl- 384.87mm (430.2mm)	Champhai- 105.48mm (359.89mm)	Saiha- 307.40 mm (507.7mm)	Kolasib- 236.00mm (428.1mm)
Lawngtlai-291.20mm (453.1mm)	Lunglei-326.00mm (465.14mm)	Mamit-204.87mm (442.80mm)	Serchhip-411.72mm (259.62mm)

Weather summary of the past three days

Maximum Tem. (°C):27-29°C
Minimum Tem. (°C):18-20°C
Maximum RH (%):97-99%
Minimum RH (%):47-84%
Wind Direction: Easterly
Cloud cover: Mainly clear
Wind speed: 2-4 km/hr


Rainfall: 08.3 mm

**03rd May– 07th May, 2017 chhunga sik leh sa
 dinhmun tur tlangpui**

Tun ni 5 chhung lo awm turah hian ruahtui tla miahlo tura beisei a ni. Khua a lum lai berin 32°C a ni ang a. A vawh lai ber in 19-23°C ni tura beisei a ni. RH san lai berin 77-99% leh a hniam lai berin 29-37% ni tur a rin niin. Thli hi darkar khatah 2-4 km vela chakin chhaklam awi zawngin a tleh rin a ni. A tlangpuiin tun ni nga chung hian khawthiang tak hmuh beisei a ni.

Weekly cumulative rainfall: 00.0mm

NDVI for Mizoram



Moderately wet mildly dry/mildly wet conditions



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



Main Crop / Animal / Fisheries	Stage	Cultural practices/ Pest/ Diseases	Agricultural / Horticultural/ animal husbandry advisories
FRUITS CROPS			
KHASI MANDARIN AND ACID LIME	A kui atanga a seng hun		- Thlasik laia thlai bul khoro lutuk tur vennis chuan hnim hnah hring tlai bul velah dahkhawm tur ani. - Thlai naupang deuah chuan chawlh kar tin a tui pek thin tur ani. - Leia tha mamawh tawh a hmuh theihna turin a hmunhma a hnim awm te thlawhfai thin tur ani. - A seng hma kar 6 chhung chu tui tha taka pek hian a rah tla tur chelh nan leh a rah than that nan te leh a rah keh tur lakah t a veng thei ani.
BANANA			
STAR FRUIT			
PLUM AND PEACH			
		Gummosis, citrus canker, citrus greening and Dieback	Temperture hniam lutuk leh hnawng vang hian natna a a tam duh a . Soil bome natna laka vennis Bordeaux past hi thing zar leh a trangah te hnawih tur ani.
		Fruit fly	Huan zau takah chuan a par tan tirh leh a rah tan tirin chawlhkar hnih chhung chu heng te hian enkawl tur ani: carbaryl 0.2 percent emaw malathion 0.15 percent suspension containing sugar or jeggery at 10 g/l.
PLANTATION CROP			
COFFEE	All stages		Nursery stage - Thlai chi thlak hma in <i>Azospirillum</i> leh <i>Phosphobacterium</i> a enkawl tur ani. - A chi hi December – January ah hmun zawl/rualrem 1.5 - 2.5 cm a in hlatin tlar mumal tak siam in chin tur ani. - Chuan a chi chu lei tlem te a chhilh a buhpawla khuh tur ani. - Nitin tui pek tur ani a, a sat lutuka loh nan niin a chhun loh nan zar hlih tur ani. - Ni 45 hnu velah a tiak thin a,chu chu bag ah an sawn chhuak leh thin ani.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			Harvesting Stage - Coffe rah hmin hi thlasik lain an seng thin a ni. - A rah hmin tha lo ho chu nuai sawm hmain an thliar hrang leh vek thin ani.
		Coffee Berry borer	- A hun takah leh fimkhur taka seng tur ani. - Hmaih neih nuaih loh tur ani. - Sneg hmaih te lakkhawma paih vek tur ani. - Seng bang zawng zawng chu uluka taka paih fai vek tur ani. - A thlai vennan a rah tlai ho chu paih vek tur ani. - Hmun dam lutukah dah loh tur. - Boruak tha taka a hmuh theih nan leh a rawn chawr no theih nan thlai chu uluk tak a hlawi tur ani. - Chuan hei hian kah tur ani. Quinalphos 25 EC @ 340 ml/200 lit emaw lamda cyhalothrin 5 EC 120 – 160 ml / 200 lit. - In leh loa sawngbawl a a ro dan tur tawkah chuan: Arabica / robusta parchment 10 %, Arabica cherry 10.5 % leh robusta cherry 11.0 %.
		Coffee Rust	- Natna hrik in a khawih tawh a hnah leh a dang te paihpaih vek tur ani. - Bordeaux mixture 0.5% in February - March (Pre-bloom) a kah phawt a, Oxycarboxin 0.03% in May - June (Pre-monsoon) ah kah chhunzawm tur ani.
CEREALS AND PULSE CROPS			
Rabi Maize	A chin hun		- Vaimim chinna tur atan lei kan let phut darh anga. - Hei hian a rawn to chhuah na tur atan a pui dawn a ni. - A chi chu kan lei leh saah chuan kan dah ang. - A chi chu Thiram @4 g/kg seed hian ak sawngbawl hmasa anga. 20-25 kg/ha vel a chi thlak hi a tawk vel viau ani. - A chi tuh hma in lei leh FYM @5-10 t/ha leh 80:60:40 kg N, P₂O₅ leh



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			K ₂ O/ha pawlh chu hman phawt tur a ni. Nitrogen dose chanve chu a chi tuh hunlaia hman tur a ni a, tichuan a bang 25% chu thla khat hnu ah ani ang a adang leh 25% chu a par hunah hman tur a ni.
Soybean, pea, lentil toria, breen gram and black gram cultivation in rice fellow	All stage	Zero tillage	- A than a that theih nan nikhat danah tui pek thin tur ani. - Lei rih vur hian thlai kung te a veng ve ani. - Thlasik lai a lei khoro lutuk tur ven nan a chungah hnim leh thildanga khuh tur ani.
Potato	Sowing stage		- Muangchang loving alu chin na tur chu buatsaih vat tur ani. - Hei hian a than hun lain natna hrikin lakah a veng dawn ani. - Lei leh hmain a hmun hma chu fai taka thlawh hmasak tur ani. - A chi thlak hma in a chi chu en fiah hmasak tur ani. - A than a that theih nan nikhat danah tui pek thin tur ani.
VEGETABLE CROP			
Tomato	Bacterial Blight disease		- Tomato bikah chuan sik leh sa hi natna an kaina tlang lawn ber ani . - Hmun hnawng leh ni hmu lo lutuk hmunah chuan natna an kai hma bik ani. - Tomato hi a uai a, a thih mai loh nan Ridomil emaw Indofil emaw Mancozeb @ 2 gm hi tui liter 1 ah pawlh a kah tur ani .
Early Cole crop	Black spot disease		- A than a that theih nan nikhat danah tui pek thin tur ani. - Thlai bul vawn hnawn nana thlai bula hnim ring vawm khawm hi tui pek zawhah dah tur ani. - Zikhlum lam chi ah chuan sik leh sa vangin a hnah ah thil dum rawn



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			awm thin a , hei hi natna tlanglawn ber ani. Thlai hna lam chi leh zikhlum lam chi reng reng enkawl nan Mancozeb @ 2gm ah tui leter 1 pawlha kah tur ani.
Onion and capsicum	Nursery stage	Poly house	- A than a that theih nan nikhat danah tui pek thin tur ani. - Thlai bul vawn hnawn nana thlai bula hnim ring vawm khawm hi tui pek zawhah dah tur ani. - Thlai chhina hmun (nursery) hi hnim a to loh nan Pendimethalin @ 3.5ml hi tui liter 1 zelah pawlh a kah hi a tha hle ani.
		Phytophthora blight	- A chi ven that nan thiram 3g/kg seed emaw Trichoderma viride 4g+ metalaxyl 4g (Apron)/ kg seed hi a tha hle ani - Hneh taka 1% Bordeaux chawhpawlh emaw 2 g captan emaw 3 copper oxychloride a tui liter 1 hi 10-15 DAS a pek hi a tha hle ani.
French bean	Sowing stage		- Tui pek a hnihnah hringa khuh tur ani a. than a that theih nan tui pek hma in lei rin pan hmasak tur ani. - A than duna theih nan leh hnim to loh na turin a kung bulah lei vur chhoh zel tur ani.
Carrot and radish	Sowing stage		- A than a that theih nan nikhat danah tui pek thin tur ani. - Tui pek hnuah thlai bul vawn hnawn na tur siam tur ani. - Zikhlum lam chi ah chuan sik leh sa vangin a hnah ah thil dum a rawn awm thina, hei hi natna tlanglawn ber ani. - Thlai hna lam chi leh zikhlum lam chi reng reng enkawl nan Mancozeb @ 2gm ah tui leter 1 pawlha kah tur ani.






GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



ANIMAL HUSBANDRY			
Pig	All stages		- Khua a vawh hian vawk hian an mahni in tih lumna tur atan chakna an mamawhna a sang bik ani. - An hriselna that leh that loh enfiah renga, a chaw ei tur tlem tlema tih tam hret hret tur ani.. - Sangha tel ah hian omega-3 hi atam em a vangin an chakna muangchanga a in siam chhoh zel theih nan a tha hle ani.
			Porcine Reproductive Respiratory Syndrome (PRRS). 1. Vawknote emaw vawk lak hran.
	Adult stage		Swine fever. 2. SF vaccines hi thla 2 hnua pek tur ani a, chumi hnuah chuan kumtin thlaruk danah pek chhonzawm tur ani.
Cattle	All age group		Hun rei tak khua a ro avanga hnim hnah hring peh tur a awm loh laia bawngin an chaw ei in buk tawk tur leh an taksa tana mamawh tur atan buh kung urea molasses hmanga sawngbawl pek tur ani.
	All age group		Kar 16 hnuah FMD vaccine pek a, chuan thla tin thla 6 chhung chhonzawm tur ani.
	Young stage		- Black Quarter Vaccine (BQV). - Vaccinne hmasa ber hi thla 6 ah emaw a hnu lamah pek tur. - Chumi hnuah chuan Vaccine hi kum tin pek tur ani.
Poultry	Litter management		- Ar te hian hmun thawl nuam tawk, chaw tha an mamawh tawk leh tui thianghlim an mamawh tawk an hmu tur ani a. - An hriselna atan enkawltha tha tawk tak pek hian natna an kai mai theih tur lak atang a venna tha ber ani.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			- Tui an in tur chhawpna tur tha /lian tha tak leh tui thianghlim tak pek tur ani. - Chaw a hmuar/thing pek loh tur ani a, an chaw eitur thlak sak thut loh tur ani.
	Preventive measures	0-3rd week	Ranikhet Disease- an pian atanga ni 1-6 ah F1 vaccine pek tur ani a, chuan a puitlingh chuan R₂B vaccine pek tur ani. - B complex with antibodies
		4th weeks	Coccidiosis- Amprolium or coccidiostat
		4-5th Weeks	Calcium tonic fortified with B₁₂
FISHERY			
	Pond preparation (Dil buatsaih)	3rd-4th weeks	Dil a leitha hman hian sangha chaw kan tih mai planktons insiam nan a tanpui thin.
			- Bawngkek hring 10 tonnes/ha/year vel dil ah hman thin a ni a; bawngkek kumkhat a kan mamawh zat hmunthum a then a hmunkhat hi dil buatsaih nan hman tur ani. A bak zawng hi tui boruak a zirin semdarh a hman thin tur ani. - Single super phosphate hi dil hectare khat zel a zauah kg 250 ang a hman thin tur ani.
			- Leitha kan hman atang a karkhat hnuah tui rawng a tangin sangha chaw inseam that leh thatloh a hriat theih a. Tui rawng eng hring deuh nghalh ah hian sangha chaw planktons te an inseam tha ang a ngaih ani. - Tui nut zawng tehma transparency pawh 30-40 cm vel ani tur ani.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD,
Guwahati)



Expert committee members:

Dr. S.B. Singh	:	Joint Director	basantasinghsoibam@rediffmail.com
Dr. Saurav Saha	:	Scientist (Agril. Physics)	sauravs.saha@gmail.com
Dr. T. Boopathi	:	Scientist (Agril Entomology)	boopathiars@gmail.com
Dr. A. Ratankumar Singh	:	Scientist (Plant Pathology)	ratantplantpatho@gmail.com
Dr. Lungmuana	:	Scientist (Soil Fertility)	lmsingson@gmail.com
Mr. P.L. Lalrinsanga	:	Scientist (Aquaculture)	viensky2@gmail.com
Dr. Dr. V. Dayal	:	Scientist (Horticulture)	Vishambhai5009@gmail.com
Dr. Samuel Lalliansanga	:	Head & Sr. Scientist	samuelpachua10@gmail.com
Mr. Samik Chowdhury	:	Technical Officer	samikchowdhury33@gmail.com
Mr. Diktea chenkual	:	Project Assistant	dikteachenkualboy@gmail.com
Dr. S.B. Singh	:	Joint Director	basantasinghsoibam@rediffmail.com

Collaborating Department:

Name of the KVK		Programme Coordinator Name and Designation	KVK Email Id	Phone no/ Mobile no
KVK Lunglei	:	Dr. Lalmuanzovi Head & Sr. Scientist	kvkhnahthial@gmail.com	9862803750 9436154614
KVK, Kolasib	:	Mr. Lalrosamga Khiantge Head & Sr. Scientist	kvkkolasib@gmail.com	9436152440
KVK, Serchhip	:	Mr. K. Laltlanmawia Head & Sr. Scientist	kvkserchhip@gmail.com	9436146115 9615389293
KVK, Champhai	:	Mrs. Lalrinawmi Renthlei Head & Sr. Scientist	kvkchawzawl@gmail.com	9436159788
KVK, Lawngtlai	:	Dr. Michel Lallawmkimi Head & Sr. Scientist	kvklawntlai@gmail.com	9436155858
KVK, Saiha	:	Dr. Vanlalhruaia Hnampe Head & Sr. Scientist	kvksaiha@gmail.com	8974656509
KVK, Mamit	:	Dr. Samuel Lalliansanga Head & Sr. Scientist	kvkmamit@gmail.com	9436147625
KVK, Aizawl	:	Dr. K. P. Chaudhary Head & Sr. Scientist	Kpchy@rediffmail.com kvkaizawl@rediffmail.com	9436351669



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
 Mizoram Centre, Kolasib- 796081, MIZORAM
(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



District: Aizawl

Period: 03 May – 07 May, 2017

Bulletin No: - 697/2017/ Bulletin/English

Date of issue: 02nd May, 2017

Parameters	03.05.2017	04.05.2017	05.05.2017	06.05.2017	07.05.2017
Rainfall (mm)	10	0	0	10	6
Max Temp (°C)	31	32	32	32	32
Min Temp (°C)	21	21	21	21	21
Cloud Coverage	Partially clear	Partially clear	Partially clear	Partially clear	Partially clear
Max RH (%)	98	92	87	92	96
Min RH (%)	43	45	42	33	38
Wind Speed (Kmph)	4	4	4	4	4
*Wind Direction	S-E	S-E	S-E	S-E	S

**Northerly- N, North-Easterly- N-E, Easterly- E, South-Easterly- S-E,
 Southerly- S, South-Westerly- S-W, Westerly- W, North-westerly- N-W.**

STATUS OF MONSOON- June 1-30, 2016 (Percent of deviation from normal in parenthesis)

Aizawl- 384.87mm (430.2mm)	Champhai- 105.48mm (359.89mm)	Saiha- 307.40 mm (507.7mm)	Kolasib- 236.00mm (428.1mm)
Lawngtlai-291.20mm (453.1mm)	Lunglei-326.00mm (465.14mm)	Mamit-204.87mm (442.80mm)	Serchhip-411.72mm (259.62mm)

Weather summary of the past three days

Maximum Tem. (°C):26-28°C
Minimum Tem. (°C):17-20°C
Maximum RH (%):97-99%
Minimum RH (%):46-52%
Wind Direction: southeasterly
Cloud cover: Mainly clear
Wind speed: 2-4 km/hr


Rainfall: 15.1 mm

Weather forecast valid from 03rd May, 2017 To 07thMay, 2017.

There are chances of moderate to light rainfall during the next 3 days. The maximum and minimum temperatures for the next 5 days may range for 31-32°C and 21°C. Maximum relative humidity is expected in the range of 87-98% and minimum may from 32-45%. Wind direction would be southeasterly to southerly with the wind speed of 4 km per hour. Partially clear sky will prevail during the next five days.

Weekly cumulative rainfall: 26.0 mm

NDVI for Mizoram



Moderately wet mildly dry/mildly wet conditions



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



Main Crop / Animal / Fisheries	Stage	Cultural practices/ Pest/ Diseases	Agricultural / Horticultural/ animal husbandry advisories
FRUITS CROPS			
KHASI MANDARIN AND ACID LIME	Seedling transplanting stage		- Fruit plant should be planted in a sunny and wind-protected area. - In the citrus belt, trees can be planted at any time; however, spring is the best time for container grown plants. - Standard-size trees should be spaced 12 to 25 feet apart and dwarf trees should be set 6 to 10 feet apart. The exact distance depends on the variety. The bigger the fruit, the farther the distance. - If the soil is not well-drained, plant the trees on a slight mound to prevent water logging. - Mulching should be done with dry grasses near the tree base to conserve soil moisture during winter. - The young fruit plant must be irrigated at weekly interval for better establishment.
BANANA			
STAR FRUIT			
PLUM AND PEACH			
		Gummosis, citrus Canker, Citrus greening, Dieback, Lamon butterfly and leaf minor	Die back- Due to low temperature and humidity disease appearance will more. Use Bordeaux past in tree trunk, twigs and branches protect healthy plant from soil borne disease. Lamon butterfly- Spray monocrotophos @0.04% @1.2 ml/lt of water. Leaf minor- Spray confidor 0.05% (0.5 ml/lit of water) at each flush emergence. Citrus Canker- Apply bacterimycin @0.6 g/lt of water.
Passion Fruit	Transplanting stage		- High yielding mother vine with good quality fruits and free of virus diseases should be selected to provide cuttings. - A cutting should contain at least 3 buds and must be planted in sand beds.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			- Immediately after planting these should be kept inside a high humid chamber made out of bamboo and polythene. Grafting: - The root stock of yellow Passion fruit is planted in polythene sleeves and the section from Rahangala hybrid is grafted using wedge or approach method of grafting.
Pineapple	Flowering stage		- Apply flowering inducing chemical (Ethrel 10 PPM+2% urea+0.04% Sodium Carbonate) should be applied in the heart of the plant. In evening and only when plants have at least 32 leaves. - The flowering emergence will come out after 55-60 days after chemical spraying. - Apply split doses of fertilizer @ 60:50:60 g per plant. - Remove all unwanted leaves, branches and weed near to the plant.
PLANTATION CROP			
COFFEE	Blooming stage		- If day temperature and prolong dry spell occur it lead to Floral abnormalities like "Star Flower" in Arabica and "Pink Flower" in Robusta. - Irrigation of plants at alternate day's interval, 6 weeks before harvesting improves fruit retention and fruit development, and reduces fruit cracking and improves the fruit quality. - Mulching should be done with dry grasses near the tree base to conserve soil moisture during winter. - The young fruit plant must be irrigated at weekly interval for better establishment. - Foliar application of Mepiquat chloride @ 1000 PPM concentration or 0.75% SSP @ 1.5 g per 200 lt of water 15 days





GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



Rubber	Nursery stage		interval. ✚ Clearing operation may be done during the month of February to April. ✚ Make fire line to protect the young tree and seedlings. ✚ 10-12 kg of well rotten organic manure and 225 gm rock phosphate should be apply at time of planting to each pit as basal dose application.
CEREALS AND PULSE CROPS			
Pre Kharif Rice	Nursery stage		✚ Use only Well filled and healthy seeds. ✚ Put the seed in 2.5% salt solution i.e 250 g of common salt in 10 lts of water. ✚ Seed treated with Bavistin 50 WP @ 0.1% (2 g/ltr) solution. ✚ Mulching is requiring for better germination in nursery.
Jhum Rice	Sowing stage		✚ Sowing in ridge and furrow. ✚ Mulching with grass for better germination. ✚ Intercropping with greengram and bunchy type frenchbean.
Maize (Jhum)	Sowing stage		✚ Seed should be treated with Thiram @4 g/kg seed. ✚ Use optimum seed rate (20-25 kg/ha) for desire plant population. ✚ Apply well decomposed FYM/pig manure @ 5-10 t/ha along with 80:60:40 kg N, P₂O₅ and K₂O/ha incorporate with soil before sowing. Half nitrogen dose will use at the time of sowing and remaining 25% after one month and 25% at flowering stage.
Rabi Maize	Cob formation stage		✚ Light irrigation on every week may be given for better establishment and smooth growth. ✚ Earthing up soil near to plant for better support. ✚ Maize rust disease will prevail due to high relative humidity with low temperature. Apply Mancozeb Kg/ha



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			for effective control. Remove the alternate host <i>Oxalis comiculata</i>.
VEGETABLE CROP			
Cowpea	Sowing stage	KOLASIB	- Plough the field properly, at least 2-3 times. - Mix fertilizer with FYM 50:60:60Kg/ha. - Sow 2-3 seed per whole. - Spacing should be 30 X 20 cm.
Okra	Sowing stage	Weeding and light irrigation in nursery bed. Provide irrigation in transplanted okra field.	- Plough the field with the help of spade. - Sow 2 seed 45 X 45 cm spacing. - Before sowing seed provide one or two irrigation. - Provide fertilizer @ 120: 60: 60 Kg/ha
Ginger and turmeric	Sowing stage	SERCHHIP	- Rhizome should be treated with Thiram @4 g/kg seed. - Use optimum seed rate (50-60 kg/ha) for desire plant population. - Apply well decomposed FYM/ pig manure @ 10-20 t/ha along with 120:80:60 kg N, P₂O₅ and K₂O/ha incorporate with soil before sowing. Half nitrogen dose will use at the time of sowing and remaining 25% after one month and 25% at flowering stage.
Colocasia	Sowing stage	LUNGLEI	- Planting is done well prepared land or pits filled up with FYM (12-15) t/ha - Sprouted corms or cormels are planted 5-7 deep at a spacing of 40-50 cm between and within rows in the pits. - Inorganic fertilizer like Urea, SSP and MOP @ 220: 375: 134 kg.
ANIMAL HUSBANDRY			
Pig	All stages	LAWNGTELA SAIHA	- As the weather gets colder, your pigs' energy requirement will increase, as they need more energy to keep warm. - Regularly monitor their level of 'fitness'



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			and increase their feed intake to maintain. ✚ Fish oils are excellent for providing slow-release energy with the added advantage of a high level of omega-3.
		Porcine Reproductive Respiratory Syndrome (PRRS).	1. Culling of positive pigs or piglets.
	Adult stage	Swine fever.	2. Vaccination of pigs with SF vaccines at 2 months and yearly interval/6 month interval
Cattle	All age group		Due to prolong dry spell there is a shortage of green grass in the field. For balanced diet and nutrition to your cattle, provide urea molasses treated paddy straw.
	All age group	Foot and Mouth Disease (FMD)	FMD vaccine at 16 week and repeat every 6 month.
	Young stage	Black Quarter (BQ)	- Black Quarter Vaccine (BQV). ❖ Primary vaccination 6 month or above ❖ Revaccination annually
Poultry	Litter management		✚ Birds require adequate space, sufficient feed to meet their nutritional requirements and an adequate supply of good-quality water. ✚ Good management and sanitation are the best ways to avoid infectious disease in poultry. ✚ Provide ample quantity of clean drinking water. ✚ Avoid feeding of mouldy feed. Don't make sudden changes in feed
	Preventive measures	0-3rd week	✚ Ranikhet Disease- F1 vaccine at (1-6) days of birth and R₂B vaccine for adult birds. ✚ B complex with antibodies
		4th weeks	✚ Coccidiosis- Amprolium or coccidiostat



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



		4-5 th Weeks	✚ Calcium tonic fortified with B ₁₂
FISHERY			
	Pond preparation	3 rd -4 th weeks	✚ Application of fertilizers/manure helps in development of plankton which serve as natural feed for the fishes.
		KOLASIB	✚ Raw cowdung should be applied in the pond at the rate of 10 tonnes/ha/year. One third of the total dose should be applied initially and the rest may be applied in a spilt doses. ✚ Single super phosphate should also be applied at the rate of 250 kg/ha in the pond.
		AIZAWL	✚ After one week of application development of planktons could be observed in the pond depending on the colour of the water. Yellowish green colour is an indicator of the good plankton development. ✚ Transparency of the water needs to be maintained at 30-40 cm.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD,
Guwahati)



Expert committee members:

Dr. S.B. Singh	:	Joint Director	basantasinghsoibam@rediffmail.com
Dr. Saurav Saha	:	Scientist (Agril. Physics)	sauravs.saha@gmail.com
Dr. T. Boopathi	:	Scientist (Agril Entomology)	boopathiars@gmail.com
Dr. A. Ratankumar Singh	:	Scientist (Plant Pathology)	ratantplantpatho@gmail.com
Dr. Lungmuana	:	Scientist (Soil Fertility)	lmsingson@gmail.com
Mr. P.L. Lalrinsanga	:	Scientist (Aquaculture)	viensky2@gmail.com
Dr. Dr. V. Dayal	:	Scientist (Horticulture)	Vishambhai5009@gmail.com
Dr. Samuel Lalliansanga	:	Head & Sr. Scientist	samuelpachua10@gmail.com
Mr. Samik Chowdhury	:	Technical Officer	samikchowdhury33@gmail.com
Mr. Diktea chenkuai	:	Project Assistant	dikteachenkuai10@gmail.com

Collaborating Department:

Name of the KVK	Programme Coordinator Name and Designation	KVK Email Id	Phone no/ Mobile no
KVK Lunglei	Dr. Lalmuanzovi Head & Sr. Scientist	kvknhathial@gmail.com	9862803750 9436154614
KVK, Kolasib	Mr. Lalrosamga Khiangte Head & Sr. Scientist	kvkkolasib@gmail.com	9436152440
KVK, Serchhip	Mr. K. Laltlanmawia Head & Sr. Scientist	kvkserchhip@gmail.com	9436146115 9615389293
KVK, Champhai	Mrs. Lalrinawmi Renthlei Head & Sr. Scientist	kvkchawzawl@gmail.com	9436159788
KVK, Lawngtlai	Dr. Michel Lallawmkimi Head & Sr. Scientist	kvklawntlai@gmail.com	9436155858
KVK, Saiha	Dr. Vanlalhraia Hnampe Head & Sr. Scientist	kvksaiha@gmail.com	8974656509
KVK, Mamit	Dr. Samuel Lalliansanga Head & Sr. Scientist	kvkmamit@gmail.com	9436147625
KVK, Aizawl	Dr. K. P. Chaudhary Head & Sr. Scientist	Kpchy@rediffmail.com kvkaizawl@rediffmail.com	9436351669



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
 Mizoram Centre, Kolasib- 796081, MIZORAM
(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



District: Aizawl

Period: 03 May – 07 May, 2017

Bulletin No: - 697/2017/ Bulletin/Mizo

Date of issue: 02nd May, 2017

Parameters	03.05.2017	04.05.2017	05.05.2017	06.05.2017	07.05.2017
Rainfall (mm)	10	0	0	10	6
Max Temp (°C)	31	32	32	32	32
Min Temp (°C)	21	21	21	21	21
Cloud Coverage	Partially clear	Partially clear	Partially clear	Partially clear	Partially clear
Max RH (%)	98	92	87	92	96
Min RH (%)	43	45	42	33	38
Wind Speed (Kmph)	4	4	4	4	4
*Wind Direction	S-E	S-E	S-E	S-E	S

**Northerly- N, North-Easterly- N-E, Easterly- E, South-Easterly- S-E,
 Southerly- S, South-Westerly- S-W, Westerly- W, North-westerly- N-W.**

STATUS OF MONSOON- June 1-30, 2016 (Percent of deviation from normal in parenthesis)

Aizawl- 384.87mm (430.2mm)	Champhai- 105.48mm (359.89mm)	Saiha- 307.40 mm (507.7mm)	Kolasib- 236.00mm (428.1mm)
Lawngtlai-291.20mm (453.1mm)	Lunglei-326.00mm (465.14mm)	Mamit-204.87mm (442.80mm)	Serchhip-411.72mm (259.62mm)

Weather summary of the past three days

Maximum Tem. (°C):26-28°C
Minimum Tem. (°C):17-20°C
Maximum RH (%):97-99%
Minimum RH (%):46-52%
Wind Direction: southeasterly
Cloud cover: Mainly clear
Wind speed: 2-4 km/hr


Rainfall: 15.1 mm

**03rd May– 07th May, 2017 chhunga sik leh sa
 dinhmun tur tlangpui**

Tun ni 3 chhung lo awm turah hian ruahtui tla miahlo tura beisei a ni. Khua a lum lai berin 31-32°C a ni ang a. A vawh lai ber in 21°C ni tura beisei a ni. RH san lai berin 87-98% leh a hniam lai berin 32-45% ni tur a rin niin. Thli hi darkar khatah 4 km vela chakin chhaklam awi zawngin a tleh rin a ni. A tlangpuiin tun ni nga chhung hian khawthiang tak hmuh beisei a ni.

Weekly cumulative rainfall: 26.0mm

NDVI for Mizoram



Moderately wet mildly dry/mildly wet conditions




GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



Main Crop/ Animal /Fisheries	Stage	Cultural practices/ Pest/ Diseases	Agricultural / Horticultural/ animal husbandry advisories
FRUITS CROPS			
KHASI MANDARIN AND ACID LIME	A kui atanga a seng hun		- Thlasik laia thlai bul khoro lutuk tur vennan chuan hnim hnah hring tlai bul velah dahkhawm tur ani. - Thlai naupang deuah chuan chawlh kar tin a tui pek thin tur ani. - Leia tha mamawh tawh a hmuh theihna turin a hmunhma a hnim awm te thlawhfai thin tur ani. - A seng hma kar 6 chhung chu tui tha taka pek hian a rah tla tur chelh nan leh a rah than that nan te leh a rah keh tur lakah t a veng thei ani.
BANANA			
STAR FRUIT			
PLUM AND PEACH			
		Gummosis, citrus canker, citrus greening and Dieback	Temperture hniam lutuk leh hnawng vang hian natna a a tam duh a . Soil bome natna laka vennan Bordeaux past hi thing zar leh a trangah te hnawih tur ani.
		Fruit fly	Huan zau takah chuan a par tan tirh leh a rah tan tirin chawlhkar hnih chhung chu heng te hian enkawl tur ani: carbaryl 0.2 percent emaw malathion 0.15 percent suspension containing sugar or jeggery at 10 g/l.
PLANTATION CROP			
COFFEE	All stages		Nursery stage - Thlai chi thlak hma in <i>Azospirillum</i> leh <i>Phosphobacterium</i> a enkawl tur ani. - A chi hi December – January ah hmun zawl/rualrem 1.5 - 2.5 cm a in hlatin tlar mumal tak siam in chin tur ani. - Chuan a chi chu lei tlem te a chhilh a buhpawla khuh tur ani. - Nitin tui pek tur ani a, a sat lutuka loh nan niin a chhun loh nan zar hlih tur ani. - Ni 45 hnu velah a tiak thin a,chu chu bag ah an sawn chhuak leh thin ani.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			Harvesting Stage - Coffe rah hmin hi thlasik lain an seng thin a ni. - A rah hmin tha lo ho chu nuai sawm hmain an thliar hrang leh vek thin ani.
		Coffee Berry borer	- A hun takah leh fimkhur taka seng tur ani. - Hmaih neih nuaih loh tur ani. - Sneg hmaih te lakkhawma paih vek tur ani. - Seng bang zawng zawng chu uluka taka paih fai vek tur ani. - A thlai vennan a rah tlai ho chu paih vek tur ani. - Hmun dam lutukah dah loh tur. - Boruak tha taka a hmuh theih nan leh a rawn chawr no theih nan thlai chu uluk tak a hlawi tur ani. - Chuan hei hian kah tur ani. Quinalphos 25 EC @ 340 ml/200 lit emaw lamda cyhalothrin 5 EC 120 – 160 ml / 200 lit. - In leh loa sawngbawl a a ro dan tur tawkah chuan: Arabica / robusta parchment 10 %, Arabica cherry 10.5 % leh robusta cherry 11.0 %.
		Coffee Rust	- Natna hrik in a khawih tawh a hnah leh a dang te paihpaih vek tur ani. - Bordeaux mixture 0.5% in February - March (Pre-bloom) a kah phawt a, Oxycarboxin 0.03% in May - June (Pre-monsoon) ah kah chhunzawm tur ani.
CEREALS AND PULSE CROPS			
Rabi Maize	A chin hun		- Vaimim chinna tur atan lei kan let phut darh anga. - Hei hian a rawn to chhuah na tur atan a pui dawn a ni. - A chi chu kan lei leh saah chuan kan dah ang. - A chi chu Thiram @4 g/kg seed hian ak sawngbawl hmasa anga. 20-25 kg/ha vel a chi thlak hi a tawk vel viau ani. - A chi tuh hma in lei leh FYM @5-10 t/ha leh 80:60:40 kg N, P₂O₅ leh



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			K ₂ O/ha pawlh chu hman phawt tur a ni. Nitrogen dose chanve chu a chi tuh hunlaia hman tur a ni a, tichuan a bang 25% chu thla khat hnu ah ani ang a adang leh 25% chu a par hunah hman tur a ni.
Soybean, pea, lentil toria, breen gram and black gram cultivation in rice fellow	All stage	Zero tillage	- A than a that theih nan nikhat danah tui pek thin tur ani. - Lei rih vur hian thlai kung te a veng ve ani. - Thlasik lai a lei khoro lutuk tur ven nan a chungah hnim leh thildanga khuh tur ani.
Potato	Sowing stage		- Muangchang loving alu chin na tur chu buatsaih vat tur ani. - Hei hian a than hun lain natna hrikin lakah a veng dawn ani. - Lei leh hmain a hmun hma chu fai taka thlawh hmasak tur ani. - A chi thlak hma in a chi chu en fiah hmasak tur ani. - A than a that theih nan nikhat danah tui pek thin tur ani.
VEGETABLE CROP			
Tomato	Bacterial Blight disease		- Tomato bikah chuan sik leh sa hi natna an kaina tlang lawn ber ani . - Hmun hnawng leh ni hmu lo lutuk hmunah chuan natna an kai hma bik ani. - Tomato hi a uai a, a thih mai loh nan Ridomil emaw Indofil emaw Mancozeb @ 2 gm hi tui liter 1 ah pawlh a kah tur ani .
Early Cole crop	Black spot disease		- A than a that theih nan nikhat danah tui pek thin tur ani. - Thlai bul vawn hnawn nana thlai bula hnim ring vawm khawm hi tui pek zawhah dah tur ani. - Zikhlum lam chi ah chuan sik leh sa vangin a hnah ah thil dum rawn



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			awm thin a , hei hi natna tlanglawn ber ani. Thlai hna lam chi leh zikhlum lam chi reng reng enkawl nan Mancozeb @ 2gm ah tui leter 1 pawlha kah tur ani.
Onion and capsicum	Nursery stage	Poly house	- A than a that theih nan nikhat danah tui pek thin tur ani. - Thlai bul vawn hnawn nana thlai bula hnim ring vawm khawm hi tui pek zawhah dah tur ani. - Thlai chhina hmun (nursery) hi hnim a to loh nan Pendimethalin @ 3.5ml hi tui liter 1 zelah pawlh a kah hi a tha hle ani.
		Phytophthora blight	- A chi ven that nan thiram 3g/kg seed emaw Trichoderma viride 4g+ metalaxyl 4g (Apron)/ kg seed hi a tha hle ani - Hneh taka 1% Bordeaux chawhpawlh emaw 2 g captan emaw 3 copper oxychloride a tui liter 1 hi 10-15 DAS a pek hi a tha hle ani.
French bean	Sowing stage		- Tui pek a hnihnah hringa khuh tur ani a. than a that theih nan tui pek hma in lei rin pan hmasak tur ani. - A than duna theih nan leh hnim to loh na turin a kung bulah lei vur chhoh zel tur ani.
Carrot and radish	Sowing stage		- A than a that theih nan nikhat danah tui pek thin tur ani. - Tui pek hnuah thlai bul vawn hnawn na tur siam tur ani. - Zikhlum lam chi ah chuan sik leh sa vangin a hnah ah thil dum a rawn awm thina, hei hi natna tlanglawn ber ani. - Thlai hna lam chi leh zikhlum lam chi reng reng enkawl nan Mancozeb @ 2gm ah tui leter 1 pawlha kah tur ani.






GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



ANIMAL HUSBANDRY			
Pig	All stages		- Khua a vawh hian vawk hian an mahni in tih lumna tur atan chakna an mamawhna a sang bik ani. - An hriselna that leh that loh enfiah renga, a chaw ei tur tlem tlema tih tam hret hret tur ani.. - Sangha tel ah hian omega-3 hi atam em a vanging an chakna muangchanga a in siam chhoh zel theih nan a tha hle ani.
			Porcine Reproductive Respiratory Syndrome (PRRS). 1. Vawknote emaw vawk lak hran.
	Adult stage		Swine fever. 2. SF vaccines hi thla 2 hnua pek tur ani a, chumi hnuah chuan kumtin thlaruk danah pek chhonzawm tur ani.
Cattle	All age group		Hun rei tak khua a ro avanga hnim hnah hring peh tur a awm loh laia bawngin an chaw ei in buk tawk tur leh an taksa tana mamawh tur atan buh kung urea molasses hmanga sawngbawl pek tur ani.
	All age group		Kar 16 hnuah FMD vaccine pek a, chuan thla tin thla 6 chhung chhonzawm tur ani.
	Young stage		- Black Quarter Vaccine (BQV). - Vaccinne hmasa ber hi thla 6 ah emaw a hnu lamah pek tur. - Chumi hnuah chuan Vaccine hi kum tin pek tur ani.
Poultry	Litter management		- Ar te hian hmun thawl nuam tawk, chaw tha an mamawh tawk leh tui thianghlim an mamawh tawk an hmu tur ani a. - An hriselna atan enkawltha tha tawk tak pek hian natna an kai mai theih tur lak atang a venna tha ber ani.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			- Tui an in tur chhawpna tur tha /lian tha tak leh tui thianghlim tak pek tur ani. - Chaw a hmuar/thing pek loh tur ani a, an chaw eitur thlak sak thut loh tur ani.
	Preventive measures	0-3rd week	Ranikhet Disease- an pian atanga ni 1-6 ah F1 vaccine pek tur ani a, chuan a puitlingh chuan R₂B vaccine pek tur ani. - B complex with antibodies
		4th weeks	Coccidiosis- Amprolium or coccidiostat
		4-5th Weeks	Calcium tonic fortified with B₁₂
FISHERY			
	Pond preparation (Dil buatsaih)	3rd-4th weeks	Dil a leitha hman hian sangha chaw kan tih mai planktons insiam nan a tanpui thin.
			- Bawngkek hring 10 tonnes/ha/year vel dil ah hman thin a ni a; bawngkek kumkhat a kan mamawh zat hmunthum a then a hmunkhat hi dil buatsaih nan hman tur ani. A bak zawng hi tui boruak a zirin semdarh a hman thin tur ani. - Single super phosphate hi dil hectare khat zel a zauah kg 250 ang a hman thin tur ani.
			- Leitha kan hman atang a karkhat hnuah tui rawng a tangin sangha chaw inseam that leh thatloh a hriat theih a. Tui rawng eng hring deuh nghalh ah hian sangha chaw planktons te an inseam tha ang a ngaih ani. - Tui nut zawng tehma transparency pawh 30-40 cm vel ani tur ani.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD,
Guwahati)



Expert committee members:

Dr. S.B. Singh	: Joint Director	basantasinghsoibam@rediffmail.com
Dr. Saurav Saha	: Scientist (Agril. Physics)	sauravs.saha@gmail.com
Dr. T. Boopathi	: Scientist (Agril Entomology)	boopathiars@gmail.com
Dr. A. Ratankumar Singh	: Scientist (Plant Pathology)	ratantplantpatho@gmail.com
Dr. Lungmuana	: Scientist (Soil Fertility)	lmsingson@gmail.com
Mr. P.L. Lalrinsanga	: Scientist (Aquaculture)	viensky2@gmail.com
Dr. Dr. V. Dayal	: Scientist (Horticulture)	Vishambhai5009@gmail.com
Dr. Samuel Lalliansanga	: Head & Sr. Scientist	samuelpachua10@gmail.com
Mr. Samik Chowdhury	: Technical Officer	samikchowdhury33@gmail.com
Mr. Diktea chenkual	: Project Assistant	dikteachenkualboy@gmail.com

Collaborating Department:

Name of the KVK	Programme Coordinator Name and Designation	KVK Email Id	Phone no/ Mobile no
KVK Lunglei	: Dr. Lalmuanzovi Head & Sr. Scientist	kvkhnahthial@gmail.com	9862803750 9436154614
KVK, Kolasib	: Mr. Lalrosamga Khiangte Head & Sr. Scientist	kvkkolasib@gmail.com	9436152440
KVK, Serchhip	: Mr. K. Laltlanmawia Head & Sr. Scientist	kvkserchhip@gmail.com	9436146115 9615389293
KVK, Champhai	: Mrs. Lalrinawmi Renthlei Head & Sr. Scientist	kvkchawzawl@gmail.com	9436159788
KVK, Lawngtlai	: Dr. Michel Lallawmkimi Head & Sr. Scientist	kvklawntlai@gmail.com	9436155858
KVK, Saiha	: Dr. Vanlalhruaia Hnampe Head & Sr. Scientist	kvksaiha@gmail.com	8974656509
KVK, Mamit	: Dr. Samuel Lalliansanga Head & Sr. Scientist	kvkmamit@gmail.com	9436147625
KVK, Aizawl	: Dr. K. P. Chaudhary Head & Sr. Scientist	Kpchy@rediffmail.com kvkaizawl@rediffmail.com	9436351669



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
 Mizoram Centre, Kolasib- 796081, MIZORAM
(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



District: Champhai

Period: 03 May – 07 May, 2017

Bulletin No: - 697/2017/ Bulletin/English

Date of issue: 02nd May, 2017

Parameters	03.05.2017	04.05.2017	05.05.2017	06.05.2017	07.05.2017
Rainfall (mm)	9	0	0	3	6
Max Temp (°C)	31	32	32	32	32
Min Temp (°C)	21	22	22	22	22
Cloud Coverage	Partially clear	Clear sky	Clear sky	Partially clear	Partially clear
Max RH (%)	91	83	76	85	92
Min RH (%)	35	37	33	28	30
Wind Speed (Kmph)	4	4	4	3	2
*Wind Direction	S-E	S-E	S	S	S

**Northerly- N, North-Easterly- N-E, Easterly- E, South-Easterly- S-E,
 Southerly- S, South-Westerly- S-W, Westerly- W, North-westerly- N-W.**

STATUS OF MONSOON- June 1-30, 2016 (Percent of deviation from normal in parenthesis)

Aizawl- 384.87mm (430.2mm)	Champhai- 105.48mm (359.89mm)	Saiha- 307.40 mm (507.7mm)	Kolasib- 236.00mm (428.1mm)
Lawngtlai-291.20mm (453.1mm)	Lunglei-326.00mm (465.14mm)	Mamit-204.87mm (442.80mm)	Serchhip-411.72mm (259.62mm)

Weather summary of the past three days

Maximum Tem. (°C):25-28°C
Minimum Tem. (°C):18-21°C
Maximum RH (%):89-94%
Minimum RH (%):35-43%
Wind Direction: Southeasterly
Cloud cover: Mainly clear
Wind Speed: 3-4 km/hr


Rainfall: 10.2 mm

Weather forecast valid from 03rd May, 2017 To 07thMay, 2017.

There are chances of light rainfall during the next 3 days. The maximum and minimum temperatures for the next 5 days may range for 31-32°C and 21-22°C. Maximum relative humidity is expected in the range of 76-92% and minimum may from 28-37%. Wind direction would be southeasterly to southerly with the wind speed of 2-4 km per hour. Partially clear sky will prevail during the next five days.

Weekly cumulative rainfall: 18.0 mm

NDVI for Mizoram



Moderately wet mildly dry/mildly wet conditions



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



Main Crop / Animal / Fisheries	Stage	Cultural practices/ Pest/ Diseases	Agricultural / Horticultural/ animal husbandry advisories
FRUITS CROPS			
KHASI MANDARIN AND ACID LIME	Seedling transplanting stage		- Fruit plant should be planted in a sunny and wind-protected area. - In the citrus belt, trees can be planted at any time, however, spring is the best time for container grown plants. - Standard-size trees should be spaced 12 to 25 feet apart and dwarf trees should be set 6 to 10 feet apart. The exact distance depends on the variety. The bigger the fruit, the farther the distance. - If the soil is not well-drained, plant the trees on a slight mound to prevent water logging. - Mulching should be done with dry grasses near the tree base to conserve soil moisture during winter. - The young fruit plant must be irrigated at weekly interval for better establishment.
BANANA			
STAR FRUIT			
PLUM AND PEACH			
		Gummosis, citrus Canker, Citrus greening, Dieback, Lamon butterfly and leaf minor	Die back- Due to low temperature and humidity disease appearance will more. Use Bordeaux past in tree trunk, twigs and branches protect healthy plant from soil borne disease. Lamon butterfly- Spray monocrotophos @0.04% @1.2 ml/lt of water. Leaf minor- Spray confidor 0.05% (0.5 ml/lit of water) at each flush emergence. Citrus Canker- Apply bacterimycin @0.6 g/lt of water.
Passion Fruit	Transplanting stage		- High yielding mother vine with good quality fruits and free of virus diseases should be selected to provide cuttings. - A cutting should contain at least 3 buds and must be planted in sand beds.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			- Immediately after planting these should be kept inside a high humid chamber made out of bamboo and polythene. Grafting: - The root stock of yellow Passion fruit is planted in polythene sleeves and the section from Rahangala hybrid is grafted using wedge or approach method of grafting.
Pineapple	Flowering stage		- Apply flowering inducing chemical (Ethrel 10 PPM+2% urea+0.04% Sodium Carbonate) should be applied in the heart of the plant. In evening and only when plants have at least 32 leaves. - The flowering emergence will come out after 55-60 days after chemical spraying. - Apply split doses of fertilizer @ 60:50:60 g per plant. - Remove all unwanted leaves, branches and weed near to the plant.
PLANTATION CROP			
COFFEE	Blooming stage		- If day temperature and prolong dry spell occur it lead to Floral abnormalities like "Star Flower" in Arabica and "Pink Flower" in Robusta. - Irrigation of plants at alternate day's interval, 6 weeks before harvesting improves fruit retention and fruit development, and reduces fruit cracking and improves the fruit quality. - Mulching should be done with dry grasses near the tree base to conserve soil moisture during winter. - The young fruit plant must be irrigated at weekly interval for better establishment. - Foliar application of Mepiquat chloride @ 1000 PPM concentration or 0.75% SSP @ 1.5 g per 200 lt of water 15 days




GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



Rubber	Nursery stage		interval. ✚ Clearing operation may be done during the month of February to April. ✚ Make fire line to protect the young tree and seedlings. ✚ 10-12 kg of well rotten organic manure and 225 gm rock phosphate should be apply at time of planting to each pit as basal dose application.
CEREALS AND PULSE CROPS			
Pre Kharif Rice	Nursery stage		✚ Use only Well filled and healthy seeds. ✚ Put the seed in 2.5% salt solution i.e 250 g of common salt in 10 lts of water. ✚ Seed treated with Bavistin 50 WP @ 0.1% (2 g/ltr) solution. ✚ Mulching is requiring for better germination in nursery.
Jhum Rice	Sowing stage		✚ Sowing in ridge and furrow. ✚ Mulching with grass for better germination. ✚ Intercropping with greengram and bunchy type frenchbean.
Maize (Jhum)	Sowing stage		✚ Seed should be treated with Thiram @4 g/kg seed. ✚ Use optimum seed rate (20-25 kg/ha) for desire plant population. ✚ Apply well decomposed FYM/pig manure @ 5-10 t/ha along with 80:60:40 kg N, P₂O₅ and K₂O/ha incorporate with soil before sowing. Half nitrogen dose will use at the time of sowing and remaining 25% after one month and 25% at flowering stage.
Rabi Maize	vegetative stage		✚ Light irrigation on every week may be given for better establishment and smooth growth. ✚ Earthing up soil near to plant for better support. ✚ Maize rust disease will prevail due to high relative humidity with low temperature. Apply Mancozeb Kg/ha



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			for effective control. Remove the alternate host <i>Oxalis comiculata</i>.
VEGETABLE CROP			
Cowpea	Sowing stage	KOLASIB	- Plough the field properly, at least 2-3 times. - Mix fertilizer with FYM 50:60:60Kg/ha. - Sow 2-3 seed per whole. - Spacing should be 30 X 20 cm.
Okra	Sowing stage	Weeding and light irrigation in nursery bed. Provide irrigation in transplanted okra field.	- Plough the field with the help of spade. - Sow 2 seed 45 X 45 cm spacing. - Before sowing seed provide one or two irrigation. - Provide fertilizer @ 120: 60: 60 Kg/ha
Ginger and turmeric	Sowing stage	SERCHHIP	- Rhizome should be treated with Thiram @4 g/kg seed. - Use optimum seed rate (50-60 kg/ha) for desire plant population. - Apply well decomposed FYM/ pig manure @ 10-20 t/ha along with 120:80:60 kg N, P₂O₅ and K₂O/ha incorporate with soil before sowing. Half nitrogen dose will use at the time of sowing and remaining 25% after one month and 25% at flowering stage.
Colocasia	Sowing stage	LUNGLEI	- Planting is done well prepared land or pits filled up with FYM (12-15) t/ha - Sprouted corms or cormels are planted 5-7 deep at a spacing of 40-50 cm between and within rows in the pits. - Inorganic fertilizer like Urea, SSP and MOP @ 220: 375: 134 kg.
ANIMAL HUSBANDRY			
Pig	All stages	LAWNGTIA	- As the weather gets colder, your pigs' energy requirement will increase, as they need more energy to keep warm. - Regularly monitor their level of 'fitness'



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			and increase their feed intake to maintain. ✚ Fish oils are excellent for providing slow-release energy with the added advantage of a high level of omega-3.
		Porcine Reproductive Respiratory Syndrome (PRRS).	1. Culling of positive pigs or piglets.
	Adult stage	Swine fever.	2. Vaccination of pigs with SF vaccines at 2 months and yearly interval/6 month interval
Cattle	All age group		Due to prolong dry spell there is a shortage of green grass in the field. For balanced diet and nutrition to your cattle, provide urea molasses treated paddy straw.
	All age group	Foot and Mouth Disease (FMD)	FMD vaccine at 16 week and repeat every 6 month.
	Young stage	Black Quarter (BQ)	- Black Quarter Vaccine (BQV). ✦ Primary vaccination 6 month or above ✦ Revaccination annually
Poultry	Litter management		✚ Birds require adequate space, sufficient feed to meet their nutritional requirements and an adequate supply of good-quality water. ✚ Good management and sanitation are the best ways to avoid infectious disease in poultry. ✚ Provide ample quantity of clean drinking water. ✚ Avoid feeding of mouldy feed. Don't make sudden changes in feed
	Preventive measures	0-3rd week	✚ Ranikhet Disease- F1 vaccine at (1-6) days of birth and R₂B vaccine for adult birds. ✚ B complex with antibodies
		4th weeks	✚ Coccidiosis- Amprolium or coccidiostat



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



		4-5 th Weeks	✚ Calcium tonic fortified with B ₁₂
FISHERY			
	Pond preparation	3 rd -4 th weeks	✚ Application of fertilizers/manure helps in development of plankton which serve as natural feed for the fishes.
		KOLASIB	✚ Raw cowdung should be applied in the pond at the rate of 10 tonnes/ha/year. One third of the total dose should be applied initially and the rest may be applied in a spilt doses. ✚ Single super phosphate should also be applied at the rate of 250 kg/ha in the pond.
		AIZAWL	✚ After one week of application development of planktons could be observed in the pond depending on the colour of the water. Yellowish green colour is an indicator of the good plankton development. ✚ Transparency of the water needs to be maintained at 30-40 cm.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD,
Guwahati)



Expert committee members:

Dr. S.B. Singh	:	Joint Director	basantasinghsoibam@rediffmail.com
Dr. Saurav Saha	:	Scientist (Agril. Physics)	sauravs.saha@gmail.com
Dr. T. Boopathi	:	Scientist (Agril Entomology)	boopathiars@gmail.com
Dr. A. Ratankumar Singh	:	Scientist (Plant Pathology)	ratantplantpatho@gmail.com
Dr. Lungmuana	:	Scientist (Soil Fertility)	lmsingson@gmail.com
Mr. P.L. Lalrinsanga	:	Scientist (Aquaculture)	viensky2@gmail.com
Dr. Dr. V. Dayal	:	Scientist (Horticulture)	Vishambhai5009@gmail.com
Dr. Samuel Lalliansanga	:	Head & Sr. Scientist	samuelpachua10@gmail.com
Mr. Samik Chowdhury	:	Technical Officer	samikchowdhury33@gmail.com
Mr. Diktea chenkuai	:	Project Assistant	dikteachenkuaiboy@gmail.com

Collaborating Department:

Name of the KVK	Programme Coordinator Name and Designation	KVK Email Id	Phone no/ Mobile no
KVK Lunglei	Dr. Lalmuanzovi Head & Sr. Scientist	kvkhnahthial@gmail.com	9862803750 9436154614
KVK, Kolasib	Mr. Lalrosamga Khiangte Head & Sr. Scientist	kvkkolasib@gmail.com	9436152440
KVK, Serchhip	Mr. K. Laltlanmawia Head & Sr. Scientist	kvkserchhip@gmail.com	9436146115 9615389293
KVK, Champhai	Mrs. Lalrinawmi Renthlei Head & Sr. Scientist	kvkchawzawl@gmail.com	9436159788
KVK, Lawngtlai	Dr. Michel Lallawmkimi Head & Sr. Scientist	kvklawntlai@gmail.com	9436155858
KVK, Saiha	Dr. Vanlalhraia Hnampe Head & Sr. Scientist	kvksaiha@gmail.com	8974656509
KVK, Mamit	Dr. Samuel Lalliansanga Head & Sr. Scientist	kvkmamit@gmail.com	9436147625
KVK, Aizawl	Dr. K. P. Chaudhary Head & Sr. Scientist	Kpchy@rediffmail.com kvkaizawl@rediffmail.com	9436351669



GRAMIN KRISHI MAUSAM SEWA

ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM
(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



District: Champhai

Period: 03 May – 07 May, 2017

Bulletin No: - 697/2017/ Bulletin/Mizo

Date of issue: 02nd May, 2017

Parameters	03.05.2017	04.05.2017	05.05.2017	06.05.2017	07.05.2017
Rainfall (mm)	9	0	0	3	6
Max Temp (°C)	31	32	32	32	32
Min Temp (°C)	21	22	22	22	22
Cloud Coverage	Partially clear	Clear sky	Clear sky	Partially clear	Partially clear
Max RH (%)	91	83	76	85	92
Min RH (%)	35	37	33	28	30
Wind Speed (Kmph)	4	4	4	3	2
*Wind Direction	S-E	S-E	S	S	S

Northerly- N, North-Easterly- N-E, Easterly- E, South-Easterly- S-E, Southerly- S, South-Westerly- S-W, Westerly- W, North-westerly- N-W.

STATUS OF MONSOON- June 1-30, 2016 (Percent of deviation from normal in parenthesis)

Aizawl- 384.87mm (430.2mm)	Champhai- 105.48mm (359.89mm)	Saiha- 307.40 mm (507.7mm)	Kolasib- 236.00mm (428.1mm)
Lawngtlai-291.20mm (453.1mm)	Lunglei-326.00mm (465.14mm)	Mamit-204.87mm (442.80mm)	Serchhip-411.72mm (259.62mm)

Weather summary of the past three days

Maximum Tem. (°C):25-28°C
Minimum Tem. (°C):18-21°C
Maximum RH (%):89-94%
Minimum RH (%):35-43%
Wind Direction: Southeasterly
Cloud cover: Mainly clear
Wind Speed: 3-4 km/hr


Rainfall: 10.2 mm

03rd May– 07th May, 2017 chhunga sik leh sa dinhmun tur tlangpui

Tun ni 3 chhung lo awm turah hian ruahtui tla miahlo tura beisei a ni. Khua a lum lai berin 31-32°C a ni ang a. A vawh lai ber in 21-22°C ni tura beisei a ni. RH san lai berin 76-92% leh a hniam lai berin 28-37% ni tur a rin niin. Thli hi darkar khatah 2-4 km vela chakin chhaklam awi zawngin a tleh rin a ni. A tlangpuiin tun ni nga chung hian khawthiang tak hmuh beisei a ni.

Weekly cumulative rainfall: 18.0mm

NDVI for Mizoram



Moderately wet mildly dry/mildly wet conditions



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



Main Crop / Animal / Fisheries	Stage	Cultural practices/ Pest/ Diseases	Agricultural / Horticultural/ animal husbandry advisories
FRUITS CROPS			
KHASI MANDARIN AND ACID LIME	A kui atanga a seng hun		- Thlasik laia thlai bul khoro lutuk tur vennan chuan hnim hnah hring tlai bul velah dahkhawm tur ani. - Thlai naupang deuah chuan chawlh kar tin a tui pek thin tur ani. - Leia tha mamawh tawh a hmuh theihna turin a hmunhma a hnim awm te thlawhfai thin tur ani. - A seng hma kar 6 chhung chu tui tha taka pek hian a rah tla tur chelh nan leh a rah than that nan te leh a rah keh tur lakah t a veng thei ani.
BANANA			
STAR FRUIT			
PLUM AND PEACH			
		Gummosis, citrus canker, citrus greening and Dieback	Temperture hniam lutuk leh hnawng vang hian natna a a tam duh a . Soil bome natna laka vennan Bordeaux past hi thing zar leh a trangah te hnawih tur ani.
		Fruit fly	Huan zau takah chuan a par tan tirh leh a rah tan tirin chawlhkar hnih chhung chu heng te hian enkawl tur ani: carbaryl 0.2 percent emaw malathion 0.15 percent suspension containing sugar or jeggery at 10 g/l.
PLANTATION CROP			
COFFEE	All stages		Nursery stage - Thlai chi thlak hma in <i>Azospirillum</i> leh <i>Phosphobacterium</i> a enkawl tur ani. - A chi hi December – January ah hmun zawl/rualrem 1.5 - 2.5 cm a in hlatin tlar mumal tak siam in chin tur ani. - Chuan a chi chu lei tlem te a chhilh a buhpawla khuh tur ani. - Nitin tui pek tur ani a, a sat lutuka loh nan niin a chhun loh nan zar hlih tur ani. - Ni 45 hnu velah a tiak thin a,chu chu bag ah an sawn chhuak leh thin ani.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			Harvesting Stage - Coffe rah hmin hi thlasik lain an seng thin a ni. - A rah hmin tha lo ho chu nuai sawm hmain an thliar hrang leh vek thin ani.
		Coffee Berry borer	- A hun takah leh fimkhur taka seng tur ani. - Hmaih neih nuaih loh tur ani. - Sneg hmaih te lakkhawma paih vek tur ani. - Seng bang zawng zawng chu uluka taka paih fai vek tur ani. - A thlai vennan a rah tlai ho chu paih vek tur ani. - Hmun dam lutukah dah loh tur. - Boruak tha taka a hmuh theih nan leh a rawn chawr no theih nan thlai chu uluk tak a hlawi tur ani. - Chuan hei hian kah tur ani. Quinalphos 25 EC @ 340 ml/200 lit emaw lamda cyhalothrin 5 EC 120 – 160 ml / 200 lit. - In leh loa sawngbawl a a ro dan tur tawkah chuan: Arabica / robusta parchment 10 %, Arabica cherry 10.5 % leh robusta cherry 11.0 %.
		Coffee Rust	- Natna hrik in a khawih tawh a hnah leh a dang te paihpaih vek tur ani. - Bordeaux mixture 0.5% in February - March (Pre-bloom) a kah phawt a, Oxycarboxin 0.03% in May - June (Pre-monsoon) ah kah chhunzawm tur ani.
CEREALS AND PULSE CROPS			
Rabi Maize	A chin hun		- Vaimim chinna tur atan lei kan let phut darh anga. - Hei hian a rawn to chhuah na tur atan a pui dawn a ni. - A chi chu kan lei leh saah chuan kan dah ang. - A chi chu Thiram @4 g/kg seed hian ak sawngbawl hmasa anga. 20-25 kg/ha vel a chi thlak hi a tawk vel viau ani. - A chi tuh hma in lei leh FYM @5-10 t/ha leh 80:60:40 kg N, P₂O₅ leh



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			K ₂ O/ha pawlh chu hman phawt tur a ni. Nitrogen dose chanve chu a chi tuh hunlaia hman tur a ni a, tichuan a bang 25% chu thla khat hnu ah ani ang a adang leh 25% chu a par hunah hman tur a ni.
Soybean, pea, lentil toria, breen gram and black gram cultivation in rice fellow	All stage	Zero tillage	- A than a that theih nan nikhat danah tui pek thin tur ani. - Lei rih vur hian thlai kung te a veng ve ani. - Thlasik lai a lei khoro lutuk tur ven nan a chungah hnim leh thildanga khuh tur ani.
Potato	Sowing stage		- Muangchang loving alu chin na tur chu buatsaih vat tur ani. - Hei hian a than hun lain natna hrikin lakah a veng dawn ani. - Lei leh hmain a hmun hma chu fai taka thlawh hmasak tur ani. - A chi thlak hma in a chi chu en fiah hmasak tur ani. - A than a that theih nan nikhat danah tui pek thin tur ani.
VEGETABLE CROP			
Tomato	Bacterial Blight disease		- Tomato bikah chuan sik leh sa hi natna an kaina tlang lawn ber ani . - Hmun hnawng leh ni hmu lo lutuk hmunah chuan natna an kai hma bik ani. - Tomato hi a uai a, a thih mai loh nan Ridomil emaw Indofil emaw Mancozeb @ 2 gm hi tui liter 1 ah pawlh a kah tur ani .
Early Cole crop	Black spot disease		- A than a that theih nan nikhat danah tui pek thin tur ani. - Thlai bul vawn hnawn nana thlai bula hnim ring vawm khawm hi tui pek zawhah dah tur ani. - Zikhlum lam chi ah chuan sik leh sa vangin a hnah ah thil dum rawn



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			awm thin a , hei hi natna tlanglawn ber ani. Thlai hna lam chi leh zikhlum lam chi reng reng enkawl nan Mancozeb @ 2gm ah tui leter 1 pawlha kah tur ani.
Onion and capsicum	Nursery stage	Poly house	- A than a that theih nan nikhat danah tui pek thin tur ani. - Thlai bul vawn hnawn nana thlai bula hnim ring vawm khawm hi tui pek zawhah dah tur ani. - Thlai chhina hmun (nursery) hi hnim a to loh nan Pendimethalin @ 3.5ml hi tui liter 1 zelah pawlh a kah hi a tha hle ani.
		Phytophthora blight	- A chi ven that nan thiram 3g/kg seed emaw Trichoderma viride 4g+ metalaxyl 4g (Apron)/ kg seed hi a tha hle ani - Hneh taka 1% Bordeaux chawhpawlh emaw 2 g captan emaw 3 copper oxychloride a tui liter 1 hi 10-15 DAS a pek hi a tha hle ani.
French bean	Sowing stage		- Tui pek a hnihnah hringa khuh tur ani a. than a that theih nan tui pek hma in lei rin pan hmasak tur ani. - A than duna theih nan leh hnim to loh na turin a kung bulah lei vur chhoh zel tur ani.
Carrot and radish	Sowing stage		- A than a that theih nan nikhat danah tui pek thin tur ani. - Tui pek hnuah thlai bul vawn hnawn na tur siam tur ani. - Zikhlum lam chi ah chuan sik leh sa vangin a hnah ah thil dum a rawn awm thina, hei hi natna tlanglawn ber ani. - Thlai hna lam chi leh zikhlum lam chi reng reng enkawl nan Mancozeb @ 2gm ah tui leter 1 pawlha kah tur ani.






GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



ANIMAL HUSBANDRY			
Pig	All stages		- Khua a vawh hian vawk hian an mahni in tih lumna tur atan chakna an mamawhna a sang bik ani. - An hriselna that leh that loh enfiah renga, a chaw ei tur tlem tlema tih tam hret hret tur ani.. - Sangha tel ah hian omega-3 hi atam em a vanging an chakna muangchanga a in siam chhoh zel theih nan a tha hle ani.
			Porcine Reproductive Respiratory Syndrome (PRRS). 1. Vawknote emaw vawk lak hran.
	Adult stage		Swine fever. 2. SF vaccines hi thla 2 hnua pek tur ani a, chumi hnuah chuan kumtin thlaruk danah pek chhonzawm tur ani.
Cattle	All age group		Hun rei tak khua a ro avanga hnim hnah hring peh tur a awm loh laia bawngin an chaw ei in buk tawk tur leh an taksa tana mamawh tur atan buh kung urea molasses hmanga sawngbawl pek tur ani.
	All age group		Kar 16 hnuah FMD vaccine pek a, chuan thla tin thla 6 chhung chhonzawm tur ani.
	Young stage		- Black Quarter Vaccine (BQV). - Vaccinne hmasa ber hi thla 6 ah emaw a hnu lamah pek tur. - Chumi hnuah chuan Vaccine hi kum tin pek tur ani.
Poultry	Litter management		- Ar te hian hmun thawl nuam tawk, chaw tha an mamawh tawk leh tui thianghlim an mamawh tawk an hmu tur ani a. - An hriselna atan enkawltha tha tawk tak pek hian natna an kai mai theih tur lak atang a venna tha ber ani.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			- Tui an in tur chhawpna tur tha /lian tha tak leh tui thianghlim tak pek tur ani. - Chaw a hmuar/thing pek loh tur ani a, an chaw eitur thlak sak thut loh tur ani.
	Preventive measures	0-3rd week	Ranikhet Disease- an pian atanga ni 1-6 ah F1 vaccine pek tur ani a, chuan a puitlingh chuan R₂B vaccine pek tur ani. - B complex with antibodies
		4th weeks	Coccidiosis- Amprolium or coccidiostat
		4-5th Weeks	Calcium tonic fortified with B₁₂
FISHERY			
	Pond preparation (Dil buatsaih)	3rd-4th weeks	Dil a leitha hman hian sangha chaw kan tih mai planktons insiam nan a tanpui thin.
			- Bawngkek hring 10 tonnes/ha/year vel dil ah hman thin a ni a; bawngkek kumkhat a kan mamawh zat hmunthum a then a hmunkhat hi dil buatsaih nan hman tur ani. A bak zawng hi tui boruak a zirin semdarh a hman thin tur ani. - Single super phosphate hi dil hectare khat zel a zauah kg 250 ang a hman thin tur ani.
			- Leitha kan hman atang a karkhat hnuah tui rawng a tangin sangha chaw inseam that leh thatloh a hriat theih a. Tui rawng eng hring deuh nghalh ah hian sangha chaw planktons te an inseam tha ang a ngaih ani. - Tui nut zawng tehma transparency pawh 30-40 cm vel ani tur ani.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD,
Guwahati)



Expert committee members:

Dr. S.B. Singh	:	Joint Director	basantasinghsoibam@rediffmail.com
Dr. Saurav Saha	:	Scientist (Agril. Physics)	sauravs.saha@gmail.com
Dr. T. Boopathi	:	Scientist (Agril Entomology)	boopathiars@gmail.com
Dr. A. Ratankumar Singh	:	Scientist (Plant Pathology)	ratantplantpatho@gmail.com
Dr. Lungmuana	:	Scientist (Soil Fertility)	lmsingson@gmail.com
Mr. P.L. Lalrinsanga	:	Scientist (Aquaculture)	viensky2@gmail.com
Dr. Dr. V. Dayal	:	Scientist (Horticulture)	Vishambhai5009@gmail.com
Dr. Samuel Lalliansanga	:	Head & Sr. Scientist	samuelpachua10@gmail.com
Mr. Samik Chowdhury	:	Technical Officer	samikchowdhury33@gmail.com
Mr. Diktea chenkuai	:	Project Assistant	dikteachenkuaiboy@gmail.com

Collaborating Department:

Name of the KVK	Programme Coordinator Name and Designation	KVK Email Id	Phone no/ Mobile no
KVK Lunglei	Dr. Lalmuanzovi Head & Sr. Scientist	kvknhathial@gmail.com	9862803750 9436154614
KVK, Kolasib	Mr. Lalrosamga Khiangte Head & Sr. Scientist	kvkkolasib@gmail.com	9436152440
KVK, Serchhip	Mr. K. Laltlanmawia Head & Sr. Scientist	kvkserchhip@gmail.com	9436146115 9615389293
KVK, Champhai	Mrs. Lalrinawmi Renthlei Head & Sr. Scientist	kvkchawzawl@gmail.com	9436159788
KVK, Lawngtlai	Dr. Michel Lallawmkimi Head & Sr. Scientist	kvklawntlai@gmail.com	9436155858
KVK, Saiha	Dr. Vanlalhraia Hnampe Head & Sr. Scientist	kvksaiha@gmail.com	8974656509
KVK, Mamit	Dr. Samuel Lalliansanga Head & Sr. Scientist	kvkmamit@gmail.com	9436147625
KVK, Aizawl	Dr. K. P. Chaudhary Head & Sr. Scientist	Kpchy@rediffmail.com kvkaizawl@rediffmail.com	9436351669



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
 Mizoram Centre, Kolasib- 796081, MIZORAM
(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



District: Kolasib

Period: 03 May – 07 May, 2017

Bulletin No: - 697/2017/ Bulletin/English

Date of issue: 02nd May, 2017

Parameters	03.05.2017	04.05.2017	05.05.2017	06.05.2017	07.05.2017
Rainfall (mm)	11	0	0	10	9
Max Temp (°C)	31	32	32	32	32
Min Temp (°C)	21	21	20	20	21
Cloud Coverage	Partially clear	Partially clear	Partially clear	Partially clear	Partially clear
Max RH (%)	97	92	89	94	96
Min RH (%)	46	49	45	38	43
Wind Speed (Kmph)	4	4	4	5	5
*Wind Direction	S-E	S-E	S-E	S-E	S

**Northerly- N, North-Easterly- N-E, Easterly- E, South-Easterly- S-E,
 Southerly- S, South-Westerly- S-W, Westerly- W, North-westerly- N-W.**

STATUS OF MONSOON- June 1-30, 2016 (Percent of deviation from normal in parenthesis)

Aizawl- 384.87mm (430.2mm)	Champhai- 105.48mm (359.89mm)	Saiha- 307.40 mm (507.7mm)	Kolasib- 236.00mm (428.1mm)
Lawngtlai-291.20mm (453.1mm)	Lunglei-326.00mm (465.14mm)	Mamit-204.87mm (442.80mm)	Serchhip-411.72mm (259.62mm)

Weather summary of the past three days

Maximum Tem. (°C):28-30°C
Minimum Tem. (°C):16-22°C
Maximum RH (%):94-99%
Minimum RH (%):61-91%
Wind Direction: southeasterly
Cloud cover: Mainly clear
Wind speed: 2-3 km/hr


Rainfall: 18.3 mm

Weather forecast valid from 03rd May, 2017 To 07thMay, 2017.

There are chances of moderate to light rainfall during the next 3 days. The maximum and minimum temperatures for the next 5 days may range for 31-32°C and 20-21°C. Maximum relative humidity is expected in the range of 89-97% and minimum may from 38-49%. Wind direction would be southeasterly to southerly with the wind speed of 4-5 km per hour. Partially clear sky will prevail during the next five days.

Weekly cumulative rainfall: 30.0 mm

NDVI for Mizoram



Moderately wet mildly dry/mildly wet conditions



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



Main Crop / Animal / Fisheries	Stage	Cultural practices/ Pest/ Diseases	Agricultural / Horticultural/ animal husbandry advisories
FRUITS CROPS			
KHASI MANDARIN AND ACID LIME	Seedling transplanting stage		- Fruit plant should be planted in a sunny and wind-protected area. - In the citrus belt, trees can be planted at any time, however, spring is the best time for container grown plants. - Standard-size trees should be spaced 12 to 25 feet apart and dwarf trees should be set 6 to 10 feet apart. The exact distance depends on the variety. The bigger the fruit, the farther the distance. - If the soil is not well-drained, plant the trees on a slight mound to prevent water logging. - Mulching should be done with dry grasses near the tree base to conserve soil moisture during winter. - The young fruit plant must be irrigated at weekly interval for better establishment.
BANANA			
STAR FRUIT			
PLUM AND PEACH			
		Gummosis, citrus Canker, Citrus greening, Dieback, Lamon butterfly and leaf minor	Die back- Due to low temperature and humidity disease appearance will more. Use Bordeaux past in tree trunk, twigs and branches protect healthy plant from soil borne disease. Lamon butterfly- Spray monocrotophos @0.04% @1.2 ml/lt of water. Leaf minor- Spray confidor 0.05% (0.5 ml/lit of water) at each flush emergence. Citrus Canker- Apply bacterimycin @0.6 g/lt of water.
Passion Fruit	Transplanting stage		- High yielding mother vine with good quality fruits and free of virus diseases should be selected to provide cuttings. - A cutting should contain at least 3 buds and must be planted in sand beds.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			- Immediately after planting these should be kept inside a high humid chamber made out of bamboo and polythene. Grafting: - The root stock of yellow Passion fruit is planted in polythene sleeves and the section from Rahangala hybrid is grafted using wedge or approach method of grafting.
Pineapple	Flowering stage		- Apply flowering inducing chemical (Ethrel 10 PPM+2% urea+0.04% Sodium Carbonate) should be applied in the heart of the plant. In evening and only when plants have at least 32 leaves. - The flowering emergence will come out after 55-60 days after chemical spraying. - Apply split doses of fertilizer @ 60:50:60 g per plant. - Remove all unwanted leaves, branches and weed near to the plant.
PLANTATION CROP			
COFFEE	Blooming stage		- If day temperature and prolong dry spell occur it lead to Floral abnormalities like "Star Flower" in Arabica and "Pink Flower" in Robusta. - Irrigation of plants at alternate day's interval, 6 weeks before harvesting improves fruit retention and fruit development, and reduces fruit cracking and improves the fruit quality. - Mulching should be done with dry grasses near the tree base to conserve soil moisture during winter. - The young fruit plant must be irrigated at weekly interval for better establishment. - Foliar application of Mepiquat chloride @ 1000 PPM concentration or 0.75% SSP @ 1.5 g per 200 lt of water 15 days




GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



Rubber	Nursery stage		interval. ✚ Clearing operation may be done during the month of February to April. ✚ Make fire line to protect the young tree and seedlings. ✚ 10-12 kg of well rotten organic manure and 225 gm rock phosphate should be apply at time of planting to each pit as basal dose application.
CEREALS AND PULSE CROPS			
Pre Kharif Rice	Nursery stage		✚ Use only Well filled and healthy seeds. ✚ Put the seed in 2.5% salt solution i.e 250 g of common salt in 10 lts of water. ✚ Seed treated with Bavistin 50 WP @ 0.1% (2 g/ltr) solution. ✚ Mulching is requiring for better germination in nursery.
Jhum Rice	Sowing stage		✚ Sowing in ridge and furrow. ✚ Mulching with grass for better germination. ✚ Intercropping with greengram and bunchy type frenchbean.
Maize (Jhum)	Sowing stage		✚ Seed should be treated with Thiram @4 g/kg seed. ✚ Use optimum seed rate (20-25 kg/ha) for desire plant population. ✚ Apply well decomposed FYM/pig manure @ 5-10 t/ha along with 80:60:40 kg N, P₂O₅ and K₂O/ha incorporate with soil before sowing. Half nitrogen dose will use at the time of sowing and remaining 25% after one month and 25% at flowering stage.
Rabi Maize	vegetative stage		✚ Light irrigation on every week may be given for better establishment and smooth growth. ✚ Earthing up soil near to plant for better support. ✚ Maize rust disease will prevail due to high relative humidity with low temperature. Apply Mancozeb Kg/ha



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			for effective control. ✚ Remove the alternate host <i>Oxalis comiculata</i> .
VEGETABLE CROP			
Cowpea	Sowing stage	KOLASIB	✚ Plough the field properly, at least 2-3 times. ✚ Mix fertilizer with FYM 50:60:60Kg/ha. ✚ Sow 2-3 seed per whole. ✚ Spacing should be 30 X 20 cm.
Okra	Sowing stage	MAMIT Weeding and light irrigation in nursery bed. Provide irrigation in transplanted okra field.	✚ Plough the field with the help of spade. ✚ Sow 2 seed 45 X 45 cm spacing. ✚ Before sowing seed provide one or two irrigation. ✚ Provide fertilizer @ 120: 60: 60 Kg/ha
Ginger and turmeric	Sowing stage	SERCHHIP	✚ Rhizome should be treated with Thiram @4 g/kg seed. ✚ Use optimum seed rate (50-60 kg/ha) for desire plant population. ✚ Apply well decomposed FYM/ pig manure @ 10-20 t/ha along with 120:80:60 kg N, P₂O₅ and K₂O/ha incorporate with soil before sowing. Half nitrogen dose will use at the time of sowing and remaining 25% after one month and 25% at flowering stage.
Colocasia	Sowing stage	LUNGLEI	✚ Planting is done well prepared land or pits filled up with FYM (12-15) t/ha ✚ Sprouted corms or cormels are planted 5-7 deep at a spacing of 40-50 cm between and within rows in the pits. ✚ Inorganic fertilizer like Urea, SSP and MOP @ 220: 375: 134 kg.
ANIMAL HUSBANDRY			
Pig	All stages	LAWNGTIA, SAIHA	✚ As the weather gets colder, your pigs' energy requirement will increase, as they need more energy to keep warm. ✚ Regularly monitor their level of 'fitness'



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			and increase their feed intake to maintain. ✚ Fish oils are excellent for providing slow-release energy with the added advantage of a high level of omega-3.
		Porcine Reproductive Respiratory Syndrome (PRRS).	1. Culling of positive pigs or piglets.
	Adult stage	Swine fever.	2. Vaccination of pigs with SF vaccines at 2 months and yearly interval/6 month interval
Cattle	All age group		Due to prolong dry spell there is a shortage of green grass in the field. For balanced diet and nutrition to your cattle, provide urea molasses treated paddy straw.
	All age group	Foot and Mouth Disease (FMD)	FMD vaccine at 16 week and repeat every 6 month.
	Young stage	Black Quarter (BQ)	- Black Quarter Vaccine (BQV). ✦ Primary vaccination 6 month or above ✦ Revaccination annually
Poultry	Litter management		✚ Birds require adequate space, sufficient feed to meet their nutritional requirements and an adequate supply of good-quality water. ✚ Good management and sanitation are the best ways to avoid infectious disease in poultry. ✚ Provide ample quantity of clean drinking water. ✚ Avoid feeding of mouldy feed. Don't make sudden changes in feed
	Preventive measures	0-3rd week	✚ Ranikhet Disease- F1 vaccine at (1-6) days of birth and R₂B vaccine for adult birds. ✚ B complex with antibodies
		4th weeks	✚ Coccidiosis- Amprolium or coccidiostat



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



		4-5 th Weeks	✚ Calcium tonic fortified with B ₁₂
FISHERY			
	Pond preparation	3 rd -4 th weeks	✚ Application of fertilizers/manure helps in development of plankton which serve as natural feed for the fishes.
		KOLASIB	✚ Raw cowdung should be applied in the pond at the rate of 10 tonnes/ha/year. One third of the total dose should be applied initially and the rest may be applied in a spilt doses. ✚ Single super phosphate should also be applied at the rate of 250 kg/ha in the pond.
		AIZAWL	✚ After one week of application development of planktons could be observed in the pond depending on the colour of the water. Yellowish green colour is an indicator of the good plankton development. ✚ Transparency of the water needs to be maintained at 30-40 cm.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD,
Guwahati)



Expert committee members:

Dr. S.B. Singh	:	Joint Director	basantasinghsoibam@rediffmail.com
Dr. Saurav Saha	:	Scientist (Agril. Physics)	sauravs.saha@gmail.com
Dr. T. Boopathi	:	Scientist (Agril Entomology)	boopathiars@gmail.com
Dr. A. Ratankumar Singh	:	Scientist (Plant Pathology)	ratantplantpatho@gmail.com
Dr. Lungmuana	:	Scientist (Soil Fertility)	lmsingson@gmail.com
Mr. P.L. Lalrinsanga	:	Scientist (Aquaculture)	viensky2@gmail.com
Dr. Dr. V. Dayal	:	Scientist (Horticulture)	Vishambhai5009@gmail.com
Dr. Samuel Lalliansanga	:	Head & Sr. Scientist	samuelpachua10@gmail.com
Mr. Samik Chowdhury	:	Technical Officer	samikchowdhury33@gmail.com
Mr. Diktea chenkuai	:	Project Assistant	dikteachenkuaiboy@gmail.com

Collaborating Department:

Name of the KVK	Programme Coordinator Name and Designation	KVK Email Id	Phone no/ Mobile no
KVK Lunglei	Dr. Lalmuanzovi Head & Sr. Scientist	kvkhnahthial@gmail.com	9862803750 9436154614
KVK, Kolasib	Mr. Lalrosamga Khiangte Head & Sr. Scientist	kvkkolasib@gmail.com	9436152440
KVK, Serchhip	Mr. K. Laltlanmawia Head & Sr. Scientist	kvkserchhip@gmail.com	9436146115 9615389293
KVK, Champhai	Mrs. Lalrinawmi Renthlei Head & Sr. Scientist	kvkchawzawl@gmail.com	9436159788
KVK, Lawngtlai	Dr. Michel Lallawmkimi Head & Sr. Scientist	kvklawntlai@gmail.com	9436155858
KVK, Saiha	Dr. Vanlalhraia Hnampe Head & Sr. Scientist	kvksaiha@gmail.com	8974656509
KVK, Mamit	Dr. Samuel Lalliansanga Head & Sr. Scientist	kvkmamit@gmail.com	9436147625
KVK, Aizawl	Dr. K. P. Chaudhary Head & Sr. Scientist	Kpchy@rediffmail.com kvkaizawl@rediffmail.com	9436351669



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
 Mizoram Centre, Kolasib- 796081, MIZORAM
(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



District: Kolasib

Period: 03 May – 07 May, 2017

Bulletin No: - 697/2017/ Bulletin/Mizo

Date of issue: 02nd May, 2017

Parameters	03.05.2017	04.05.2017	05.05.2017	06.05.2017	07.05.2017
Rainfall (mm)	4	5	6	5	3
Max Temp (°C)	30	30	31	31	31
Min Temp (°C)	21	20	20	21	21
Cloud Coverage	Partially clear	Partially clear	Partially clear	Partially clear	Partially clear
Max RH (%)	98	99	99	98	99
Min RH (%)	73	61	65	50	52
Wind Speed (Kmph)	4	2	3	2	3
*Wind Direction	S	S	S	S-E	S-E

**Northerly- N, North-Easterly- N-E, Easterly- E, South-Easterly- S-E,
 Southerly- S, South-Westerly- S-W, Westerly- W, North-westerly- N-W.**

STATUS OF MONSOON- June 1-30, 2016 (Percent of deviation from normal in parenthesis)

Aizawl- 384.87mm (430.2mm)	Champhai- 105.48mm (359.89mm)	Saiha- 307.40 mm (507.7mm)	Kolasib- 236.00mm (428.1mm)
Lawngtlai-291.20mm (453.1mm)	Lunglei-326.00mm (465.14mm)	Mamit-204.87mm (442.80mm)	Serchhip-411.72mm (259.62mm)

Weather summary of the past three days

Maximum Tem. (°C):28-30°C
Minimum Tem. (°C):16-22°C
Maximum RH (%):94-99%
Minimum RH (%):61-91%
Wind Direction: southeasterly
Cloud cover: Mainly clear
Wind speed: 2-3 km/hr


Rainfall: 18.3 mm

**03rd May– 07th May, 2017 chhunga sik leh sa
 dinhmun tur tlangpui**

Tun ni 3 chhung lo awm turah hian ruahtui tla miahlo tura beisei a ni. Khua a lum lai berin 31-32°C a ni ang a. A vawh lai ber in 20-21°C ni tura beisei a ni. RH san lai berin 89-97% leh a hniam lai berin 38-49% ni tur a rin niin. Thli hi darkar khatah 4-5 km vela chakin chhaklam awi zawngin a tleh rin a ni. A tlangpuiin tun ni nga chung hian khawthiang tak hmuh beisei a ni.

Weekly cumulative rainfall: 30.0mm

NDVI for Mizoram



Moderately wet mildly dry/mildly wet conditions



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



Main Crop / Animal / Fisheries	Stage	Cultural practices/ Pest/ Diseases	Agricultural / Horticultural/ animal husbandry advisories
FRUITS CROPS			
KHASI MANDARIN AND ACID LIME	A kui atanga a seng hun		- Thlasik laia thlai bul khoro lutuk tur vennan chuan hnim hnah hring tlai bul velah dahkhawm tur ani. - Thlai naupang deuah chuan chawlh kar tin a tui pek thin tur ani. - Leia tha mamawh tawh a hmuh theihna turin a hmunhma a hnim awm te thlawhfai thin tur ani. - A seng hma kar 6 chhung chu tui tha taka pek hian a rah tla tur chelh nan leh a rah than that nan te leh a rah keh tur lakah t a veng thei ani.
BANANA			
STAR FRUIT			
PLUM AND PEACH			
		Gummosis, citrus canker, citrus greening and Dieback	Temperture hniam lutuk leh hnawng vang hian natna a a tam duh a . Soil bome natna laka vennan Bordeaux past hi thing zar leh a trangah te hnawih tur ani.
		Fruit fly	Huan zau takah chuan a par tan tirh leh a rah tan tirin chawlhkar hnih chhung chu heng te hian enkawl tur ani: carbaryl 0.2 percent emaw malathion 0.15 percent suspension containing sugar or jeggery at 10 g/l.
PLANTATION CROP			
COFFEE	All stages		Nursery stage - Thlai chi thlak hma in <i>Azospirillum</i> leh <i>Phosphobacterium</i> a enkawl tur ani. - A chi hi December – January ah hmun zawl/rualrem 1.5 - 2.5 cm a in hlatin tlar mumal tak siam in chin tur ani. - Chuan a chi chu lei tlem te a chhilh a buhpawla khuh tur ani. - Nitin tui pek tur ani a, a sat lutuka loh nan niin a chhun loh nan zar hlih tur ani. - Ni 45 hnu velah a tiak thin a,chu chu bag ah an sawn chhuak leh thin ani.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			Harvesting Stage - Coffe rah hmin hi thlasik lain an seng thin a ni. - A rah hmin tha lo ho chu nuai sawm hmain an thliar hrang leh vek thin ani.
		Coffee Berry borer	- A hun takah leh fimkhur taka seng tur ani. - Hmaih neih nuaih loh tur ani. - Sneg hmaih te lakkhawma paih vek tur ani. - Seng bang zawng zawng chu uluka taka paih fai vek tur ani. - A thlai vennan a rah tlai ho chu paih vek tur ani. - Hmun dam lutukah dah loh tur. - Boruak tha taka a hmuh theih nan leh a rawn chawr no theih nan thlai chu uluk tak a hlawi tur ani. - Chuan hei hian kah tur ani. Quinalphos 25 EC @ 340 ml/200 lit emaw lamda cyhalothrin 5 EC 120 – 160 ml / 200 lit. - In leh loa sawngbawln a a ro dan tur tawkah chuan: Arabica / robusta parchment 10 %, Arabica cherry 10.5 % leh robusta cherry 11.0 %.
		Coffee Rust	- Natna hrik in a khawih tawh a hnah leh a dang te paihpaih vek tur ani. - Bordeaux mixture 0.5% in February - March (Pre-bloom) a kah phawt a, Oxycarboxin 0.03% in May - June (Pre-monsoon) ah kah chhunzawm tur ani.
CEREALS AND PULSE CROPS			
Rabi Maize	A chin hun		- Vaimim chinna tur atan lei kan let phut darh anga. - Hei hian a rawn to chhuah na tur atan a pui dawn a ni. - A chi chu kan lei leh saah chuan kan dah ang. - A chi chu Thiram @4 g/kg seed hian ak sawngbawl hmasa anga. 20-25 kg/ha vel a chi thlak hi a tawk vel viau ani. - A chi tuh hma in lei leh FYM @5-10 t/ha leh 80:60:40 kg N, P₂O₅ leh



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			K ₂ O/ha pawlh chu hman phawt tur a ni. Nitrogen dose chanve chu a chi tuh hunlaia hman tur a ni a, tichuan a bang 25% chu thla khat hnu ah ani ang a adang leh 25% chu a par hunah hman tur a ni.
Soybean, pea, lentil toria, breen gram and black gram cultivation in rice fellow	All stage	Zero tillage	- A than a that theih nan nikhat danah tui pek thin tur ani. - Lei rih vur hian thlai kung te a veng ve ani. - Thlasik lai a lei khoro lutuk tur ven nan a chungah hnim leh thildanga khuh tur ani.
Potato	Sowing stage		- Muangchang loving alu chin na tur chu buatsaih vat tur ani. - Hei hian a than hun lain natna hrikin lakah a veng dawn ani. - Lei leh hmain a hmun hma chu fai taka thlawh hmasak tur ani. - A chi thlak hma in a chi chu en fiah hmasak tur ani. - A than a that theih nan nikhat danah tui pek thin tur ani.
VEGETABLE CROP			
Tomato	Bacterial Blight disease		- Tomato bikah chuan sik leh sa hi natna an kaina tlang lawn ber ani . - Hmun hnawng leh ni hmu lo lutuk hmunah chuan natna an kai hma bik ani. - Tomato hi a uai a, a thih mai loh nan Ridomil emaw Indofil emaw Mancozeb @ 2 gm hi tui liter 1 ah pawlh a kah tur ani .
Early Cole crop	Black spot disease		- A than a that theih nan nikhat danah tui pek thin tur ani. - Thlai bul vawn hnawn nana thlai bula hnim ring vawm khawm hi tui pek zawhah dah tur ani. - Zikhlum lam chi ah chuan sik leh sa vangin a hnah ah thil dum rawn



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			awm thin a , hei hi natna tlanglawn ber ani. Thlai hna lam chi leh zikhlum lam chi reng reng enkawl nan Mancozeb @ 2gm ah tui leter 1 pawlha kah tur ani.
Onion and capsicum	Nursery stage	Poly house	- A than a that theih nan nikhat danah tui pek thin tur ani. - Thlai bul vawn hnawn nana thlai bula hnim ring vawm khawm hi tui pek zawhah dah tur ani. - Thlai chhina hmun (nursery) hi hnim a to loh nan Pendimethalin @ 3.5ml hi tui liter 1 zelah pawlh a kah hi a tha hle ani.
		Phytophthora blight	- A chi ven that nan thiram 3g/kg seed emaw Trichoderma viride 4g+ metalaxyl 4g (Apron)/ kg seed hi a tha hle ani - Hneh taka 1% Bordeaux chawhpawlh emaw 2 g captan emaw 3 copper oxychloride a tui liter 1 hi 10-15 DAS a pek hi a tha hle ani.
French bean	Sowing stage		- Tui pek a hnihnah hringa khuh tur ani a. than a that theih nan tui pek hma in lei rin pan hmasak tur ani. - A than duna theih nan leh hnim to loh na turin a kung bulah lei vur chhoh zel tur ani.
Carrot and radish	Sowing stage		- A than a that theih nan nikhat danah tui pek thin tur ani. - Tui pek hnuah thlai bul vawn hnawn na tur siam tur ani. - Zikhlum lam chi ah chuan sik leh sa vangin a hnah ah thil dum a rawn awm thina, hei hi natna tlanglawn ber ani. - Thlai hna lam chi leh zikhlum lam chi reng reng enkawl nan Mancozeb @ 2gm ah tui leter 1 pawlha kah tur ani.






GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



ANIMAL HUSBANDRY			
Pig	All stages		- Khua a vawh hian vawk hian an mahni in tih lumna tur atan chakna an mamawhna a sang bik ani. - An hriselna that leh that loh enfiah renga, a chaw ei tur tlem tlema tih tam hret hret tur ani.. - Sangha tel ah hian omega-3 hi atam em a vangin an chakna muangchanga a in siam chhoh zel theih nan a tha hle ani.
			Porcine Reproductive Respiratory Syndrome (PRRS). 1. Vawknote emaw vawk lak hran.
	Adult stage		Swine fever. 2. SF vaccines hi thla 2 hnua pek tur ani a, chumi hnuah chuan kumtin thlaruk danah pek chhonzawm tur ani.
Cattle	All age group		Hun rei tak khua a ro avanga hnim hnah hring peh tur a awm loh laia bawngin an chaw ei in buk tawk tur leh an taksa tana mamawh tur atan buh kung urea molasses hmanga sawngbawl pek tur ani.
	All age group		Kar 16 hnuah FMD vaccine pek a, chuan thla tin thla 6 chhung chhonzawm tur ani.
	Young stage		- Black Quarter Vaccine (BQV). - Vaccinne hmasa ber hi thla 6 ah emaw a hnu lamah pek tur. - Chumi hnuah chuan Vaccine hi kum tin pek tur ani.
Poultry	Litter management		- Ar te hian hmun thawl nuam tawk, chaw tha an mamawh tawk leh tui thianghlim an mamawh tawk an hmu tur ani a. - An hriselna atan enkawltha tha tawk tak pek hian natna an kai mai theih tur lak atang a venna tha ber ani.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			- Tui an in tur chhawpna tur tha /lian tha tak leh tui thianghlim tak pek tur ani. - Chaw a hmuar/thing pek loh tur ani a, an chaw eitur thlak sak thut loh tur ani.
	Preventive measures	0-3rd week	Ranikhet Disease- an pian atanga ni 1-6 ah F1 vaccine pek tur ani a, chuan a puitlingh chuan R₂B vaccine pek tur ani. - B complex with antibodies
		4th weeks	Coccidiosis- Amprolium or coccidiostat
		4-5th Weeks	Calcium tonic fortified with B₁₂
FISHERY			
	Pond preparation (Dil buatsaih)	3rd-4th weeks	Dil a leitha hman hian sangha chaw kan tih mai planktons insiam nan a tanpui thin.
			- Bawngkek hring 10 tonnes/ha/year vel dil ah hman thin a ni a; bawngkek kumkhat a kan mamawh zat hmunthum a then a hmunkhat hi dil buatsaih nan hman tur ani. A bak zawng hi tui boruak a zirin semdarh a hman thin tur ani. - Single super phosphate hi dil hectare khat zel a zauah kg 250 ang a hman thin tur ani.
			- Leitha kan hman atang a karkhat hnuah tui rawng a tangin sangha chaw inseam that leh thatloh a hriat theih a. Tui rawng eng hring deuh nghalh ah hian sangha chaw planktons te an inseam tha ang a ngaih ani. - Tui nut zawng tehma transparency pawh 30-40 cm vel ani tur ani.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD,
Guwahati)



Expert committee members:

Dr. S.B. Singh	:	Joint Director	basantasinghsoibam@rediffmail.com
Dr. Saurav Saha	:	Scientist (Agril. Physics)	sauravs.saha@gmail.com
Dr. T. Boopathi	:	Scientist (Agril Entomology)	boopathiars@gmail.com
Dr. A. Ratankumar Singh	:	Scientist (Plant Pathology)	ratantplantpatho@gmail.com
Dr. Lungmuana	:	Scientist (Soil Fertility)	lmsingson@gmail.com
Mr. P.L. Lalrinsanga	:	Scientist (Aquaculture)	viensky2@gmail.com
Dr. Dr. V. Dayal	:	Scientist (Horticulture)	Vishambhai5009@gmail.com
Dr. Samuel Lalliansanga	:	Head & Sr. Scientist	samuelpachua10@gmail.com
Mr. Samik Chowdhury	:	Technical Officer	samikchowdhury33@gmail.com
Mr. Diktea chenkuai	:	Project Assistant	dikteachenkuaiboy@gmail.com

Collaborating Department:

Name of the KVK	Programme Coordinator Name and Designation	KVK Email Id	Phone no/ Mobile no
KVK Lunglei	Dr. Lalmuanzovi Head & Sr. Scientist	kvkhnahthial@gmail.com	9862803750 9436154614
KVK, Kolasib	Mr. Lalrosamga Khiangte Head & Sr. Scientist	kvkkolasib@gmail.com	9436152440
KVK, Serchhip	Mr. K. Laltlanmawia Head & Sr. Scientist	kvkserchhip@gmail.com	9436146115 9615389293
KVK, Champhai	Mrs. Lalrinawmi Renthlei Head & Sr. Scientist	kvkchawzawl@gmail.com	9436159788
KVK, Lawngtlai	Dr. Michel Lallawmkimi Head & Sr. Scientist	kvklawntlai@gmail.com	9436155858
KVK, Saiha	Dr. Vanlalhraia Hnampe Head & Sr. Scientist	kvksaiha@gmail.com	8974656509
KVK, Mamit	Dr. Samuel Lalliansanga Head & Sr. Scientist	kvkmamit@gmail.com	9436147625
KVK, Aizawl	Dr. K. P. Chaudhary Head & Sr. Scientist	Kpchy@rediffmail.com kvkaizawl@rediffmail.com	9436351669



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
 Mizoram Centre, Kolasib- 796081, MIZORAM
(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



District: Lawntlai

Period: 03 May – 07 May, 2017

Bulletin No: - 697/2017/ Bulletin/English

Date of issue: 02nd May, 2017

Parameters	03.05.2017	04.05.2017	05.05.2017	06.05.2017	07.05.2017
Rainfall (mm)	0	0	0	0	0
Max Temp (°C)	32	33	35	34	33
Min Temp (°C)	20	22	21	21	21
Cloud Coverage	Mainly clear	Mainly clear	Mainly clear	Mainly clear	Mainly clear
Max RH (%)	97	81	88	83	87
Min RH (%)	35	32	32	30	36
Wind Speed (Kmph)	4	4	4	5	4
*Wind Direction	E	E	S-E	S-E	S-E

**Northerly- N, North-Easterly- N-E, Easterly- E, South-Easterly- S-E,
 Southerly- S, South-Westerly- S-W, Westerly- W, North-westerly- N-W.**

STATUS OF MONSOON- June 1-30, 2016 (Percent of deviation from normal in parenthesis)

Aizawl- 384.87mm (430.2mm)	Champhai- 105.48mm (359.89mm)	Saiha- 307.40 mm (507.7mm)	Kolasib- 236.00mm (428.1mm)
Lawngtlai-291.20mm (453.1mm)	Lunglei-326.00mm (465.14mm)	Mamit-204.87mm (442.80mm)	Serchhip-411.72mm (259.62mm)

Weather summary of the past three days

Maximum Tem. (°C):28-31°C
Minimum Tem. (°C):18-21°C
Maximum RH (%):91-98%
Minimum RH (%):55-82%
Wind Direction: southeasterly
Cloud cover: Mainly clear
Wind speed: 2-3 km/hr


Rainfall: 00.0 mm

Weather forecast valid from 03rd May, 2017 To 07thMay, 2017.

There are no chances of rainfall during the next 5 days. The maximum and minimum temperatures for the next 5 days may range for 32-35°C and 20-22°C. Maximum relative humidity is expected in the range of 81-97% and minimum may from 30-36%. Wind direction would be easterly to southeasterly with the wind speed of 4-5 km per hour. Mainly clear sky will prevail during the next five days.

Weekly cumulative rainfall: 00.0 mm

NDVI for Mizoram



Moderately wet mildly dry/mildly wet conditions



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



Main Crop / Animal / Fisheries	Stage	Cultural practices/ Pest/ Diseases	Agricultural / Horticultural/ animal husbandry advisories
FRUITS CROPS			
KHASI MANDARIN AND ACID LIME	Seedling transplanting stage		- Fruit plant should be planted in a sunny and wind-protected area. - In the citrus belt, trees can be planted at any time, however, spring is the best time for container grown plants. - Standard-size trees should be spaced 12 to 25 feet apart and dwarf trees should be set 6 to 10 feet apart. The exact distance depends on the variety. The bigger the fruit, the farther the distance. - If the soil is not well-drained, plant the trees on a slight mound to prevent water logging. - Mulching should be done with dry grasses near the tree base to conserve soil moisture during winter. - The young fruit plant must be irrigated at weekly interval for better establishment.
BANANA			
STAR FRUIT			
PLUM AND PEACH			
		Gummosis, citrus Canker, Citrus greening, Dieback, Lamon butterfly and leaf minor	Die back- Due to low temperature and humidity disease appearance will more. Use Bordeaux past in tree trunk, twigs and branches protect healthy plant from soil borne disease. Lamon butterfly- Spray monocrotophos @0.04% @1.2 ml/lt of water. Leaf minor- Spray confidor 0.05% (0.5 ml/lit of water) at each flush emergence. Citrus Canker- Apply bacterimycin @0.6 g/lt of water.
Passion Fruit	Transplanting stage		- High yielding mother vine with good quality fruits and free of virus diseases should be selected to provide cuttings. - A cutting should contain at least 3 buds and must be planted in sand beds.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			- Immediately after planting these should be kept inside a high humid chamber made out of bamboo and polythene. Grafting: - The root stock of yellow Passion fruit is planted in polythene sleeves and the section from Rahangala hybrid is grafted using wedge or approach method of grafting.
Pineapple	Flowering stage		- Apply flowering inducing chemical (Ethrel 10 PPM+2% urea+0.04% Sodium Carbonate) should be applied in the heart of the plant. In evening and only when plants have at least 32 leaves. - The flowering emergence will come out after 55-60 days after chemical spraying. - Apply split doses of fertilizer @ 60:50:60 g per plant. - Remove all unwanted leaves, branches and weed near to the plant.
PLANTATION CROP			
COFFEE	Blooming stage		- If day temperature and prolong dry spell occur it lead to Floral abnormalities like "Star Flower" in Arabica and "Pink Flower" in Robusta. - Irrigation of plants at alternate day's interval, 6 weeks before harvesting improves fruit retention and fruit development, and reduces fruit cracking and improves the fruit quality. - Mulching should be done with dry grasses near the tree base to conserve soil moisture during winter. - The young fruit plant must be irrigated at weekly interval for better establishment. - Foliar application of Mepiquat chloride @ 1000 PPM concentration or 0.75% SSP @ 1.5 g per 200 lt of water 15 days




GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



Rubber	Nursery stage		interval. ✚ Clearing operation may be done during the month of February to April. ✚ Make fire line to protect the young tree and seedlings. ✚ 10-12 kg of well rotten organic manure and 225 gm rock phosphate should be apply at time of planting to each pit as basal dose application.
CEREALS AND PULSE CROPS			
Pre Kharif Rice	Nursery stage		✚ Use only Well filled and healthy seeds. ✚ Put the seed in 2.5% salt solution i.e 250 g of common salt in 10 lts of water. ✚ Seed treated with Bavistin 50 WP @ 0.1% (2 g/ltr) solution. ✚ Mulching is requiring for better germination in nursery.
Jhum Rice	Sowing stage		✚ Sowing in ridge and furrow. ✚ Mulching with grass for better germination. ✚ Intercropping with greengram and bunchy type frenchbean.
Maize (Jhum)	Sowing stage		✚ Seed should be treated with Thiram @4 g/kg seed. ✚ Use optimum seed rate (20-25 kg/ha) for desire plant population. ✚ Apply well decomposed FYM/pig manure @ 5-10 t/ha along with 80:60:40 kg N, P₂O₅ and K₂O/ha incorporate with soil before sowing. Half nitrogen dose will use at the time of sowing and remaining 25% after one month and 25% at flowering stage.
Rabi Maize	vegetative stage		✚ Light irrigation on every week may be given for better establishment and smooth growth. ✚ Earthing up soil near to plant for better support. ✚ Maize rust disease will prevail due to high relative humidity with low temperature. Apply Mancozeb Kg/ha



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			for effective control. Remove the alternate host <i>Oxalis comiculata</i>.
VEGETABLE CROP			
Cowpea	Sowing stage	KOLASIB	- Plough the field properly, at least 2-3 times. - Mix fertilizer with FYM 50:60:60Kg/ha. - Sow 2-3 seed per whole. - Spacing should be 30 X 20 cm.
Okra	Sowing stage	Weeding and light irrigation in nursery bed. Provide irrigation in transplanted okra field.	- Plough the field with the help of spade. - Sow 2 seed 45 X 45 cm spacing. - Before sowing seed provide one or two irrigation. - Provide fertilizer @ 120: 60: 60 Kg/ha
Ginger and turmeric	Sowing stage	SERCHHIP	- Rhizome should be treated with Thiram @4 g/kg seed. - Use optimum seed rate (50-60 kg/ha) for desire plant population. - Apply well decomposed FYM/ pig manure @ 10-20 t/ha along with 120:80:60 kg N, P₂O₅ and K₂O/ha incorporate with soil before sowing. Half nitrogen dose will use at the time of sowing and remaining 25% after one month and 25% at flowering stage.
Colocasia	Sowing stage	LUNGLEI	- Planting is done well prepared land or pits filled up with FYM (12-15) t/ha - Sprouted corms or cormels are planted 5-7 deep at a spacing of 40-50 cm between and within rows in the pits. - Inorganic fertilizer like Urea, SSP and MOP @ 220: 375: 134 kg.
ANIMAL HUSBANDRY			
Pig	All stages	LAWNGTIA	- As the weather gets colder, your pigs' energy requirement will increase, as they need more energy to keep warm. - Regularly monitor their level of 'fitness'



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



			and increase their feed intake to maintain. ✚ Fish oils are excellent for providing slow-release energy with the added advantage of a high level of omega-3.
		Porcine Reproductive Respiratory Syndrome (PRRS).	1. Culling of positive pigs or piglets.
	Adult stage	Swine fever.	2. Vaccination of pigs with SF vaccines at 2 months and yearly interval/6 month interval
Cattle	All age group		Due to prolong dry spell there is a shortage of green grass in the field. For balanced diet and nutrition to your cattle, provide urea molasses treated paddy straw.
	All age group	Foot and Mouth Disease (FMD)	FMD vaccine at 16 week and repeat every 6 month.
	Young stage	Black Quarter (BQ)	- Black Quarter Vaccine (BQV). ✦ Primary vaccination 6 month or above ✦ Revaccination annually
Poultry	Litter management		✚ Birds require adequate space, sufficient feed to meet their nutritional requirements and an adequate supply of good-quality water. ✚ Good management and sanitation are the best ways to avoid infectious disease in poultry. ✚ Provide ample quantity of clean drinking water. ✚ Avoid feeding of mouldy feed. Don't make sudden changes in feed
	Preventive measures	0-3rd week	✚ Ranikhet Disease- F1 vaccine at (1-6) days of birth and R₂B vaccine for adult birds. ✚ B complex with antibodies
		4th weeks	✚ Coccidiosis- Amprolium or coccidiostat



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD, Guwahati)



		4-5 th Weeks	✚ Calcium tonic fortified with B ₁₂
FISHERY			
	Pond preparation	3 rd -4 th weeks	✚ Application of fertilizers/manure helps in development of plankton which serve as natural feed for the fishes.
		KOLASIB	✚ Raw cowdung should be applied in the pond at the rate of 10 tonnes/ha/year. One third of the total dose should be applied initially and the rest may be applied in a spilt doses. ✚ Single super phosphate should also be applied at the rate of 250 kg/ha in the pond.
		AIZAWL	✚ After one week of application development of planktons could be observed in the pond depending on the colour of the water. Yellowish green colour is an indicator of the good plankton development. ✚ Transparency of the water needs to be maintained at 30-40 cm.



GRAMIN KRISHI MAUSAM SEWA ICAR RESEARCH COMPLEX FOR NEH REGION

Mizoram Centre, Kolasib- 796081, MIZORAM

(Prepared based on District wise Weather Forecast received from IMD,
Guwahati)



Expert committee members:

Dr. S.B. Singh	:	Joint Director	basantasinghsoibam@rediffmail.com
Dr. Saurav Saha	:	Scientist (Agril. Physics)	sauravs.saha@gmail.com
Dr. T. Boopathi	:	Scientist (Agril Entomology)	boopathiars@gmail.com
Dr. A. Ratankumar Singh	:	Scientist (Plant Pathology)	ratantplantpatho@gmail.com
Dr. Lungmuana	:	Scientist (Soil Fertility)	lmsingson@gmail.com
Mr. P.L. Lalrinsanga	:	Scientist (Aquaculture)	viensky2@gmail.com
Dr. Dr. V. Dayal	:	Scientist (Horticulture)	Vishambhai5009@gmail.com
Dr. Samuel Lalliansanga	:	Head & Sr. Scientist	samuelpachua10@gmail.com
Mr. Samik Chowdhury	:	Technical Officer	samikchowdhury33@gmail.com
Mr. Diktea chenkuai	:	Project Assistant	dikteachenkuaiboy@gmail.com

Collaborating Department:

Name of the KVK	Programme Coordinator Name and Designation	KVK Email Id	Phone no/ Mobile no
KVK Lunglei	Dr. Lalmuanzovi Head & Sr. Scientist	kvknhathial@gmail.com	9862803750 9436154614
KVK, Kolasib	Mr. Lalrosamga Khiangte Head & Sr. Scientist	kvkkolasib@gmail.com	9436152440
KVK, Serchhip	Mr. K. Laltlanmawia Head & Sr. Scientist	kvkserchhip@gmail.com	9436146115 9615389293
KVK, Champhai	Mrs. Lalrinawmi Renthlei Head & Sr. Scientist	kvkchawzawl@gmail.com	9436159788
KVK, Lawngtlai	Dr. Michel Lallawmkimi Head & Sr. Scientist	kvklawntlai@gmail.com	9436155858
KVK, Saiha	Dr. Vanlalhruaia Hnampe Head & Sr. Scientist	kvksaiha@gmail.com	8974656509
KVK, Mamit	Dr. Samuel Lalliansanga Head & Sr. Scientist	kvkmamit@gmail.com	9436147625
KVK, Aizawl	Dr. K. P. Chaudhary Head & Sr. Scientist	Kpchy@rediffmail.com kvkaizawl@rediffmail.com	9436351669