



Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 06-08-2019

Weather Forecast of SAIHA(Mizoram) Issued On : 2019-08-06(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-08-07	16.0	29.0	23.0	99	65	4.9	113	8
2019-08-08	10.0	28.0	24.0	98	92	5.0	110	7
2019-08-09	8.0	29.0	23.0	99	71	5.1	94	5
2019-08-10	14.0	28.0	24.0	100	62	3.0	112	7
2019-08-11	41.0	28.0	24.0	100	80	2.4	207	8

Weather Summary/Alert:

Moderate to heavy rainfall occurred last 3 days. The maximum and minimum temperatures for the last 3 days range for 25-28 C and 16-19 C. Maximum relative humidity was expected in the range of 92-98% and minimum from 62-84%. Wind direction was easterly southeasterly to northeasterly with the wind speed of 2.2-4.6 km per hour. Partially clear sky was prevailed last three days. Weekly cumulative rainfall: 47 mm

Tun nithum (3 days) chhung khan ruahtui vantlang tanga tam lam dawn a ni. Khua a lumlai ber in 25-28 C a ni a, a vawhlai ber erawh chuan 16-19 C a ni. Boruak hnawn lai berin 92-98% a nih lain a hul lai erawh chuan 62-84% a ni thung. Thli hi darker khatah 2.2-4.6 a chak in chhaklam atanga tannin chhim lamah chuan hmarchhak lamah awileh in chhim lamah a tleh thlaleh a ni. Tun nithum chhung khan van ah chum hmuh tur a tam hle. Kartluana ruahtui tlingkhawm: 47 mm

General Advisory:

Possibility of rainfall will be less and temperature will be high. Following general measures should be taken by farmers. Harvest all mature crops and proper sun dry for preservation of seed. Application of fertilizers/manure can be done. Check of vaccination schedule in poultry/livestock.

Ruahtui a tlak tlem rin a nih avangin boruak a lum deuh beisei a ni. Heng a hnuaia enkawl dan tarlan tehi huan neitu tehian zawm a hman theuh tur a ni. A kung puitling tawh techu seng tur, nisa ah phoro a a chi atana dahthat tur a ni. Leitha pek theih a ni. Ran neite tan vaccine pek hun tur (date) ennawn thin ni se.

SMS Advisory:

Pig- Swine fever- Vaccination of pigs @ 1 ml/pig with SF vaccines at 2 months interval.

Vawk natna-Swine fever enkawl dan- SF vaccines chu thla 2 danah vawk pakhat chiu nan 1 ml zel hman thin tur a ni.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
JHUM RICE(LOCAL)	☐ Drainage should be maintain properly ☐ Use split dose of fertilizer @ 40 kg/ha in clear/dry day. ☐ Keep the surrounding clean/ weed free to avoid disease/pest harboring. ☐ Spray tricyclozole 0.6 g/lit of water against leaf blast of rice in clear/dry day. ☐ Tuitling paihchhuahna tha taka siam tur ☐ Khaw that laiin leitha pek tur te chu split dose hmangin hectare khatah kg 40 zel hman tur a ni. ☐ A bul leh a chhehvel te tihfai a hnim hnah emaw rannung emaw natna thlen thei chi lakah enkawl tur a ni. ☐ Buh natna leaf blast laka vennis tricyclozole chu tui litre khatah 0.6 g pawlh chu khaw that laiin kah tur a ni.
MAIZE(LOCAL)	☐ To manage this northern leaf blight disease spray aproach (Methoxy-acrylates) @ 2 g/li of water in clear/ dry day. ☐ Crop rotation with legume. ☐ Drainage should be maintain properly in heavy rainfall ☐ He Northern leaf blight natna enkawl nan hian Methoxy-acrylates gram 2 chu tui litre khatah chawhpawlh chu khaw that laia kah tur a ni. ☐ Thlai (Legume) ho nen chin pawlh ni se. ☐ Ruahtui tling paihchhuahna tur tha taka siam tur.
GREENGRAM(TRC M 151)	☐ Drainage should be maintain properly to avoid water stagnation problem. ☐ Intercrop with maize every 4-5 lines interval. ☐ Keep the surrounding clean/ weed free to avoid disease/pest harboring. ☐ Tuitling paihchhuahna tha taka siam tur. ☐ Vaimim nen line 4-5 dan zelah chin pawlh tur. ☐ A bul leh a chhehvel te tihfai a hnim hnah emaw rannung emaw natna thlen thei chi lakah enkawl tur a ni.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
BHINDI(ARKA ANAMIKA)	☐ Pick tender and immature pods when they are approximately 3-4 inch in length. ☐ Harvest mature pod every alternate days. ☐ Harvest over matured pod for seed production purpose ☐ Seed should be dry properly and keep it for next year. ☐ A kawm (rah) la no leh la changlo techu inches 3-4 inkar an nih in lawh tur. ☐ A kawm (rah) hmin tawh techu ni khat danah lawh thin tur. ☐ A kawm (rah) hmin leh chang techu lawh a a chi atana dah that tur. ☐ A rah lawhsa tawh techu hmun hul ah dah ro a a kum leh atan dah that tur.
CUCURBITACEAE CROP(LOCAL)	☐ Drainage should be maintain properly ☐ Keep the surrounding clean/ weed free to avoid disease/pest harboring. ☐ Harvest all mature fruits and store dry seed for next year ☐ Tuitling paihchhuahna tha taka siam tur. ☐ A bul leh a chhehvel te tihfai a hnim hnah emaw rannung emaw natna thlen thei chi lakah enkawl tur a ni. ☐ A rah hmin tawh techu seng a, pho ro a, a kum leh a tuh atan dahthat tur a ni.

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
COW(LOCAL)	Balanced feeding of dairy animals and allow calf suckling immediately after parturition. Full hand milking should be practiced. Frequent incomplete milking at interval of 4hrs should be practiced after parturition for minimum 1 week. Afterwards complete milking in a day is recommended. A pui in nau a neih hnu ah a innmil tawkin a naute chu hnute tui leh chaw pek tur a ni. A hnute sawr dan thin full hand milking chu hman tur a ni. Nau a neih hnu atangin a loh ber ah kar khat tal chu darkar li danah a hnute chu vawi khata sawr zo vek lo in sawr zeuh zeuh thin tur a ni. Chumi hnu ah chuan a vaia sawr zawh thin chauh tur a ni.
PIG(LOCAL)	The bedding is always. If possible wooden sleeping box should be provided. Provide artificial heat source to new born piglets.

Livestock(Varieties)	Livestock Specific Advisory
	An mut hmun hi a pawimawh hle. A theih hram chuan an mutna chhuat chu thing a siam ni thei se. A no piang hlim tan an taksa tih lumna tur ngaihtuahsak tur a ni.



Gramin Krishi Mausam Sewa
India Meteorological Department
ICAR Complex Branch
Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 06-08-2019

Weather Forecast of SERCHHIP(Mizoram) Issued On : 2019-08-06(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-08-07	10.0	29.0	23.0	99	51	6.6	88	8
2019-08-08	17.0	28.0	24.0	99	76	4.0	104	8
2019-08-09	6.0	29.0	23.0	100	58	4.1	97	7
2019-08-10	42.0	28.0	24.0	100	53	3.4	173	7
2019-08-11	41.0	28.0	24.0	100	86	1.8	191	8

Weather Summary/Alert:

Moderate to heavy rainfall occurred last 3 days. The maximum and minimum temperatures for the last 3 days range for 28-29 C and 18-21 C. Maximum relative humidity was expected in the range of 94-99% and minimum from 76-82%. Wind direction was easterly to southeasterly and northeasterly with the wind speed of 1.9-3.4 km per hour. Partially clear sky was prevailed last three days. Weekly cumulative rainfall: 40 mm

Tun nithum (3 days) chhung khan ruahtui vantlang tanga tam lam dawn a ni. Khua a lum lai ber 28-29 C a ni a, a vawh lai ber erawh chuan 18-21 C a ni. Boruak hnawn lai berin 94-99% a nih laiin a hul lai erawh chuan 76-82% a ni thung. Thli hi darker khatah 2.2-4.6 a chak in chhaklam atanga tannin chhim lamah chuan hmarchhak lamah awileh in chhim lamah a tleh thlaleh a ni. Tun nithum chhung khan van ah chum hmuh tur a tam hle. Kartluana ruahtui tlingkhawm: 40 mm

General Advisory:

Possibility of rainfall will be less and temperature will be high. Following general measures should be taken by farmers. Harvest all mature crops and proper sun dry for preservation of seed. Application of fertilizers/manure can be done. Check of vaccination schedule in poultry/livestock.

Ruahtui a tlak tlem rin a nih avangin boruak a lum deuh beisei a ni. Heng a hnuaia enkawl dan tarlan tehi huan neitu tehian zawm a hman theuh tur a ni. A kung puitling tawh techu seng tur, nisa ah phoro a a chi atana dahthat tur a ni. Leitha pek theih a ni. Ran neite tan vaccine pek hun tur (date) ennawn thin ni se.

SMS Advisory:

Pig- Swine fever- Vaccination of pigs @ 1 ml/pig with SF vaccines at 2 months interval.

Vawk natna-Swine fever enkawl dan- SF vaccines chu thla 2 danah vawk pakhat chiu nan 1 ml zel hman thin tur a ni.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
JHUM RICE(LOCAL)	☐ Drainage should be maintain properly ☐ Use split dose of fertilizer @ 40 kg/ha in clear/dry day. ☐ Keep the surrounding clean/ weed free to avoid disease/pest harboring. ☐ Spray tricyclozole 0.6 g/lit of water against leaf blast of rice in clear/dry day. ☐ Tuitling paihchhuahna tha taka siam tur ☐ Khaw that laiin leitha pek tur te chu split dose hmangin hectare khatah kg 40 zel hman tur a ni. ☐ A bul leh a chhehvel te tihfai a hnim hnah emaw rannung emaw natna thlen thei chi lakah enkawl tur a ni. ☐ Buh natna leaf blast laka vennis tricyclozole chu tui litre khatah 0.6 g pawlh chu khaw that laiin kah tur a ni.
RICE(GOMATI)	☐ Plough 2-3 time properly for reducing weed population. ☐ Apply fertilizer @40:60:40NPK kg and properly mix with soil. ☐ Spacing should be maintain 20x15 cm ☐ Hnim hnah a to tlem nan lei chu wawi 2-3 vel chawhphut tur. ☐ Leitha NPK 40:60:40 kg chu lei nen tha taka chawhpawlh tur a ni. ☐ A inkar hlat zawng hi 20x15 cm a ni ang.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
CUCURBITACEAE CROP(LOCAL)	☐ Drainage should be maintain properly ☐ Keep the surrounding clean/ weed free to avoid disease/pest harboring. ☐ Harvest all mature fruits and store dry seed for next year ☐ Tuitling paihchhuahna tha taka siam tur. ☐ A bul leh a chhehvel te tihfai a hnim hnah emaw rannung emaw natna thlen thei chi lakah enkawl tur a ni. ☐ A rah hmin tawh techu seng a, pho ro a, a kum leh a tuh atan dahthat tur a ni.
BHINDI(ARKA ANAMIKA)	☐ Pick tender and immature pods when they are approximately 3-4 inch in length. ☐ Harvest mature pod every alternate days. ☐ Harvest over matured pod for seed production purpose ☐ Seed should be dry properly and keep it for next year. ☐ A kawm (rah) la no leh la changlo techu inches 3-4 inkar an nih in lawh tur. ☐ A kawm (rah) hmin tawh techu ni khat danah lawh thin tur. ☐ A kawm (rah) hmin leh chang techu lawh a a chi atana dah that tur. ☐ A rah lawhsa tawh techu hmun hul ah dah ro a a kum leh atan dah that tur.

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG(LOCAL)	The bedding is always. If possible wooden sleeping box should be provided. Provide artificial heat source to new born piglets. An mut hmun hi a pawimawh hle. A theih hram chuan an mutna chhuat chu thing a siam ni thei se. A no piang hlim tan an taksa tih lumna tur ngaihtuhsak tur a ni.

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
FISH(COMPOSITE FISH)	To avoid bacterial disease liming should be done @10kg/Bigha. FYM 2400 kg and Lime 80 kg per bigha may be applied in new constructed pond. Bacteria natna laka an him theih nan chinai al @10kg/Bigha hman tur a ni FYM 2400 kg leh Lime 80 kg per bigha chu tuikhuah siam thar ah chuan chhun luh tur a ni.



Gramin Krishi Mausam Sewa
India Meteorological Department
ICAR Complex Branch
Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 06-08-2019

Weather Forecast of AIZAWL(Mizoram) Issued On : 2019-08-06(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-08-07	7.0	29.0	23.0	96	53	5.7	109	5
2019-08-08	33.0	28.0	24.0	99	65	3.5	115	8
2019-08-09	22.0	29.0	23.0	99	59	3.3	96	7
2019-08-10	43.0	28.0	24.0	100	56	2.8	113	8
2019-08-11	38.0	28.0	24.0	99	89	2.2	122	8

Weather Summary/Alert:

Moderate to heavy rainfall occurred last 3 days. The maximum and minimum temperatures for the last 3 days range for 29-30 C and 18-21 C. Maximum relative humidity was expected in the range of 95-95% and minimum from 75-89%. Wind direction was easterly to southeasterly with the wind speed of 2.9-4.1 km per hour. Partially clear sky was prevailed last three days. Weekly cumulative rainfall: 60 mm

Tun nithum (3 days) chhung khan ruahtui vantlang tanga tam lam dawn a ni. Khua a lum lai ber in 29-30 C a ni a, a vawh lai ber erawh chuan 18-21 C a ni. Boruak hnawn lai berin 95-95% a nih lain a hul lai erawh chuan 75-89% a ni thung. Thli hi darker khatah 2.9-4.1 a chak in chhaklam atanga tannin chhimlamah chuan hmarchhak lamah awileh in chhim lamah a tleh thlaleh a ni. Tun nithum chhung khan van ah chum hmuh tur a tam hle. Kartluana ruahtui tlingkhawm: 60 mm

General Advisory:

Possibility of rainfall will be less and temperature will be high. Following general measures should be taken by farmers. Harvest all mature crops and proper sun dry for preservation of seed. Application of fertilizers/manure can be done. Check of vaccination schedule in poultry/livestock.

Ruahtui a tlak tlem rin a nih avangin boruak a lum deuh beisei a ni. Heng a hnuaia enkawl dan tarlan tehi huan neitu tehian zawm a hman theuh tur a ni. A kung puitling tawh techu seng tur, nisa ah phoro a a chi atana dahthat tur a ni. Leitha pek theih a ni. Ran neite tan vaccine pek hun tur (date) ennawn thin ni se.

SMS Advisory:

Pig- Swine fever- Vaccination of pigs @ 1 ml/pig with SF vaccines at 2 months interval.

Vawk natna-Swine fever enkawl dan- SF vaccines chu thla 2 danah vawk pakhat chiu nan 1 ml zel hman thin tur a ni.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
JHUM RICE(LOCAL)	☐ Drainage should be maintain properly ☐ Use split dose of fertilizer @ 40 kg/ha in clear/dry day. ☐ Keep the surrounding clean/ weed free to avoid disease/pest harboring. ☐ Spray tricyclozole 0.6 g/lit of water against leaf blast of rice in clear/dry day. ☐ Tuitling paihchhuahna tha taka siam tur ☐ Khaw that laiin leitha pek tur te chu split dose hmangin hectare khatah kg 40 zel hman tur a ni. ☐ A bul leh a chhehvel te tihfai a hnim hnah emaw rannung emaw natna thlen thei chi lakah enkawl tur a ni. ☐ Buh natna leaf blast laka vennis tricyclozole chu tui litre khatah 0.6 g pawlh chu khaw that laiin kah tur a ni.
MAIZE(LOCAL)	☐ To manage this northern leaf blight disease spray aproach (Methoxy-acrylates) @ 2 g/li of water in clear/ dry day. ☐ Crop rotation with legume. ☐ Drainage should be maintain properly in heavy rainfall ☐ He Northern leaf blight natna enkawl nan hian Methoxy-acrylates gram 2 chu tui litre khatah chawhpawlh chu khaw that laia kah tur a ni. ☐ Thlai (Legume) ho nen chin pawlh ni se. ☐ Ruahtui tling paihchhuahna tur tha taka siam tur.
GREENGRAM(TRC M 151)	☐ Drainage should be maintain properly to avoid water stagnation problem. ☐ Intercrop with maize every 4-5 lines interval. ☐ Keep the surrounding clean/ weed free to avoid disease/pest harboring. ☐ Tuitling paihchhuahna tha taka siam tur. ☐ Vaimim nen line 4-5 dan zelah chin pawlh tur. ☐ A bul leh a chhehvel te tihfai a hnim hnah emaw rannung emaw natna thlen thei chi lakah enkawl tur a ni.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
BHINDI(ARKA ANAMIKA)	☐ Pick tender and immature pods when they are approximately 3-4 inch in length. ☐ Harvest mature pod every alternate days. ☐ Harvest over matured pod for seed production purpose ☐ Seed should be dry properly and keep it for next year. ☐ A kawm (rah) la no leh la changlo techu inches 3-4 inkar an nih in lawh tur. ☐ A kawm (rah) hmin tawh techu ni khat danah lawh thin tur. ☐ A kawm (rah) hmin leh chang techu lawh a a chi atana dah that tur. ☐ A rah lawhsa tawh techu hmun hul ah dah ro a a kum leh atan dah that tur.
CUCURBITACEAE CROP(LOCAL)	☐ Drainage should be maintain properly ☐ Keep the surrounding clean/ weed free to avoid disease/pest harboring. ☐ Harvest all mature fruits and store dry seed for next year ☐ Tuitling paihchhuahna tha taka siam tur. ☐ A bul leh a chhehvel te tihfai a hnim hnah emaw rannung emaw natna thlen thei chi lakah enkawl tur a ni. ☐ A rah hmin tawh techu seng a, pho ro a, a kum leh a tuh atan dahthat tur a ni.
GINGER(LOCAL)	☐ Drainage should be maintain properly to avoid water stagnation problem. ☐ Keep the surrounding clean/ weed free to avoid disease/pest harboring. ☐ Use split dose of fertilizer @ 40 kg/ha. ☐ Tuitling paihchhuahna tha taka siam tur. ☐ A bul leh a chhehvel te tihfai a hnim hnah emaw rannung emaw natna thlen thei chi lakah enkawl tur a ni. ☐ Leitha pek tur techu split dose hmangin hectare khatah kg 40 zel hman tur a ni.

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG(LOCAL)	The bedding is always. If possible wooden sleeping box should be provided. Provide artificial heat source to new born piglets. An mut hmun hi a pawimawh hle. A theih hram chuan an mutna chhuat chu thing a siam ni thei se. A no piang him tan an taksa tih lumna tur ngaituhsak tur a ni.
Livestock(Varieties)	Livestock Specific Advisory

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
FISH(COMPOSITE FISH)	To avoid bacterial disease liming should be done @10kg/Bigha. FYM 2400 kg and Lime 80 kg per bigha may be applied in new constructed pond.
	Bacteria natna laka an him theih nan chinai al @10kg/Bigha hman tur a ni FYM 2400 kg leh Lime 80 kg per bigha chu tuikhuah siam thar ah chuan chhun luh tur a ni.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICKEN(GIRIRAJA)	☐ Vaccination against Ranikhet at 4 days of birth and boost dose after 1 month of age
	☐ An pianni atanga ni 4 hnu ah Ranikhet natna laka invennan chiu tur a ni. Thla khat an nih hnu ah an taksa chak nan vaccine ngai tho lak leh tur.



Gramin Krishi Mausam Sewa
India Meteorological Department
ICAR Complex Branch
Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 06-08-2019

Weather Forecast of CHAMPHAI(Mizoram) Issued On : 2019-08-06(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-08-07	1.0	29.0	23.0	97	64	6.0	95	5
2019-08-08	27.0	28.0	24.0	99	68	3.1	116	7
2019-08-09	11.0	29.0	23.0	100	61	2.7	116	7
2019-08-10	41.0	28.0	24.0	100	52	2.5	203	7
2019-08-11	39.0	28.0	24.0	100	87	2.0	67	8

Weather Summary/Alert:

Moderate to heavy rainfall occurred last 3 days. The maximum and minimum temperatures for the last 3 days range for 27-26 C and 19-22 C. Maximum relative humidity was expected in the range of 89-95% and minimum from 72-85%. Wind direction was southeasterly to easterly to southeasterly and westerly with the wind speed of 2.4-4.1 km per hour. Partially clearsky was prevailed last three days. Weekly cumulative rainfall: 50 mm

Tun nithum (3 days) chhung khan ruahtui vantlang tanga tam lam dawn a ni. Khua a lum lai ber in 27-26 C a ni a, a vawhlai ber erawh chuan 19-22 C a ni. Boruak hnawn lai berin 89-95% a nih laiin a hul lai erawh chuan 72-85% a ni thung. Thli hi darker khatah 2.4-4.1 a chak in chhaklam atanga tannin chhimlam ah chuan hmarchhak lamah awileh in chhim lamah a tleh thlaleh a ni. Tun nithum chhung khan van ah chum hmuh tur a tam hle. Kartluana ruahtui tlingkhawm: 50 mm

General Advisory:

Possibility of rainfall will be less and temperature will be high. Following general measures should be taken by farmers. Harvest all mature crops and proper sun dry for preservation of seed. Application of fertilizers/manure can be done. Check of vaccination schedule in poultry/livestock.

Ruahtui a tlak tlem rin a nih avangin boruak a lum deuh beisei a ni. Heng a hnuaia enkawl dan tarlan tehi huan neitu tehian zawm a hman theuh tur a ni. A kung puitling tawh techu seng tur, nisa ah phoro a a chi atana dahthat tur a ni. Leitha pek theih a ni. Ran neite tan vaccine pek hun tur (date) ennawn thin ni se.

SMS Advisory:

Pig- Swine fever- Vaccination of pigs @ 1 ml/pig with SF vaccines at 2 months interval.

Vawk natna-Swine fever enkawl dan- SF vaccines chu thla 2 danah vawk pakhat chiu nan 1 ml zel hman thin tur a ni.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
MAIZE(LOCAL)	☐ To manage this northern leaf blight disease spray aproach (Methoxy-acrylates) @ 2 g/li of water in clear/ dry day. ☐ Crop rotation with legume. ☐ Drainage should be maintain properly in heavy rainfall ☐ He Northern leaf blight natna enkawl nan hian Methoxy-acrylates gram 2 chu tui litre khatah chawhpawlh chu khaw that laia kah tur a ni. ☐ Thlai (Legume) ho nen chin pawlh ni se. ☐ Ruahtui tling paihchhuahna tur tha taka siam tur.
JHUM RICE(LOCAL)	☐ Drainage should be maintain properly ☐ Use split dose of fertilizer @ 40 kg/ha in clear/dry day. ☐ Keep the surrounding clean/ weed free to avoid disease/pest harboring. ☐ Spray tricyclozole 0.6 g/lit of water against leaf blast of rice in clear/dry day. ☐ Tuitling paihchhuahna tha taka siam tur ☐ Khaw that laiin leitha pek tur te chu split dose hmangin hectare khatah kg 40 zel hman tur a ni. ☐ A bul leh a chhehvel te tihfai a hnim hnah emaw rannung emaw natna thlen thei chi lakah enkawl tur a ni. ☐ Buh natna leaf blast laka vennan tricyclozole chu tui litre khatah 0.6 g pawlh chu khaw that laiin kah tur a ni.
GREENGRAM(TRC M 151)	☐ Drainage should be maintain properly to avoid water stagnation problem. ☐ Intercrop with maize every 4-5 lines interval. ☐ Keep the surrounding clean/ weed free to avoid disease/pest harboring. ☐ Tuitling paihchhuahna tha taka siam tur. ☐ Vaimim nen line 4-5 dan zelah chin pawlh tur. ☐ A bul leh a chhehvel te tihfai a hnim hnah emaw rannung emaw natna thlen thei chi lakah enkawl tur a ni.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
BHINDI(ARKA ANAMIKA)	☐ Pick tender and immature pods when they are approximately 3-4 inch in length. ☐ Harvest mature pod every alternate days. ☐ Harvest over matured pod for seed production purpose ☐ Seed should be dry properly and keep it for next year. ☐ A kawm (rah) la no leh la changlo techu inches 3-4 inkar an nih in lawh tur. ☐ A kawm (rah) hmin tawh techu ni khat danah lawh thin tur. ☐ A kawm (rah) hmin leh chang techu lawh a a chi atana dah that tur. ☐ A rah lawhsa tawh techu hmun hul ah dah ro a a kum leh atan dah that tur.
CUCURBITACEAE CROP(LOCAL)	☐ Drainage should be maintain properly ☐ Keep the surrounding clean/ weed free to avoid disease/pest harboring. ☐ Harvest all mature fruits and store dry seed for next year ☐ Tuitling paihchhuahna tha taka siam tur. ☐ A bul leh a chhehvel te tihfai a hnim hnah emaw rannung emaw natna thlen thei chi lakah enkawl tur a ni. ☐ A rah hmin tawh techu seng a, pho ro a, a kum leh a tuh atan dahthat tur a ni.
GINGER(LOCAL)	☐ Drainage should be maintain properly to avoid water stagnation problem. ☐ Keep the surrounding clean/ weed free to avoid disease/pest harboring. ☐ Use split dose of fertilizer @ 40 kg/ha. ☐ Tuitling paihchhuahna tha taka siam tur. ☐ A bul leh a chhehvel te tihfai a hnim hnah emaw rannung emaw natna thlen thei chi lakah enkawl tur a ni. ☐ Leitha pek tur techu split dose hmangin hectare khatah kg 40 zel hman tur a ni.

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
COW(LOCAL)	Balanced feeding of dairy animals and allow calf suckling immediately after parturition. Full hand milking should be practiced. Frequent incomplete milking at interval of 4hrs should be practiced after parturition for minimum 1 week. Afterwards complete milking in a day is recommended.

Livestock(Varieties)	Livestock Specific Advisory
	A pui in nau a neih hnu ah a inmil tawkin a naute chu hnute tui leh chaw pek tur a ni. A hnute sawr dan thin full hand milking chu hman tur a ni. Nau a neih hnu atangin a loh ber ah kar khat tal chu darkar li danah a hnute chu vawi khata sawr zo vek lo in sawr zeuh zeuh thin tur a ni. Chumi hnu ah chuan a vaia sawr zawh thin chauh tur a ni.
PIG(LOCAL)	The bedding is always. If possible wooden sleeping box should be provided. Provide artificial heat source to new born piglets.
	An mut hmun hi a pawimawh hle. A theih hram chuan an mutna chhuat chu thing a siam ni thei se. A no piang hlim tan an taksa tih lumna tur ngaihtuahsak tur a ni.

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
FISH(COMPOSITE FISH)	To avoid bacterial disease liming should be done @10kg/Bigha. FYM 2400 kg and Lime 80 kg per bigha may be applied in new constructed pond.
	Bacteria natna laka an him theih nan chinai al @10kg/Bigha hman tur a ni FYM 2400 kg leh Lime 80 kg per bigha chu tuikhuah siam thar ah chuan chhun luh tur a ni.



Gramin Krishi Mausam Sewa
India Meteorological Department
ICAR Complex Branch
Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 06-08-2019

Weather Forecast of KOLASIB(Mizoram) Issued On : 2019-08-06(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-08-07	1.0	32.0	25.0	97	54	6.2	101	5
2019-08-08	22.0	30.0	26.0	99	56	3.4	105	8
2019-08-09	17.0	31.0	26.0	99	61	3.0	81	8
2019-08-10	31.0	30.0	25.0	99	56	2.8	50	8
2019-08-11	10.0	30.0	26.0	98	75	2.7	129	8

Weather Summary/Alert:

Moderate to heavy rainfall occurred last 3 days. The maximum and minimum temperatures for the last 3 days range for 30-31 C and 22-24 C. Maximum relative humidity was expected in the range of 94-98% and minimum from 85-92%. Wind direction was easterly to northwesterly with the wind speed of 1.6-3.2 km per hour. Partially clear sky was prevailed last three days. Weekly cumulative rainfall: 65 mm

Tun nithum (3 days) chhung khan ruahtui vantlang tanga tam lam dawn a ni. Khua a lum lai ber 30-31 C a ni a, a vawh lai ber erawh chuan 22-24 C a ni. Boruak hnawn lai berin 94-98% a nih laiin a hul lai erawh chuan 85-92% a ni thung. Thli hi darker khatah 1.6-3.2 a chak in chhaklam atanga tannin chhimlamah chuan hmarchhak lamah awileh in chhimlamah a tleh thlaleh a ni. Tun nithum chhung khan van ah chum hmuh tur a tam hle. Kartluana ruahtui tlingkhawm: 65 mm

General Advisory:

Possibility of rainfall will be less and temperature will be high. Following general measures should be taken by farmers. Harvest all mature crops and proper sun dry for preservation of seed. Application of fertilizers/manure can be done. Check of vaccination schedule in poultry/livestock.

Ruahtui a tlak tlem rin a nih avangin boruak a lum deuh beisei a ni. Heng a hnuaia enkawl dan tarlan tehi huan neitu tehian zawm a hman theuh tur a ni. A kung puitling tawh techu seng tur, nisa ah phoro a a chi atana dahthat tur a ni. Leitha pek theih a ni. Ran neite tan vaccine pek hun tur (date) ennawn thin ni se.

SMS Advisory:

Pig- Swine fever- Vaccination of pigs @ 1 ml/pig with SF vaccines at 2 months interval

Vawk natna-Swine fever enkawl dan- SF vaccines chu thla 2 danah vawk pakhat chiu nan 1 ml zel hman thin tur a ni.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
JHUM RICE(LOCAL)	☐ Drainage should be maintain properly ☐ Use split dose of fertilizer @ 40 kg/ha in clear/dry day. ☐ Keep the surrounding clean/ weed free to avoid disease/pest harboring. ☐ Spray tricyclozole 0.6 g/lit of water against leaf blast of rice in clear/dry day. ☐ Tuitling paihchhuahna tha taka siam tur ☐ Khaw that laiin leitha pek tur te chu split dose hmangin hectare khatah kg 40 zel hman tur a ni. ☐ A bul leh a chhehvel te tihfai a hnim hnah emaw rannung emaw natna thlen thei chi lakah enkawl tur a ni. ☐ Buh natna leaf blast laka vennis tricyclozole chu tui litre khatah 0.6 g pawlh chu khaw that laiin kah tur a ni.
RICE(GOMATI)	☐ Plough 2-3 time properly for reducing weed population. ☐ Apply fertilizer @40:60:40NPK kg and properly mix with soil. ☐ Spacing should be maintain 20x15 cm ☐ Hnim hnah a to tlem nan lei chu wawi 2-3 vel chawhphut tur. ☐ Leitha NPK 40:60:40 kg chu lei nen tha taka chawhpawlh tur a ni. ☐ A inkar hlat zawng hi 20x15 cm a ni ang.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
BHINDI(ARKA ANAMIKA)	☐ Pick tender and immature pods when they are approximately 3-4 inch in length. ☐ Harvest mature pod every alternate days. ☐ Harvest over matured pod for seed production purpose ☐ Seed should be dry properly and keep it for next year. ☐ A kawm (rah) la no leh la changlo techu inches 3-4 inkar an nih in lawh tur. ☐ A kawm (rah) hmin tawh techu ni khat danah lawh thin tur. ☐ A kawm (rah) hmin leh chang techu lawh a a chi atana dah that tur. ☐ A rah lawhsa tawh techu hmun hul ah dah ro a a kum leh atan dah that tur.
CUCURBITACEAE CROP(LOCAL)	☐ Drainage should be maintain properly ☐ Keep the surrounding clean/ weed free to avoid disease/pest harboring. ☐ Harvest all mature fruits and store dry seed for next year ☐ Tuitling paihchhuahna tha taka siam tur. ☐ A bul leh a chhehvel te tihfai a hnim hnah emaw rannung emaw natna thlen thei chi lakah enkawl tur a ni. ☐ A rah hmin tawh techu seng a, pho ro a, a kum leh a tuh atan dahthat tur a ni.

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG(LOCAL)	The bedding is always. If possible wooden sleeping box should be provided. Provide artificial heat source to new born piglets. An mut hmun hi a pawimawh hle. A theih hram chuan an mutna chhuat chu thing a siam ni thei se. A no piang hlim tan an taksa tih lumna tur ngaihtuhsak tur a ni.

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
FISH(COMPOSITE FISH)	To avoid bacterial disease liming should be done @10kg/Bigha. FYM 2400 kg and Lime 80 kg per bigha may be applied in new constructed pond. Bacteria natna laka an him theih nan chinai al @10kg/Bigha hman tur a ni FYM 2400 kg leh Lime 80 kg per bigha chu tuikhuah siam thar ah chuan chhun luh tur a ni.



Gramin Krishi Mausam Sewa
India Meteorological Department
ICAR Complex Branch
Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 06-08-2019

Weather Forecast of LAWNGTLAI(Mizoram) Issued On : 2019-08-06(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-08-07	14.0	29.0	23.0	98	60	5.9	119	8
2019-08-08	7.0	28.0	24.0	96	83	5.8	114	5
2019-08-09	8.0	29.0	23.0	98	68	5.8	100	5
2019-08-10	13.0	28.0	24.0	99	52	3.4	132	7
2019-08-11	27.0	28.0	24.0	99	68	2.1	175	8

Weather Summary/Alert:

Moderate to heavy rainfall occurred last 3 days. The maximum and minimum temperatures for the last 3 days range for 29-26 C and 21-23 C. Maximum relative humidity was expected in the range of 88-95% and minimum from 72-83%. Wind direction was northeasterly to southeasterly and westerly with the wind speed of 2.1-3.3 km per hour. Partially clear sky was prevailed last three days. Weekly cumulative rainfall: 40 mm

Tun nithum (3 days) chhung khan ruahtui vantlang tanga tam lam dawn a ni. Khua a lumlai ber in 26-29 C a ni a, a vawhlai ber erawh chuan 21-23 C a ni. Boruak hnawn lai berin 88-95% a nih lain a hul lai erawh chuan 72-83% a ni thung. Thli hi darker khatah 2.1-3.3 a chak in chhaklam atanga tannin chhimlamah chuan hmarchhak lamah awileh in chhimlamah a tleh thlaleh a ni. Tun nithum chhung khan van ah chum hmuh tur a tam hle. Kartluana ruahtui tlingkhawm: 40 mm

General Advisory:

Possibility of rainfall will be less and temperature will be high. Following general measures should be taken by farmers. Harvest all mature crops and proper sun dry for preservation of seed. Application of fertilizers/manure can be done. Check of vaccination schedule in poultry/livestock.

Ruahtui a tlak tlem rin a nih avangin boruak a lum deuh beisei a ni. Heng a hnuaia enkawl dan tarlan tehi huan neitu tehian zawm a hman theuh tur a ni. A kung puitling tawh techu seng tur, nisa ah phoro a a chi atana dahthat tur a ni. Leitha pek theih a ni. Ran neite tan vaccine pek hun tur (date) ennawn thin ni se.

SMS Advisory:

Pig- Swine fever- Vaccination of pigs @ 1 ml/pig with SF vaccines at 2 months interval.

Vawk natna-Swine fever enkawl dan- SF vaccines chu thla 2 danah vawk pakhat chiu nan 1 ml zel hman thin tur a ni.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
MAIZE(LOCAL)	☐ To manage this northern leaf blight disease spray aproach (Methoxy-acrylates) @ 2 g/li of water in clear/ dry day. ☐ Crop rotation with legume. ☐ Drainage should be maintain properly in heavy rainfall ☐ He Northern leaf blight natna enkawl nan hian Methoxy-acrylates gram 2 chu tui litre khatah chawhpawlh chu khaw that laia kah tur a ni. ☐ Thlai (Legume) ho nen chin pawlh ni se. ☐ Ruahtui tling paihchhuahna tur tha taka siam tur.
JHUM RICE(LOCAL)	☐ Drainage should be maintain properly ☐ Use split dose of fertilizer @ 40 kg/ha in clear/dry day. ☐ Keep the surrounding clean/ weed free to avoid disease/pest harboring. ☐ Spray tricyclozole 0.6 g/lit of water against leaf blast of rice in clear/dry day. ☐ Tuitling paihchhuahna tha taka siam tur ☐ Khaw that laiin leitha pek tur te chu split dose hmangin hectare khatah kg 40 zel hman tur a ni. ☐ A bul leh a chhehvel te tihfai a hnim hnah emaw rannung emaw natna thlen thei chi lakah enkawl tur a ni. ☐ Buh natna leaf blast laka vennis tricyclozole chu tui litre khatah 0.6 g pawlh chu khaw that laiin kah tur a ni.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
CUCURBITACEAE CROP(LOCAL)	☐ Drainage should be maintain properly ☐ Keep the surrounding clean/ weed free to avoid disease/pest harboring. ☐ Harvest all mature fruits and store dry seed for next year ☐ Tuitling paihchhuahna tha taka siam tur. ☐ A bul leh a chhehvel te tihfai a hnim hnah emaw rannung emaw natna thlen thei chi lakah enkawl tur a ni. ☐ A rah hmin tawh techu seng a, pho ro a, a kum leh a tuh atan dahthat tur a ni.
BHINDI(ARKA ANAMIKA)	☐ Pick tender and immature pods when they are approximately 3-4 inch in length. ☐ Harvest mature pod every alternate days. ☐ Harvest over matured pod for seed production purpose ☐ Seed should be dry properly and keep it for next year. ☐ A kawm (rah) la no leh la changlo techu inches 3-4 inkar an nih in lawh tur. ☐ A kawm (rah) hmin tawh techu ni khat danah lawh thin tur. ☐ A kawm (rah) hmin leh chang techu lawh a a chi atana dah that tur. ☐ A rah lawhsa tawh techu hmun hul ah dah ro a a kum leh atan dah that tur.

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG(LOCAL)	The bedding is always. If possible wooden sleeping box should be provided. Provide artificial heat source to new born piglets. An mut hmun hi a pawimawh hle. A theih hram chuan an mutna chhuat chu thing a siam ni thei se. A no piang hlim tan an taksa tih lumna tur ngaihtuhsak tur a ni.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICKEN(LOCAL)	☐ Vaccination against Ranikhet at 4 days of birth and boost dose after 1 month of age ☐ An pianni atanga ni 4 hnu ah Ranikhet natna laka invennis chiu tur a ni. Thla khat an nih hnu ah an taksa chak nan vaccine ngai tho lak leh tur.



Gramin Krishi Mausam Sewa
India Meteorological Department
ICAR Complex Branch
Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 06-08-2019

Weather Forecast of LUNGLEI(Mizoram) Issued On : 2019-08-06(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-08-07	16.0	31.0	25.0	98	54	5.7	100	8
2019-08-08	13.0	31.0	25.0	96	82	5.5	102	7
2019-08-09	8.0	32.0	26.0	99	62	5.2	91	6
2019-08-10	13.0	32.0	25.0	99	53	3.3	133	6
2019-08-11	45.0	31.0	26.0	99	74	2.1	209	8

Weather Summary/Alert:

Moderate to heavy rainfall occurred last 3 days. The maximum and minimum temperatures for the last 3 days range for 24-27 C and 17-19 C. Maximum relative humidity was expected in the range of 94-98% and minimum from 71-85%. Wind direction was easterly to southeasterly and westerly with the wind speed of 2.5-3.8 km per hour. Partially clear sky was prevailed last three days. Weekly cumulative rainfall: 50 mm

Tun nithum (3 days) chhung khan ruahtui vantlang tanga tam lam dawn a ni. Khua a lumlai ber in 24-27 C a ni a, a vawhlai ber erawh chuan 17-19 C a ni. Boruak hnawn lai berin 94-98% a nih laiin a hul lai erawh chuan 71-85% a ni thung. Thli hi darker khatah 2.5-3.8 a chak in chhaklam atanga tannin chhimlamah chuan hmarchhak lamah awileh in chhim lamah a tleh thlaleh a ni. Tun nithum chhung khan van ah chum hmuh tur a tam hle. Kartluana ruahtui tlingkhawm: 50 mm

General Advisory:

Possibility of rainfall will be less and temperature will be high. Following general measures should be taken by farmers. Harvest all mature crops and proper sun dry for preservation of seed. Application of fertilizers/manure can be done. Check of vaccination schedule in poultry/livestock.

Ruahtui a tlak tlem rin a nih avangin boruak a lum deuh beisei a ni. Heng a hnuaia enkawl dan tarlan tehi huan neitu tehian zawm a hman theuh tur a ni. A kung puitling tawh techu seng tur, nisa ah phoro a a chi atana dahthat tur a ni. Leitha pek theih a ni. Ran neite tan vaccine pek hun tur (date) ennawn thin ni se.

SMS Advisory:

Pig- Swine fever- Vaccination of pigs @ 1 ml/pig with SF vaccines at 2 months interval.

Vawk natna-Swine fever enkawl dan- SF vaccines chu thla 2 danah vawk pakhat chiu nan 1 ml zel hman thin tur a ni.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
MAIZE(LOCAL)	☐ To manage this northern leaf blight disease spray aproach (Methoxy-acrylates) @ 2 g/li of water in clear/ dry day. ☐ Crop rotation with legume. ☐ Drainage should be maintain properly in heavy rainfall ☐ He Northern leaf blight natna enkawl nan hian Methoxy-acrylates gram 2 chu tui litre khatah chawhpawlh chu khaw that laia kah tur a ni. ☐ Thlai (Legume) ho nen chin pawlh ni se. ☐ Ruahtui tling paihchhuahna tur tha taka siam tur.
JHUM RICE(LOCAL)	☐ Drainage should be maintain properly ☐ Use split dose of fertilizer @ 40 kg/ha in clear/dry day. ☐ Keep the surrounding clean/ weed free to avoid disease/pest harboring. ☐ Spray tricyclozole 0.6 g/lit of water against leaf blast of rice in clear/dry day. ☐ Tuitling paihchhuahna tha taka siam tur ☐ Khaw that laiin leitha pek tur te chu split dose hmangin hectare khatah kg 40 zel hman tur a ni. ☐ A bul leh a chhehvel te tihfai a hnim hnah emaw rannung emaw natna thlen thei chi lakah enkawl tur a ni. ☐ Buh natna leaf blast laka vennis tricyclozole chu tui litre khatah 0.6 g pawlh chu khaw that laiin kah tur a ni.
RICE(GOMATI)	☐ Plough 2-3 time properly for reducing weed population. ☐ Apply fertilizer @40:60:40NPK kg and properly mix with soil. ☐ Spacing should be maintain 20x15 cm ☐ Hnim hnah a to tlem nan lei chu vawi 2-3 vel chawhphut tur. ☐ Leitha NPK 40:60:40 kg chu lei nen tha taka chawhpawlh tur a ni. ☐ A inkar hlat zawng hi 20x15 cm a ni ang.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
CUCURBITACEAE CROP(LOCAL)	☐ Drainage should be maintain properly ☐ Keep the surrounding clean/ weed free to avoid disease/pest harboring. ☐ Harvest all mature fruits and store dry seed for next year ☐ Tuitling paihchhuahna tha taka siam tur. ☐ A bul leh a chhehvel te tihfai a hnim hnah emaw rannung emaw natna thlen thei chi lakah enkawl tur a ni. ☐ A rah hmin tawh techu seng a, pho ro a, a kum leh a tuh atan dahthat tur a ni.
BHINDI(ARKA ANAMIKA)	☐ Pick tender and immature pods when they are approximately 3-4 inch in length. ☐ Harvest mature pod every alternate days. ☐ Harvest over matured pod for seed production purpose ☐ Seed should be dry properly and keep it for next year. ☐ A kawm (rah) la no leh la changlo techu inches 3-4 inkar an nih in lawh tur. ☐ A kawm (rah) hmin tawh techu ni khat danah lawh thin tur. ☐ A kawm (rah) hmin leh chang techu lawh a a chi atana dah that tur. ☐ A rah lawhsa tawh techu hmun hul ah dah ro a a kum leh atan dah that tur.

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG(LOCAL)	The bedding is always. If possible wooden sleeping box should be provided. Provide artificial heat source to new born piglets. An mut hmun hi a pawimawh hle. A theih hram chuan an mutna chhuat chu thing a siam ni thei se. A no piang hlim tan an taksa tih lumna tur ngaihtuhsak tur a ni.

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
FISH(COMPOSITE FISH)	To avoid bacterial disease liming should be done @10kg/Bigha. FYM 2400 kg and Lime 80 kg per bigha may be applied in new constructed pond.

Fisheries(Varieties)	Fisheries Specific Advisory
	Bacteria natna laka an him theih nan chinai al @10kg/Bigha hman tur a ni FYM 2400 kg leh Lime 80 kg per bigha chu tuikhuah siam thar ah chuan chhun luh tur a ni.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICKEN(LOCAL)	☐ Vaccination against Ranikhet at 4 days of birth and boost dose after 1 month of age ☐ An pianni atanga ni 4 hnu ah Ranikhet natna laka invennan chiu tur a ni. Thla khat an nih hnu ah an taksa chak nan vaccine ngai tho lak leh tur.



Gramin Krishi Mausam Sewa
India Meteorological Department
ICAR Complex Branch
Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 06-08-2019

Weather Forecast of MAMIT(Mizoram) Issued On : 2019-08-06(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-08-07	8.0	29.0	23.0	98	50	5.1	105	5
2019-08-08	15.0	28.0	24.0	96	70	4.7	97	8
2019-08-09	6.0	30.0	24.0	98	54	5.1	87	5
2019-08-10	24.0	28.0	24.0	99	53	3.8	139	7
2019-08-11	36.0	28.0	24.0	99	55	2.9	158	8

Weather Summary/Alert:

Moderate to heavy rainfall occurred last 3 days. The maximum and minimum temperatures for the last 3 days range for 30-32 C and 22-24 C. Maximum relative humidity was expected in the range of 92-98% and minimum from 71-85%. Wind direction was easterly to northeasterly and southeasterly with the wind speed of 2.4-3.5km per hour. Partially clear sky was prevailed last three days. Weekly cumulative rainfall: 45 mm

Tun nithum (3 days) chhung khan ruahtui vantlang tanga tam lam dawn a ni. Khua a lumlai ber 30-32 C a ni a, a vawh lai ber erawh chuan 22-24 C a ni. Boruak hnawn lai berin 92-98% a nih lain a hul lai erawh chuan 71-85% a ni thung. Thli hi darker khatah 2.4-3.5 a chak in chhaklam atanga tannin chhimlamah chuan hmarchhak lamah awileh in chhim lamah a tleh thlaleh a ni. Tun nithum chhung khan van ah chum hmuh tur a tam hle. Kartluana ruahtui tlingkhawm: 45 mm

General Advisory:

Possibility of rainfall will be less and temperature will be high. Following general measures should be taken by farmers. Harvest all mature crops and proper sun dry for preservation of seed. Application of fertilizers/manure can be done. Check of vaccination schedule in poultry/livestock.

Ruahtui a tlak tlem rin a nih avangin boruak a lum deuh beisei a ni. Heng a hnuaia enkawl dan tarlan tehi huan neitu tehian zawm a hman theuh tur a ni. A kung puitling tawh techu seng tur, nisa ah phoro a a chi atana dahthat tur a ni. Leitha pek theih a ni. Ran neite tan vaccine pek hun tur (date) ennawn thin ni se.

SMS Advisory:

Pig- Swine fever- Vaccination of pigs @ 1 ml/pig with SF vaccines at 2 months interval.

Vawk natna-Swine fever enkawl dan- SF vaccines chu thla 2 danah vawk pakhat chiu nan 1 ml zel hman thin tur a ni.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
MAIZE(LOCAL)	☐ To manage this northern leaf blight disease spray aproach (Methoxy-acrylates) @ 2 g/li of water in clear/ dry day. ☐ Crop rotation with legume. ☐ Drainage should be maintain properly in heavy rainfall ☐ He Northern leaf blight natna enkawl nan hian Methoxy-acrylates gram 2 chu tui litre khatah chawhpawlh chu khaw that laia kah tur a ni. ☐ Thlai (Legume) ho nen chin pawlh ni se. ☐ Ruahtui tling paihchhuahna tur tha taka siam tur.
JHUM RICE(LOCAL)	☐ Drainage should be maintain properly ☐ Use split dose of fertilizer @ 40 kg/ha in clear/dry day. ☐ Keep the surrounding clean/ weed free to avoid disease/pest harboring. ☐ Spray tricyclozole 0.6 g/lit of water against leaf blast of rice in clear/dry day. ☐ Tuitling paihchhuahna tha taka siam tur ☐ Khaw that laiin leitha pek tur te chu split dose hmangin hectare khatah kg 40 zel hman tur a ni. ☐ A bul leh a chhehvel te tihfai a hnim hnah emaw rannung emaw natna thlen thei chi lakah enkawl tur a ni. ☐ Buh natna leaf blast laka vennis tricyclozole chu tui litre khatah 0.6 g pawlh chu khaw that laiin kah tur a ni.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
BHINDI(ARKA ANAMIKA)	☐ Pick tender and immature pods when they are approximately 3-4 inch in length. ☐ Harvest mature pod every alternate days. ☐ Harvest over matured pod for seed production purpose ☐ Seed should be dry properly and keep it for next year. ☐ A kawm (rah) la no leh la changlo techu inches 3-4 inkar an nih in lawh tur. ☐ A kawm (rah) hmin tawh techu ni khat danah lawh thin tur. ☐ A kawm (rah) hmin leh chang techu lawh a a chi atana dah that tur. ☐ A rah lawhsa tawh techu hmun hul ah dah ro a a kum leh atan dah that tur.
CUCURBITACEAE CROP(LOCAL)	☐ Drainage should be maintain properly ☐ Keep the surrounding clean/ weed free to avoid disease/pest harboring. ☐ Harvest all mature fruits and store dry seed for next year ☐ Tuitling paihchhuahna tha taka siam tur. ☐ A bul leh a chhehvel te tihfai a hnim hnah emaw rannung emaw natna thlen thei chi lakah enkawl tur a ni. ☐ A rah hmin tawh techu seng a, pho ro a, a kum leh a tuh atan dahthat tur a ni.

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG(LOCAL)	The bedding is always. If possible wooden sleeping box should be provided. Provide artificial heat source to new born piglets. An mut hmun hi a pawimawh hle. A theih hram chuan an mutna chhuat chu thing a siam ni thei se. A no piang hlim tan an taksa tih lumna tur ngaihtuhsak tur a ni.

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
FISH(COMPOSITE FISH)	To avoid bacterial disease liming should be done @10kg/Bigha. FYM 2400 kg and Lime 80 kg per bigha may be applied in new constructed pond. Bacteria natna laka an him theih nan chinai al @10kg/Bigha hman tur a ni FYM 2400 kg leh Lime 80 kg per bigha chu tuikhuah siam thar ah chuan chhun luh tur a ni.