



Integrated Agromet Advisory Service Bulletin 11th to 15th Dec'13

Longleng District

Bulletin No:93/2013

Weather summary of the preceding week

No rain occurred the past week.
Maximum and minimum temperatures ranged from 22°C to 23°C and 6°C to 7°C, respectively.
Relative humidity varied from 38% to 95%.
Wind speed ranged from 2 to 3 kmph.

Weather forecast valid upto 15th Dec '13

There is possibility of no rain the coming week.
Max temp is expected to be around 20°C - 21°C and min temp is expected to be around 6°C to 7°C.
Possibility of mainly clear sky the coming week
Relative Humidity will range from 38% to 90%
The wind speed 3 –4 kmph
Wind direction will blow from southerly

Main Crops

Stage

Pest/ diseases

Agro-meteorological Advisories

Field crops

Grain storage

Storage

-

Rice should be stored in gunny sacks on wooden platforms that allow free passage of air between grain sacks and the floor. Low temperature and low moisture content are necessary for storing rice grain for long period

Maize

Tasseling stage

Stalk
borer

Farmers are advised to monitor the maize plant against stalk borer. The larvae may remain inside the whole and start damaging the tassel stalk. Application of Neem oil 2 % would minimize the infestation.

Mustard

Vegetative stage

-

Earthing up, gap filling along with weeding should be carried out.

Horticultural crops

Probability no rain will occur the coming 3 or 4 days, therefore the farmers are advised to irrigate the crops daily or at an interval depending on the availability of water.

Cabbage

Head formation
stage

-

In case of cabbage, irrigation is very important for good growth and development.

Tomato

Vegetative stage

-

Frequent inter tillage and cultivation should be done in the field to keep it free from weeds.
Proper pruning and staking should be carried out so as better and large tomatoes can be produced.

Livestock

Regular cleaning and disinfection of cattle shed and its premises should be carried out.

Fishery

Feed your fish at the same place every day, so that you will know whether your fish are eating well or not. If you give them too much, the food that is not eaten will stay on the bottom.






Integrated Agromet Advisory Service Bulletin 11th to 15th Dec'13

Mokokchung District

Bulletin No: 93/2013

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 15th Dec '13</i>
No rain occurred the past week Maximum and minimum temperatures ranged 19 ⁰ C to 22 ⁰ C and 10 ⁰ C to 11 ⁰ C, respectively. Relative humidity varied from 39% to 97%. Wind speed ranged from 2 to 3 kmph.			There is possibility of no rain the coming week. Max temp is expected to be around 19 ⁰ C - 20 ⁰ C and min temp is expected to be around 3 ⁰ C to 6 ⁰ C. Possibility of mainly cloudy to clear sky the coming week Relative Humidity will range from 39% to 96% and Wind speed 3 – 4 kmph Wind direction will blow from southerly.
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Grain storage	Storage	-	Rice should be stored in gunny sacks on wooden platforms that allow free passage of air between grain sacks and the floor. Low temperature and low moisture content are necessary for storing rice grain for long period
Maize	Tasseling stage	Stalk borer	Farmers are advised to monitor the maize plant against stalk borer. The larvae may remain inside the whole and start damaging the tassel stalk. Application of Neem oil 2 % would minimize the infestation.
Mustard	Vegetative stage	-	Earthing up, gap filling along with weeding should be carried out.
Horticultural crops			
Probability no rain will occur the coming 3 or 4 days, therefore the farmers are advised to irrigate the crops daily or at an interval depending on the availability of water.			
Rajmash	Vegetative stage	wilt	Symptoms are premature yellowing and whitening of leaves and finally the diving of the entire plant. Plant may continue to wilt in patches in quick succession till the maturity period. Disease affected plants should be pulled out.
Broccoli	Vegetative stage	-	Broccoli requires sufficient moisture in the soil for uniform and continuous growth. Dry conditions adversely affect the quality and yield of shoots by being more fibrous
Cabbage	Head formation stage	-	In case of cabbage, irrigation is very important for good growth and development.
Potato	Vegetative stage	-	Proper cultural practices should be maintained, infested vines should be removed from the field for controlling the pest. Farmers are advised to carry out intercultural operations like hand weeding and

				earthingup to destroy weeds, to keep the soil loose for proper development of tubers and to cover the tubers with soil.
	Tomato	Vegetative stage	-	Frequent inter tillage and cultivation should be done in the field to keep it free from weeds. Proper pruning and staking should be carried out so as better and large tomatoes can be produced.
	Livestock	Regular cleaning and disinfection of cattle shed and its premises should be carried out.		
	Fishery	Feed your fish at the same place every day, so that you will know whether your fish are eating well or not. If you give them too much, the food that is not eaten will stay on the bottom.		



Integrated Agromet Advisory Service Bulletin from 11th to 15th Dec'13 Mon District

Bulletin No:93/2013


Weather summary of the preceding week

- No rain occurred the past week
- **Maximum** and **minimum temperatures** ranged 21⁰C to 22⁰C and 4⁰C to 5⁰C, respectively.
- **Relative humidity** varied from 37% to 95%.
- **Wind speed** ranged from 2 to 3 kmph.

Weather forecast valid upto 15th Dec '13

- Possibility of no rain the coming week.
- The **max temp** is likely to be 20⁰C - 21⁰C and the **min temp** 4⁰C - 5⁰C.
- **Sky is likely to be mainly cloudy to clear the coming week**
- **Relative Humidity** is likely to range from 38% to 94%
- **Wind speed** may reach upto 2- 3kmph
- **Wind direction** will be from southeasterly

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Grain storage	Storage	-	Rice should be stored in gunny sacks on wooden platforms that allow free passage of air between grain sacks and the floor. Low temperature and low moisture content are necessary for storing rice grain for long period
Maize	Tasseling stage	Stalk borer	Farmers are advised to monitor the maize plant against stalk borer. The larvae may remain inside the whole and start damaging the tassel stalk. Application of Neem oil 2 % would minimize the infestation.
Mustard	Vegetative stage	-	Earthing up, gap filling along with weeding should be carried out.
Horticultural crops			
Probability no rain will occur the coming 3 or 4 days, therefore the farmers are advised to irrigate the crops daily or at an interval depending on the availability of water.			
Cabbage	Head formation stage	-	In case of cabbage, irrigation is very important for good growth and development.
French bean	Vegetative stage	wilt	Symptoms are premature yellowing and whitening of leaves and finally the diving of the entire plant. Plant may continue to wilt in patches in quick succession till the maturity period. Disease affected plants should be pulled out.
Pea	Vegetative stage	-	When the crops are about two months old bamboo stick should be given as a support. Delay in this operation will reduce the yield considerably.
Potato	Vegetative stage	-	Proper cultural practices should be maintained, infested vines should be removed from the field for controlling the pest. Farmers are advised to carry out intercultural operations like hand weeding and earthingup to destroy weeds, to keep the soil loose for proper

				development of tubers and to cover the tubers with soil.
	Tomato	Vegetative stage	-	Frequent inter tillage and cultivation should be done in the field to keep it free from weeds. Proper pruning and staking should be carried out so as better and large tomatoes can be produced.
	Livestock	Regular cleaning and disinfection of cattle shed and its premises should be carried out.		
	Fishery	Feed your fish at the same place every day, so that you will know whether your fish are eating well or not. If you give them too much, the food that is not eaten will stay on the bottom.		



Integrated Agromet Advisory Service Bulletin 11th to 15th Dec'13

Peren District

Bulletin No:93/2013

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 15th Dec '13</i>
<p>No rain occurred the past week. Maximum and minimum temperature range from 23⁰C to 24⁰C and 7⁰C to 8⁰C, respectively. Relative humidity varied from 38% to 98%. Wind speed ranged from 3 to 4 kmph.</p>			<p>There is possibility of no rain the coming week. Max temp is expected to be around 22⁰C - 24⁰C and min temp is expected to be around 7⁰C to 8⁰C. Possibility of mainly clear sky the coming week Relative Humidity will range from 35% to 97% The wind speed may reach upto 3 - 4 kmph Wind direction will blow from easterly</p>
Main Crops	Stage	Pest/ diseases	Agro-meteorological Advisories
Field crops			
Grain storage	Storage	-	Rice should be stored in gunny sacks on wooden platforms that allow free passage of air between grain sacks and the floor. Low temperature and low moisture content are necessary for storing rice grain for long period
Maize	Tasseling stage	Stalk borer	Farmers are advised to monitor the maize plant against stalk borer. The larvae may remain inside the whole and start damaging the tassel stalk. Application of Neem oil 2 % would minimize the infestation.
Mustard	Vegetative stage	-	Earthing up, gap filling along with weeding should be carried out.
Horticultural crops			
Probability no rain will occur the coming 3 or 4 days, therefore the farmers are advised to irrigate the crops daily or at an interval depending on the availability of water.			
Cabbage	Head formation stage	-	In case of cabbage, irrigation is very important for good growth and development.
Tomato	Vegetative stage	-	Frequent inter tillage and cultivation should be done in the field to keep it free from weeds. Proper pruning and staking should be carried out so as better and large tomatoes can be produced.
Bean	Vegetative stage	wilt	Symptoms are premature yellowing and whitening of leaves and finally the diving of the entire plant. Plant may continue to wilt in patches in quick succession till the maturity period. Disease affected plants should be pulled out.



<i>Livestock</i>	Regular cleaning and disinfection of cattle shed and its premises should be carried out.
<i>Fishery</i>	Feed your fish at the same place every day, so that you will know whether your fish are eating well or not. If you give them too much, the food that is not eaten will stay on the bottom.



Integrated Agromet Advisory Service Bulletin 11th to 15th Dec'13

Phek District

Bulletin No: 93/2013

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 15th Dec '13</i>
No rain occurred the past week Maximum and minimum temperatures ranged from 20 ⁰ C to 21 ⁰ C and 5 ⁰ C to 9 ⁰ C, respectively. Relative humidity varied from 39% to 98%. Wind speed ranged from 1 to 2 kmph.			There is possibility of no rain the coming week. Max temp is expected to be around 19 ⁰ C - 20 ⁰ C and min temp is expected to be around 8 ⁰ C to 9 ⁰ C. Possibility of mainly cloudy to clear sky the coming week Relative Humidity will range from 39% to 98% The wind speed 1- 2 kmph Wind direction will blow from southerly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Field crops</i>			
<i>Grain storage</i>	Storage	-	Rice should be stored in gunny sacks on wooden platforms that allow free passage of air between grain sacks and the floor. Low temperature and low moisture content are necessary for storing rice grain for long period
<i>Maize</i>	Tasseling stage	Stalk borer	Farmers are advised to monitor the maize plant against stalk borer. The larvae may remain inside the whole and start damaging the tassel stalk. Application of Neem oil 2 % would minimize the infestation.
<i>Mustard</i>	Vegetative stage	-	Earthing up, gap filling along with weeding should be carried out.
<i>Horticultural crops</i>			
Probability no rain will occur the coming 3 or 4 days, therefore the farmers are advised to irrigate the crops daily or at an interval depending on the availability of water.			
<i>Cabbage</i>	Head formation stage	-	In case of cabbage, irrigation is very important for good growth and development.
<i>Tomato</i>	Vegetative stage	-	Frequent inter tillage and cultivation should be done in the field to keep it free from weeds. Proper pruning and staking should be carried out so as better and large tomatoes can be produced.
<i>Livestock</i>	Regular cleaning and disinfection of cattle shed and its premises should be carried out.		

***Fishery***

Feed your fish at the same place every day, so that you will know whether your fish are eating well or not. If you give them too much, the food that is not eaten will stay on the bottom.




Integrated Agromet Advisory Service Bulletin 11th to 15th Dec'13

Tuensang District

Bulletin No:93/2013

<i>Weather summary of the preceding week</i>	<i>Weather forecast valid upto 15th Dec '13</i>
No rain occurred the past week Maximum and minimum temperatures ranged 17 ⁰ C to 19 ⁰ C and 3 ⁰ C to 5 ⁰ C, respectively. Relative humidity varied from 38% to 97%. Wind speed ranged from 1 to 2 kmph.	There is possibility of no rain the coming week. Max temp is expected to be around 16 ⁰ C - 18 ⁰ C and min temp is expected to be around 2 ⁰ C to 3 ⁰ C. Possibility of partly cloudy sky the coming week Relative humidity will range from 37% to 97% Wind speed 1 – 2 kmph Wind direction will blow from westerly

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Grain storage	Storage	-	Rice should be stored in gunny sacks on wooden platforms that allow free passage of air between grain sacks and the floor. Low temperature and low moisture content are necessary for storing rice grain for long period
Maize	Tasseling stage	Stalk borer	Farmers are advised to monitor the maize plant against stalk borer. The larvae may remain inside the whole and start damaging the tassel stalk. Application of Neem oil 2 % would minimize the infestation.
Mustard	Vegetative stage	-	Earthing up, gap filling along with weeding should be carried out.
Horticultural crops			
Probability no rain will occur the coming 3 or 4 days, therefore the farmers are advised to irrigate the crops daily or at an interval depending on the availability of water.			
Rajmash	Vegetative stage	wilt	Symptoms are premature yellowing and whitening of leaves and finally the diving of the entire plant. Plant may continue to wilt in patches in quick succession till the maturity period. Disease affected plants should be pulled out.
Cabbage	Head formation stage	-	In case of cabbage, irrigation is very important for good growth and development.
Potato	Vegetative stage	-	Proper cultural practices should be maintained, infested vines should be removed from the field for controlling the pest. Farmers are advised to carry out intercultural operations like hand weeding and

				earthingup to destroy weeds, to keep the soil loose for proper development of tubers and to cover the tubers with soil.
	Tomato	Vegetative stage	-	Frequent inter tillage and cultivation should be done in the field to keep it free from weeds. Proper pruning and staking should be carried out so as better and large tomatoes can be produced.
	Livestock	Regular cleaning and disinfection of cattle shed and its premises should be carried out.		
	Fishery	Feed your fish at the same place every day, so that you will know whether your fish are eating well or not. If you give them too much, the food that is not eaten will stay on the bottom.		



Integrated Agromet Advisory Service Bulletin 11th to 15th Dec'13 Wokha District

Bulletin No:93/2013


Weather summary of the preceding week

No rain occurred the past week.
Maximum and minimum temperatures ranged from 21⁰C to 23⁰C and 9⁰C to 10⁰C, respectively.
Relative humidity varied from 40% to 97%.
Wind speed ranged from 2 to 3 kmph

Weather forecast valid upto 15th Dec '13

There is possibility of no rain the coming week.
Max temp is expected to be around 20⁰C - 21⁰C and min temp is expected to be around 9⁰C to 10⁰C.
Possibility of mainly cloudy sky the coming week
Relative Humidity will range from 40% to 98%
The wind speed 2– 3 kmph
Wind direction will blow from southeasterly

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Field crops</i>			
<i>Grain storage</i>	Storage	-	Rice should be stored in gunny sacks on wooden platforms that allow free passage of air between grain sacks and the floor. Low temperature and low moisture content are necessary for storing rice grain for long period
<i>Maize</i>	Tasseling stage	Stalk borer	Farmers are advised to monitor the maize plant against stalk borer. The larvae may remain inside the whole and start damaging the tassel stalk. Application of Neem oil 2 % would minimize the infestation.
<i>Mustard</i>	Vegetative stage	-	Earthing up, gap filling along with weeding should be carried out.
<i>Horticultural crops</i>			
Probability no rain will occur the coming 3 or 4 days, therefore the farmers are advised to irrigate the crops daily or at an interval depending on the availability of water.			
<i>Cabbage</i>	Head formation stage	-	In case of cabbage, irrigation is very important for good growth and development.
<i>Bean</i>	Vegetative stage	wilt	Symptoms are premature yellowing and whitening of leaves and finally the diving of the entire plant. Plant may continue to wilt in patches in quick succession till the maturity period. Disease affected plants should be pulled out.
<i>Tomato</i>	Vegetative stage	-	Frequent inter tillage and cultivation should be done in the field to keep it free from weeds.

				Proper pruning and staking should be carried out so as better and large tomatoes can be produced.
	Livestock	Regular cleaning and disinfection of cattle shed and its premises should be carried out.		
	Fishery	Feed your fish at the same place every day, so that you will know whether your fish are eating well or not. If you give them too much, the food that is not eaten will stay on the bottom.		




--	--



Integrated Agromet Advisory Service Bulletin 11th to 15th Dec'13 Zunheboto District

Bulletin No:93/2013

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 15th Dec '13</i>
<p>No rain occurred the past week Maximum and minimum temperatures ranged from 21⁰C to 22⁰C and 3⁰C to 7⁰C, respectively. Relative humidity varied from 39% to 95%. Wind speed ranged from 1 to 2 kmph.</p>			<p>There is possibility of no rain the coming week. Max temp is expected to be around 19⁰C - 21⁰C and min temp is expected to be around 3⁰C to 4⁰C. Possibility of mainly clear to clear sky the coming week Relative Humidity will range from 39% to 95% The wind speed 2- 3 kmph Wind direction will blow southerly</p>
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Grain storage	Storage	-	Rice should be stored in gunny sacks on wooden platforms that allow free passage of air between grain sacks and the floor. Low temperature and low moisture content are necessary for storing rice grain for long period
Maize	Tasseling stage	Stalk borer	Farmers are advised to monitor the maize plant against stalk borer. The larvae may remain inside the whole and start damaging the tassel stalk. Application of Neem oil 2 % would minimize the infestation.
Mustard	Vegetative stage	-	Earthing up, gap filling along with weeding should be carried out.
Horticultural crops			
Probability no rain will occur the coming 3 or 4 days, therefore the farmers are advised to irrigate the crops daily or at an interval depending on the availability of water.			
Cabbage	Head formation stage	-	In case of cabbage, irrigation is very important for good growth and development.
Potato	Vegetative stage	-	Proper cultural practices should be maintained, infested vines should be removed from the field for controlling the pest. Farmers are advised to carry out intercultural operations like hand weeding and earthingup to destroy weeds, to keep the soil loose for proper development of tubers and to cover the tubers with soil.
Tomato	Vegetative	-	Frequent inter tillage and cultivation should be done in the field to keep it free

		stage	from weeds. Proper pruning and staking should be carried out so as better and large tomatoes can be produced.
	Livestock	Regular cleaning and disinfection of cattle shed and its premises should be carried out.	
	Fishery	Feed your fish at the same place every day, so that you will know whether your fish are eating well or not. If you give them too much, the food that is not eaten will stay on the bottom.	



Integrated Agromet Advisory Service Bulletin 11th to 15th Dec'13

Dimapur District

Bulletin No:93/2013


Weather summary of the preceding week

No rain occurred the past week.
Maximum and minimum temperatures ranged 24⁰C to 26⁰C and 10⁰C to 12⁰C, respectively.
Relative humidity varied from 41% to 96%.
Wind speed ranged from 2 to 4 kmph.

Weather forecast valid upto 15th Dec '13

There is possibility of no rain the coming week.
Max temp is expected to be around 22⁰C - 25⁰C and min temp is expected to be around 7⁰C to 8⁰C.
Possibility of mainly cloudy sky the coming week
Relative Humidity will range from 41% to 96% and
Wind speed 3– 4 kmph
Wind direction will be from easterly

Main Crops	Stage	Pest/ diseases	Agro-meteorological Advisories
Field crops			
Grain storage	Storage	-	Rice should be stored in gunny sacks on wooden platforms that allow free passage of air between grain sacks and the floor. Low temperature and low moisture content are necessary for storing rice grain for long period
Maize	Tasseling stage	Stalk borer	Farmers are advised to monitor the maize plant against stalk borer. The larvae may remain inside the whole and start damaging the tassel stalk. Application of Neem oil 2 % would minimize the infestation.
Mustard	Vegetative stage	-	Earthing up, gap filling along with weeding should be carried out.
Horticultural crops			
Probability no rain will occur the coming 3 or 4 days, therefore the farmers are advised to irrigate the crops daily or at an interval depending on the availability of water.			
Cabbage	Head formation stage	-	In case of cabbage, irrigation is very important for good growth and development.
Tomato	Vegetative stage	-	Frequent inter tillage and cultivation should be done in the field to keep it free from weeds. Proper pruning and staking should be carried out so as better and large tomatoes can be produced.
Broccoli	Vegetative stage	-	Broccoli requires sufficient moisture in the soil for uniform and continuous growth. Dry conditions adversely affect the quality and yield of shoots by being more fibrous
Bean	Vegetative stage	wilt	Symptoms are premature yellowing and whitening of leaves and finally the diving of the entire plant. Plant may continue to wilt in patches in quick succession till the maturity period. Disease affected plants should be pulled out.

	<i>Livestock</i>	Regular cleaning and disinfection of cattle shed and its premises should be carried out.
	<i>Fishery</i>	Feed your fish at the same place every day, so that you will know whether your fish are eating well or not. If you give them too much, the food that is not eaten will stay on the bottom.




Integrated Agromet Advisory Service Bulletin 11th to 15th Dec'13

Kiphire District

Bulletin No:93/2013

Weather summary of the preceding week			Weather forecast valid upto 15 th Dec '13
<p>No rain occurred the past week. Maximum and minimum temperatures ranged from 20⁰C to 23⁰C and 3⁰C to 9⁰C, respectively. Relative humidity varied from 38% to 98%. Wind speed ranged from 1 to 2 kmph.</p>			<p>There is possibility of no rain the coming week. Max temp is expected to be around 22⁰C - 23⁰C and min temp is expected to be around 8⁰C to 9⁰C. Possibility of mainly cloudy to clear sky the coming week Relative Humidity will range from 37% to 98% The wind speed 1- 2 kmph Wind direction will blow from westerly</p>
Main Crops	Stage	Pest/ diseases	Agro-meteorological Advisories
Field crops			
Grain storage	Storage	-	Rice should be stored in gunny sacks on wooden platforms that allow free passage of air between grain sacks and the floor. Low temperature and low moisture content are necessary for storing rice grain for long period
Maize	Tasseling stage	Stalk borer	Farmers are advised to monitor the maize plant against stalk borer. The larvae may remain inside the whole and start damaging the tassel stalk. Application of Neem oil 2 % would minimize the infestation.
Mustard	Vegetative stage	-	Earthing up, gap filling along with weeding should be carried out.
Horticultural crops			
Probability no rain will occur the coming 3 or 4 days, therefore the farmers are advised to irrigate the crops daily or at an interval depending on the availability of water.			
Rajmash	Vegetative stage	wilt	Symptoms are premature yellowing and whitening of leaves and finally the diving of the entire plant. Plant may continue to wilt in patches in quick succession till the maturity period. Disease affected plants should be pulled out.
Cabbage	Head formation stage	-	In case of cabbage, irrigation is very important for good growth and development.
Potato	Vegetative stage	-	Proper cultural practices should be maintained, infested vines should be removed from the field for controlling the pest. Farmers are advised to carry out intercultural operations like hand weeding and earthingup to destroy weeds, to keep the soil loose for proper development of tubers and to cover the tubers with soil.

	Tomato	Vegetative stage	-	<p>Frequent inter tillage and cultivation should be done in the field to keep it free from weeds.</p> <p>Proper pruning and staking should be carried out so as better and large tomatoes can be produced.</p>
	Livestock	Regular cleaning and disinfection of cattle shed and its premises should be carried out.		
	Fishery	Feed your fish at the same place every day, so that you will know whether your fish are eating well or not. If you give them too much, the food that is not eaten will stay on the bottom.		



Integrated Agromet Advisory Service Bulletin 11th to 15th Dec'13 Kohima District

Bulletin No:93/2013


Weather summary of the preceding week

No rain occurred the past week.
Maximum and minimum temperatures ranged from 22⁰C to 24⁰C and 7⁰C to 8⁰C, respectively.
Relative humidity varied from 37% to 98%.
Wind speed ranged from 3 to 4 kmph.

Weather forecast valid upto 15th Dec '13

There is possibility of no rain the coming week.
Max temp is expected to be around 21⁰C - 24⁰C and min temp is expected to be around 7⁰C to 8⁰C.
Possibility of mainly cloudy sky the coming week
Relative Humidity will range from 34% to 98%
Wind speed 2- 3 kmph
Wind direction will blow from easterly

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Grain storage	Storage	-	Rice should be stored in gunny sacks on wooden platforms that allow free passage of air between grain sacks and the floor. Low temperature and low moisture content are necessary for storing rice grain for long period
Maize	Tasseling stage	Stalk borer	Farmers are advised to monitor the maize plant against stalk borer. The larvae may remain inside the whole and start damaging the tassel stalk. Application of Neem oil 2 % would minimize the infestation.
Mustard	Vegetative stage	-	Earthing up, gap filling along with weeding should be carried out.
Horticultural crops			
Probability no rain will occur the coming 3 or 4 days, therefore the farmers are advised to irrigate the crops daily or at an interval depending on the availability of water.			
Cabbage	Head formation stage	-	In case of cabbage, irrigation is very important for good growth and development.
Potato	Vegetative stage	-	Proper cultural practices should be maintained, infested vines should be removed from the field for controlling the pest. Farmers are advised to carry out intercultural operations like hand weeding and earthingup to destroy weeds, to keep the soil loose for proper development of tubers and to cover the tubers with soil.
Broccoli	Vegetative stage	-	Broccoli requires sufficient moisture in the soil for uniform and continuous growth. Dry conditions adversely affect the quality and yield of shoots by being more fibrous

	<i>Tomato</i>	Vegetative stage	-	Frequent inter tillage and cultivation should be done in the field to keep it free from weeds. Proper pruning and staking should be carried out so as better and large tomatoes can be produced.
	<i>Livestock</i>	Regular cleaning and disinfection of cattle shed and its premises should be carried out.		
	<i>Fishery</i>	Feed your fish at the same place every day, so that you will know whether your fish are eating well or not. If you give them too much, the food that is not eaten will stay on the bottom.		