



## Integrated Agromet Advisory Service Bulletin 29<sup>th</sup> Sept to 3<sup>rd</sup> Oct '12 Dimapur District

**Bulletin No:74/2012**

Weather summary of the preceding week			Weather forecast valid upto 3 <sup>rd</sup> Oct '12
<p>Light rain occurred the past week. The <b>maximum</b> and the <b>minimum temperatures</b> ranged 29<sup>0</sup>C to 31<sup>0</sup>C and 23<sup>0</sup>C to 24<sup>0</sup>C, respectively. The <b>Relative humidity</b> varied from 75% to 93%. <b>Wind speed</b> ranged from 2 to 4 kmph.</p>			<p>There will be light to medium rain the coming week (4 – 15 mm) The <b>max temp</b> range from 28<sup>0</sup>C - 30<sup>0</sup>C and the <b>min temp</b> 23<sup>0</sup>C - 24<sup>0</sup>C. <b>Sky will be partly cloudy the coming days.</b> <b>Relative Humidity</b> will range from 74% to 93% and <b>Wind speed</b> 2 – 5 kmph <b>Wind direction</b> will be from westerly</p>
Main Crops	Stage	Pest/ diseases	Agricultural Advisories
<b>Wetland paddy</b>	Dough development stage	Gundhi bug	<b>Traditional method:</b> Placing a rotten frog/ dry fish in the field during milky stage, the foul smell of these attract bugs and keep busy on feeding carcass
<b>Tomato</b>	Nursery	-	The best soil for tomato is a fertile loam soil with more sand in the surface layer, and clay in the sub-surface layers and the most favourable range of soil pH is 6.0–7.0. The suitable height for tomato plant for transplanting is 7.5 cm to 10 cm with a good sturdy stem
<b>Papaya</b>	Growth stage	-	Earthing up should be carried out so as to improve drainage.
<b>Banana</b>	Vegetative stage	-	Unwanted suckers should be removed for reducing internal competition with the main plant. Small suckers should be removed on regular basis upto 7-8 months.
<b>Piggery</b>	Find out what diseases are prevalent in your area and arrange for vaccination and treatment if necessary. Internal parasites are a constant problem, and animals should be dewormed from time to time.		
<b>Duckery</b>			
<b>Poultry</b>	Plentiful of clean and fresh water is very much essential. All waterers should be cleaned daily.		



<b><i>Poultry</i></b>	Plentiful of clean and fresh water is very much essential. All waterers should be cleaned daily.
<b><i>Fishery</i></b>	Feed your fish at the same place every day, so that you will know whether your fish are eating well or not. If you give them too much, the food that is not eaten will stay on the bottom



## Integrated Agromet Advisory Service Bulletin 29<sup>th</sup> Sept to 3<sup>rd</sup> Oct '12 Kiphire District

**Bulletin No:74/2012**

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 3<sup>rd</sup> Oct '12</i>
<p>Light rain occurred the past week. The <b>maximum</b> and the <b>minimum temperatures</b> ranged from 25<sup>0</sup>C to 26<sup>0</sup>C and 18<sup>0</sup>C to 19<sup>0</sup>C, respectively. The <b>Relative humidity</b> varied from 72% to 98%. <b>Wind speed</b> ranged from 1 to 2 kmph.</p>			<p>There will be light to medium rain the coming week (3 to 14 mm) The <b>maximum temp</b> range 24<sup>0</sup>C - 27<sup>0</sup>C and the <b>minimum temp</b> 18<sup>0</sup>C to 19<sup>0</sup>C. <b>Sky will be mainly cloudy the coming days.</b> <b>Relative Humidity</b> will range from 78% to 96% The <b>wind speed</b> 1- 2 kmph Wind direction will blow from southerly.</p>
Main Crops	Stage	Pest/ diseases	Agricultural Advisories
<b>Jhum paddy</b>	Harvesting stage	-	As the grain is to be harvested, the store house should be fumigated; store bins cleaned and should be weather proof, rodent proof and damp proof.
<b>Wetland Paddy</b>	Dough development stage	Gundhi bug	<b>Traditional method:</b> Placing a rotten frog/ dry fish in the field during milky stage, the foul smell of these attract bugs and keep busy on feeding carcass
<b>Tomato</b>	Nursery	-	The best soil for tomato is a fertile loam soil with more sand in the surface layer, and clay in the sub-surface layers and the most favourable range of soil pH is 6.0–7.0. The suitable height for tomato plant for transplanting is 7.5 cm to 10 cm with a good sturdy stem
<b>French bean</b>	Vegetative stage	-	Second weeding will be essential 50 days later after the first weeding.
<b>Piggery</b>	Find out what diseases are prevalent in your area and arrange for vaccination and treatment if necessary. Internal parasites are a constant problem, and animals should be dewormed from time to time.		
<b>Poultry</b>	Plentiful of clean and fresh water is very much essential. All waterers should be cleaned daily.		



<b><i>Fishery</i></b>	Feed your fish at the same place every day, so that you will know whether your fish are eating well or not. If you give them too much, the food that is not eaten will stay on the bottom



भा क अतुप - पु प सं  
ICAR - RCNEH

ICAR RESEARCH COMPLEX FOR NEH REGION  
(Indian Council of Agricultural Research)  
NAGALAND CENTER, JHARNAPANI, MEDZIPHEMA



# Integrated Agromet Advisory Service Bulletin 29<sup>th</sup> Sept to 3<sup>rd</sup> Oct '12

## Kohima District

**Bulletin No:74/2012**

Weather summary of the preceding week			Weather forecast valid upto 3 <sup>rd</sup> Oct '12
<p>Light rain occurred the past week. The <b>maximum</b> and the <b>minimum temperatures</b> ranged from 27<sup>0</sup>C to 28<sup>0</sup>C and 21<sup>0</sup>C to 22<sup>0</sup>C, respectively. The <b>Relative humidity</b> varied from 85% to 97%. <b>Wind speed</b> ranged from 2 to 4 kmph.</p>			<p>There will be light rain the coming week (5 to 15 mm) The <b>max temp</b> ranged 25<sup>0</sup>C - 28<sup>0</sup>C and the <b>min temp</b> 21<sup>0</sup>C - 22<sup>0</sup>C. <b>Sky will be cloudy the coming week.</b> <b>Relative Humidity</b> will range from 91% to 98% <b>Wind speed</b> 2 – 5 kmph <b>Wind direction</b> will blow from easterly to southerly</p>
Main Crops	Stage	Pest/ Diseases	Agricultural Advisories
<b>Wetland Paddy</b>	Dough development stage	Gundhi bug	<b>Traditional method:</b> Placing a rotten frog/ dry fish in the field during milky stage, the foul smell of these attract bugs and keep busy on feeding carcass
<b>Cabbage</b>	Nursery		Four to six weeks old seedlings should be used for transplantation.
<b>Tomato</b>	Nursery	-	The best soil for tomato is a fertile loam soil with more sand in the surface layer, and clay in the sub-surface layers and the most favourable range of soil pH is 6.0–7.0. The suitable height for tomato plant for transplanting is 7.5 cm to 10 cm with a good sturdy stem
<b>Colocasia</b>	Vegetative stage	-	Keeping the field clean and free of weeds is essential for better growth.
<b>Piggery</b>	Find out what diseases are prevalent in your area and arrange for vaccination and treatment if necessary. Internal parasites are a constant problem, and animals should be dewormed from time to time.		
<b>Poultry</b>	Plentiful of clean and fresh water is very much essential. All waterers should be cleaned daily.		



***Fishery***

Feed your fish at the same place every day, so that you will know whether your fish are eating well or not. If you give them too much, the food that is not eaten will stay on the bottom



# Integrated Agromet Advisory Service Bulletin 29<sup>th</sup> Sept to 3<sup>rd</sup> Oct '12 Longleng District

**Bulletin No:74/2012**

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 3<sup>rd</sup> Oct '12</i>
<p>Light rain occurred the past week. The <b>maximum</b> and the <b>minimum temperatures</b> ranged from 27<sup>0</sup>C to 29<sup>0</sup>C and 22<sup>0</sup>C to 23<sup>0</sup>C, respectively. The <b>Relative humidity</b> varied from 78% to 95%. <b>Wind speed</b> ranged from 2 to 4 kmph.</p>			<p>There will be light to medium rain the coming week (9 to 15 mm) The <b>maximum temp</b> range 26<sup>0</sup>C - 29<sup>0</sup>C and the <b>minimum temp</b> 22<sup>0</sup>C to 23<sup>0</sup>C. <b>Sky will be cloudy the coming days.</b> <b>Relative Humidity</b> will range from 85% to 93% The <b>wind speed</b> 2 – 4 kmph Wind direction will blow from southerly</p>
Main Crops	Stage	Pest/ diseases	Agricultural Advisories
<i>Jhum paddy</i>	Harvesting stage	-	As the grain is to be harvested, the store house should be fumigated; store bins cleaned and should be weather proof, rodent proof and damp proof.
<i>Wetland paddy</i>	Dough development stage	Gundhi bug	<b>Traditional method:</b> Placing a rotten frog/ dry fish in the field during milky stage, the foul smell of these attract bugs and keep busy on feeding carcass
	Inflorescence stage	-	Monitor the plants pest and diseases.
<i>Banana</i>	Vegetative stage	-	Unwanted suckers should be removed for reducing internal competition with the main plant. Small suckers should be removed on regular basis upto 7-8 months.
<i>Tomato</i>	Nursery	-	The best soil for tomato is a fertile loam soil with more sand in the surface layer, and clay in the sub-surface layers and the most favourable range of soil pH is 6.0–7.0. The suitable height for tomato plant for transplanting is 7.5 cm to 10 cm with a good sturdy stem
<i>Piggery</i>	Find out what diseases are prevalent in your area and arrange for vaccination and treatment if necessary. Internal parasites are a constant problem, and animals should be dewormed from time to time.		



<b><i>Poultry</i></b>	Plentiful of clean and fresh water is very much essential. All waterers should be cleaned daily.
<b><i>Fishery</i></b>	Feed your fish at the same place every day, so that you will know whether your fish are eating well or not. If you give them too much, the food that is not eaten will stay on the bottom





# Integrated Agromet Advisory Service Bulletin 29<sup>th</sup> Sept to 3<sup>rd</sup> Oct '12

## Mokokchung District

Bulletin No:74/2012

Weather summary of the preceding week			Weather forecast valid upto 3 <sup>rd</sup> Oct '12
<p>Light rain occurred the past week</p> <p>The <b>maximum and the minimum temperatures</b> ranged 26<sup>0</sup>C to 29<sup>0</sup>C and 21<sup>0</sup>C to 23<sup>0</sup>C, respectively.</p> <p>The <b>Relative humidity</b> varied from 82% to 98%.</p> <p><b>Wind speed</b> ranged from 2 to 4 kmph.</p>			<p>There will be light to medium rain the coming week (9 to 13 mm)</p> <p>The <b>max temp</b> range 26<sup>0</sup>C - 29<sup>0</sup>C and the <b>min temp</b> 21<sup>0</sup>C to 22<sup>0</sup>C.</p> <p><b>Sky will be partly cloudy the coming days.</b></p> <p><b>Relative Humidity</b> will range from 85% to 93% and</p> <p><b>Wind speed</b> 2 - 4 kmph</p> <p>Wind direction will blow from southerly</p>
Main Crops	Stage	Pest/ Diseases	Agricultural Advisories
<b>Jhum paddy</b>	Harvesting stage	-	As the grain is to be harvested, the store house should be fumigated; store bins cleaned and should be weather proof, rodent proof and damp proof.
<b>Wetland Paddy</b>	Milk development stage	Gundhi bug	<b>Traditional method:</b> Placing a rotten frog/ dry fish in the field during milky stage, the foul smell of these attract bugs and keep busy on feeding carcass
<b>Tomato</b>	Nursery	-	The best soil for tomato is a fertile loam soil with more sand in the surface layer, and clay in the sub-surface layers and the most favourable range of soil pH is 6.0–7.0. The suitable height for tomato plant for transplanting is 7.5 cm to 10 cm with a good sturdy stem
<b>French bean</b>	Vegetative stage	-	Give support to the plant
<b>Cabbage</b>	Nursery		Four to six weeks old seedlings should be used for transplantation.
<b>Piggery</b>	Find out what diseases are prevalent in your area and arrange for vaccination and treatment if necessary. Internal parasites are a constant problem, and animals should be dewormed from time to time.		
<b>Poultry</b>	Plentiful of clean and fresh water is very much essential. All waterers should be cleaned daily.		
<b>Fishery</b>	Feed your fish at the same place every day, so that you will know whether your fish are eating well or not. If you give them too much, the food that is not eaten will stay on the bottom		



# Integrated Agromet Advisory Service Bulletin from 29<sup>th</sup> Sept to 3<sup>rd</sup> Oct'12

## Mon District

**Bulletin No:74/2012**

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 3<sup>rd</sup> Oct '12</i>
Light rain occurred the past week. The <b>maximum</b> and the <b>minimum temperatures</b> ranged 25 <sup>0</sup> C to 29 <sup>0</sup> C and 21 <sup>0</sup> C to 22 <sup>0</sup> C, respectively. The <b>Relative humidity</b> varied from 84% to 96%. <b>Wind speed</b> ranged from 2 to 4 kmph.			There will be light rain the coming week (7 – 10 mm) The <b>max temp</b> range from 25 <sup>0</sup> C - 27 <sup>0</sup> C and the <b>min temp</b> 21 <sup>0</sup> C - 22 <sup>0</sup> C. <b>Sky will be partly cloudy the coming days.</b> <b>Relative Humidity</b> will range from 87% to 95% and <b>Wind speed</b> 2 – 4 kmph <b>Wind direction</b> will be from southerly
Main Crops	Stage	Pest/ Diseases	Agricultural Advisories
<i>Jhum paddy</i>	Harvesting stage	-	As the grain is to be harvested, the store house should be fumigated; store bins cleaned and should be weather proof, rodent proof and damp proof.
<i>Wetland paddy</i>	Milk development stage	Gundhi bug	<b>Traditional method:</b> Placing a rotten frog/ dry fish in the field during milky stage, the foul smell of these attract bugs and keep busy on feeding carcass
<i>Soybean</i>	Vegetative stage	-	Crop rotation should be carried out as it break disease and insect cycles and increase yield
<i>Piggery</i>	Find out what diseases are prevalent in your area and arrange for vaccination and treatment if necessary. Internal parasites are a constant problem, and animals should be dewormed from time to time.		
<i>Poultry</i>	Plentiful of clean and fresh water is very much essential. All waterers should be cleaned daily.		
<i>Fishery</i>	Feed your fish at the same place every day, so that you will know whether your fish are eating well or not. If you give them too much, the food that is not eaten will stay on the bottom.		



## Integrated Agromet Advisory Service Bulletin 29<sup>th</sup> Sept to 3<sup>rd</sup> Oct '12 Peren District

**Bulletin No:74/2012**

<i>Weather summary of the preceding week</i>	<i>Weather forecast valid upto 3<sup>rd</sup> Oct '12</i>
<p>Light rain occurred the past week. The <b>maximum</b> and the <b>minimum temperatures</b> ranged from 26<sup>0</sup>C to 29<sup>0</sup>C and 21<sup>0</sup>C to 22<sup>0</sup>C, respectively. The <b>Relative humidity</b> varied from 83% to 96%. <b>Wind speed</b> ranged from 2 to 5 kmph.</p>	<p>There will be light rain the coming week (3 to 10 mm) The <b>maximum temp</b> range 25<sup>0</sup>C - 27<sup>0</sup>C and the <b>minimum temp</b> 21<sup>0</sup>C to 22<sup>0</sup>C. <b>Sky will be cloudy the coming days.</b> <b>Relative Humidity</b> will range from 87% to 95% The <b>wind speed</b> 2- 4 kmph Wind direction will blow from easterly to southeasterly.</p>

Main Crops	Stage	Pest/ diseases	Agricultural Advisories
<b>Wetland paddy</b>	Booting stage	Stem borer	Release of the egg parasitoid <i>Trichogramma japonicum</i> @ 50,000 adults /ha 5 times starting from 30 days after transplanting for control of stem borers.
<b>Tomato</b>	Nursery	-	The best soil for tomato is a fertile loam soil with more sand in the surface layer, and clay in the sub-surface layers and the most favourable range of soil pH is 6.0–7.0. The suitable height for tomato plant for transplanting is 7.5 cm to 10 cm with a good sturdy stem
<b>Papaya</b>	Growth stage	-	Earthing up should be carried out so as to improve drainage.
<b>Banana</b>	Vegetative stage	-	Unwanted suckers should be removed for reducing internal competition with the main plant. Small suckers should be removed on regular basis upto 7-8 months.
<b>Piggery</b>	Find out what diseases are prevalent in your area and arrange for vaccination and treatment if necessary. Internal parasites are a constant problem, and animals should be dewormed from time to time.		
<b>Poultry</b>	Plentiful of clean and fresh water is very much essential. All waterers should be cleaned daily.		



<b><i>Fishery</i></b>	Feed your fish at the same place every day, so that you will know whether your fish are eating well or not. If you give them too much, the food that is not eaten will stay on the bottom



# Integrated Agromet Advisory Service Bulletin 29<sup>th</sup> Sept to 3<sup>rd</sup> Oct '12 Phek District

**Bulletin No:74/2012**

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 3<sup>rd</sup> Oct '12</i>
<p>Light rain occurred the past week. The <b>maximum</b> and the <b>minimum temperatures</b> ranged from 24<sup>0</sup>C to 25<sup>0</sup>C and 18<sup>0</sup>C to 19<sup>0</sup>C, respectively. The <b>Relative humidity</b> varied from 83% to 97%. <b>Wind speed</b> ranged from 1 to 2 kmph.</p>			<p>There will be light rain the coming week (3 to 9 mm) The <b>maximum temp</b> range 22<sup>0</sup>C - 25<sup>0</sup>C and the <b>minimum temp</b> 17<sup>0</sup>C to 18<sup>0</sup>C. <b>Sky will be partly cloudy the coming days.</b> <b>Relative Humidity</b> will range from 88% to 95% The <b>wind speed</b> 2 - 5 kmph Wind direction will blow from southeasterly.</p>
Main Crops	Stage	Pest/ Diseases	Agricultural Advisories
<b>Jhum Paddy</b>	Harvesting stage	-	As the grain is to be harvested, the store house should be fumigated; store bins cleaned and should be weather proof, rodent proof and damp proof.
<b>Wetland Paddy</b>	Milk development stage	Gundhi bug	<b>Traditional method:</b> Placing a rotten frog/ dry fish in the field during milky stage, the foul smell of these attract bugs and keep busy on feeding carcass
<b>Colocasia</b>	Vegetative stage	-	Keeping the field clean and free of weeds is essential for better growth.
<b>Tomato</b>	Nursery	-	The best soil for tomato is a fertile loam soil with more sand in the surface layer, and clay in the sub-surface layers and the most favourable range of soil pH is 6.0–7.0. The suitable height for tomato plant for transplanting is 7.5 cm to 10 cm with a good sturdy stem
<b>Cabbage</b>	Nursery		Four to six weeks old seedlings should be used for transplantation.
<b>Fishery</b>	Feed your fish at the same place every day, so that you will know whether your fish are eating well or not. If you give them too much, the food that is not eaten will stay on the bottom		
<b>Piggery</b>	Find out what diseases are prevalent in your area and arrange for vaccination and treatment if necessary. Internal parasites are a constant problem, and animals should be dewormed from time to time.		
<b>Poultry</b>	Plentiful of clean and fresh water is very much essential. All waterers should be cleaned daily.		



# Integrated Agromet Advisory Service Bulletin 29<sup>th</sup> Sept to 3<sup>rd</sup> Oct '12 Tuensang District

**Bulletin No:74/2012**

## *Weather summary of the preceding week*

Light rain occurred the past week  
The **maximum** and the **minimum temperatures** ranged 25<sup>0</sup>C to 26<sup>0</sup>C and 18<sup>0</sup>C to 19<sup>0</sup>C, respectively.  
The **Relative humidity** varied from 80% to 97%.  
**Wind speed** ranged from 2 to 4 kmph.

## *Weather forecast valid upto 3<sup>rd</sup> Oct '12*

There will be light to medium rain the coming week (7 to 10 mm)  
The **max temp** range 24<sup>0</sup>C - 27<sup>0</sup>C and the **min temp** 18<sup>0</sup>C to 19<sup>0</sup>C.  
**Sky will be partly cloudy the coming days.**  
**Relative Humidity** will range from 86% to 97% and  
**Wind speed** 2 - 3 kmph  
Wind direction will blow from southerly

Main Crops	Stage	Pest/ Diseases	Agricultural Advisories
<i>Jhum paddy</i>	Harvesting stage		As the grain is to be harvested, the store house should be fumigated; store bins cleaned and should be weather proof, rodent proof and damp proof.
<i>Wetland paddy</i>	Dough development stage		
<i>Colocasia</i>	Vegetative stage	-	Keeping the field clean and free of weeds is essential for better growth.
<i>Pea</i>	Sowing stage	-	It should be sown in light soil, does not thrive well on highly acidic soil. FYM should be applied during field preparation.
<i>Cabbage</i>	Sowing stage	-	Organic manure 25 t/ha is recommended
<i>Piggery</i>	Find out what diseases are prevalent in your area and arrange for vaccination and treatment if necessary. Internal parasites are a constant problem, and animals should be dewormed from time to time.		
<i>Poultry</i>	Plentiful of clean and fresh water is very much essential. All waterers should be cleaned daily.		





<b><i>Poultry</i></b>	Plentiful of clean and fresh water is very much essential. All waterers should be cleaned daily.
<b><i>Fishery</i></b>	Feed your fish at the same place every day, so that you will know whether your fish are eating well or not. If you give them too much, the food that is not eaten will stay on the bottom



# Integrated Agromet Advisory Service Bulletin 29<sup>th</sup> Sept to 3<sup>rd</sup> Oct '12 Wokha District

**Bulletin No:74/2012**

## Weather summary of the preceding week

Light rain occurred the past week.  
The **maximum** and the **minimum temperatures** ranged from 27<sup>0</sup>C to 29<sup>0</sup>C and 21<sup>0</sup>C to 22<sup>0</sup>C, respectively.  
The **Relative humidity** varied from 81% to 95%.  
**Wind speed** ranged from 2 to 3 kmph

## Weather forecast valid upto 3<sup>rd</sup> Oct '12

There will be light rain the coming week (9 to 19 mm)  
The **max temp** 25<sup>0</sup>C - 28<sup>0</sup>C and **min temp** 21<sup>0</sup>C - 22<sup>0</sup>C.  
**Sky will be partly cloudy the coming days.**  
**Relative Humidity** will range from 86% to 95%  
The **wind speed** 2 – 4 kmph  
Wind direction will blow from southeasterly

Main Crops	Stage	Pest/ Diseases	Agricultural Advisories
<b>Jhum paddy</b>	Dough development stage	Gundhi bug	<b>Traditional method:</b> Placing a rotten frog/ dry fish in the field during milky stage, the foul smell of these attract bugs and keep busy on feeding carcass
<b>Wetland paddy</b>	Inflorescence stage	-	Proper monitoring should be done against pest and diseases
<b>Colocasia</b>	Vegetative stage	-	Keeping the field clean and free of weeds is essential for better growth.
<b>Tomato</b>	Nursery	-	The best soil for tomato is a fertile loam soil with more sand in the surface layer, and clay in the sub-surface layers and the most favourable range of soil pH is 6.0–7.0. The suitable height for tomato plant for transplanting is 7.5 cm to 10 cm with a good sturdy stem
<b>Fishery</b>	Feed your fish at the same place every day, so that you will know whether your fish are eating well or not. If you give them too much, the food that is not eaten will stay on the bottom		
<b>Piggery</b>	Find out what diseases are prevalent in your area and arrange for vaccination and treatment if necessary. Internal parasites are a constant problem, and animals should be dewormed from time to time.		
<b>Poultry</b>	Plentiful of clean and fresh water is very much essential. All waterers should be cleaned daily.		







--	--



## Integrated Agromet Advisory Service Bulletin 29<sup>th</sup> Sept to 3<sup>rd</sup> Oct '12 Zunheboto District

**Bulletin No:74/2012**

Weather summary of the preceding week			Weather forecast valid upto 3 <sup>rd</sup> Oct '12
<p>Light rain occurred the past week The <b>maximum</b> and the <b>minimum temperatures</b> ranged from 24<sup>0</sup>C to 25<sup>0</sup>C and 19<sup>0</sup>C to 20<sup>0</sup>C, respectively. The <b>Relative humidity</b> varied from 86% to 97%. <b>Wind speed</b> ranged from 2 to 3 kmph.</p>			<p>There will be light rain the coming week (4 to 10 mm) The <b>maximum temp</b> range 23<sup>0</sup>C - 25<sup>0</sup>C and the <b>minimum temp</b> 19<sup>0</sup>C to 20<sup>0</sup>C. <b>Sky will be partly cloudy the coming days.</b> <b>Relative Humidity</b> will range from 89% to 93% The <b>wind speed</b> 2- 4 kmph Wind direction will blow from southeasterly</p>
Main Crops	Stage	Pest/ Diseases	Agricultural Advisories
<b>Jhum Paddy</b>	Harvesting stage	-	As the grain is to be harvested, the store house should be fumigated; store bins cleaned and should be weather proof, rodent proof and damp proof.
<b>Wetland Paddy</b>	Milk development stage	Gundhi bug	<b>Traditional method:</b> Placing a rotten frog/ dry fish in the field during milky stage, the foul smell of these attract bugs and keep busy on feeding carcass
<b>Tomato</b>	Nursery	-	The best soil for tomato is a fertile loam soil with more sand in the surface layer, and clay in the sub-surface layers and the most favourable range of soil pH is 6.0–7.0. The suitable height for tomato plant for transplanting is 7.5 cm to 10 cm with a good sturdy stem
<b>Soybean</b>	Vegetative stage	-	Crop rotation should be carried out as it break disease and insect cycles and increase yield
<b>Papaya</b>	Growth stage	-	Earthing up should be carried out so as to improve drainage.
<b>Piggery</b>	Find out what diseases are prevalent in your area and arrange for vaccination and treatment if necessary. Internal parasites are a constant problem, and animals should be dewormed from time to time.		
<b>Poultry</b>	Plentiful of clean and fresh water is very much essential. All waterers should be cleaned daily.		



<b><i>Poultry</i></b>	Plentiful of clean and fresh water is very much essential. All waterers should be cleaned daily.
<b><i>Fishery</i></b>	Feed your fish at the same place every day, so that you will know whether your fish are eating well or not. If you give them too much, the food that is not eaten will stay on the bottom