




Integrated Agromet Advisory Service Bulletin 18th to 22nd Jan'14 Kohima District

Bulletin No:05/2014

Weather summary of the preceding week			Weather forecast valid upto 22 nd Jan'14
<p>No rain occurred the past week. Maximum and minimum temperatures ranged from 20⁰C to 22⁰C and 2⁰C to 8⁰C, respectively. Relative humidity varied from 27% to 96%. Wind speed ranged from 2 to 3 kmph.</p>			<p>There is possibility of no rain the coming week. Max temp is expected to be around 21⁰C - 22⁰C and min temp is expected to be around 7⁰C to 8⁰C. Possibility of clear sky the coming week Relative Humidity will range from 29% to 97% Wind speed 3- 4 kmph Wind direction will blow from easterly</p>
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Rice	Storage	-	<p>It should be ensured that the bags are clean, dry and well stitched, to keep the risk of loss as low as possible right from the start. Care should be taken as high proportion of broken grains encourages the development of microorganisms, mites and beetles Rice is particularly likely to be infested with storage pests if it has been stored for more than two months.</p>
Maize	Post -harvesting stage	-	<p>Provide protection from common storage loss agents such as insect pests, rodents, moulds, birds etc</p>
Mustard	Podding stage	Aphid	<p>Aphids suck sap as a result of it the plants remain stunted, flowers dry up and seeds do not develop. So proper monitoring should be carried out.</p>
Horticultural crops			
<p>Probability no rain will occur the coming 3 or 4 days, therefore the farmers are advised to irrigate the crops daily or at an interval depending on the availability of water.</p>			
Citrus	Nursery stage	-	<p>In citrus nursery, diseases may appear any time of plant growth through contaminated water, soil and even through nursery workers and implements. Therefore a regular monitoring should be done for infection and in case of infection; the infected/contaminated plants should immediately be uprooted and destroyed to keep the nursery totally free.</p>
Vegetables	Harvesting stage	-	<p>Harvest vegetables at their right stage and storing them properly will help extend their shelf life</p>

	<i>Cabbage</i>	Heading stage		Harvesting should be done when the heads are large and solid
	<i>Ginger</i>	Harvesting stage		Harvesting should be done when the foliage is completely dried out.
	<i>Fishery</i>	Never over feed your fish. Excess food will pollute the pond water, so only feed amounts that will be completely consumed by your fish within 2 minutes.		
	<i>Piggery</i>	Good housekeeping practices such as removing manure, washing down pens, disinfecting and generally cleaning up the facility are essential to maintaining a healthy herd.		
	<i>Poultry</i>	During cold weather, cool outside air should be prevented from contacting birds.		



Integrated Agromet Advisory Service Bulletin 18th to 22nd Jan'14

Longleng District

Bulletin No:05/2014


Weather summary of the preceding week	Weather forecast valid upto 22 nd Jan'14
<p>No rain occurred the past week. Maximum and minimum temperatures ranged from 20⁰C to 23⁰C and 3⁰C to 8⁰C, respectively. Relative humidity varied from 32% to 86%. Wind speed ranged from 2 to 4 kmph.</p>	<p>There is possibility of no rain the coming week. Max temp is expected to be around 22⁰C - 23⁰C and min temp is expected to be around 7⁰C to 8⁰C. Possibility of clear sky the coming week Relative Humidity will range from 29% to 80% The wind speed 3 –4 kmph Wind direction will blow from southeasterly</p>

Main Crops	Stage	Pest/ diseases	Agro-meteorological Advisories
Field crops			
Rice	Storage	-	<p>It should be ensured that the bags are clean, dry and well stitched, to keep the risk of loss as low as possible right from the start. Care should be taken as high proportion of broken grains encourages the development of microorganisms, mites and beetles Rice is particularly likely to be infested with storage pests if it has been stored for more than two months.</p>
Maize	Post -harvesting stage	-	Provide protection from common storage loss agents such as insect pests, rodents, moulds, birds etc
Mustard	Podding stage	Aphid	Aphids suck sap as a result of it the plants remain stunted, flowers dry up and seeds do not develop. So proper monitoring should be carried out.

Horticultural crops

Probability no rain will occur the coming 3 or 4 days, therefore the farmers are advised to irrigate the crops daily or at an interval depending on the availability of water.

Citrus	Nursery stage	-	In citrus nursery, diseases may appear any time of plant growth through contaminated water, soil and even through nursery workers and implements. Therefore a regular monitoring should be done for infection and in case of infection; the infected/contaminated plants should immediately be uprooted and destroyed to keep the nursery totally free.
Vegetables	Harvesting stage	-	Harvest vegetables at their right stage and storing them properly will help extend their shelf life
Cabbage	Heading stage		Harvesting should be done when the heads are large and solid

	<i>Ginger</i>	Harvesting stage		Harvesting should be done when the foliage is completely dried out.
	<i>Fishery</i>	Never over feed your fish. Excess food will pollute the pond water, so only feed amounts that will be completely consumed by your fish within 2 minutes.		
	<i>Piggery</i>	Good housekeeping practices such as removing manure, washing down pens, disinfecting and generally cleaning up the facility are essential to maintaining a healthy herd.		
	<i>Poultry</i>	During cold weather, cool outside air should be prevented from contacting birds.		




Integrated Agromet Advisory Service Bulletin 18th to 22nd Jan'14

Mokokchung District

Bulletin No: 05/2014

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 22nd Jan'14</i>
<p>No rain occurred the past week Maximum and minimum temperatures ranged 15⁰C to 24⁰C and 3⁰C to 8⁰C, respectively. Relative humidity varied from 33% to 92%. Wind speed ranged from 2 to 4 kmph.</p>			<p>There is possibility of no rain the coming week. Max temp is expected to be around 23⁰C - 24⁰C and min temp is expected to be around 7⁰C to 8⁰C. Possibility of clear sky the coming week Relative Humidity will range from 30% to 84% and Wind speed 3– 4 kmph Wind direction will blow from southeasterly.</p>
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Rice	Storage	-	<p>It should be ensured that the bags are clean, dry and well stitched, to keep the risk of loss as low as possible right from the start. Care should be taken as high proportion of broken grains encourages the development of microorganisms, mites and beetles Rice is particularly likely to be infested with storage pests if it has been stored for more than two months.</p>
Maize	Post -harvesting stage	-	<p>Provide protection from common storage loss agents such as insect pests, rodents, moulds, birds etc</p>
Mustard	Podding stage	Aphid	<p>Aphids suck sap as a result of it the plants remain stunted, flowers dry up and seeds do not develop. So proper monitoring should be carried out.</p>
Horticultural crops			
<p>Probability no rain will occur the coming 3 or 4 days, therefore the farmers are advised to irrigate the crops daily or at an interval depending on the availability of water.</p>			
Citrus	Nursery stage	-	<p>In citrus nursery, diseases may appear any time of plant growth through contaminated water, soil and even through nursery workers and implements. Therefore a regular monitoring should be done for infection and in case of infection; the infected/contaminated plants should immediately be uprooted and destroyed to keep the nursery totally free.</p>
Vegetables	Harvesting stage	-	<p>Harvest vegetables at their right stage and storing them properly will help extend their shelf life</p>
Cabbage	Heading stage		<p>Harvesting should be done when the heads are large and solid</p>

	<i>Ginger</i>	Harvesting stage		Harvesting should be done when the foliage is completely dried out.
	<i>Fishery</i>	Never over feed your fish. Excess food will pollute the pond water, so only feed amounts that will be completely consumed by your fish within 2 minutes.		
	<i>Piggery</i>	Good housekeeping practices such as removing manure, washing down pens, disinfecting and generally cleaning up the facility are essential to maintaining a healthy herd.		
	<i>Poultry</i>	During cold weather, cool outside air should be prevented from contacting birds.		



Integrated Agromet Advisory Service Bulletin from 18th to 22nd Jan'14 Mon District

Bulletin No:05/2014


Weather summary of the preceding week

- No rain occurred the past week
- **Maximum** and **minimum temperatures** ranged 18⁰C to 24⁰C and 4⁰C to 10⁰C, respectively.
- **Relative humidity** varied from 32% to 89%.
- **Wind speed** ranged from 2 to 4 kmph.

Weather forecast valid upto 22nd Jan'14

- Possibility of no rain the coming week.
- The **max temp** is likely to be 23⁰C - 24⁰C and the **min temp** 9⁰C - 10⁰C.
- **Sky is likely to be clear the coming week**
- **Relative Humidity** is likely to range from 30% to 80%
- **Wind speed** may reach upto 3- 4kmph
- **Wind direction** will be from southerly

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Rice	Storage	-	It should be ensured that the bags are clean, dry and well stitched, to keep the risk of loss as low as possible right from the start. Care should be taken as high proportion of broken grains encourages the development of microorganisms, mites and beetles Rice is particularly likely to be infested with storage pests if it has been stored for more than two months.
Maize	Post -harvesting stage	-	Provide protection from common storage loss agents such as insect pests, rodents, moulds, birds etc
Mustard	Podding stage	Aphid	Aphids suck sap as a result of it the plants remain stunted, flowers dry up and seeds do not develop. So proper monitoring should be carried out.
Horticultural crops			
Probability no rain will occur the coming 3 or 4 days too, therefore the farmers are advised to irrigate the crops daily or at an interval depending on the availability of water.			
Citrus	Nursery stage	-	In citrus nursery, diseases may appear any time of plant growth through contaminated water, soil and even through nursery workers and implements. Therefore a regular monitoring should be done for infection and in case of infection; the infected/contaminated plants should immediately be uprooted and destroyed to keep the nursery totally free.
Vegetables	Harvesting stage	-	Harvest vegetables at their right stage and storing them properly will help extend their shelf life
Cabbage	Heading stage		Harvesting should be done when the heads are large and solid

	<i>Ginger</i>	Harvesting stage		Harvesting should be done when the foliage is completely dried out.
	<i>Fishery</i>	Never over feed your fish. Excess food will pollute the pond water, so only feed amounts that will be completely consumed by your fish within 2 minutes.		
	<i>Piggery</i>	Good housekeeping practices such as removing manure, washing down pens, disinfecting and generally cleaning up the facility are essential to maintaining a healthy herd.		
	<i>Poultry</i>	During cold weather, cool outside air should be prevented from contacting birds.		



Integrated Agromet Advisory Service Bulletin 18th to 22nd Jan'14

Peren District

Bulletin No:05/2014

Weather summary of the preceding week

No rain occurred the past week.
Maximum and **minimum temperature** range from 21⁰C to 24⁰C and 5⁰C to 11⁰C, respectively.
Relative humidity varied from 29% to 97%.
Wind speed ranged from 2 to 4 kmph.

Weather forecast valid upto 22nd Jan'14

There is possibility of no rain the coming week.
Max temp is expected to be around 23⁰C - 24⁰C and **min temp is expected to be around** 10⁰C to 11⁰C.
Possibility of clear sky the coming week
Relative Humidity will range from 30% to 97%
The **wind speed** may reach upto 3 - 4 kmph
Wind direction will blow from easterly

Main Crops

Stage

Pest/ diseases

Agro-meteorological Advisories


Field crops

Rice	Storage	-	It should be ensured that the bags are clean, dry and well stitched, to keep the risk of loss as low as possible right from the start. Care should be taken as high proportion of broken grains encourages the development of microorganisms, mites and beetles Rice is particularly likely to be infested with storage pests if it has been stored for more than two months.
Maize	Post -harvesting stage	-	Provide protection from common storage loss agents such as insect pests, rodents, moulds, birds etc
Mustard	Podding stage	Aphid	Aphids suck sap as a result of it the plants remain stunted, flowers dry up and seeds do not develop. So proper monitoring should be carried out.

Horticultural crops

Probability no rain will occur the coming 3 or 4 days, therefore the farmers are advised to irrigate the crops daily or at an interval depending on the availability of water.

Citrus	Nursery stage	-	In citrus nursery, diseases may appear any time of plant growth through contaminated water, soil and even through nursery workers and implements. Therefore a regular monitoring should be done for infection and in case of infection; the infected/contaminated plants should immediately be uprooted and destroyed to keep the nursery totally free.
Vegetables	Harvesting stage	-	Harvest vegetables at their right stage and storing them properly will help extend their shelf life

	<i>Cabbage</i>	Heading stage		Harvesting should be done when the heads are large and solid
	<i>Ginger</i>	Harvesting stage		Harvesting should be done when the foliage is completely dried out.
	<i>Fishery</i>	Never over feed your fish. Excess food will pollute the pond water, so only feed amounts that will be completely consumed by your fish within 2 minutes.		
	<i>Piggery</i>	Good housekeeping practices such as removing manure, washing down pens, disinfecting and generally cleaning up the facility are essential to maintaining a healthy herd.		
	<i>Poultry</i>	During cold weather, cool outside air should be prevented from contacting birds.		




Integrated Agromet Advisory Service Bulletin 18th to 22nd Jan'14

Phek District

Bulletin No: 05/2014

Weather summary of the preceding week			Weather forecast valid upto 22 nd Jan'14
No rain occurred the past week Maximum and minimum temperatures ranged from 19 ⁰ C to 23 ⁰ C and 5 ⁰ C to 9 ⁰ C, respectively. Relative humidity varied from 31% to 96%. Wind speed ranged from 1 to 2 kmph.			There is possibility of no rain the coming week. Max temp is expected to be around 22 ⁰ C - 23 ⁰ C and min temp is expected to be around 8 ⁰ C to 9 ⁰ C. Possibility of clear sky the coming week Relative Humidity will range from 32% to 97% The wind speed 1- 2 kmph Wind direction will blow from southerly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Rice	Storage	-	It should be ensured that the bags are clean, dry and well stitched, to keep the risk of loss as low as possible right from the start. Care should be taken as high proportion of broken grains encourages the development of microorganisms, mites and beetles Rice is particularly likely to be infested with storage pests if it has been stored for more than two months.
Maize	Post - harvesting stage	-	Provide protection from common storage loss agents such as insect pests, rodents, moulds, birds etc
Mustard	Podding stage	Aphid	Aphids suck sap as a result of it the plants remain stunted, flowers dry up and seeds do not develop. So proper monitoring should be carried out.
Horticultural crops			
Probability no rain will occur the coming 3 or 4 days, therefore the farmers are advised to irrigate the crops daily or at an interval depending on the availability of water.			
Citrus	Nursery stage	-	In citrus nursery, diseases may appear any time of plant growth through contaminated water, soil and even through nursery workers and implements. Therefore a regular monitoring should be done for infection and in case of infection; the infected/contaminated plants should immediately be uprooted and destroyed to keep the nursery totally free.

	Vegetables	Harvesting stage	-	Harvest vegetables at their right stage and storing them properly will help extend their shelf life
	Cabbage	Heading stage		Harvesting should be done when the heads are large and solid
	Ginger	Harvesting stage		Harvesting should be done when the foliage is completely dried out.
	Fishery	Never over feed your fish. Excess food will pollute the pond water, so only feed amounts that will be completely consumed by your fish within 2 minutes.		
	Piggery	Good housekeeping practices such as removing manure, washing down pens, disinfecting and generally cleaning up the facility are essential to maintaining a healthy herd.		
	Poultry	During cold weather, cool outside air should be prevented from contacting birds.		



Integrated Agromet Advisory Service Bulletin 18th to 22nd Jan'14 Tuensang District

Bulletin No:05/2014


Weather summary of the preceding week

No rain occurred the past week
Maximum and minimum temperatures ranged 15⁰C to 24⁰C and 5⁰C to 7⁰C, respectively.
Relative humidity varied from 30% to 97%.
Wind speed ranged from 2 to 4 kmph.

Weather forecast valid upto 22nd Jan'14

There is possibility of no rain the coming week.
Max temp is expected to be around 23⁰C - 24⁰C and min temp is expected to be around 7⁰C to 8⁰C.
Possibility of clear sky the coming week
Relative humidity will range from 33% to 97%
Wind speed 2 – 3kmph
Wind direction will blow from southwesterly

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Field crops</i>			
<i>Rice</i>	Storage	-	It should be ensured that the bags are clean, dry and well stitched, to keep the risk of loss as low as possible right from the start. Care should be taken as high proportion of broken grains encourages the development of microorganisms, mites and beetles Rice is particularly likely to be infested with storage pests if it has been stored for more than two months.
<i>Maize</i>	Post - harvesting stage	-	Provide protection from common storage loss agents such as insect pests, rodents, moulds, birds etc
<i>Mustard</i>	Podding stage	Aphid	Aphids suck sap as a result of it the plants remain stunted, flowers dry up and seeds do not develop. So proper monitoring should be carried out.
<i>Horticultural crops</i>			
Probability no rain will occur the coming 3 or 4 days, therefore the farmers are advised to irrigate the crops daily or at an interval depending on the availability of water.			
<i>Citrus</i>	Nursery stage	-	In citrus nursery, diseases may appear any time of plant growth through contaminated water, soil and even through nursery workers and implements. Therefore a regular monitoring should be done for infection and in case of infection; the infected/contaminated plants should immediately be uprooted and destroyed to keep the nursery totally free.
<i>Vegetables</i>	Harvesting	-	Harvest vegetables at their right stage and storing them properly will help extend their shelf life

		stage		
	<i>Cabbage</i>	Heading stage		Harvesting should be done when the heads are large and solid
	<i>Ginger</i>	Harvesting stage		Harvesting should be done when the foliage is completely dried out.
	<i>Fishery</i>	Never over feed your fish. Excess food will pollute the pond water, so only feed amounts that will be completely consumed by your fish within 2 minutes.		
	<i>Piggery</i>	Good housekeeping practices such as removing manure, washing down pens, disinfecting and generally cleaning up the facility are essential to maintaining a healthy herd.		
	<i>Poultry</i>	During cold weather, cool outside air should be prevented from contacting birds.		



Integrated Agromet Advisory Service Bulletin 18th to 22nd Jan'14 Wokha District

Bulletin No:05/2014


Weather summary of the preceding week

No rain occurred the past week.
Maximum and minimum temperatures ranged from 16⁰C to 23⁰C and 5⁰C to 8⁰C, respectively.
Relative humidity varied from 33% to 97%.
Wind speed ranged from 2 to 3 kmph

Weather forecast valid upto 22nd Jan'14

There is possibility of no rain the coming week.
Max temp is expected to be around 22⁰C - 23⁰C and min temp is expected to be around 7⁰C to 8⁰C.
Possibility of clear sky the coming week
Relative Humidity will range from 31% to 94%
The wind speed 2- 3 kmph
Wind direction will blow from southerly

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Field crops</i>			
<i>Rice</i>	Storage	-	It should be ensured that the bags are clean, dry and well stitched, to keep the risk of loss as low as possible right from the start. Care should be taken as high proportion of broken grains encourages the development of microorganisms, mites and beetles Rice is particularly likely to be infested with storage pests if it has been stored for more than two months.
<i>Maize</i>	Post - harvesting stage	-	Provide protection from common storage loss agents such as insect pests, rodents, moulds, birds etc
<i>Mustard</i>	Podding stage	Aphid	Aphids suck sap as a result of it the plants remain stunted, flowers dry up and seeds do not develop. So proper monitoring should be carried out.
<i>Horticultural crops</i>			
Probability no rain will occur the coming 3 or 4 days, therefore the farmers are advised to irrigate the crops daily or at an interval depending on the availability of water.			
<i>Citrus</i>	Nursery stage	-	In citrus nursery, diseases may appear any time of plant growth through contaminated water, soil and even through nursery workers and implements. Therefore a regular monitoring should be done for infection and in case of infection; the infected/contaminated plants should immediately be uprooted and destroyed to keep the nursery totally free.

	Vegetables	Harvesting stage	-	Harvest vegetables at their right stage and storing them properly will help extend their shelf life
	Cabbage	Heading stage		Harvesting should be done when the heads are large and solid
	Ginger	Harvesting stage		Harvesting should be done when the foliage is completely dried out.
	Fishery	Never over feed your fish. Excess food will pollute the pond water, so only feed amounts that will be completely consumed by your fish within 2 minutes.		
	Piggery	Good housekeeping practices such as removing manure, washing down pens, disinfecting and generally cleaning up the facility are essential to maintaining a healthy herd.		
	Poultry	During cold weather, cool outside air should be prevented from contacting birds.		




--	--



Integrated Agromet Advisory Service Bulletin 18th to 22nd Jan'14 Zunheboto District

Bulletin No:05/2014

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 22nd Jan'14</i>
<p>No rain occurred the past week Maximum and minimum temperatures ranged from 19⁰C to 22⁰C and 5⁰C to 7⁰C, respectively. Relative humidity varied from 32% to 96%. Wind speed ranged from 2 to 3 kmph.</p>			<p>There is possibility of no rain the coming week. Max temp is expected to be around 21⁰C - 22⁰C and min temp is expected to be around 6⁰C to 7⁰C. Possibility of clear sky the coming week Relative Humidity will range from 31% to 95% The wind speed 2- 3 kmph Wind direction will blow southerly</p>
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Rice	Storage	-	<p>It should be ensured that the bags are clean, dry and well stitched, to keep the risk of loss as low as possible right from the start. Care should be taken as high proportion of broken grains encourages the development of microorganisms, mites and beetles Rice is particularly likely to be infested with storage pests if it has been stored for more than two months.</p>
Maize	Post - harvesting stage	-	<p>Provide protection from common storage loss agents such as insect pests, rodents, moulds, birds etc</p>
Mustard	Podding stage	Aphid	<p>Aphids suck sap as a result of it the plants remain stunted, flowers dry up and seeds do not develop. So proper monitoring should be carried out.</p>
Horticultural crops			
<p>Probability no rain will occur the coming 3 or 4 days, therefore the farmers are advised to irrigate the crops daily or at an interval depending on the availability of water.</p>			
Citrus	Nursery stage	-	<p>In citrus nursery, diseases may appear any time of plant growth through contaminated water, soil and even through nursery workers and implements. Therefore a regular monitoring should be done for infection and in case of infection; the infected/contaminated plants should immediately be uprooted and destroyed to keep the nursery totally free.</p>
Vegetables	Harvesting	-	<p>Harvest vegetables at their right stage and storing them properly will help</p>

		stage		extend their shelf life
	<i>Cabbage</i>	Heading stage		Harvesting should be done when the heads are large and solid
	<i>Ginger</i>	Harvesting stage		Harvesting should be done when the foliage is completely dried out.
	<i>Fishery</i>	Never over feed your fish. Excess food will pollute the pond water, so only feed amounts that will be completely consumed by your fish within 2 minutes.		
	<i>Piggery</i>	Good housekeeping practices such as removing manure, washing down pens, disinfecting and generally cleaning up the facility are essential to maintaining a healthy herd.		
	<i>Poultry</i>	During cold weather, cool outside air should be prevented from contacting birds.		




Integrated Agromet Advisory Service Bulletin 18th to 22nd Jan'14

Dimapur District

Bulletin No:05/2014

Weather summary of the preceding week			Weather forecast valid upto 22 nd Jan'14
No rain occurred the past week. Maximum and minimum temperatures ranged 22 ⁰ C to 24 ⁰ C and 5 ⁰ C to 6 ⁰ C, respectively. Relative humidity varied from 34% to 96%. Wind speed ranged from 2 to 4 kmph.			There is possibility of no rain the coming week. Max temp is expected to be around 23 ⁰ C - 24 ⁰ C and min temp is expected to be around 9 ⁰ C to 10 ⁰ C. Possibility of clear sky the coming week Relative Humidity will range from 30% to 95% and Wind speed 2– 3 kmph Wind direction will be from easterly
Main Crops	Stage	Pest/ diseases	Agro-meteorological Advisories
Field crops			
Rice	Storage	-	It should be ensured that the bags are clean, dry and well stitched, to keep the risk of loss as low as possible right from the start. Care should be taken as high proportion of broken grains encourages the development of microorganisms, mites and beetles Rice is particularly likely to be infested with storage pests if it has been stored for more than two months.
Maize	Post -harvesting stage	-	Provide protection from common storage loss agents such as insect pests, rodents, moulds, birds etc
Mustard	Podding stage	Aphid	Aphids suck sap as a result of it the plants remain stunted, flowers dry up and seeds do not develop. So proper monitoring should be carried out.
Horticultural crops			
Probability no rain will occur the coming 3 or 4 days, therefore the farmers are advised to irrigate the crops daily or at an interval depending on the availability of water.			
Citrus	Nursery stage	-	In citrus nursery, diseases may appear any time of plant growth through contaminated water, soil and even through nursery workers and implements. Therefore a regular monitoring should be done for infection and in case of infection; the infected/contaminated plants should immediately be uprooted and destroyed to keep the nursery totally free.
Vegetables	Harvesting stage	-	Harvest vegetables at their right stage and storing them properly will help extend their shelf life
Cabbage	Heading stage		Harvesting should be done when the heads are large and solid

	<i>Ginger</i>	Harvesting stage		Harvesting should be done when the foliage is completely dried out.
	<i>Fishery</i>	Never over feed your fish. Excess food will pollute the pond water, so only feed amounts that will be completely consumed by your fish within 2 minutes.		
	<i>Piggery</i>	Good housekeeping practices such as removing manure, washing down pens, disinfecting and generally cleaning up the facility are essential to maintaining a healthy herd.		
	<i>Poultry</i>	During cold weather, cool outside air should be prevented from contacting birds.		




Integrated Agromet Advisory Service Bulletin 18th to 22nd Jan'14

Kiphire District

Bulletin No:05/2014

Weather summary of the preceding week			Weather forecast valid upto 22 nd Jan'14
<p>No rain occurred the past week. Maximum and minimum temperatures ranged from 17⁰C to 23⁰C and 5⁰C to 10⁰C, respectively. Relative humidity varied from 30% to 98%. Wind speed ranged from 2 to 4 kmph.</p>			<p>There is possibility of no rain the coming week. Max temp is expected to be around 22⁰C - 23⁰C and min temp is expected to be around 9⁰C to 10⁰C. Possibility of clear sky the coming week Relative Humidity will range from 33% to 97% The wind speed 2- 3 kmph Wind direction will blow from southwesterly</p>
Main Crops	Stage	Pest/ diseases	Agro-meteorological Advisories
Field crops			
Rice	Storage	-	<p>It should be ensured that the bags are clean, dry and well stitched, to keep the risk of loss as low as possible right from the start. Care should be taken as high proportion of broken grains encourages the development of microorganisms, mites and beetles Rice is particularly likely to be infested with storage pests if it has been stored for more than two months.</p>
Maize	Post -harvesting stage	-	<p>Provide protection from common storage loss agents such as insect pests, rodents, moulds, birds etc</p>
Mustard	Podding stage	Aphid	<p>Aphids suck sap as a result of it the plants remain stunted, flowers dry up and seeds do not develop. So proper monitoring should be carried out.</p>
Horticultural crops			
<p>Probability no rain will occur the coming 3 or 4 days, therefore the farmers are advised to irrigate the crops daily or at an interval depending on the availability of water.</p>			
Citrus	Nursery stage	-	<p>In citrus nursery, diseases may appear any time of plant growth through contaminated water, soil and even through nursery workers and implements. Therefore a regular monitoring should be done for infection and in case of infection; the infected/contaminated plants should immediately be uprooted and destroyed to keep the nursery totally free.</p>
Vegetables	Harvesting stage	-	<p>Harvest vegetables at their right stage and storing them properly will help extend their shelf life</p>

	<i>Cabbage</i>	Heading stage		Harvesting should be done when the heads are large and solid
	<i>Ginger</i>	Harvesting stage		Harvesting should be done when the foliage is completely dried out.
	<i>Fishery</i>	Never over feed your fish. Excess food will pollute the pond water, so only feed amounts that will be completely consumed by your fish within 2 minutes.		
	<i>Piggery</i>	Good housekeeping practices such as removing manure, washing down pens, disinfecting and generally cleaning up the facility are essential to maintaining a healthy herd.		
	<i>Poultry</i>	During cold weather, cool outside air should be prevented from contacting birds.		