




Integrated Agromet Advisory Service Bulletin 1st to 5th March'14

Tuensang District

Bulletin No:16/2014

<i>Weather summary of the preceding week</i>		<i>Weather forecast valid upto 5th March'14</i>	
No rain occurred the past week Maximum and minimum temperatures ranged 16 ⁰ C to 17 ⁰ C and 6 ⁰ C to 7 ⁰ C, respectively. Relative humidity varied from 35% to 94%. Wind speed ranged from 2 to 4 kmph.		There is possibility of no rain the coming week. Max temp is expected to be around 16 ⁰ C-17 ⁰ C and min temp is expected to be around 6 ⁰ C to 7 ⁰ C. Possibility of mainly clear sky the coming week Relative humidity will range from 32% to 94% Wind speed 3 – 4kmph Wind direction will blow from westerly	
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
<i>Paddy field</i>	-	-	Crop rotation should be carried out as it may interrupt the relationship between pests and host plant. Rice based cropping systems involving pre-kharif green-manure crop followed by kharif rice and non- rice rabi crop directly benefit the rice crop.
<i>Mustard</i>	Maturing stage	-	If harvesting is going to be carried out in some places then it should be postponed to next week as there is a probability of light rain the coming week
Horticultural crops			
<i>Khasi mandarin (Orange)</i>	-	-	During this season, light earth work should be carried out in the plant basin and mulching should be carried out with locally available dry grasses/ leaves/ straw near the base of the trunk to conserve soil moisture.
<i>Vegetables</i>	Harvesting stage (Tomato,broccoli, bottlegourd,carrot)	-	Harvest during the coolest part of the day that is early morning or late afternoon; do not harvest produce when it is wet from dew as it will be more likely to decay. Protect harvested produce in the field by putting it under open-sided shade when transport is not immediately available.
<i>Fishery</i>	If you have enough water to fill the pond again, it is best to drain all of the water and clean the bottom of your pond but if you have only enough water to fill part of your pond, do not drain out all of the water, and drain only part of it.		
<i>Piggery</i>	Animals should be culled if it is suffering from incurable and chronic infections.		

 The logo of the Indian Council of Agricultural Research (ICAR) is located in the top-left corner. It features a green emblem with a central leaf-like shape and radiating lines, set within a circular frame. Below the emblem, the text 'भारतीय' is written in Devanagari script, and 'ICAR' is written in English capital letters.	<i>Poultry</i>	<p>For prevention of diseases, some general measures should be followed. Provide ample quantity of clean drinking water, don't make sudden changes in feed, maintain good ventilation avoid overcrowding and dispose dead birds by burning or burial.</p>
---	-----------------------	---



Integrated Agromet Advisory Service Bulletin 1st to 5th March'14 Wokha District

Bulletin No:16/2014

Weather summary of the preceding week

No rain occurred the past week.
Maximum and minimum temperatures ranged from 20⁰C to 21⁰C and 8⁰C to 9⁰C, respectively.
Relative humidity varied from 33% to 95%.
Wind speed ranged from 2 to 4 kmph

Weather forecast valid upto 5th March'14

There is possibility of no rain the coming week.
Max temp is expected to be around 20⁰C - 21⁰C and min temp is expected to be around 8⁰C to 9⁰C.
Possibility of partly cloudy sky the coming week
Relative Humidity will range from 30% to 92%
The wind speed 1– 2 kmph
Wind direction will blow from southeasterly

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Field crops</i>			
<i>Paddy field</i>	-	-	Crop rotation should be carried out as it may interrupt the relationship between pests and host plant. Rice based cropping systems involving pre-kharif green-manure crop followed by kharif rice and non- rice rabi crop directly benefit the rice crop.
<i>Mustard</i>	Maturing stage	-	If harvesting is going to be carried out in some places then it should be postponed to next week as there is a probability of light rain the coming week
<i>Horticultural crops</i>			
<i>Khasi mandarin (Orange)</i>	-	-	During this season, light earth work should be carried out in the plant basin and mulching should be carried out with locally available dry grasses/ leaves/ straw near the base of the trunk to conserve soil moisture.
<i>Vegetables</i>	Harvesting stage (Tomato, broccoli, bottlegourd, carrot)	-	Harvest during the coolest part of the day that is early morning or late afternoon; do not harvest produce when it is wet from dew as it will be more likely to decay. Protect harvested produce in the field by putting it under open-sided shade when transport is not immediately available.
<i>Fishery</i>	If you have enough water to fill the pond again, it is best to drain all of the water and clean the bottom of your pond but if you have only enough water to fill part of your pond, do not drain out all of the water, and drain only part of it.		



<i>Piggery</i>	Animals should be culled if it is suffering from incurable and chronic infections.
<i>Poultry</i>	For prevention of diseases, some general measures should be followed. Provide ample quantity of clean drinking water, don't make sudden changes in feed, maintain good ventilation avoid overcrowding and dispose dead birds by burning or burial.



--	--



Integrated Agromet Advisory Service Bulletin 1st to 5th March'14 Zunheboto District

Bulletin No:16/2014

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 5th March'14</i>
No rain occurred the past week Maximum and minimum temperatures ranged from 21 ⁰ C to 22 ⁰ C and 6 ⁰ C to 8 ⁰ C, respectively. Relative humidity varied from 31% to 90%. Wind speed ranged from 2 to 4 kmph.			There is possibility of no rain the coming week. Max temp is expected to be around 21 ⁰ C - 23 ⁰ C and min temp is expected to be around 7 ⁰ C to 8 ⁰ C. Possibility of partly cloudy the coming week Relative Humidity will range from 31% to 95% The wind speed 2- 3 kmph Wind direction will blow southerly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
<i>Paddy field</i>	-	-	Crop rotation should be carried out as it may interrupt the relationship between pests and host plant. Rice based cropping systems involving pre-kharif green-manure crop followed by kharif rice and non- rice rabi crop directly benefit the rice crop.
<i>Mustard</i>	Maturing stage	-	If harvesting is going to be carried out in some places then it should be postponed to next week as there is a probability of light rain the coming week
Horticultural crops			
<i>Khasi mandarin (Orange)</i>	-	-	During this season, light earth work should be carried out in the plant basin and mulching should be carried out with locally available dry grasses/ leaves/ straw near the base of the trunk to conserve soil moisture.
<i>Vegetables</i>	Harvesting stage (Tomato,broccoli, bottlegourd,carrot)	-	Harvest during the coolest part of the day that is early morning or late afternoon; do not harvest produce when it is wet from dew as it will be more likely to decay. Protect harvested produce in the field by putting it under open-sided shade when transport is not immediately available.
<i>Fishery</i>	If you have enough water to fill the pond again, it is best to drain all of the water and clean the bottom of your pond but if you have only enough water to fill part of your pond, do not drain out all of the water, and drain only part of it.		
<i>Piggery</i>	Animals should be culled if it is suffering from incurable and chronic infections.		



Poultry

For prevention of diseases, some general measures should be followed.
Provide ample quantity of clean drinking water, don't make sudden changes in feed, maintain good ventilation avoid overcrowding and dispose dead birds by burning or burial.



Integrated Agromet Advisory Service Bulletin 1st to 5th March'14

Dimapur District

Bulletin No:16/2014

Weather summary of the preceding week			Weather forecast valid upto 5 th March'14
<p>No rain occurred the past week. Maximum and minimum temperatures ranged 25⁰C to 26⁰C and 10⁰C to 12⁰C, respectively. Relative humidity varied from 30% to 97%. Wind speed ranged from 2 to 4 kmph.</p>			<p>There is possibility of no rain the coming week. Max temp is expected to be around 23⁰C - 26⁰C and min temp is expected to be around 11⁰C to 12⁰C. Possibility of partly cloudy sky the coming week Relative Humidity will range from 28% to 95% Wind speed 1– 2 kmph Wind direction will be from easterly</p>
Main Crops	Stage	Pest/ diseases	Agro-meteorological Advisories
Field crops			
Paddy field	-	-	<p>Crop rotation should be carried out as it may interrupt the relationship between pests and host plant. Rice based cropping systems involving pre-kharif green-manure crop followed by kharif rice and non- rice rabi crop directly benefit the rice crop.</p>
Mustard	Maturing stage	-	<p>If harvesting is going to be carried out in some places then it should be postponed to next week as there is a probability of light rain the coming week</p>
Horticultural crops			
Khasi mandarin (Orange)	-	-	<p>During this season, light earth work should be carried out in the plant basin and mulching should be carried out with locally available dry grasses/ leaves/ straw near the base of the trunk to conserve soil moisture.</p>
Vegetables	Harvesting stage (Tomato,broccoli, bottlegourd,carrot)	-	<p>Harvest during the coolest part of the day that is early morning or late afternoon; do not harvest produce when it is wet from dew as it will be more likely to decay. Protect harvested produce in the field by putting it under open-sided shade when transport is not immediately available.</p>
Fishery	<p>If you have enough water to fill the pond again, it is best to drain all of the water and clean the bottom of your pond but if you have only enough water to fill part of your pond, do not drain out all of the water, and drain only part of it.</p>		
Piggery	<p>Animals should be culled if it is suffering from incurable and chronic infections.</p>		



Poultry

For prevention of diseases, some general measures should be followed.

Provide ample quantity of clean drinking water, don't make sudden changes in feed, maintain good ventilation avoid overcrowding and dispose dead birds by burning or burial.



Integrated Agromet Advisory Service Bulletin 1st to 5th March'14

Kiphire District

Bulletin No:16/2014

Weather summary of the preceding week			Weather forecast valid upto 5 th March'14
<p>No rain occurred the past week. Maximum and minimum temperatures ranged from 22^oC to 23^oC and 7^oC to 8^oC, respectively. Relative humidity varied from 35% to 93%. Wind speed ranged from 2 to 4 kmph.</p>			<p>There is possibility of light rain the coming week. Max temp is expected to be around 22^oC - 23^oC and min temp is expected to be around 7^oC to 8^oC. Possibility of mainly clear sky the coming week Relative Humidity will range from 32% to 95% The wind speed 3- 4 kmph Wind direction will blow from westerly</p>
Main Crops	Stage	Pest/ diseases	Agro-meteorological Advisories
Field crops			
Paddy field	-	-	<p>Crop rotation should be carried out as it may interrupt the relationship between pests and host plant. Rice based cropping systems involving pre-kharif green-manure crop followed by kharif rice and non- rice rabi crop directly benefit the rice crop.</p>
Mustard	Maturing stage	-	<p>If harvesting is going to be carried out in some places then it should be postponed to next week as there is a probability of light rain the coming week</p>
Horticultural crops			
Khasi mandarin (Orange)	-	-	<p>During this season, light earth work should be carried out in the plant basin and mulching should be carried out with locally available dry grasses/ leaves/ straw near the base of the trunk to conserve soil moisture.</p>
Vegetables	Harvesting stage (Tomato,broccoli, bottlegourd,carrot)	-	<p>Harvest during the coolest part of the day that is early morning or late afternoon; do not harvest produce when it is wet from dew as it will be more likely to decay. Protect harvested produce in the field by putting it under open-sided shade when transport is not immediately available.</p>
Cabbage	Harvesting stage	-	<p>Harvesting should be done when the heads are well developed and firm.</p>
Fishery	<p>If you have enough water to fill the pond again, it is best to drain all of the water and clean the bottom of your pond but if you have only enough water to fill part of your pond, do not drain out all of the water, and drain only part of it.</p>		
Piggery	<p>Animals should be culled if it is suffering from incurable and chronic infections.</p>		



Poultry

For prevention of diseases, some general measures should be followed.

Provide ample quantity of clean drinking water, don't make sudden changes in feed, maintain good ventilation avoid overcrowding and dispose dead birds by burning or burial.



Integrated Agromet Advisory Service Bulletin 1st to 5th March'14

Kohima District

Bulletin No:16/2014


Weather summary of the preceding week

No rain occurred the past week.
Maximum and minimum temperatures ranged from 20⁰C to 21⁰C and 6⁰C to 9⁰C, respectively.
Relative humidity varied from 23% to 97%.
Wind speed ranged from 2 to 3 kmph.

Weather forecast valid upto 5th March'14

There is possibility of no rain the coming week.
Max temp is expected to be around 20⁰C - 21⁰C and min temp is expected to be around 8⁰C to 9⁰C.
Possibility of partly cloudy sky the coming week
Relative Humidity will range from 23% to 95%
Wind speed 2-4 kmph
Wind direction will blow from southerly

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Paddy field	-	-	Crop rotation should be carried out as it may interrupt the relationship between pests and host plant. Rice based cropping systems involving pre-kharif green-manure crop followed by kharif rice and non- rice rabi crop directly benefit the rice crop.
Mustard	Maturing stage	-	If harvesting is going to be carried out in some places then it should be postponed to next week as there is a probability of light rain the coming week
Horticultural crops			
Khasi mandarin (Orange)	-	-	During this season, light earth work should be carried out in the plant basin and mulching should be carried out with locally available dry grasses/ leaves/ straw near the base of the trunk to conserve soil moisture.
Vegetables	Harvesting stage (Tomato,broccoli, bottlegourd,carrot)	-	Harvest during the coolest part of the day that is early morning or late afternoon; do not harvest produce when it is wet from dew as it will be more likely to decay. Protect harvested produce in the field by putting it under open-sided shade when transport is not immediately available.
Fishery	If you have enough water to fill the pond again, it is best to drain all of the water and clean the bottom of your pond but if you have only enough water to fill part of your pond, do not drain out all of the water, and drain only part of it.		

	<i>Piggery</i>	Animals should be culled if it is suffering from incurable and chronic infections.
	<i>Poultry</i>	<p>For prevention of diseases, some general measures should be followed.</p> <p>Provide ample quantity of clean drinking water, don't make sudden changes in feed, maintain good ventilation avoid overcrowding and dispose dead birds by burning or burial.</p>



Integrated Agromet Advisory Service Bulletin 1st to 5th March'14

Longleng District

Bulletin No:16/2014

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 5th March'14</i>
<p>No rain occurred the past week. Maximum and minimum temperatures ranged from 22⁰C to 23⁰C and 7⁰C to 8⁰C, respectively. Relative humidity varied from 32% to 79%. Wind speed ranged from 3 to 4 kmph.</p>			<p>There is possibility of no rain the coming week. Max temp is expected to be around 22⁰C - 23⁰C and min temp is expected to be around 7⁰C to 8⁰C. Possibility of partly cloudy sky the coming week Relative Humidity will range from 28% to 95% The wind speed 2 –4 kmph Wind direction will blow from southerly</p>
Main Crops	Stage	Pest/ diseases	Agro-meteorological Advisories
Field crops			
Paddy field	-	-	<p>Crop rotation should be carried out as it may interrupt the relationship between pests and host plant. Rice based cropping systems involving pre-kharif green-manure crop followed by kharif rice and non- rice rabi crop directly benefit the rice crop.</p>
Mustard	Maturing stage	-	<p>If harvesting is going to be carried out in some places then it should be postponed to next week as there is a probability of light rain the coming week</p>
Horticultural crops			
Khasi mandarin (Orange)	-	-	<p>During this season, light earth work should be carried out in the plant basin and mulching should be carried out with locally available dry grasses/ leaves/ straw near the base of the trunk to conserve soil moisture.</p>
Vegetables	Harvesting stage (Tomato,broccoli, bottlegourd,carrot)	-	<p>Harvest during the coolest part of the day that is early morning or late afternoon; do not harvest produce when it is wet from dew as it will be more likely to decay. Protect harvested produce in the field by putting it under open-sided shade when transport is not immediately available.</p>
Cabbage	Harvesting stage	-	<p>Harvesting should be done when the heads are well developed and firm.</p>
Fishery	<p>If you have enough water to fill the pond again, it is best to drain all of the water and clean the bottom of your pond but if you have only enough water to fill part of your pond, do not drain out all of the water, and drain only part of it.</p>		



	<i>Piggery</i>	Animals should be culled if it is suffering from incurable and chronic infections.
	<i>Poultry</i>	For prevention of diseases, some general measures should be followed. Provide ample quantity of clean drinking water, don't make sudden changes in feed, maintain good ventilation avoid overcrowding and dispose dead birds by burning or burial.



Integrated Agromet Advisory Service Bulletin 1st to 5th March'14

Mokokchung District

Bulletin No: 16/2014

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 5th March'14</i>
<p>No rain occurred the past week Maximum and minimum temperatures ranged 17⁰C to 18⁰C and 6⁰C to 8⁰C, respectively. Relative humidity varied from 33% to 87%. Wind speed ranged from 3 to 4 kmph.</p>			<p>There is possibility of no rain the coming week. Max temp is expected to be around 18⁰C - 19⁰C and min temp is expected to be around 7⁰C to 8⁰C. Possibility of clear sky the coming week Relative Humidity will range from 29% to 95% and Wind speed 2– 3 kmph Wind direction will blow from southerly.</p>
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
<i>Paddy field</i>	-	-	<p>Crop rotation should be carried out as it may interrupt the relationship between pests and host plant. Rice based cropping systems involving pre-kharif green-manure crop followed by kharif rice and non- rice rabi crop directly benefit the rice crop.</p>
<i>Mustard</i>	Maturing stage	-	<p>If harvesting is going to be carried out in some places then it should be postponed to next week as there is a probability of light rain the coming week</p>
Horticultural crops			
<i>Khasi mandarin (Orange)</i>	-	-	<p>During this season, light earth work should be carried out in the plant basin and mulching should be carried out with locally available dry grasses/ leaves/ straw near the base of the trunk to conserve soil moisture.</p>
<i>Vegetables</i>	Harvesting stage (Tomato,broccoli, bottlegourd,carrot)	-	<p>Harvest during the coolest part of the day that is early morning or late afternoon; do not harvest produce when it is wet from dew as it will be more likely to decay. Protect harvested produce in the field by putting it under open-sided shade when transport is not immediately available.</p>
<i>Fishery</i>	<p>If you have enough water to fill the pond again, it is best to drain all of the water and clean the bottom of your pond but if you have only enough water to fill part of your pond, do not drain out all of the water, and drain only part of it.</p>		
<i>Piggery</i>	<p>Animals should be culled if it is suffering from incurable and chronic infections.</p>		



<i>Poultry</i>	For prevention of diseases, some general measures should be followed. Provide ample quantity of clean drinking water, don't make sudden changes in feed, maintain good ventilation avoid overcrowding and dispose dead birds by burning or burial.
-----------------------	---



Integrated Agromet Advisory Service Bulletin from 1st to 5th March'14 Mon District

Bulletin No:16/2014


Weather summary of the preceding week

- No rain occurred the past week
- **Maximum** and **minimum temperatures** ranged 21⁰C to 22⁰C and 7⁰C to 8⁰C, respectively.
- **Relative humidity** varied from 32% to 82%.
- **Wind speed** ranged from 3 to 4 kmph.

Weather forecast valid upto 5th March'14

- Possibility of no rain the coming week.
- The **max temp** is likely to be 19⁰C - 22⁰C and the **min temp** 7⁰C - 8⁰C.
- **Sky is likely to be partly cloudy the coming week**
- **Relative Humidity** is likely to range from 28% to 95%
- **Wind speed** may reach upto 2- 4kmph
- **Wind direction** will be from southeasterly

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Paddy field	-	-	Crop rotation should be carried out as it may interrupt the relationship between pests and host plant. Rice based cropping systems involving pre-kharif green-manure crop followed by kharif rice and non- rice rabi crop directly benefit the rice crop.
Mustard	Maturing stage	-	If harvesting is going to be carried out in some places then it should be postponed to next week as there is a probability of light rain the coming week
Horticultural crops			
Khasi mandarin (Orange)	-	-	During this season, light earth work should be carried out in the plant basin and mulching should be carried out with locally available dry grasses/ leaves/ straw near the base of the trunk to conserve soil moisture.
Beans	Vegetative stage	-	Earthing up should be carried out.
Vegetables	Harvesting stage (Tomato,broccoli, bottlegourd,carrot, knolkhol)	-	Harvest during the coolest part of the day that is early morning or late afternoon; do not harvest produce when it is wet from dew as it will be more likely to decay. Protect harvested produce in the field by putting it under open-sided shade when transport is not immediately available.
Fishery	If you have enough water to fill the pond again, it is best to drain all of the water and clean the bottom of your pond but if you have only enough water to fill part of your pond, do not drain out all of the water, and drain only part of it.		

	<i>Piggery</i>	Animals should be culled if it is suffering from incurable and chronic infections.
	<i>Poultry</i>	<p>For prevention of diseases, some general measures should be followed.</p> <p>Provide ample quantity of clean drinking water, don't make sudden changes in feed, maintain good ventilation avoid overcrowding and dispose dead birds by burning or burial.</p>



Integrated Agromet Advisory Service Bulletin 1st to 5th March'14

Peren District

Bulletin No:16/2014

Weather summary of the preceding week

No rain occurred the past week.
Maximum and minimum temperature range from 23⁰C to 24⁰C and 5⁰C to 7⁰C, respectively.
Relative humidity varied from 25% to 97%.
Wind speed ranged from 2 to 3 kmph.

Weather forecast valid upto 5th March'14

There is possibility of no rain the coming week.
Max temp is expected to be around 24⁰C - 25⁰C and min temp is expected to be around 6⁰C to 7⁰C.
Possibility of partly cloudy sky the coming week
Relative Humidity will range from 25% to 94%
The wind speed may reach upto 2 - 4 kmph
Wind direction will blow from southerly

Main Crops

Stage

Pest/ diseases

Agro-meteorological Advisories

Field crops

Paddy field

-

-

Crop rotation should be carried out as it may interrupt the relationship between pests and host plant.
Rice based cropping systems involving **pre-kharif green-manure crop** followed by **kharif rice and non- rice rabi crop** directly benefit the rice crop.

Mustard

Maturing stage

-

If harvesting is going to be carried out in some places then it should be postponed to next week as there is a probability of light rain the coming week

Horticultural crops

Khasi mandarin (Orange)

-

-

During this season, light earth work should be carried out in the plant basin and mulching should be carried out with locally available dry grasses/ leaves/ straw near the base of the trunk to conserve soil moisture.

Vegetables

Harvesting stage
(Tomato,broccoli
,
bottlegourd,carrot
)

-

Harvest during the coolest part of the day that is early morning or late afternoon; do not harvest produce when it is wet from dew as it will be more likely to decay. Protect harvested produce in the field by putting it under open-sided shade when transport is not immediately available.

Fishery

If you have enough water to fill the pond again, it is best to drain all of the water and clean the bottom of your pond but if you have only enough water to fill part of your pond, do not drain out all of the water, and drain only part of it.

Piggery

Animals should be culled if it is suffering from incurable and chronic infections.



<i>Poultry</i>	For prevention of diseases, some general measures should be followed. Provide ample quantity of clean drinking water, don't make sudden changes in feed, maintain good ventilation avoid overcrowding and dispose dead birds by burning or burial.
-----------------------	---



Integrated Agromet Advisory Service Bulletin 1st to 5th March'14

Phek District

Bulletin No: 16/2014

Weather summary of the preceding week

No rain occurred the past week
Maximum and **minimum temperatures** ranged from 16⁰C to 17⁰C and 6⁰C to 7⁰C, respectively.
Relative humidity varied from 30% to 90%.
Wind speed ranged from 2 to 4 kmph.

Weather forecast valid upto 5th March'14

There is possibility of no rain the coming week.
Max temp is expected to be around 16⁰C - 17⁰C and **min temp is expected to be around 6⁰C to 7⁰C**.
Possibility of partly cloudy sky the coming week
Relative Humidity will range from 30% to 97%
The **wind speed** 1- 2 kmph
Wind direction will blow from southerly

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Field crops</i>			
<i>Paddy field</i>	-	-	Crop rotation should be carried out as it may interrupt the relationship between pests and host plant. Rice based cropping systems involving pre-kharif green-manure crop followed by kharif rice and non- rice rabi crop directly benefit the rice crop.
<i>Mustard</i>	Maturing stage	-	If harvesting is going to be carried out in some places then it should be postponed to next week as there is a probability of light rain the coming week
<i>Horticultural crops</i>			
<i>Khasi mandarin (Orange)</i>	-	-	During this season, light earth work should be carried out in the plant basin and mulching should be carried out with locally available dry grasses/ leaves/ straw near the base of the trunk to conserve soil moisture.
<i>Vegetables</i>	Harvesting stage (Tomato, broccoli, bottlegourd, carrot)	-	Harvest during the coolest part of the day that is early morning or late afternoon; do not harvest produce when it is wet from dew as it will be more likely to decay. Protect harvested produce in the field by putting it under open-sided shade when transport is not immediately available.
<i>Fishery</i>	If you have enough water to fill the pond again, it is best to drain all of the water and clean the bottom of your pond but if you have only enough water to fill part of your pond, do not drain out all of the water, and drain only part of it.		
<i>Piggery</i>	Animals should be culled if it is suffering from incurable and chronic infections.		



Poultry

For prevention of diseases, some general measures should be followed.
Provide ample quantity of clean drinking water, don't make sudden changes in feed, maintain good ventilation avoid overcrowding and dispose dead birds by burning or burial.