



Integrated Agromet Advisory Service Bulletin 5th to 9th March'14

Kiphire District

Bulletin No:17/2014

Weather summary of the preceding week

No rain occurred the past week.
Maximum and **minimum temperatures** ranged from 22^oC to 23^oC and 7^oC to 8^oC, respectively.
Relative humidity varied from 32% to 92%.
Wind speed ranged from 3 to 4 kmph.

Weather forecast valid upto 9th March'14

There is possibility of no rain the coming week.
Max temp is expected to be around 22^oC - 23^oC and **min temp** is expected to be around 8^oC to 9^oC.
Possibility of mainly clear sky the coming week
Relative Humidity will range from 25% to 94%
The **wind speed** 3- 4 kmph
Wind direction will blow from westerly

Main Crops

Stage

Pest/ diseases

Agro-meteorological Advisories

Field crops

Paddy field

-

-

Crop rotation should be carried out as it may interrupt the relationship between pests and host plant.
Rice based cropping systems involving **pre-kharif green-manure crop** followed by **kharif rice and non- rice rabi crop** directly benefit the rice crop.

Horticultural crops

Citrus dieback (Mandarin orange)

-

-

Leaves become yellow and fall with the beginning of summer and die-back of twig starts leading to drying of branches and declining – death of trees. Transmission of disease take place through infected planting materials, mechanically through contaminated pruning knife and sometimes from soil borne pathogen. So, famers should take precautionary measures.

Summer vegetables

-

-

Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. as growing of early summer vegetables crops will fetch more income.

Potato

Sowing stage

-

At the time of planting, the tubers should closely be inspected and all suspicious looking tubers should be rejected.

Fishery

Stocking should not be delayed once a new pond has filled. As soon as the pond has adequate water in it, contamination by unwanted fish is possible before a good fish population develops. Prevent livestock from entering the pond and degrading the banks of the pond.

Piggery

Deworming of the pigs should be done routinely and clean, fresh water — and lots of it — must always be available

to your pigs.



Poultry


During this season, plenty of cool, clean fresh drinking water should be provided as the water consumption increases during summer and feed should be provided during cooler parts of the day such as early morning or late evening.



Integrated Agromet Advisory Service Bulletin 5th to 9th March'14 Kohima District

Bulletin No:17/2014

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 9th March'14</i>
No rain occurred the past week. Maximum and minimum temperatures ranged from 20 ⁰ C to 21 ⁰ C and 8 ⁰ C to 9 ⁰ C, respectively. Relative humidity varied from 23% to 93%. Wind speed ranged from 1 to 2 kmph.			There is possibility of no rain the coming week. Max temp is expected to be around 20 ⁰ C - 21 ⁰ C and min temp is expected to be around 8 ⁰ C to 9 ⁰ C. Possibility of mainly clear sky the coming week Relative Humidity will range from 16% to 92% Wind speed 1-2 kmph Wind direction will blow from easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Field crops</i>			
<i>Paddy field</i>	-	-	Crop rotation should be carried out as it may interrupt the relationship between pests and host plant. Rice based cropping systems involving pre-kharif green-manure crop followed by kharif rice and non- rice rabi crop directly benefit the rice crop.
<i>Horticultural crops</i>			
<i>Citrus dieback (Mandarin orange)</i>	-	-	Leaves become yellow and fall with the beginning of summer and die-back of twig starts leading to drying of branches and declining – death of trees. Transmission of disease take place through infected planting materials, mechanically through contaminated pruning knife and sometimes from soil borne pathogen. So, famers should take precautionary measures.
<i>Summer vegetables</i>	-	-	Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. as growing of early summer vegetables crops will fetch more income.
<i>Potato</i>	Sowing stage	-	At the time of planting, the tubers should closely be inspected and all suspicious looking tubers should be rejected.
<i>Fishery</i>	Stocking should not be delayed once a new pond has filled. As soon as the pond has adequate water in it, contamination by unwanted fish is possible before a good fish population develops. Prevent livestock from entering the pond and degrading the banks of the pond.		

	<i>Piggery</i>	Deworming of the pigs should be done routinely and clean, fresh water — and lots of it — must always be available to your pigs.
	<i>Poultry</i>	During this season, plenty of cool, clean fresh drinking water should be provided as the water consumption increases during summer and feed should be provided during cooler parts of the day such as early morning or late evening.



Integrated Agromet Advisory Service Bulletin 5th to 9th March'14

Longleng District

Bulletin No:17/2014

Weather summary of the preceding week

No rain occurred the past week.
Maximum and **minimum temperatures** ranged from 22^oC to 23^oC and 7^oC to 8^oC, respectively.
Relative humidity varied from 31% to 95%.
Wind speed ranged from 2 to 4 kmph.

Weather forecast valid upto 9th March'14

There is possibility of no rain the coming week.
Max temp is expected to be around 22^oC - 23^oC and **min temp is expected to be around 8^oC to 9^oC**.
Possibility of mainly clear sky the coming week
Relative Humidity will range from 22% to 80%
The wind speed 2 –4 kmph
Wind direction will blow from southerly

Main Crops

Stage

Pest/ diseases

Agro-meteorological Advisories

Field crops

Paddy field

-

-

Crop rotation should be carried out as it may interrupt the relationship between pests and host plant.
Rice based cropping systems involving **pre-kharif green-manure crop** followed by **kharif rice and non- rice rabi crop** directly benefit the rice crop.

Horticultural crops

Citrus dieback (Mandarin orange)

-

-

Leaves become yellow and fall with the beginning of summer and die-back of twig starts leading to drying of branches and declining – death of trees. Transmission of disease take place through infected planting materials, mechanically through contaminated pruning knife and sometimes from soil borne pathogen. So, famers should take precautionary measures.

Summer vegetables

-

-

Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. as growing of early summer vegetables crops will fetch more income.

Potato

Sowing stage

-

At the time of planting, the tubers should closely be inspected and all suspicious looking tubers should be rejected.

Fishery

Stocking should not be delayed once a new pond has filled. As soon as the pond has adequate water in it, contamination by unwanted fish is possible before a good fish population develops. Prevent livestock from entering the pond and degrading the banks of the pond.



	<i>Piggery</i>	Deworming of the pigs should be done routinely and clean, fresh water — and lots of it — must always be available to your pigs.
	<i>Poultry</i>	During this season, plenty of cool, clean fresh drinking water should be provided as the water consumption increases during summer and feed should be provided during cooler parts of the day such as early morning or late evening.



Integrated Agromet Advisory Service Bulletin 5th to 9th March'14

Mokokchung District

Bulletin No: 17/2014

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 9th March'14</i>
<p>No rain occurred the past week Maximum and minimum temperatures ranged 17⁰C to 18⁰C and 7⁰C to 8⁰C, respectively. Relative humidity varied from 32% to 95%. Wind speed ranged from 2 to 4 kmph.</p>			<p>There is possibility of no rain the coming week. Max temp is expected to be around 19⁰C - 20⁰C and min temp is expected to be around 11⁰C to 12⁰C. Possibility of clear sky the coming week Relative Humidity will range from 23% to 85% and Wind speed 3– 4 kmph Wind direction will blow from southerly.</p>
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Field crops</i>			
<i>Paddy field</i>	-	-	<p>Crop rotation should be carried out as it may interrupt the relationship between pests and host plant. Rice based cropping systems involving pre-kharif green-manure crop followed by kharif rice and non- rice rabi crop directly benefit the rice crop.</p>
<i>Horticultural crops</i>			
<i>Citrus dieback (Mandarin orange)</i>	-	-	<p>Leaves become yellow and fall with the beginning of summer and die-back of twig starts leading to drying of branches and declining – death of trees. Transmission of disease take place through infected planting materials, mechanically through contaminated pruning knife and sometimes from soil borne pathogen. So, famers should take precautionary measures.</p>
<i>Summer vegetables</i>	-	-	<p>Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. as growing of early summer vegetables crops will fetch more income.</p>
<i>Potato</i>	Sowing stage	-	<p>At the time of planting, the tubers should closely be inspected and all suspicious looking tubers should be rejected.</p>
<i>Fishery</i>	<p>Stocking should not be delayed once a new pond has filled. As soon as the pond has adequate water in it, contamination by unwanted fish is possible before a good fish population develops. Prevent livestock from entering the pond and degrading the banks of the pond.</p>		
<i>Piggery</i>	<p>Deworming of the pigs should be done routinely and clean, fresh water — and lots of it — must always be available to your pigs.</p>		



Poultry

During this season, plenty of cool, clean fresh drinking water should be provided as the water consumption increases during summer and feed should be provided during cooler parts of the day such as early morning or late evening.



Integrated Agromet Advisory Service Bulletin 5th to 9th March'14

Peren District

Bulletin No:17/2014

Weather summary of the preceding week

No rain occurred the past week.
Maximum and **minimum temperature** range from 23⁰C to 24⁰C and 6⁰C to 7⁰C, respectively.
Relative humidity varied from 23% to 94%.
Wind speed ranged from 2 to 3 kmph.

Weather forecast valid upto 9th March'14

There is possibility of no rain the coming week.
Max temp is expected to be around 24⁰C - 25⁰C and **min temp** is expected to be around 7⁰C to 9⁰C.
Possibility of mainly clear sky the coming week
Relative Humidity will range from 17% to 90%
The wind speed may reach upto 1 - 2 kmph
Wind direction will blow from easterly

Main Crops

Stage

**Pest/
diseases**

Agro-meteorological Advisories

Field crops

Paddy field

-

-

Crop rotation should be carried out as it may interrupt the relationship between pests and host plant.
Rice based cropping systems involving **pre-kharif green-manure crop** followed by **kharif rice and non- rice rabi crop** directly benefit the rice crop.

Horticultural crops

***Citrus dieback
(Mandarin
orange)***

-

-

Leaves become yellow and fall with the beginning of summer and die-back of twig starts leading to drying of branches and declining – death of trees. Transmission of disease take place through infected planting materials, mechanically through contaminated pruning knife and sometimes from soil borne pathogen. So, famers should take precautionary measures.

***Summer
vegetables***

-

-

Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. as growing of early summer vegetables crops will fetch more income.

Potato

Sowing stage

-

At the time of planting, the tubers should closely be inspected and all suspicious looking tubers should be rejected.

Fishery

Stocking should not be delayed once a new pond has filled. As soon as the pond has adequate water in it, contamination by unwanted fish is possible before a good fish population develops. Prevent livestock from entering the pond and degrading the banks of the pond.



ICAR RESEARCH COMPLEX FOR NEH REGION

(Indian Council of Agricultural Research)

NAGALAND CENTER, JHARNAPANI, MEDZIPHEMA

<i>Piggery</i>	Deworming of the pigs should be done routinely and clean, fresh water — and lots of it — must always be available to your pigs.
<i>Poultry</i>	During this season, plenty of cool, clean fresh drinking water should be provided as the water consumption increases during summer and feed should be provided during cooler parts of the day such as early morning or late evening.



Integrated Agromet Advisory Service Bulletin 5th to 9th March'14

Phek District

Bulletin No: 17/2014


Weather summary of the preceding week

No rain occurred the past week
Maximum and **minimum temperatures** ranged from 16⁰C to 17⁰C and 5⁰C to 6⁰C, respectively.
Relative humidity varied from 30% to 96%.
Wind speed ranged from 2 to 4 kmph.

Weather forecast valid upto 9th March'14

There is possibility of no rain the coming week.
Max temp is expected to be around 17⁰C - 18⁰C and **min temp is expected to be around 6⁰C to 7⁰C**.
Possibility of mainly clear sky the coming week
Relative Humidity will range from 23% to 92%
The wind speed 2- 4 kmph
Wind direction will blow from southerly

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Field crops</i>			
<i>Paddy field</i>	-	-	Crop rotation should be carried out as it may interrupt the relationship between pests and host plant. Rice based cropping systems involving pre-kharif green-manure crop followed by kharif rice and non- rice rabi crop directly benefit the rice crop.
<i>Horticultural crops</i>			
<i>Citrus dieback (Mandarin orange)</i>	-	-	Leaves become yellow and fall with the beginning of summer and die-back of twig starts leading to drying of branches and declining – death of trees. Transmission of disease take place through infected planting materials, mechanically through contaminated pruning knife and sometimes from soil borne pathogen. So, famers should take precautionary measures.
<i>Summer vegetables</i>	-	-	Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. as growing of early summer vegetables crops will fetch more income.
<i>Potato</i>	Sowing stage	-	At the time of planting, the tubers should closely be inspected and all suspicious looking tubers should be rejected.
<i>Fishery</i>	Stocking should not be delayed once a new pond has filled. As soon as the pond has adequate water in it, contamination by unwanted fish is possible before a good fish population develops. Prevent livestock from entering the pond and degrading the banks of the pond.		

	<i>Piggery</i>	Deworming of the pigs should be done routinely and clean, fresh water — and lots of it — must always be available to your pigs.
	<i>Poultry</i>	During this season, plenty of cool, clean fresh drinking water should be provided as the water consumption increases during summer and feed should be provided during cooler parts of the day such as early morning or late evening.





Integrated Agromet Advisory Service Bulletin 5th to 9th March'14 Tuensang District

Bulletin No:17/2014


Weather summary of the preceding week

No rain occurred the past week
Maximum and minimum temperatures ranged 16^oC to 17^oC and 6^oC to 7^oC, respectively.
Relative humidity varied from 33% to 97%.
Wind speed ranged from 3 to 4 kmph.

Weather forecast valid upto 9th March'14

There is possibility of no rain the coming week.
Max temp is expected to be around 17^oC-19^oC and **min temp is expected to be around 8^oC to 9^oC**.
Possibility of mainly clear sky the coming week
Relative humidity will range from 25% to 94%
Wind speed 3 – 4kmph
 Wind direction will blow from westerly

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Field crops</i>			
<i>Paddy field</i>	-	-	Crop rotation should be carried out as it may interrupt the relationship between pests and host plant. Rice based cropping systems involving pre-kharif green-manure crop followed by kharif rice and non- rice rabi crop directly benefit the rice crop.
<i>Horticultural crops</i>			
<i>Citrus dieback (Mandarin orange)</i>	-	-	Leaves become yellow and fall with the beginning of summer and die-back of twig starts leading to drying of branches and declining – death of trees. Transmission of disease take place through infected planting materials, mechanically through contaminated pruning knife and sometimes from soil borne pathogen. So, famers should take precautionary measures.
<i>Summer vegetables</i>	-	-	Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. as growing of early summer vegetables crops will fetch more income.
<i>Potato</i>	Sowing stage	-	At the time of planting, the tubers should closely be inspected and all suspicious looking tubers should be rejected.
<i>Fishery</i>	Stocking should not be delayed once a new pond has filled. As soon as the pond has adequate water in it, contamination by unwanted fish is possible before a good fish population develops. Prevent livestock from entering the pond and degrading the banks of the pond.		

	<i>Piggery</i>	Deworming of the pigs should be done routinely and clean, fresh water — and lots of it — must always be available to your pigs.
 The logo of the Indian Council of Agricultural Research (ICAR) is located in the bottom-left corner of the second row. It features a green emblem with a central leaf and wheat stalks, surrounded by a circular border. Below the emblem, the text 'भारतीय कृषि अनुसंधान परिषद' is written in Devanagari script, and 'ICAR' is written in English below that.	<i>Poultry</i>	During this season, plenty of cool, clean fresh drinking water should be provided as the water consumption increases during summer and feed should be provided during cooler parts of the day such as early morning or late evening.



Integrated Agromet Advisory Service Bulletin 5th to 9th March'14

Wokha District

Bulletin No:17/2014

Weather summary of the preceding week

No rain occurred the past week.
Maximum and minimum temperatures ranged from 20⁰C to 21⁰C and 8⁰C to 9⁰C, respectively.
Relative humidity varied from 32% to 95%.
Wind speed ranged from 2 to 3 kmph

Weather forecast valid upto 9th March'14


There is possibility of no rain the coming week.
Max temp is expected to be around 20⁰C - 24⁰C and **min temp** is expected to be around 9⁰C to 10⁰C.
Possibility of mainly clear sky the coming week
Relative Humidity will range from 23% to 90%
The wind speed 2– 3 kmph
Wind direction will blow from southerly

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Field crops</i>			
<i>Paddy field</i>	-	-	Crop rotation should be carried out as it may interrupt the relationship between pests and host plant. Rice based cropping systems involving pre-kharif green-manure crop followed by kharif rice and non- rice rabi crop directly benefit the rice crop.
<i>Horticultural crops</i>			
<i>Citrus dieback (Mandarin orange)</i>	-	-	Leaves become yellow and fall with the beginning of summer and die-back of twig starts leading to drying of branches and declining – death of trees. Transmission of disease take place through infected planting materials, mechanically through contaminated pruning knife and sometimes from soil borne pathogen. So, famers should take precautionary measures.
<i>Summer vegetables</i>	-	-	Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. as growing of early summer vegetables crops will fetch more income.
<i>Potato</i>	Sowing stage	-	At the time of planting, the tubers should closely be inspected and all suspicious looking tubers should be rejected.
<i>Fishery</i>	Stocking should not be delayed once a new pond has filled. As soon as the pond has adequate water in it, contamination by unwanted fish is possible before a good fish population develops. Prevent livestock from entering the pond and degrading the banks of the pond.		

ICAR RESEARCH COMPLEX FOR NEH REGION
 (Indian Council of Agricultural Research)
 NAGALAND CENTER, JHARNAPANI, MEDZIPHEMA



<i>Piggery</i>	Deworming of the pigs should be done routinely and clean, fresh water — and lots of it — must always be available to your pigs.
<i>Poultry</i>	During this season, plenty of cool, clean fresh drinking water should be provided as the water consumption increases during summer and feed should be provided during cooler parts of the day such as early morning or late evening.





भारत कृषि अनुसंधान परिषद
ICAR - RCNEH


ICAR RESEARCH COMPLEX FOR NEH REGION
(Indian Council of Agricultural Research)
NAGALAND CENTER, JHARNAPANI, MEDZIPHEMA

Integrated Agromet Advisory Service Bulletin 5th to 9th March'14 Zunheboto District

Bulletin No:17/2014

<i>Weather summary of the preceding week</i>	<i>Weather forecast valid upto 9th March'14</i>
<p>No rain occurred the past week Maximum and minimum temperatures ranged from 21⁰C to 22⁰C and 7⁰C to 8⁰C, respectively. Relative humidity varied from 30% to 96%. Wind speed ranged from 2 to 4 kmph.</p>	<p>There is possibility of no rain the coming week. Max temp is expected to be around 21⁰C - 23⁰C and min temp is expected to be around 8⁰C to 9⁰C. Possibility of mainly clear the coming week Relative Humidity will range from 31% to 95% The wind speed 2- 4 kmph Wind direction will blow southerly</p>

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Paddy field	-	-	<p>Crop rotation should be carried out as it may interrupt the relationship between pests and host plant. Rice based cropping systems involving pre-kharif green-manure crop followed by kharif rice and non- rice rabi crop directly benefit the rice crop.</p>
Horticultural crops			
Citrus dieback (Mandarin orange)	-	-	<p>Leaves become yellow and fall with the beginning of summer and die-back of twig starts leading to drying of branches and declining – death of trees. Transmission of disease take place through infected planting materials, mechanically through contaminated pruning knife and sometimes from soil borne pathogen. So, famers should take precautionary measures.</p>
Summer vegetables	-	-	<p>Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. as growing of early summer vegetables crops will fetch more income.</p>
Potato	Sowing stage	-	<p>At the time of planting, the tubers should closely be inspected and all suspicious looking tubers should be rejected.</p>
Fishery	<p>Stocking should not be delayed once a new pond has filled. As soon as the pond has adequate water in it, contamination by unwanted fish is possible before a good fish population develops. Prevent livestock from entering the pond and degrading the banks of the pond.</p>		

	<i>Piggery</i>	Deworming of the pigs should be done routinely and clean, fresh water — and lots of it — must always be available to your pigs.
 The logo of the Indian Council of Agricultural Research (ICAR) is located in the bottom-left corner of the second row. It features a stylized green plant with three leaves inside a circular frame, with the text 'भारतीय कृषि अनुसंधान परिषद' in Hindi and 'ICAR' in English below it.	<i>Poultry</i>	During this season, plenty of cool, clean fresh drinking water should be provided as the water consumption increases during summer and feed should be provided during cooler parts of the day such as early morning or late evening.



Integrated Agromet Advisory Service Bulletin 5th to 9th March'14

Dimapur District

Bulletin No:17/2014

Weather summary of the preceding week

No rain occurred the past week.
Maximum and **minimum temperatures** ranged 25⁰C to 26⁰C and 11⁰C to 12⁰C, respectively.
Relative humidity varied from 27% to 95%.
Wind speed ranged from 2 to 3 kmph.

Weather forecast valid upto 9th March'14

There is possibility of no rain the coming week.
Max temp is expected to be around 23⁰C - 26⁰C and **min temp** is expected to be around 11⁰C to 12⁰C.
Possibility of mainly clear sky the coming week
Relative Humidity will range from 21% to 87%
Wind speed 1– 2 kmph
Wind direction will be from easterly

Main Crops	Stage	Pest/ diseases	Agro-meteorological Advisories
<i>Field crops</i>			
<i>Paddy field</i>	-	-	Crop rotation should be carried out as it may interrupt the relationship between pests and host plant. Rice based cropping systems involving pre-kharif green-manure crop followed by kharif rice and non- rice rabi crop directly benefit the rice crop.
<i>Horticultural crops</i>			
<i>Citrus dieback (Mandarin orange)</i>	-	-	Leaves become yellow and fall with the beginning of summer and die-back of twig starts leading to drying of branches and declining – death of trees. Transmission of disease take place through infected planting materials, mechanically through contaminated pruning knife and sometimes from soil borne pathogen. So, famers should take precautionary measures.
<i>Summer vegetables</i>	-	-	Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. as growing of early summer vegetables crops will fetch more income.
<i>Fishery</i>	Stocking should not be delayed once a new pond has filled. As soon as the pond has adequate water in it, contamination by unwanted fish is possible before a good fish population develops. Prevent livestock from entering the pond and degrading the banks of the pond.		
<i>Piggery</i>	Deworming of the pigs should be done routinely and clean, fresh water — and lots of it — must always be available to your pigs.		

ICAR RESEARCH COMPLEX FOR NEH REGION
 (Indian Council of Agricultural Research)
 NAGALAND CENTER, JHARNAPANI, MEDZIPHEMA



Poultry

During this season, plenty of cool, clean fresh drinking water should be provided as the water consumption increases during summer and feed should be provided during cooler parts of the day such as early morning or late evening.