



# Integrated Agromet Advisory Service Bulletin 12<sup>th</sup> to 16<sup>th</sup> March'14

## Longleng District

**Bulletin No:19/2014**

### *Weather summary of the preceding week*

No rain occurred the past week.  
**Maximum** and **minimum temperatures** ranged from 22<sup>o</sup>C to 25<sup>o</sup>C and 8<sup>o</sup>C to 9<sup>o</sup>C, respectively.  
**Relative humidity** varied from 22% to 80%.  
**Wind speed** ranged from 3 to 4 kmph.

### *Weather forecast valid upto 16<sup>th</sup> March'14*

There is possibility of no rain the coming week.  
**Max temp** is expected to be around 25<sup>o</sup>C - 27<sup>o</sup>C and **min temp** is expected to be around 8<sup>o</sup>C to 9<sup>o</sup>C.  
**Possibility of mainly clear sky** the coming week  
**Relative Humidity** will range from 20% to 63%  
The **wind speed** 3 –4 kmph  
Wind direction will blow from southerly

### **Main Crops**

### **Stage**

### **Pest/ diseases**

### **Agro-meteorological Advisories**

### *Field crops*

#### *Cereal crops*

-

-

Cultivate cereal crops like maize, paddy in rotation with legume crops like soybean, peas, beans etc to escape disease and for sustainable yield.

### *Horticultural crops*

#### *Mandarin orange*

-

Citrus dieback

Leaves become yellow and fall with the beginning of summer and die-back of twig starts leading to drying of branches and declining – death of trees. Transmission of disease take place through infected planting materials, mechanically through contaminated pruning knife and sometimes from soil borne pathogen. So, famers should take precautionary measures - proper drainage should be made, certified planting material should be use and good cultural practices should be carried out.

#### *Summer vegetables*

-

-

Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. and certified seeds should be use to produce good yield.

#### *Water harvesting*

-

-

Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price.

#### *Potato*

Sowing stage

Early blight

Proper sanitation should be maintained to prevent early blight. Weeds should not be allowed to grow on borders of the field.



<b><i>Fishery</i></b>	As most of the pond is dry, take up renovation work like repairing of dyke and drainage system and bottom of the pond should be cleaned.
<b><i>Piggery</i></b>	Deworming (feeding of anti worm drugs) should be provided to the piglets' right after weaning. Separate piglets from the mother (weaning) after 2 months of age for better growth of piglets
<b><i>Poultry</i></b>	During this season, avoid overcrowding of birds and floor space should be increased by at least 10%.



# Integrated Agromet Advisory Service Bulletin 12<sup>th</sup> to 16<sup>th</sup> March'14

## Mokokchung District

**Bulletin No: 19/2014**

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 16<sup>th</sup> March'14</i>
<p>No rain occurred the past week  <b>Maximum and minimum temperatures</b> ranged 20<sup>0</sup>C to 23<sup>0</sup>C and 10<sup>0</sup>C to 11<sup>0</sup>C, respectively.  <b>Relative humidity</b> varied from 24% to 85%.  <b>Wind speed</b> ranged from 3 to 4 kmph.</p>			<p>There is possibility of no rain the coming week.  <b>Max temp</b> is expected to be around 22<sup>0</sup>C - 24<sup>0</sup>C and <b>min temp</b> is expected to be around 10<sup>0</sup>C to 11<sup>0</sup>C.  <b>Possibility of clear sky the coming week</b>  <b>Relative Humidity</b> will range from 21% to 69% and  <b>Wind speed</b> 3– 4 kmph            Wind direction will blow from southerly.</p>
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<b>Field crops</b>			
<i>Cereal crops</i>	-	-	Cultivate cereal crops like maize, paddy in rotation with legume crops like soybean, peas, beans etc to escape disease and for sustainable yield.
<b>Horticultural crops</b>			
<i>Mandarin orange</i>	-	Citrus dieback	Leaves become yellow and fall with the beginning of summer and die-back of twig starts leading to drying of branches and declining – death of trees. Transmission of disease take place through infected planting materials, mechanically through contaminated pruning knife and sometimes from soil borne pathogen. So, famers should take precautionary measures - proper drainage should be made, certified planting material should be use and good cultural practices should be carried out.
<i>Summer vegetables</i>	-	-	Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. and certified seeds should be use to produce good yield.
<i>Water harvesting</i>	-	-	Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price.
<i>Potato</i>	Sowing stage	Early blight	Proper sanitation should be maintained to prevent early blight. Weeds should not be allowed to grow on borders of the field.
<i>Fishery</i>	As most of the pond is dry, take up renovation work like repairing of dyke and drainage system and bottom of the pond should be cleaned.		
<i>Piggery</i>	Deworming (feeding of anti worm drugs) should be provided to the piglets' right after weaning. Separate piglets from the mother (weaning) after 2 months of age for better growth of piglets		



***Poultry***

During this season, avoid overcrowding of birds and floor space should be increased by at least 10%.



# Integrated Agromet Advisory Service Bulletin from 12<sup>th</sup> to 16<sup>th</sup> March'14 Mon District

**Bulletin No:19/2014**


### *Weather summary of the preceding week*

- No rain occurred the past week
- **Maximum** and **minimum temperatures** ranged 22<sup>o</sup>C to 24<sup>o</sup>C and 7<sup>o</sup>C to 8<sup>o</sup>C, respectively.
- **Relative humidity** varied from 22% to 83%.
- **Wind speed** ranged from 3 to 4 kmph.

### *Weather forecast valid upto 16<sup>th</sup> March'14*

- Possibility of no rain the coming week.
- The **max temp** is likely to be 23<sup>o</sup>C - 25<sup>o</sup>C and the **min temp** 6<sup>o</sup>C - 7<sup>o</sup>C.
- **Sky is likely to be mainly clear the coming week**
- **Relative Humidity** is likely to range from 21% to 66%
- **Wind speed** may reach upto 3- 4kmph
- **Wind direction** will be from southerly

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<b><i>Field crops</i></b>			
<b><i>Cereal crops</i></b>	-	-	Cultivate cereal crops like maize, paddy in rotation with legume crops like soybean, peas, beans etc to escape disease and for sustainable yield.
<b><i>Horticultural crops</i></b>			
<b><i>Mandarin orange</i></b>	-	Citrus dieback	Leaves become yellow and fall with the beginning of summer and die-back of twig starts leading to drying of branches and declining – death of trees. Transmission of disease take place through infected planting materials, mechanically through contaminated pruning knife and sometimes from soil borne pathogen. So, famers should take precautionary measures - proper drainage should be made, certified planting material should be use and good cultural practices should be carried out.
<b><i>Summer vegetables</i></b>	-	-	Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. and certified seeds should be use to produce good yield.
<b><i>Water harvesting</i></b>	-	-	Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price.
<b><i>Potato</i></b>	Sowing stage	Early blight	Proper sanitation should be maintained to prevent early blight. Weeds should not be allowed to grow on borders of the field.
<b><i>Fishery</i></b>	As most of the pond is dry, take up renovation work like repairing of dyke and drainage system and bottom of the pond should be cleaned.		

	<b><i>Piggery</i></b>	Deworming (feeding of anti worm drugs) should be provided to the piglets' right after weaning. Separate piglets from the mother (weaning) after 2 months of age for better growth of piglets
	<b><i>Poultry</i></b>	During this season, avoid overcrowding of birds and floor space should be increased by at least 10%.





# Integrated Agromet Advisory Service Bulletin 12<sup>th</sup> to 16<sup>th</sup> March'14

## Peren District

**Bulletin No:19/2014**

### *Weather summary of the preceding week*

No rain occurred the past week.  
**Maximum** and **minimum temperature** range from 23<sup>0</sup>C to 24<sup>0</sup>C and 7<sup>0</sup>C to 9<sup>0</sup>C, respectively.  
**Relative humidity** varied from 19% to 90%.  
**Wind speed** ranged from 1 to 2 kmph.

### *Weather forecast valid upto 16<sup>th</sup> March'14*

There is possibility of no rain the coming week.  
**Max temp** is expected to be around 24<sup>0</sup>C - 25<sup>0</sup>C and **min temp** is expected to be around 8<sup>0</sup>C to 9<sup>0</sup>C.  
**Possibility of mainly clear sky the coming week**  
**Relative Humidity** will range from 18% to 76%  
**The wind speed** may reach upto 1 - 2 kmph  
**Wind direction** will blow from easterly

<b>Main Crops</b>	<b>Stage</b>	<b>Pest/ diseases</b>	<b>Agro-meteorological Advisories</b>
<b><i>Field crops</i></b>			
<b><i>Cereal crops</i></b>	-	-	Cultivate cereal crops like maize, paddy in rotation with legume crops like soybean, peas, beans etc to escape disease and for sustainable yield.
<b><i>Horticultural crops</i></b>			
<b><i>Mandarin orange</i></b>	-	Citrus dieback	Leaves become yellow and fall with the beginning of summer and die-back of twig starts leading to drying of branches and declining – death of trees. Transmission of disease take place through infected planting materials, mechanically through contaminated pruning knife and sometimes from soil borne pathogen. So, famers should take precautionary measures - proper drainage should be made, certified planting material should be use and good cultural practices should be carried out.
<b><i>Summer vegetables</i></b>	-	-	Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. and certified seeds should be use to produce good yield.
<b><i>Water harvesting</i></b>	-	-	Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price.
<b><i>Potato</i></b>	Sowing stage	Early blight	Proper sanitation should be maintained to prevent early blight. Weeds should not be allowed to grow on borders of the field.



<b><i>Fishery</i></b>	As most of the pond is dry, take up renovation work like repairing of dyke and drainage system and bottom of the pond should be cleaned.
<b><i>Piggery</i></b>	Deworming (feeding of anti worm drugs) should be provided to the piglets' right after weaning. Separate piglets from the mother (weaning) after 2 months of age for better growth of piglets
<b><i>Poultry</i></b>	During this season, avoid overcrowding of birds and floor space should be increased by at least 10%.





भा क अनुप - पु प सं  
ICAR - RCNEH

CAR RESEARCH COMPLEX FOR NEH REGION

(Indian Council of Agricultural Research)


NAGALAND CENTER, JHARNAPANI, MEDZIPHEMA

## Integrated Agromet Advisory Service Bulletin 12<sup>th</sup> to 16<sup>th</sup> March'14 Phek District

**Bulletin No: 19/2014**

<i>Weather summary of the preceding week</i>	<i>Weather forecast valid upto 16<sup>th</sup> March'14</i>
<p>No rain occurred the past week  <b>Maximum</b> and <b>minimum temperatures</b> ranged from 17<sup>0</sup>C to 23<sup>0</sup>C and 7<sup>0</sup>C to 9<sup>0</sup>C, respectively.  <b>Relative humidity</b> varied from 25% to 93%.  <b>Wind speed</b> ranged from 2 to 4 kmph.</p>	<p>There is possibility of no rain the coming week.  <b>Max temp is expected to be around 22<sup>0</sup>C - 24<sup>0</sup>C</b> and <b>min temp is expected to be around 8<sup>0</sup>C to 9<sup>0</sup>C</b>.  <b>Possibility of mainly clear sky the coming week</b>  <b>Relative Humidity</b> will range from 21% to 78%  <b>The wind speed 3- 4 kmph</b>  <b>Wind direction</b> will blow from southerly</p>

<b>Main Crops</b>	<b>Stage</b>	<b>Pest/ Diseases</b>	<b>Agro-meteorological Advisories</b>
<b><i>Field crops</i></b>			
<b><i>Cereal crops</i></b>	-	-	Cultivate cereal crops like maize, paddy in rotation with legume crops like soybean, peas, beans etc to escape disease and for sustainable yield.
<b><i>Horticultural crops</i></b>			
<b><i>Mandarin orange</i></b>	-	Citrus dieback	Leaves become yellow and fall with the beginning of summer and die-back of twig starts leading to drying of branches and declining – death of trees. Transmission of disease take place through infected planting materials, mechanically through contaminated pruning knife and sometimes from soil borne pathogen. So, famers should take precautionary measures - proper drainage should be made, certified planting material should be use and good cultural practices should be carried out.
<b><i>Summer vegetables</i></b>	-	-	Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. and certified seeds should be use to produce good yield.
<b><i>Water harvesting</i></b>	-	-	Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price.
<b><i>Potato</i></b>	Sowing stage	Early blight	Proper sanitation should be maintained to prevent early blight. Weeds should not be allowed to grow on borders of the field.
<b><i>Fishery</i></b>	As most of the pond is dry, take up renovation work like repairing of dyke and drainage system and bottom of the pond should be cleaned.		

	<b><i>Piggery</i></b>	Deworming (feeding of anti worm drugs) should be provided to the piglets' right after weaning. Separate piglets from the mother (weaning) after 2 months of age for better growth of piglets
 <p>भारत ICAR</p>	<b><i>Poultry</i></b>	During this season, avoid overcrowding of birds and floor space should be increased by at least 10%.



# Integrated Agromet Advisory Service Bulletin 12<sup>th</sup> to 16<sup>th</sup> March'14

## Tuensang District

**Bulletin No:19/2014**

### *Weather summary of the preceding week*


No rain occurred the past week  
**Maximum and minimum temperatures** ranged 19<sup>0</sup>C to 20<sup>0</sup>C and 8<sup>0</sup>C to 9<sup>0</sup>C, respectively.  
**Relative humidity** varied from 26% to 94%.  
**Wind speed** ranged from 3 to 4 kmph.

### *Weather forecast valid upto 16<sup>th</sup> March'14*

There is possibility of no rain the coming week.  
**Max temp is expected to be around 19<sup>0</sup>C-20<sup>0</sup>C** and **min temp is expected to be around 8<sup>0</sup>C to 9<sup>0</sup>C**.  
**Possibility of mainly clear sky the coming week**  
**Relative humidity** will range from 24% to 86%  
**Wind speed** 2 – 4kmph  
 Wind direction will blow from westerly

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<b><i>Field crops</i></b>			
<b><i>Cereal crops</i></b>	-	-	Cultivate cereal crops like maize, paddy in rotation with legume crops like soybean, peas, beans etc to escape disease and for sustainable yield.
<b><i>Horticultural crops</i></b>			
<b><i>Mandarin orange</i></b>	-	Citrus dieback	Leaves become yellow and fall with the beginning of summer and die-back of twig starts leading to drying of branches and declining – death of trees. Transmission of disease take place through infected planting materials, mechanically through contaminated pruning knife and sometimes from soil borne pathogen. So, famers should take precautionary measures - proper drainage should be made, certified planting material should be use and good cultural practices should be carried out.
<b><i>Summer vegetables</i></b>	-	-	Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. and certified seeds should be use to produce good yield.
<b><i>Water harvesting</i></b>	-	-	Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price.
<b><i>Potato</i></b>	Sowing stage	Early blight	Proper sanitation should be maintained to prevent early blight. Weeds should not be allowed to grow on borders of the field.
<b><i>Fishery</i></b>	As most of the pond is dry, take up renovation work like repairing of dyke and drainage system and bottom of the pond should be cleaned.		

ICAR RESEARCH COMPLEX FOR NEH REGION  
 (Indian Council of Agricultural Research)  
 NAGALAND CENTER, JHARNAPANI, MEDZIPHEMA

	<b><i>Piggery</i></b>	Deworming (feeding of anti worm drugs) should be provided to the piglets' right after weaning. Separate piglets from the mother (weaning) after 2 months of age for better growth of piglets
 <p>भारत ICAR</p>	<b><i>Poultry</i></b>	During this season, avoid overcrowding of birds and floor space should be increased by at least 10%.



# Integrated Agromet Advisory Service Bulletin 12<sup>th</sup> to 16<sup>th</sup> March'14

## Wokha District

**Bulletin No:19/2014**

### *Weather summary of the preceding week*

No rain occurred the past week.  
**Maximum and minimum temperatures** ranged from 20<sup>o</sup>C to 23<sup>o</sup>C and 9<sup>o</sup>C to 10<sup>o</sup>C, respectively.  
**Relative humidity** varied from 24% to 91%.  
**Wind speed** ranged from 2 to 3 kmph

### *Weather forecast valid upto 16<sup>th</sup> March'14*

There is possibility of no rain the coming week.  
**Max temp is expected to be around 23<sup>o</sup>C - 24<sup>o</sup>C and min temp is expected to be around 9<sup>o</sup>C to 10<sup>o</sup>C.**  
**Possibility of clear sky the coming week**  
**Relative Humidity** will range from 19% to 68%  
**The wind speed** 3– 4 kmph  
**Wind direction** will blow from southerly


Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<b><i>Field crops</i></b>			
<b><i>Cereal crops</i></b>	-	-	Cultivate cereal crops like maize, paddy in rotation with legume crops like soybean, peas, beans etc to escape disease and for sustainable yield.
<b><i>Horticultural crops</i></b>			
<b><i>Mandarin orange</i></b>	-	Citrus dieback	Leaves become yellow and fall with the beginning of summer and die-back of twig starts leading to drying of branches and declining – death of trees. Transmission of disease take place through infected planting materials, mechanically through contaminated pruning knife and sometimes from soil borne pathogen. So, famers should take precautionary measures - proper drainage should be made, certified planting material should be use and good cultural practices should be carried out.
<b><i>Summer vegetables</i></b>	-	-	Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. and certified seeds should be use to produce good yield.
<b><i>Water harvesting</i></b>	-	-	Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price.
<b><i>Potato</i></b>	Sowing stage	Early blight	Proper sanitation should be maintained to prevent early blight. Weeds should not be allowed to grow on borders of the field.

**ICAR RESEARCH COMPLEX FOR NEH REGION**  
 (Indian Council of Agricultural Research)  
**NAGALAND CENTER, JHARNAPANI, MEDZIPHEMA**



<b><i>Fishery</i></b>	As most of the pond is dry, take up renovation work like repairing of dyke and drainage system and bottom of the pond should be cleaned.
<b><i>Piggery</i></b>	Deworming (feeding of anti worm drugs) should be provided to the piglets' right after weaning. Separate piglets from the mother (weaning) after 2 months of age for better growth of piglets
<b><i>Poultry</i></b>	During this season, avoid overcrowding of birds and floor space should be increased by at least 10%.










भा क अनुप - पु प सं  
ICAR - RCNEH

ICAR RESEARCH COMPLEX FOR NEH REGION  
(Indian Council of Agricultural Research)  
NAGALAND CENTER, JHARNAPANI, MEDZIPHEMA

## Integrated Agromet Advisory Service Bulletin 12<sup>th</sup> to 16<sup>th</sup> March'14 Zunheboto District

**Bulletin No:19/2014**

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 16<sup>th</sup> March'14</i>
<p>No rain occurred the past week  <b>Maximum and minimum temperatures</b> ranged from 22<sup>o</sup>C to 24<sup>o</sup>C and 8<sup>o</sup>C to 9<sup>o</sup>C, respectively.  <b>Relative humidity</b> varied from 25% to 92%.  <b>Wind speed</b> ranged from 3 to 4 kmph.</p>			<p>There is possibility of no rain the coming week.  <b>Max temp</b> is expected to be around 22<sup>o</sup>C - 24<sup>o</sup>C and <b>min temp</b> is expected to be around 8<sup>o</sup>C to 9<sup>o</sup>C.  <b>Possibility of clear sky the coming week</b>  <b>Relative Humidity</b> will range from 21% to 75%  <b>The wind speed</b> 3- 4 kmph  <b>Wind direction</b> will blow southerly</p>
<b>Main Crops</b>	<b>Stage</b>	<b>Pest/ Diseases</b>	<b>Agro-meteorological Advisories</b>
<b>Field crops</b>			
<b>Cereal crops</b>	-	-	Cultivate cereal crops like maize, paddy in rotation with legume crops like soybean, peas, beans etc to escape disease and for sustainable yield.
<b>Horticultural crops</b>			
<b>Mandarin orange</b>	-	Citrus dieback	Leaves become yellow and fall with the beginning of summer and die-back of twig starts leading to drying of branches and declining – death of trees. Transmission of disease take place through infected planting materials, mechanically through contaminated pruning knife and sometimes from soil borne pathogen. So, famers should take precautionary measures - proper drainage should be made, certified planting material should be use and good cultural practices should be carried out.
<b>Summer vegetables</b>	-	-	Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. and certified seeds should be use to produce good yield.
<b>Water harvesting</b>	-	-	Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price.
<b>Potato</b>	Sowing stage	Early blight	Proper sanitation should be maintained to prevent early blight. Weeds should not be allowed to grow on borders of the field.
<b>Fishery</b>	As most of the pond is dry, take up renovation work like repairing of dyke and drainage system and bottom of the pond should be cleaned.		

	<b><i>Piggery</i></b>	Deworming (feeding of anti worm drugs) should be provided to the piglets' right after weaning. Separate piglets from the mother (weaning) after 2 months of age for better growth of piglets
 <p>भारतीय ICAR</p>	<b><i>Poultry</i></b>	During this season, avoid overcrowding of birds and floor space should be increased by at least 10%.





# Integrated Agromet Advisory Service Bulletin 12<sup>th</sup> to 16<sup>th</sup> March'14

## Dimapur District

**Bulletin No:19/2014**

### *Weather summary of the preceding week*

No rain occurred the past week.  
**Maximum** and **minimum temperatures** ranged 25<sup>0</sup>C to 26<sup>0</sup>C and 11<sup>0</sup>C to 12<sup>0</sup>C, respectively.  
**Relative humidity** varied from 22% to 26%.  
**Wind speed** ranged from 2 to 3 kmph.

### *Weather forecast valid upto 16<sup>th</sup> March'14*

There is possibility of no rain the coming week.  
**Max temp** is expected to be around 25<sup>0</sup>C - 26<sup>0</sup>C and **min temp** is expected to be around 11<sup>0</sup>C to 12<sup>0</sup>C.  
**Possibility of mainly clear sky the coming week**  
**Relative Humidity** will range from 17% to 63%  
**Wind speed** 2– 4 kmph  
**Wind direction** will be from easterly

<b>Main Crops</b>	<b>Stage</b>	<b>Pest/ diseases</b>	<b>Agro-meteorological Advisories</b>
<b><i>Field crops</i></b>			
<b><i>Cereal crops</i></b>	-	-	Cultivate cereal crops like maize, paddy in rotation with legume crops like soybean, peas, beans etc to escape disease and for sustainable yield.
<b><i>Horticultural crops</i></b>			
<b><i>Citrus dieback (Mandarin orange)</i></b>	-	-	Leaves become yellow and fall with the beginning of summer and die-back of twig starts leading to drying of branches and declining – death of trees. Transmission of disease take place through infected planting materials, mechanically through contaminated pruning knife and sometimes from soil borne pathogen. So, famers should take precautionary measures.
<b><i>Summer vegetables</i></b>	-	-	Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. as growing of early summer vegetables crops will fetch more income.
<b><i>Water harvesting</i></b>	-	-	Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price.
<b><i>Fishery</i></b>	As most of the pond is dry, take up renovation work like repairing of dyke and drainage system and bottom of the pond should be cleaned.		
<b><i>Piggery</i></b>	Deworming (feeding of anti worm drugs) should be provided to the piglets' right after weaning. Separate piglets from the mother (weaning) after 2 months of age for better growth of piglets		



***Poultry***

During this season, avoid overcrowding of birds and floor space should be increased by at least 10%.



# Integrated Agromet Advisory Service Bulletin 12<sup>th</sup> to 16<sup>th</sup> March'14

## Kiphire District

**Bulletin No:19/2014**


### *Weather summary of the preceding week*

No rain occurred the past week.  
**Maximum** and **minimum temperatures** ranged from 22<sup>o</sup>C to 23<sup>o</sup>C and 8<sup>o</sup>C to 9<sup>o</sup>C, respectively.  
**Relative humidity** varied from 26% to 92%.  
**Wind speed** ranged from 3 to 4 kmph.

### *Weather forecast valid upto 16<sup>th</sup> March'14*

There is possibility of no rain the coming week.  
**Max temp** is expected to be around 22<sup>o</sup>C - 23<sup>o</sup>C and **min temp** is expected to be around 8<sup>o</sup>C to 9<sup>o</sup>C.  
**Possibility of clear sky** the coming week  
**Relative Humidity** will range from 23% to 85%  
 The **wind speed** 3- 4 kmph  
 Wind direction will blow from westerly

Main Crops	Stage	Pest/ diseases	Agro-meteorological Advisories
<b><i>Field crops</i></b>			
<b><i>Cereal crops</i></b>	-	-	Cultivate cereal crops like maize, paddy in rotation with legume crops like soybean, peas, beans etc to escape disease and for sustainable yield.
<b><i>Horticultural crops</i></b>			
<b><i>Mandarin orange</i></b>	-	Citrus dieback	Leaves become yellow and fall with the beginning of summer and die-back of twig starts leading to drying of branches and declining – death of trees. Transmission of disease take place through infected planting materials, mechanically through contaminated pruning knife and sometimes from soil borne pathogen. So, famers should take precautionary measures - proper drainage should be made, certified planting material should be use and good cultural practices should be carried out.
<b><i>Summer vegetables</i></b>	-	-	Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. and certified seeds should be use to produce good yield.
<b><i>Water harvesting</i></b>	-	-	Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price.
<b><i>Potato</i></b>	Sowing stage	Early blight	Proper sanitation should be maintained to prevent early blight. Weeds should not be allowed to grow on borders of the field.
<b><i>Fishery</i></b>	As most of the pond is dry, take up renovation work like repairing of dyke and drainage system and bottom of the pond should be cleaned.		

	<b><i>Piggery</i></b>	Deworming (feeding of anti worm drugs) should be provided to the piglets' right after weaning. Separate piglets from the mother (weaning) after 2 months of age for better growth of piglets
 <p>ICAR</p>	<b><i>Poultry</i></b>	During this season, avoid overcrowding of birds and floor space should be increased by at least 10%.




## Integrated Agromet Advisory Service Bulletin 12<sup>th</sup> to 16<sup>th</sup> March'14

### Kohima District

**Bulletin No:19/2014**

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 16<sup>th</sup> March'14</i>
No rain occurred the past week. <b>Maximum</b> and <b>minimum temperatures</b> ranged from 20 <sup>0</sup> C to 21 <sup>0</sup> C and 7 <sup>0</sup> C to 8 <sup>0</sup> C, respectively. <b>Relative humidity</b> varied from 16% to 92%. <b>Wind speed</b> ranged from 1 to 2 kmph.			There is possibility of no rain the coming week. <b>Max temp</b> is expected to be around 20 <sup>0</sup> C - 21 <sup>0</sup> C and <b>min temp</b> is expected to be around 8 <sup>0</sup> C to 9 <sup>0</sup> C. <b>Possibility of mainly clear sky</b> the coming week <b>Relative Humidity</b> will range from 18% to 82% <b>Wind speed</b> 2-3 kmph <b>Wind direction</b> will blow from easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<b><i>Field crops</i></b>			
<b><i>Cereal crops</i></b>	-	-	Cultivate cereal crops like maize, paddy in rotation with legume crops like soybean, peas, beans etc to escape disease and for sustainable yield.
<b><i>Horticultural crops</i></b>			
<b><i>Mandarin orange</i></b>	-	Citrus dieback	Leaves become yellow and fall with the beginning of summer and die-back of twig starts leading to drying of branches and declining – death of trees. Transmission of disease take place through infected planting materials, mechanically through contaminated pruning knife and sometimes from soil borne pathogen. So, famers should take precautionary measures - proper drainage should be made, certified planting material should be use and good cultural practices should be carried out.
<b><i>Summer vegetables</i></b>	-	-	Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. and certified seeds should be use to produce good yield.
<b><i>Water harvesting</i></b>	-	-	Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price.
<b><i>Potato</i></b>	Sowing stage	Early blight	Proper sanitation should be maintained to prevent early blight. Weeds should not be allowed to grow on borders of the field.
<b><i>Fishery</i></b>	As most of the pond is dry, take up renovation work like repairing of dyke and drainage system and bottom of the pond should be cleaned.		



	<b><i>Piggery</i></b>	Deworming (feeding of anti worm drugs) should be provided to the piglets' right after weaning. Separate piglets from the mother (weaning) after 2 months of age for better growth of piglets
 The logo of the Indian Council of Agricultural Research (ICAR) is located in the bottom-left corner of the table. It features a stylized green plant with three leaves and a circular emblem above it, all enclosed within a white circle. Below the emblem, the text 'भारतीय' and 'ICAR' is written in black. भारतीय ICAR	<b><i>Poultry</i></b>	During this season, avoid overcrowding of birds and floor space should be increased by at least 10%.