



Integrated Agromet Advisory Service Bulletin 19th to 23rd March'14 Dimapur District

Bulletin No:21/2014

Weather summary of the preceding week			Weather forecast valid upto 23 rd March'14
<p>No rain occurred the past week. Maximum and minimum temperatures ranged 25⁰C to 26⁰C and 11⁰C to 12⁰C, respectively. Relative humidity varied from 17% to 66%. Wind speed ranged from 3 to 4 kmph.</p>			<p>There is possibility of light rain the coming week. Max temp is expected to be around 25⁰C - 26⁰C and min temp is expected to be around 11⁰C to 12⁰C. Possibility of partly cloudy sky the coming week Relative Humidity will range from 18% to 85% Wind speed 3– 4 kmph Wind direction will be from easterly</p>
Main Crops	Stage	Pest/ diseases	Agro-meteorological Advisories
Field crops			
Cereal crops	-	-	Cultivate cereal crops like maize, paddy in rotation with legume crops like soybean, peas, beans etc to escape disease and for sustainable yield.
Horticultural crops			
Citrus dieback (Mandarin orange)	-	-	Leaves become yellow and fall with the beginning of summer and die-back of twig starts leading to drying of branches and declining – death of trees. Transmission of disease take place through infected planting materials, mechanically through contaminated pruning knife and sometimes from soil borne pathogen. So, famers should take precautionary measures.
Summer vegetables	-	-	Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. as growing of early summer vegetables crops will fetch more income.
Water harvesting	-	-	Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price.
Fishery	As most of the pond is dry, take up renovation work like repairing of dyke and drainage system and bottom of the pond should be cleaned.		
Piggery	Deworming (feeding of anti worm drugs) should be provided to the piglets' right after weaning. Separate piglets from the mother (weaning) after 2 months of age for better growth of piglets		



Poultry

During this season, avoid overcrowding of birds and floor space should be increased by at least 10%.




Integrated Agromet Advisory Service Bulletin 19th to 23rd March'14

Kiphire District

Bulletin No:21/2014

Weather summary of the preceding week			Weather forecast valid upto 23 rd March'14
<p>No rain occurred the past week. Maximum and minimum temperatures ranged from 23⁰C to 26⁰C and 9⁰C to 11⁰C, respectively. Relative humidity varied from 23% to 85%. Wind speed ranged from 3 to 4 kmph.</p>			<p>There is possibility of light rain the coming week. Max temp is expected to be around 25⁰C - 26⁰C and min temp is expected to be around 10⁰C to 11⁰C. Possibility of partly cloudy sky the coming week Relative Humidity will range from 22% to 89% The wind speed 2- 4 kmph Wind direction will blow from westerly</p>
Main Crops	Stage	Pest/ diseases	Agro-meteorological Advisories
Field crops			
Cereal crops	-	-	Cultivate cereal crops like maize, paddy in rotation with legume crops like soybean, peas, beans etc to escape disease and for sustainable yield.
Horticultural crops			
Mandarin orange	-	Citrus dieback	Leaves become yellow and fall with the beginning of summer and die-back of twig starts leading to drying of branches and declining – death of trees. Transmission of disease take place through infected planting materials, mechanically through contaminated pruning knife and sometimes from soil borne pathogen. So, famers should take precautionary measures - proper drainage should be made, certified planting material should be use and good cultural practices should be carried out.
Summer vegetables	-	-	Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. and certified seeds should be use to produce good yield.
Water harvesting	-	-	Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price.
Potato	Sowing stage	Early blight	Proper sanitation should be maintained to prevent early blight. Weeds should not be allowed to grow on borders of the field.
Fishery	As most of the pond is dry, take up renovation work like repairing of dyke and drainage system and bottom of the pond should be cleaned.		


	<i>Piggery</i>	Deworming (feeding of anti worm drugs) should be provided to the piglets' right after weaning. Separate piglets from the mother (weaning) after 2 months of age for better growth of piglets
	<i>Poultry</i>	During this season, avoid overcrowding of birds and floor space should be increased by at least 10%.



Integrated Agromet Advisory Service Bulletin 19th to 23rd March'14 Kohima District

Bulletin No:21/2014

Weather summary of the preceding week			Weather forecast valid upto 23 rd March'14
<p>No rain occurred the past week. Maximum and minimum temperatures ranged from 20⁰C to 23⁰C and 7⁰C to 8⁰C, respectively. Relative humidity varied from 18% to 77%. Wind speed ranged from 1 to 2 kmph.</p>			<p>There is possibility of light rain the coming week. Max temp is expected to be around 22⁰C - 23⁰C and min temp is expected to be around 8⁰C to 9⁰C. Possibility of partly cloudy sky the coming week Relative Humidity will range from 17% to 85% Wind speed 1-2 kmph Wind direction will blow from easterly</p>
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Cereal crops	-	-	Cultivate cereal crops like maize, paddy in rotation with legume crops like soybean, peas, beans etc to escape disease and for sustainable yield.
Horticultural crops			
Mandarin orange	-	Citrus dieback	Leaves become yellow and fall with the beginning of summer and die-back of twig starts leading to drying of branches and declining – death of trees. Transmission of disease take place through infected planting materials, mechanically through contaminated pruning knife and sometimes from soil borne pathogen. So, famers should take precautionary measures - proper drainage should be made, certified planting material should be use and good cultural practices should be carried out.
Summer vegetables	-	-	Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. and certified seeds should be use to produce good yield.
Water harvesting	-	-	Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price.
Potato	Sowing stage	Early blight	Proper sanitation should be maintained to prevent early blight. Weeds should not be allowed to grow on borders of the field.
Fishery	As most of the pond is dry, take up renovation work like repairing of dyke and drainage system and bottom of the pond should be cleaned.		

	<i>Piggery</i>	Deworming (feeding of anti worm drugs) should be provided to the piglets' right after weaning. Separate piglets from the mother (weaning) after 2 months of age for better growth of piglets
	<i>Poultry</i>	During this season, avoid overcrowding of birds and floor space should be increased by at least 10%.



Integrated Agromet Advisory Service Bulletin 19th to 23rd March'14 Longleng District

Bulletin No:21/2014

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 23rd March'14</i>
<p>No rain occurred the past week. Maximum and minimum temperatures ranged from 25⁰C to 26⁰C and 8⁰C to 9⁰C, respectively. Relative humidity varied from 22% to 63%. Wind speed ranged from 3 to 4 kmph.</p>			<p>There is possibility of light rain the coming week. Max temp is expected to be around 25⁰C - 27⁰C and min temp is expected to be around 8⁰C to 9⁰C. Possibility of partly cloudy sky the coming week Relative Humidity will range from 22% to 79% The wind speed 3 –4 kmph Wind direction will blow from southerly</p>
Main Crops	Stage	Pest/ diseases	Agro-meteorological Advisories
Field crops			
Cereal crops	-	-	Cultivate cereal crops like maize, paddy in rotation with legume crops like soybean, peas, beans etc to escape disease and for sustainable yield.
Horticultural crops			
Mandarin orange	-	Citrus dieback	Leaves become yellow and fall with the beginning of summer and die-back of twig starts leading to drying of branches and declining – death of trees. Transmission of disease take place through infected planting materials, mechanically through contaminated pruning knife and sometimes from soil borne pathogen. So, famers should take precautionary measures - proper drainage should be made, certified planting material should be use and good cultural practices should be carried out.
Summer vegetables	-	-	Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. and certified seeds should be use to produce good yield.
Water harvesting	-	-	Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price.
Potato	Sowing stage	Early blight	Proper sanitation should be maintained to prevent early blight. Weeds should not be allowed to grow on borders of the field.



<i>Fishery</i>	As most of the pond is dry, take up renovation work like repairing of dyke and drainage system and bottom of the pond should be cleaned.
<i>Piggery</i>	Deworming (feeding of anti worm drugs) should be provided to the piglets' right after weaning. Separate piglets from the mother (weaning) after 2 months of age for better growth of piglets
<i>Poultry</i>	During this season, avoid overcrowding of birds and floor space should be increased by at least 10%.



Integrated Agromet Advisory Service Bulletin 19th to 23rd March'14

Mokokchung District

Bulletin No: 21/2014

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 23rd March'14</i>
<p>No rain occurred the past week Maximum and minimum temperatures ranged 22⁰C to 24⁰C and 10⁰C to 11⁰C, respectively. Relative humidity varied from 22% to 72%. Wind speed ranged from 4 to 6 kmph.</p>			<p>There is possibility of light rain the coming week. Max temp is expected to be around 22⁰C - 24⁰C and min temp is expected to be around 12⁰C to 13⁰C. Possibility of partly cloudy sky the coming week Relative Humidity will range from 22% to 82% and Wind speed 4– 6 kmph Wind direction will blow from southerly.</p>
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Cereal crops	-	-	Cultivate cereal crops like maize, paddy in rotation with legume crops like soybean, peas, beans etc to escape disease and for sustainable yield.
Horticultural crops			
Mandarin orange	-	Citrus dieback	Leaves become yellow and fall with the beginning of summer and die-back of twig starts leading to drying of branches and declining – death of trees. Transmission of disease take place through infected planting materials, mechanically through contaminated pruning knife and sometimes from soil borne pathogen. So, famers should take precautionary measures - proper drainage should be made, certified planting material should be use and good cultural practices should be carried out.
Summer vegetables	-	-	Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. and certified seeds should be use to produce good yield.
Water harvesting	-	-	Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price.
Potato	Sowing stage	Early blight	Proper sanitation should be maintained to prevent early blight. Weeds should not be allowed to grow on borders of the field.
Fishery	As most of the pond is dry, take up renovation work like repairing of dyke and drainage system and bottom of the pond should be cleaned.		
Piggery	Deworming (feeding of anti worm drugs) should be provided to the piglets' right after weaning. Separate piglets from the mother (weaning) after 2 months of age for better growth of piglets		



<i>Poultry</i>	During this season, avoid overcrowding of birds and floor space should be increased by at least 10%.
-----------------------	--



भा क अयुष - पु प स
ICAR - RCNEH

ICAR RESEARCH COMPLEX FOR NEH REGION
(Indian Council of Agricultural Research)
NAGALAND CENTER, JHARNAPANI, MEDZIPHEMA

Integrated Agromet Advisory Service Bulletin from 19th to 23rd March'14 Mon District

Bulletin No:21/2014

Weather summary of the preceding week

- No rain occurred the past week
- **Maximum** and **minimum temperatures** ranged 23⁰C to 25⁰C and 6⁰C to 7⁰C, respectively.
- **Relative humidity** varied from 22% to 70%.
- **Wind speed** ranged from 3 to 4 kmph.

Weather forecast valid upto 23rd March'14

- Possibility of light rain the coming week.
- The **max temp** is likely to be 23⁰C - 26⁰C and the **min temp** 6⁰C - 8⁰C.
- **Sky is likely to be partly cloudy the coming week**
- **Relative Humidity** is likely to range from 22% to 80%
- **Wind speed** may reach upto 3- 4kmph
- **Wind direction** will be from southerly

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Cereal crops	-	-	Cultivate cereal crops like maize, paddy in rotation with legume crops like soybean, peas, beans etc to escape disease and for sustainable yield.
Horticultural crops			
Mandarin orange	-	Citrus dieback	Leaves become yellow and fall with the beginning of summer and die-back of twig starts leading to drying of branches and declining – death of trees. Transmission of disease take place through infected planting materials, mechanically through contaminated pruning knife and sometimes from soil borne pathogen. So, famers should take precautionary measures - proper drainage should be made, certified planting material should be use and good cultural practices should be carried out.
Summer vegetables	-	-	Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. and certified seeds should be use to produce good yield.
Water harvesting	-	-	Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price.
Potato	Sowing stage	Early blight	Proper sanitation should be maintained to prevent early blight. Weeds should not be allowed to grow on borders of the field.
Fishery	As most of the pond is dry, take up renovation work like repairing of dyke and drainage system and bottom of the pond should be cleaned.		

	<i>Piggery</i>	Deworming (feeding of anti worm drugs) should be provided to the piglets' right after weaning. Separate piglets from the mother (weaning) after 2 months of age for better growth of piglets
	<i>Poultry</i>	During this season, avoid overcrowding of birds and floor space should be increased by at least 10%.





Integrated Agromet Advisory Service Bulletin 19th to 23rd March'14 Peren District

Bulletin No:21/2014

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 23rd March'14</i>
<p>No rain occurred the past week. Maximum and minimum temperature range from 23⁰C to 24⁰C and 8⁰C to 9⁰C, respectively. Relative humidity varied from 18% to 70%. Wind speed ranged from 1 to 2 kmph.</p>			<p>There is possibility of light rain the coming week. Max temp is expected to be around 24⁰C - 25⁰C and min temp is expected to be around 8⁰C to 9⁰C. Possibility of partly cloudy sky the coming week Relative Humidity will range from 18% to 85% The wind speed may reach upto 2 - 4 kmph Wind direction will blow from easterly</p>
Main Crops	Stage	Pest/ diseases	Agro-meteorological Advisories
Field crops			
Cereal crops	-	-	Cultivate cereal crops like maize, paddy in rotation with legume crops like soybean, peas, beans etc to escape disease and for sustainable yield.
Horticultural crops			
Mandarin orange	-	Citrus dieback	Leaves become yellow and fall with the beginning of summer and die-back of twig starts leading to drying of branches and declining – death of trees. Transmission of disease take place through infected planting materials, mechanically through contaminated pruning knife and sometimes from soil borne pathogen. So, famers should take precautionary measures - proper drainage should be made, certified planting material should be use and good cultural practices should be carried out.
Summer vegetables	-	-	Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. and certified seeds should be use to produce good yield.
Water harvesting	-	-	Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price.
Potato	Sowing stage	Early blight	Proper sanitation should be maintained to prevent early blight. Weeds should not be allowed to grow on borders of the field.



<i>Fishery</i>	As most of the pond is dry, take up renovation work like repairing of dyke and drainage system and bottom of the pond should be cleaned.
<i>Piggery</i>	Deworming (feeding of anti worm drugs) should be provided to the piglets' right after weaning. Separate piglets from the mother (weaning) after 2 months of age for better growth of piglets
<i>Poultry</i>	During this season, avoid overcrowding of birds and floor space should be increased by at least 10%.




Integrated Agromet Advisory Service Bulletin 19th to 23rd March'14

Phek District

Bulletin No: 21/2014

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 23rd March'14</i>
No rain occurred the past week Maximum and minimum temperatures ranged from 22 ⁰ C to 24 ⁰ C and 9 ⁰ C to 10 ⁰ C, respectively. Relative humidity varied from 21% to 80%. Wind speed ranged from 3 to 4 kmph.			There is possibility of light rain the coming week. Max temp is expected to be around 22 ⁰ C - 24 ⁰ C and min temp is expected to be around 9 ⁰ C to 10 ⁰ C. Possibility of partly cloudy sky the coming week Relative Humidity will range from 21% to 86% The wind speed 2- 4 kmph Wind direction will blow from southerly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Cereal crops	-	-	Cultivate cereal crops like maize, paddy in rotation with legume crops like soybean, peas, beans etc to escape disease and for sustainable yield.
Horticultural crops			
Mandarin orange	-	Citrus dieback	Leaves become yellow and fall with the beginning of summer and die-back of twig starts leading to drying of branches and declining – death of trees. Transmission of disease take place through infected planting materials, mechanically through contaminated pruning knife and sometimes from soil borne pathogen. So, famers should take precautionary measures - proper drainage should be made, certified planting material should be use and good cultural practices should be carried out.
Summer vegetables	-	-	Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. and certified seeds should be use to produce good yield.
Water harvesting	-	-	Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price.
Potato	Sowing stage	Early blight	Proper sanitation should be maintained to prevent early blight. Weeds should not be allowed to grow on borders of the field.
Fishery	As most of the pond is dry, take up renovation work like repairing of dyke and drainage system and bottom of the pond should be cleaned.		

	<i>Piggery</i>	Deworming (feeding of anti worm drugs) should be provided to the piglets' right after weaning. Separate piglets from the mother (weaning) after 2 months of age for better growth of piglets
	<i>Poultry</i>	During this season, avoid overcrowding of birds and floor space should be increased by at least 10%.



Integrated Agromet Advisory Service Bulletin 19th to 23rd March'14 Tuensang District

Bulletin No:21/2014


Weather summary of the preceding week

No rain occurred the past week
Maximum and minimum temperatures ranged 19⁰C to 20⁰C and 8⁰C to 9⁰C, respectively.
Relative humidity varied from 24% to 85%.
Wind speed ranged from 3 to 4 kmph.

Weather forecast valid upto 23rd March'14

There is possibility of light rain the coming week.
Max temp is expected to be around 19⁰C-20⁰C and min temp is expected to be around 8⁰C to 9⁰C.
Possibility of partly cloudy sky the coming week
Relative humidity will range from 23% to 81%
Wind speed 2 – 4kmph
Wind direction will blow from westerly

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Field crops</i>			
<i>Cereal crops</i>	-	-	Cultivate cereal crops like maize, paddy in rotation with legume crops like soybean, peas, beans etc to escape disease and for sustainable yield.
<i>Horticultural crops</i>			
<i>Mandarin orange</i>	-	Citrus dieback	Leaves become yellow and fall with the beginning of summer and die-back of twig starts leading to drying of branches and declining – death of trees. Transmission of disease take place through infected planting materials, mechanically through contaminated pruning knife and sometimes from soil borne pathogen. So, famers should take precautionary measures - proper drainage should be made, certified planting material should be use and good cultural practices should be carried out.
<i>Summer vegetables</i>	-	-	Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. and certified seeds should be use to produce good yield.
<i>Water harvesting</i>	-	-	Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price.
<i>Potato</i>	Sowing stage	Early blight	Proper sanitation should be maintained to prevent early blight. Weeds should not be allowed to grow on borders of the field.
<i>Fishery</i>	As most of the pond is dry, take up renovation work like repairing of dyke and drainage system and bottom of the pond should be cleaned.		

	<i>Piggery</i>	Deworming (feeding of anti worm drugs) should be provided to the piglets' right after weaning. Separate piglets from the mother (weaning) after 2 months of age for better growth of piglets
	<i>Poultry</i>	During this season, avoid overcrowding of birds and floor space should be increased by at least 10%.



Integrated Agromet Advisory Service Bulletin 19th to 23rd March '14 Wokha District

Bulletin No:21/2014

Weather summary of the preceding week

No rain occurred the past week.
Maximum and minimum temperatures ranged from 23⁰C to 24⁰C and 10⁰C to 13⁰C, respectively.
Relative humidity varied from 19% to 71%.
Wind speed ranged from 3 to 4 kmph

Weather forecast valid upto 23rd March '14

There is possibility of light rain the coming week.
Max temp is expected to be around 23⁰C - 24⁰C and min temp is expected to be around 12⁰C to 13⁰C.
Possibility of partly cloudy sky the coming week
Relative Humidity will range from 20% to 80%
The wind speed 4- 5 kmph
Wind direction will blow from southerly

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Cereal crops	-	-	Cultivate cereal crops like maize, paddy in rotation with legume crops like soybean, peas, beans etc to escape disease and for sustainable yield.
Horticultural crops			
Mandarin orange	-	Citrus dieback	Leaves become yellow and fall with the beginning of summer and die-back of twig starts leading to drying of branches and declining – death of trees. Transmission of disease take place through infected planting materials, mechanically through contaminated pruning knife and sometimes from soil borne pathogen. So, famers should take precautionary measures - proper drainage should be made, certified planting material should be use and good cultural practices should be carried out.
Summer vegetables	-	-	Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. and certified seeds should be use to produce good yield.
Water harvesting	-	-	Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price.
Potato	Sowing stage	Early blight	Proper sanitation should be maintained to prevent early blight. Weeds should not be allowed to grow on borders of the field.



<i>Fishery</i>	As most of the pond is dry, take up renovation work like repairing of dyke and drainage system and bottom of the pond should be cleaned.
<i>Piggery</i>	Deworming (feeding of anti worm drugs) should be provided to the piglets' right after weaning. Separate piglets from the mother (weaning) after 2 months of age for better growth of piglets
<i>Poultry</i>	During this season, avoid overcrowding of birds and floor space should be increased by at least 10%.




--	--



Integrated Agromet Advisory Service Bulletin 19th to 23rd March'14 Zunheboto District

Bulletin No:21/2014

Weather summary of the preceding week			Weather forecast valid upto 23 rd March'14
No rain occurred the past week Maximum and minimum temperatures ranged from 22°C to 24°C and 8°C to 9°C, respectively. Relative humidity varied from 21% to 78%. Wind speed ranged from 4 to 5 kmph.			There is possibility of light rain the coming week. Max temp is expected to be around 23°C - 24°C and min temp is expected to be around 8°C to 9°C. Possibility of partly cloudy sky the coming week Relative Humidity will range from 21% to 85% The wind speed 4- 5 kmph Wind direction will blow southerly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Cereal crops	-	-	Cultivate cereal crops like maize, paddy in rotation with legume crops like soybean, peas, beans etc to escape disease and for sustainable yield.
Horticultural crops			
Mandarin orange	-	Citrus dieback	Leaves become yellow and fall with the beginning of summer and die-back of twig starts leading to drying of branches and declining – death of trees. Transmission of disease take place through infected planting materials, mechanically through contaminated pruning knife and sometimes from soil borne pathogen. So, famers should take precautionary measures - proper drainage should be made, certified planting material should be use and good cultural practices should be carried out.
Summer vegetables	-	-	Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. and certified seeds should be use to produce good yield.
Water harvesting	-	-	Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price.
Potato	Sowing stage	Early blight	Proper sanitation should be maintained to prevent early blight. Weeds should not be allowed to grow on borders of the field.
Fishery	As most of the pond is dry, take up renovation work like repairing of dyke and drainage system and bottom of the pond should be cleaned.		

	<i>Piggery</i>	Deworming (feeding of anti worm drugs) should be provided to the piglets' right after weaning. Separate piglets from the mother (weaning) after 2 months of age for better growth of piglets
	<i>Poultry</i>	During this season, avoid overcrowding of birds and floor space should be increased by at least 10%.

