



Integrated Agromet Advisory Service Bulletin 17th to 21st Sept'14

Kiphire District

Bulletin No:72/2014

Weather summary of the preceding week			Weather forecast valid upto 21 st Sept'14
<p>Light rain occurred the past week. Maximum and minimum temperatures ranged from 22⁰C to 23⁰C and 15⁰C to 18⁰C, respectively. Relative humidity varied from 82% to 97%. Wind speed ranged from 2 to 3 kmph.</p>			<p>There is possibility of light rain the coming week. Max temp is expected to be around 19⁰C - 25⁰C and min temp is expected to be around 17⁰C to 18⁰C. Possibility of partly cloudy sky the coming week Relative Humidity will range from 75% to 97% The wind speed 2-4 kmph Wind direction will blow from westerly</p>
Main Crops	Stage	Pest/diseases	Agro-meteorological Advisories
Field crops			
Jhum paddy	Dough stage	Brown spot	Stubbles should be burnt after crop harvest where the disease incidence is severe.
TRC paddy	Panicle initiation stage	Gundhi bug	They feed on all parts of the plant. Proper water management and sanitation practices should be done.
Soybean	Flowering stage	-	Prevailing weather will have a positive impact on flowering.
Horticultural crops			
Mandarin orange	-	-	Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).
Ginger	Vegetative stage		Soil around the plants should be worked with the help of <i>khurpi</i> . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.
Colocasia	Flowering stage	-	Maintain proper sanitation in the field.
Chilli	Flowering to fruiting stage		Constant monitoring of chilli plant is advised.
Cucurbits	Harvesting stage	-	Avoid injuries during harvesting and handling.
Poultry	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date, Always ensure that any wet bedding is removed and replaced with fresh clean dry bedding.		
Piggery	Drinking water must be available at all times. It should be fresh and free from contamination. Pigs should be checked daily for signs of disease, injury or non-competitiveness (for food).		



Fishery

Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond.




Integrated Agromet Advisory Service Bulletin 17th to 21st Sept'14

Kohima District

Bulletin No:72/2014

Weather summary of the preceding week			Weather forecast valid upto 21 st Sept'14
<p>Light rain occurred the past week. Maximum and minimum temperatures ranged from 25⁰C to 28⁰C and 15⁰C to 16⁰C, respectively. Relative humidity varied from 54% to 97%. Wind speed ranged from 2 to 3 kmph.</p>			<p>There is possibility of light rain the coming week. Max temp is expected to be around 24⁰C - 25⁰C and min temp is expected to be around 15⁰C to 16⁰C. Possibility of mainly cloudy sky the coming week Relative Humidity will range from 56% to 95% Wind speed 2- 4 kmph Wind direction will blow from westerly</p>
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Jhum paddy	Dough stage	Brown spot	Stubbles should be burnt after crop harvest where the disease incidence is severe.
TRC paddy	Panicle initiation stage	Gundhi bug	They feed on all parts of the plant. Proper water management and sanitation practices should be done.
Soybean	Flowering stage	-	Prevailing weather will have a positive impact on flowering.
Horticultural crops			
Mandarin orange	-	-	Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).
Ginger	Vegetative stage		Soil around the plants should be worked with the help of <i>khurpi</i> . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.
Colocasia	Flowering stage	-	Maintain proper sanitation in the field.
Chilli	Flowering to fruiting stage		Constant monitoring of chilli plant is advised.
Cucurbits	Harvesting stage	-	Avoid injuries during harvesting and handling.
Poultry	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date, Always ensure that any wet bedding is removed and replaced with fresh clean dry bedding.		

	<i>Piggery</i>	Drinking water must be available at all times. It should be fresh and free from contamination. Pigs should be checked daily for signs of disease, injury or non-competitiveness (for food).
	<i>Fishery</i>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond.



Integrated Agromet Advisory Service Bulletin 17th to 21st Sept'14

Longleng District

Bulletin No:72/2014

Weather summary of the preceding week			Weather forecast valid upto 21 st Sept'14
<p>Light rain occurred the past week. Maximum and minimum temperatures ranged from 28⁰C to 29⁰C and 19⁰C to 20⁰C, respectively. Relative humidity varied from 65% to 95%. Wind speed ranged from 2 to 3 kmph.</p>			<p>There is possibility of light rain the coming week. Max temp is expected to be around 24⁰C - 30⁰C and min temp is expected to be around 19⁰C to 20⁰C. Possibility of partly cloudy sky the coming week Relative Humidity will range from 61% to 95% The wind speed 2 –3 kmph Wind direction will blow from southeasterly</p>
Main Crops	Stage	Pest/ diseases	Agro-meteorological Advisories
Field crops			
<i>Jhum paddy</i>	Dough stage	Brown spot	Stubbles should be burnt after crop harvest where the disease incidence is severe.
<i>TRC paddy</i>	Panicle initiation stage	Gundhi bug	They feed on all parts of the plant. Proper water management and sanitation practices should be done.
<i>Soybean</i>	Flowering stage	-	Prevailing weather will have a positive impact on flowering.
Horticultural crops			
<i>Mandarin orange</i>	-	-	Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).
<i>Ginger</i>	Vegetative stage		Soil around the plants should be worked with the help of <i>khurpi</i> . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.
<i>Colocasia</i>	Flowering stage	-	Maintain proper sanitation in the field.
<i>Chilli</i>	Flowering to fruiting stage		Constant monitoring of chilli plant is advised.
<i>Cucurbits</i>	Harvesting stage	-	Avoid injuries during harvesting and handling.
<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date, Always ensure that any wet bedding is		



		removed and replaced with fresh clean dry bedding.
	<i>Piggery</i>	Drinking water must be available at all times. It should be fresh and free from contamination. Pigs should be checked daily for signs of disease, injury or non-competitiveness (for food).
	<i>Fishery</i>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond.



Integrated Agromet Advisory Service Bulletin 17th to 21st Sept'14

Mokokchung District

Bulletin No: 72/2014

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 21st Sept'14</i>
Light rain occurred the past week Maximum and minimum temperatures ranged 27 ⁰ C to 29 ⁰ C and 19 ⁰ C to 20 ⁰ C, respectively. Relative humidity varied from 65% to 93%. Wind speed ranged from 2 to 4 kmph.			There is possibility of light rain the coming week. Max temp is expected to be around 23 ⁰ C - 30 ⁰ C and min temp is expected to be around 19 ⁰ C to 20 ⁰ C. Possibility of mainly cloudy sky the coming week Relative Humidity will range from 61% to 95% and Wind speed 2 - 4 kmph Wind direction will blow from southerly.
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
<i>Jhum paddy</i>	Dough stage	Brown spot	Stubbles should be burnt after crop harvest where the disease incidence is severe.
<i>TRC paddy</i>	Panicle initiation stage	Gundhi bug	They feed on all parts of the plant. Proper water management and sanitation practices should be done.
<i>Soybean</i>	Flowering stage	-	Prevailing weather will have a positive impact on flowering.
Horticultural crops			
<i>Mandarin orange</i>	-	-	Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).
<i>Ginger</i>	Vegetative stage		Soil around the plants should be worked with the help of <i>khurpi</i> . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.
<i>Colocasia</i>	Flowering stage	-	Maintain proper sanitation in the field.
<i>Chilli</i>	Flowering to fruiting stage		Constant monitoring of chilli plant is advised.
<i>Cucurbits</i>	Harvesting stage	-	Avoid injuries during harvesting and handling.
<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date, Always ensure that any wet bedding is removed and replaced with fresh clean dry bedding.		
<i>Piggery</i>	Drinking water must be available at all times. It should be fresh and free from contamination.		




		Pigs should be checked daily for signs of disease, injury or non-competitiveness (for food).
<i>Fishery</i>		Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond.



Integrated Agromet Advisory Service Bulletin from 17th to 21st Sept'14 Mon District

Bulletin No:72/2014

Weather summary of the preceding week			Weather forecast valid upto 21 st Sept'14
<ul style="list-style-type: none"> Light rain occurred the past week Maximum and minimum temperatures ranged 26⁰C to 27⁰C and 19⁰C to 20⁰C, respectively. Relative humidity varied from 69% to 95%. Wind speed ranged from 1 to 2 kmph 			<ul style="list-style-type: none"> Possibility of light rain the coming week. The max temp is likely to be 23⁰C - 28⁰C and the min temp 19⁰C - 20⁰C. Sky is likely to be mainly cloudy the coming week Relative Humidity is likely to range from 65% to 95% Wind speed may reach upto 2- 4 kmph Wind direction will be from southerly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Jhum paddy	Harvesting stage	Stored grain rice weevil	Keeping neem leaves and branches over the stored rice helps in controlling the pest.
TRC paddy	Panicle initiation stage	Gundhi bug	They feed on all parts of the plant. Proper water management and sanitation practices should be done.
Soybean	Flowering stage	-	Prevailing weather will have a positive impact on flowering.
Horticultural crops			
Mandarin orange	-	-	Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).
Ginger	Vegetative stage		Soil around the plants should be worked with the help of <i>khurpi</i> . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.
Colocasia	Flowering stage	-	Maintain proper sanitation in the field.
Chilli	Flowering to fruiting stage		Constant monitoring of chilli plant is advised.
Cucurbits	Harvesting stage	-	Avoid injuries during harvesting and handling.
Poultry	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date, Always ensure that any wet bedding is removed and replaced with fresh clean dry bedding.		


	<i>Piggery</i>	Drinking water must be available at all times. It should be fresh and free from contamination. Pigs should be checked daily for signs of disease, injury or non-competitiveness (for food).
	<i>Fishery</i>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond.



Integrated Agromet Advisory Service Bulletin 17th to 21st Sept'14 Peren District

Bulletin No:72/2014

Weather summary of the preceding week			Weather forecast valid upto 21 st Sept'14
<p>Light rain occurred the past week. Maximum and minimum temperature range from 29^oC to 31^oC and 18^oC to 19^oC, respectively. Relative humidity varied from 52% to 97%. Wind speed ranged from 2 to 3 kmph.</p>			<p>There is possibility of light rain the coming week. Max temp is expected to be around 24^oC - 32^oC and min temp is expected to be around 18^oC to 19^oC. Possibility of partly cloudy sky the coming week Relative Humidity will range from 54% to 95% The wind speed may reach upto 2 - 3 kmph Wind direction will blow from westerly</p>
Main Crops	Stage	Pest/ diseases	Agro-meteorological Advisories
Field crops			
Jhum paddy	Hard dough development stage	Brown spot	Stubbles should be burnt after crop harvest where the disease incidence is severe.
Lowland paddy	Tillering stage	Leaf folder	Farmers should do daily monitoring against leaf folder as infestation is more in maximum tillering stage. Cloudy and humid weather, shady locations, closer spacing favour pest build up.
Soybean	Flowering stage	-	Prevailing weather will have a positive impact on flowering.
Horticultural crops			
Mandarin orange	-	-	Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).
Ginger	Vegetative stage		Soil around the plants should be worked with the help of <i>khurpi</i> . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.
Colocasia	Flowering stage	-	Maintain proper sanitation in the field.
Chilli	Flowering to fruiting stage		Field sanitation should be maintained. Collect and burn the infected plant debris of chilli and other malformed plant parts lying in the field.
Cucurbits	Harvesting stage	-	Avoid injuries during harvesting and handling.

	<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date, Always ensure that any wet bedding is removed and replaced with fresh clean dry bedding.
	<i>Piggery</i>	Drinking water must be available at all times. It should be fresh and free from contamination. Pigs should be checked daily for signs of disease, injury or non-competitiveness (for food).
	<i>Fishery</i>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond.




Integrated Agromet Advisory Service Bulletin 17th to 21st Sept'14

Phek District

Bulletin No: 72/2014

Weather summary of the preceding week			Weather forecast valid upto 21 st Sept'14
Light rain occurred the past week Maximum and minimum temperatures ranged from 25 ⁰ C to 27 ⁰ C and 18 ⁰ C to 19 ⁰ C, respectively. Relative humidity varied from 68% to 95%. Wind speed ranged from 3 to 4 kmph.			There is possibility of light rain the coming week. Max temp is expected to be around 22 ⁰ C - 28 ⁰ C and min temp is expected to be around 18 ⁰ C to 19 ⁰ C. Possibility of partly cloudy sky the coming week Relative Humidity will range from 66% to 95% The wind speed 2- 3 kmph Wind direction will blow from southwesterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
<i>Jhum paddy</i>	Dough stage	Brown spot	Stubbles should be burnt after crop harvest where the disease incidence is severe.
<i>TRC paddy</i>	Panicle initiation stage	Gundhi bug	They feed on all parts of the plant. Proper water management and sanitation practices should be done.
<i>Soybean</i>	Flowering stage	-	Prevailing weather will have a positive impact on flowering.
Horticultural crops			
<i>Mandarin orange</i>	-	-	Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).
<i>Ginger</i>	Vegetative stage		Soil around the plants should be worked with the help of <i>khurpi</i> . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.
<i>Colocasia</i>	Flowering stage	-	Maintain proper sanitation in the field.
<i>Chilli</i>	Flowering to fruiting stage		Constant monitoring of chilli plant is advised.
<i>Cucurbits</i>	Harvesting stage	-	Avoid injuries during harvesting and handling.
<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date, Always ensure that any wet bedding is removed and		

		replaced with fresh clean dry bedding.
	<i>Piggery</i>	Drinking water must be available at all times. It should be fresh and free from contamination. Pigs should be checked daily for signs of disease, injury or non-competitiveness (for food).
	<i>Fishery</i>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond.




Integrated Agromet Advisory Service Bulletin 17th to 21st Sept'14

Tuensang District

Bulletin No:72/2014

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 21st Sept'14</i>
Light rain occurred the past week Maximum and minimum temperatures ranged 18 ⁰ C to 20 ⁰ C and 15 ⁰ C to 16 ⁰ C, respectively. Relative humidity varied from 85% to 98%. Wind speed ranged from 1 to 2 kmph.			There is possibility of light to medium rain the coming week. Max temp is expected to be around 17 ⁰ C - 23 ⁰ C and min temp is expected to be around 16 ⁰ C to 17 ⁰ C. Possibility of partly cloudy sky the coming week Relative humidity will range from 78% to 95% Wind speed 2– 3 kmph Wind direction will blow from westerly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
<i>Jhum paddy</i>	Dough stage	Brown spot	Stubbles should be burnt after crop harvest where the disease incidence is severe.
<i>TRC paddy</i>	Panicle initiation stage	Gundhi bug	They feed on all parts of the plant. Proper water management and sanitation practices should be done.
<i>Soybean</i>	Flowering stage	-	Prevailing weather will have a positive impact on flowering.
Horticultural crops			
<i>Mandarin orange</i>	-	-	Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).
<i>Ginger</i>	Vegetative stage		Soil around the plants should be worked with the help of <i>khurpi</i> . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.
<i>Colocasia</i>	Flowering stage	-	Maintain proper sanitation in the field.
<i>Chilli</i>	Flowering to fruiting stage		Constant monitoring of chilli plant is advised.
<i>Cucurbits</i>	Harvesting stage	-	Avoid injuries during harvesting and handling.
<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date, Always ensure that any wet bedding is removed and replaced with fresh clean dry bedding.		

	<i>Piggery</i>	Drinking water must be available at all times. It should be fresh and free from contamination. Pigs should be checked daily for signs of disease, injury or non-competitiveness (for food).
	<i>Fishery</i>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond.



Integrated Agromet Advisory Service Bulletin 17th to 21st Sept'14 Wokha District

Bulletin No:72/2014


Weather summary of the preceding week

Light rain occurred the past week.
Maximum and minimum temperatures ranged from 29⁰C to 30⁰C and 19⁰C to 20⁰C, respectively.
Relative humidity varied from 57% to 91%.
Wind speed ranged from 3 to 4 kmph

Weather forecast valid upto 21st Sept'14

There is possibility of light rain the coming week.
Max temp is expected to be around 24⁰C - 31⁰C and min temp is expected to be around 19⁰C to 20⁰C.
Possibility of partly cloudy sky the coming week
Relative Humidity will range from 56% to 95%
The wind speed 2– 4 kmph
Wind direction will blow from southerly

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Field crops</i>			
<i>Jhum paddy</i>	Dough stage	Brown spot	Stubbles should be burnt after crop harvest where the disease incidence is severe.
<i>TRC paddy</i>	Panicle initiation stage	Gundhi bug	They feed on all parts of the plant. Proper water management and sanitation practices should be done.
<i>Soybean</i>	Flowering stage	-	Prevailing weather will have a positive impact on flowering.
<i>Horticultural crops</i>			
<i>Mandarin orange</i>	-	-	Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).
<i>Ginger</i>	Vegetative stage		Soil around the plants should be worked with the help of <i>khurpi</i> . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.
<i>Colocasia</i>	Flowering stage	-	Maintain proper sanitation in the field.
<i>Chilli</i>	Flowering to fruiting stage		Constant monitoring of chilli plant is advised.
<i>Cucurbits</i>	Harvesting stage	-	Avoid injuries during harvesting and handling.

	<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date, Always ensure that any wet bedding is removed and replaced with fresh clean dry bedding.
	<i>Piggery</i>	Drinking water must be available at all times. It should be fresh and free from contamination. Pigs should be checked daily for signs of disease, injury or non-competitiveness (for food).
	<i>Fishery</i>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond.



--	--




Integrated Agromet Advisory Service Bulletin 17th to 21st Sept'14

Zunheboto District

Bulletin No:72/2014

Weather summary of the preceding week			Weather forecast valid upto 21 st Sept'14
Light rain occurred the past week Maximum and minimum temperatures ranged from 25 ^o C to 27 ^o C and 18 ^o C to 19 ^o C, respectively. Relative humidity varied from 70% to 97%. Wind speed ranged from 2 to 3 kmph.			There is possibility of light rain the coming week. Max temp is expected to be around 22 ^o C - 29 ^o C and min temp is expected to be around 18 ^o C to 19 ^o C. Possibility of mainly cloudy sky the coming week Relative Humidity will range from 65% to 95% The wind speed 2- 4 kmph Wind direction will blow southerly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
<i>Jhum paddy</i>	Dough stage	Brown spot	Stubbles should be burnt after crop harvest where the disease incidence is severe.
<i>TRC paddy</i>	Panicle initiation stage	Gundhi bug	They feed on all parts of the plant. Proper water management and sanitation practices should be done.
<i>Soybean</i>	Flowering stage	-	Prevailing weather will have a positive impact on flowering.
Horticultural crops			
<i>Mandarin orange</i>	-	-	Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).
<i>Ginger</i>	Vegetative stage		Soil around the plants should be worked with the help of <i>khurpi</i> . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.
<i>Colocasia</i>	Flowering stage	-	Maintain proper sanitation in the field.
<i>Chilli</i>	Flowering to fruiting stage		Constant monitoring of chilli plant is advised.
<i>Cucurbits</i>	Harvesting stage	-	Avoid injuries during harvesting and handling.

	<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date, Always ensure that any wet bedding is removed and replaced with fresh clean dry bedding.
	<i>Piggery</i>	Drinking water must be available at all times. It should be fresh and free from contamination. Pigs should be checked daily for signs of disease, injury or non-competitiveness (for food).
	<i>Fishery</i>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond.




Integrated Agromet Advisory Service Bulletin 17th to 21st Sept'14

Dimapur District

Bulletin No:72/2014

Weather summary of the preceding week			Weather forecast valid upto 21 st Sept'14
<p>Light rain occurred the past week. Maximum and minimum temperatures ranged 32⁰C to 33⁰C and 20⁰C to 21⁰C, respectively. Relative humidity varied from 47% to 92%. Wind speed ranged from 2 to 4 kmph.</p>			<p>There is possibility of light rain the coming week. Max temp is expected to be around 26⁰C - 34⁰C and min temp is expected to be around 20⁰C to 21⁰C. Possibility of mainly cloudy sky the coming week Relative Humidity will range from 47 to 95% Wind speed 2– 4kmph Wind direction will be from southerly</p>
Main Crops	Stage	Pest/ diseases	Agro-meteorological Advisories
Field crops			
Jhum paddy	Hard dough development stage	Brown spot	Stubbles should be burnt after crop harvest where the disease incidence is severe.
Lowland paddy	Tillering stage	Leaf folder	Farmers should do daily monitoring against leaf folder as infestation is more in maximum tillering stage. Cloudy and humid weather, shady locations, closer spacing favour pest build up.
Soybean	Flowering stage	-	Prevailing weather will have a positive impact on flowering.
Horticultural crops			
Mandarin orange	-	-	Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).
Ginger	Vegetative stage		Soil around the plants should be worked with the help of <i>khurpi</i> . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.
Colocasia	Flowering stage	-	Maintain proper sanitation in the field.
Chilli	Flowering to fruiting stage		Constant monitoring of chilli plant is advised.
Cucurbits	Harvesting stage	-	Avoid injuries during harvesting and handling.
Poultry	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date, Always ensure that any wet bedding is removed and		

		replaced with fresh clean dry bedding.
	<i>Piggery</i>	Drinking water must be available at all times. It should be fresh and free from contamination. Pigs should be checked daily for signs of disease, injury or non-competitiveness (for food).
	<i>Fishery</i>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond.