



Integrated Agromet Advisory Service Bulletin 3rd to 7th Oct'15 Dimapur District

Bulletin No:75/2015

Weather summary of the preceding week			Weather forecast valid upto 7 th Oct '15
			<ul style="list-style-type: none"> Probability of no rain the coming week. The max temp is likely to be 31⁰C - 33⁰C and the min temp 20⁰C - 21⁰C. Sky is likely to be mainly cloudy sky the coming week Relative Humidity is likely to range from 47% to 92% Wind speed 2 kmph Wind direction will be mostly southeasterly
Main Crops	Stage	Pest/ diseases	Agro-meteorological Advisories
Field crops			
Lowland paddy	Milk development stage	Gundi bug	<i>They feed on all parts of the plant. Proper water management and sanitation practices should be done.</i>
Jhum paddy	Harvesting stage		<i>As the grain is to be harvested, the store house should be fumigated; store bins cleaned and should be weather proof, rodent proof and damp proof.</i>
Maize	Harvesting stage	-	<i>For composite and hybrid, Do not wait for stalks and leaves to dry because mostly they remain green</i>
Groundnut	Fruiting stage	-	<i>Proper earthing up should be done.</i>
Horticultural crops			
Citrus	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
Chilli	Harvesting stage	-	<i>Avoid injuries during harvesting.</i>
Ginger	Vegetative stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>
Piggery	Avoid stagnant water in and around the sty, maintain proper hygienic condition.		
Poultry	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date		

***Fishery***

Chemicals found in or used on a pond's watershed can enter the pond and cause fish kills. Runoff following a heavy rain can transport a variety of unwanted chemicals into the pond including pesticides, herbicides, fertilizers, petroleum products.

Avoid areas where the likelihood of chemical runoff is high and avoid using chemicals on the watershed



Integrated Agromet Advisory Service Bulletin 3rd to 7th Oct'15

Kiphire District

Bulletin No:75/2015

Weather summary of the preceding week			Weather forecast valid upto 7 th Oct '15
			<p>Probability of no rain the coming week</p> <p>Max temp is expected to be around 26⁰C - 27⁰C and min temp is expected to be around 12⁰C to 13⁰C.</p> <p>Possibility of mainly cloudy sky the coming week</p> <p>Relative Humidity will range from 52% to 95%</p> <p>The wind speed 1- 2 kmph</p> <p>Wind direction will blow mostly from easterly</p>
Main Crops	Stage	Pest/ diseases	Agro-meteorological Advisories
Field crops			
Terrace/ Wetland paddy	Milk development stage	Gundi bug	<i>They feed on all parts of the plant. Proper water management and sanitation practices should be done.</i>
Jhum paddy	Harvesting stage		<i>As the grain is to be harvested, the store house should be fumigated; store bins cleaned and should be weather proof, rodent proof and damp proof.</i>
Maize	Harvesting stage	-	<i>For composite and hybrid, Do not wait for stalks and leaves to dry because mostly they remain green</i>
Groundnut	Fruiting stage	-	<i>Proper earthing up should be done.</i>
Horticultural crops			
Citrus	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
Chilli	Harvesting stage	-	<i>Avoid injuries during harvesting.</i>
Ginger	Vegetative stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>
Piggery	Avoid stagnant water in and around the sty, maintain proper hygienic condition.		
Poultry	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date		



Fishery

Chemicals found in or used on a pond's watershed can enter the pond and cause fish kills. Runoff following a heavy rain can transport a variety of unwanted chemicals into the pond including pesticides, herbicides, fertilizers, petroleum products.

Avoid areas where the likelihood of chemical runoff is high and avoid using chemicals on the watershed




Integrated Agromet Advisory Service Bulletin 3rd to 7th Oct'15

Kohima District

Bulletin No:75/2015

Weather summary of the preceding week			Weather forecast valid upto 7 th Oct '15
			Probability of light rain the coming week Max temp is expected to be around 24 ⁰ C - 25 ⁰ C and min temp is expected to be around 14 ⁰ C to 15 ⁰ C. Possibility of mainly cloudy sky the coming week Relative Humidity will range from 44% to 94% Wind speed 2 kmph Wind direction will be mostly easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Terrace/ Wetland paddy	Milk development stage	Gundi bug	<i>They feed on all parts of the plant. Proper water management and sanitation practices should be done.</i>
Jhum paddy	Harvesting stage		<i>As the grain is to be harvested, the store house should be fumigated; store bins cleaned and should be weather proof, rodent proof and damp proof.</i>
Maize	Harvesting stage	-	<i>For composite and hybrid, Do not wait for stalks and leaves to dry because mostly they remain green</i>
Groundnut	Fruiting stage	-	<i>Proper earthing up should be done.</i>
Horticultural crops			
Citrus	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
Chilli	Harvesting stage	-	<i>Avoid injuries during harvesting.</i>
Ginger	Vegetative stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>
Piggery	Avoid stagnant water in and around the sty, maintain proper hygienic condition.		


	<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<i>Fishery</i>	<p>Chemicals found in or used on a pond's watershed can enter the pond and cause fish kills. Runoff following a heavy rain can transport a variety of unwanted chemicals into the pond including pesticides, herbicides, fertilizers, petroleum products.</p> <p>Avoid areas where the likelihood of chemical runoff is high and avoid using chemicals on the watershed</p>



Integrated Agromet Advisory Service Bulletin 3rd to 7th Oct'15 Longleng District

Bulletin No:75/2015

Weather summary of the preceding week			Weather forecast valid upto 7 th Oct '15
			<p>Probability of no rain the coming week Max temp is expected to be around 30⁰C - 31⁰C and min temp is expected to be around 16⁰C to 17⁰C. Possibility of cloudy sky the coming week Relative Humidity will range from 52% to 96% The wind speed 1 - 2 kmph Wind direction will be mostly easterly</p>
Main Crops	Stage	Pest/ diseases	Agro-meteorological Advisories
Field crops			
Terrace/ Wetland paddy	Milk development stage	Gundi bug	<i>They feed on all parts of the plant. Proper water management and sanitation practices should be done.</i>
Jhum paddy	Harvesting stage		<i>As the grain is to be harvested, the store house should be fumigated; store bins cleaned and should be weather proof, rodent proof and damp proof.</i>
Maize	Harvesting stage	-	<i>For composite and hybrid, Do not wait for stalks and leaves to dry because mostly they remain green</i>
Groundnut	Fruiting stage	-	<i>Proper earthing up should be done.</i>
Horticultural crops			
Citrus	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
Chilli	Harvesting stage	-	<i>Avoid injuries during harvesting.</i>
Ginger	Vegetative stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>
Piggery	Avoid stagnant water in and around the sty, maintain proper hygienic condition.		

	<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<i>Fishery</i>	<p>Chemicals found in or used on a pond's watershed can enter the pond and cause fish kills. Runoff following a heavy rain can transport a variety of unwanted chemicals into the pond including pesticides, herbicides, fertilizers, petroleum products.</p> <p>Avoid areas where the likelihood of chemical runoff is high and avoid using chemicals on the watershed</p>



Integrated Agromet Advisory Service Bulletin 3rd to 7th Oct'15

Mokokchung District

Bulletin No: 75/2015

Weather summary of the preceding week

Weather forecast valid upto 7th Oct '15

Probability no rain will occur the coming week
 Max temp is expected to be around 29⁰C - 30⁰C and min temp is expected to be around 16⁰C to 18⁰C.
 Possibility of mainly cloudy sky the coming week
 Relative Humidity will range from 51% to 95% and
 Wind speed 2 kmph
 Wind direction will be mostly easterly

Main Crops

Stage

Pest/ Diseases

Agro-meteorological Advisories

Field crops

Terrace/ Wetland paddy	Milk development stage	Gundhi bug	<i>They feed on all parts of the plant. Proper water management and sanitation practices should be done.</i>
Jhum paddy	Harvesting stage		<i>As the grain is to be harvested, the store house should be fumigated; store bins cleaned and should be weather proof, rodent proof and damp proof.</i>
Maize	Harvesting stage	-	<i>For composite and hybrid, Do not wait for stalks and leaves to dry because mostly they remain green</i>
Groundnut	Fruiting stage	-	<i>Proper earthing up should be done.</i>

Horticultural crops

Citrus	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
Chilli	Harvesting stage	-	<i>Avoid injuries during harvesting.</i>
Ginger	Vegetative stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>
Piggery	Avoid stagnant water in and around the sty, maintain proper hygienic condition.		
Poultry	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date		



Fishery

Chemicals found in or used on a pond's watershed can enter the pond and cause fish kills. Runoff following a heavy rain can transport a variety of unwanted chemicals into the pond including pesticides, herbicides, fertilizers, petroleum products.


Avoid areas where the likelihood of chemical runoff is high and avoid using chemicals on the watershed



Integrated Agromet Advisory Service Bulletin from 3rd to 7th Oct'15 Mon District

Bulletin No:75/2015

Weather summary of the preceding week			Weather forecast valid upto 7 th Oct '15
			<ul style="list-style-type: none"> ▪ Probability of no rain will occur the coming week. ▪ The max temp is likely to be 29⁰C - 30⁰C and the min temp 15⁰C - 16⁰C. ▪ Sky is likely to be mainly cloudy the coming week ▪ Relative Humidity is likely to range from 52% to 92% ▪ Wind speed may reach upto 1- 2 kmph ▪ Wind direction will be mostly easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Terrace/ Wetland paddy	Milk development stage	Gundi bug	<i>They feed on all parts of the plant. Proper water management and sanitation practices should be done.</i>
Jhum paddy	Harvesting stage		<i>As the grain is to be harvested, the store house should be fumigated; store bins cleaned and should be weather proof, rodent proof and damp proof.</i>
Maize	Harvesting stage	-	<i>For composite and hybrid, Do not wait for stalks and leaves to dry because mostly they remain green</i>
Groundnut	Fruiting stage	-	<i>Proper earthing up should be done.</i>
Horticultural crops			
Citrus	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
Chilli	Harvesting stage	-	<i>Avoid injuries during harvesting.</i>
Ginger	Vegetative stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>
Piggery	Avoid stagnant water in and around the sty, maintain proper hygienic condition.		

	<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<i>Fishery</i>	<p>Chemicals found in or used on a pond's watershed can enter the pond and cause fish kills. Runoff following a heavy rain can transport a variety of unwanted chemicals into the pond including pesticides, herbicides, fertilizers, petroleum products.</p> <p>Avoid areas where the likelihood of chemical runoff is high and avoid using chemicals on the watershed</p>




Integrated Agromet Advisory Service Bulletin 3rd to 7th Oct'15

Peren District

Bulletin No:75/2015

Weather summary of the preceding week			Weather forecast valid upto 7 th Oct '15
			<p>Probability of light rain the coming week. Max temp is expected to be around 31⁰C - 32⁰C and min temp is expected to be around 17⁰C to 19⁰C. Possibility of mainly cloudy sky the coming week Relative Humidity will range from 42% to 95% The wind speed 2 kmph Wind direction will be easterly</p>
Main Crops	Stage	Pest/ diseases	Agro-meteorological Advisories
Field crops			
Lowland paddy	Milk development stage	Gundi bug	They feed on all parts of the plant. Proper water management and sanitation practices should be done.
Jhum paddy	Harvesting stage		As the grain is to be harvested, the store house should be fumigated; store bins cleaned and should be weather proof, rodent proof and damp proof.
Maize	Harvesting stage	-	For composite and hybrid, Do not wait for stalks and leaves to dry because mostly they remain green
Groundnut	Fruiting stage	-	Proper earthing up should be done.
Horticultural crops			
Citrus	-		Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).
Chilli	Harvesting stage	-	Avoid injuries during harvesting.
Ginger	Vegetative stage		Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.

	<i>Piggery</i>	Avoid stagnant water in and around the sty, maintain proper hygienic condition.
	<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<i>Fishery</i>	Chemicals found in or used on a pond's watershed can enter the pond and cause fish kills. Runoff following a heavy rain can transport a variety of unwanted chemicals into the pond including pesticides, herbicides, fertilizers, petroleum products. Avoid areas where the likelihood of chemical runoff is high and avoid using chemicals on the watershed




Integrated Agromet Advisory Service Bulletin 3rd to 7th Oct'15

Phek District

Bulletin No:75/2015

CAR RESEARCH COMPLEX FOR NEH REGION
(Indian Council of Agricultural Research)
NAGALAND CENTER, JHARNAPANI, MEDZIPHEMA

Weather summary of the preceding week			Weather forecast valid upto 7 th Oct '15
			<p>Probability there will be light rain the coming week</p> <p>Max temp is expected to be around 27⁰C - 29⁰C and min temp is expected to be around 14⁰C to 15⁰C.</p> <p>Possibility of mainly cloudy sky the coming week</p> <p>Relative Humidity will range from 51% to 95%</p> <p>The wind speed 1-2 kmph</p> <p>Wind direction will blow mostly from easterly</p>
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Terrace/ Wetland paddy	Milk development stage	Gundi bug	<i>They feed on all parts of the plant. Proper water management and sanitation practices should be done.</i>
Jhum paddy	Harvesting stage		<i>As the grain is to be harvested, the store house should be fumigated; store bins cleaned and should be weather proof, rodent proof and damp proof.</i>
Maize	Harvesting stage	-	<i>For composite and hybrid, Do not wait for stalks and leaves to dry because mostly they remain green</i>
Groundnut	Fruiting stage	-	<i>Proper earthing up should be done.</i>
Horticultural crops			
Citrus	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
Chilli	Harvesting stage	-	<i>Avoid injuries during harvesting.</i>
Ginger	Vegetative stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>

	<i>Piggery</i>	Avoid stagnant water in and around the sty, maintain proper hygienic condition.
	<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<i>Fishery</i>	Chemicals found in or used on a pond's watershed can enter the pond and cause fish kills. Runoff following a heavy rain can transport a variety of unwanted chemicals into the pond including pesticides, herbicides, fertilizers, petroleum products. Avoid areas where the likelihood of chemical runoff is high and avoid using chemicals on the watershed




Integrated Agromet Advisory Service Bulletin 3rd to 7th Oct'15

Tuensang District

Bulletin No:75/2015

Weather summary of the preceding week			Weather forecast valid upto 7 th Oct '15
			Probability there will be light rain the coming week Max temp is expected to be around 24 ⁰ C - 25 ⁰ C and min temp is expected to be around 9 ⁰ C to 10 ⁰ C. Possibility of mainly cloudy sky the coming week Relative humidity will range from 53% to 93% Wind speed 1-2 kmph Wind direction will be mostly south easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Terrace/ Wetland paddy	Milk development stage	Gundi bug	<i>They feed on all parts of the plant. Proper water management and sanitation practices should be done.</i>
Jhum paddy	Harvesting stage		<i>As the grain is to be harvested, the store house should be fumigated; store bins cleaned and should be weather proof, rodent proof and damp proof.</i>
Maize	Harvesting stage	-	<i>For composite and hybrid, Do not wait for stalks and leaves to dry because mostly they remain green</i>
Groundnut	Fruiting stage	-	<i>Proper earthing up should be done.</i>
Horticultural crops			
Citrus	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
Chilli	Harvesting stage	-	<i>Avoid injuries during harvesting.</i>
Ginger	Vegetative stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>
Piggery	Avoid stagnant water in and around the sty, maintain proper hygienic condition.		


	<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<i>Fishery</i>	<p>Chemicals found in or used on a pond's watershed can enter the pond and cause fish kills. Runoff following a heavy rain can transport a variety of unwanted chemicals into the pond including pesticides, herbicides, fertilizers, petroleum products.</p> <p>Avoid areas where the likelihood of chemical runoff is high and avoid using chemicals on the watershed</p>



Integrated Agromet Advisory Service Bulletin 3rd to 7th Oct'15 Wokha District

Bulletin No:75/2015

Weather summary of the preceding week			Weather forecast valid upto 7 th Oct '15
			<p>Probability no rain will occur the coming week</p> <p>Max temp is expected to be around 28⁰C - 31⁰C and min temp is expected to be around 17⁰C to 19⁰C.</p> <p>Possibility of mainly cloudy sky the coming week</p> <p>Relative Humidity will range from 52% to 95%</p> <p>The wind speed 1- 2 kmph</p> <p>Wind direction will blow mostly from easterly</p>
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Terrace/ Wetland paddy	Milk development stage	Gundi bug	<i>They feed on all parts of the plant. Proper water management and sanitation practices should be done.</i>
Jhum paddy	Harvesting stage		<i>As the grain is to be harvested, the store house should be fumigated; store bins cleaned and should be weather proof, rodent proof and damp proof.</i>
Maize	Harvesting stage	-	<i>For composite and hybrid, Do not wait for stalks and leaves to dry because mostly they remain green</i>
Groundnut	Fruiting stage	-	<i>Proper earthing up should be done.</i>
Horticultural crops			
Citrus	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
Chilli	Harvesting stage	-	<i>Avoid injuries during harvesting.</i>
Ginger	Vegetative stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>

	<i>Piggery</i>	Avoid stagnant water in and around the sty, maintain proper hygienic condition.
	<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<i>Fishery</i>	Chemicals found in or used on a pond's watershed can enter the pond and cause fish kills. Runoff following a heavy rain can transport a variety of unwanted chemicals into the pond including pesticides, herbicides, fertilizers, petroleum products. Avoid areas where the likelihood of chemical runoff is high and avoid using chemicals on the watershed




--	--



Integrated Agromet Advisory Service Bulletin 3rd to 7th Oct'15 Zunheboto District

Bulletin No:75/2015

Weather summary of the preceding week			Weather forecast valid upto 7 th Oct '15
			<p>Probability light rain will occur the coming week</p> <p>Max temp is expected to be around 27⁰C - 29⁰C and min temp is expected to be around 14⁰C to 15⁰C.</p> <p>Possibility of mainly cloudy sky the coming week</p> <p>Relative Humidity will range from 51% to 91%</p> <p>The wind speed 2 kmph</p> <p>Wind direction will blow mostly from easterly</p>
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Terrace/ Wetland paddy	Milk development stage	Gundi bug	<i>They feed on all parts of the plant. Proper water management and sanitation practices should be done.</i>
Jhum paddy	Harvesting stage		<i>As the grain is to be harvested, the store house should be fumigated; store bins cleaned and should be weather proof, rodent proof and damp proof.</i>
Maize	Harvesting stage	-	<i>For composite and hybrid, Do not wait for stalks and leaves to dry because mostly they remain green</i>
Groundnut	Fruiting stage	-	<i>Proper earthing up should be done.</i>
Horticultural crops			
Citrus	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
Chilli	Harvesting stage	-	<i>Avoid injuries during harvesting.</i>
Ginger	Vegetative stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>
Piggery	Avoid stagnant water in and around the sty, maintain proper hygienic condition.		

	<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<i>Fishery</i>	<p>Chemicals found in or used on a pond's watershed can enter the pond and cause fish kills. Runoff following a heavy rain can transport a variety of unwanted chemicals into the pond including pesticides, herbicides, fertilizers, petroleum products.</p> <p>Avoid areas where the likelihood of chemical runoff is high and avoid using chemicals on the watershed</p>

