




## Integrated Agromet Advisory Service Bulletin 7<sup>th</sup> to 11<sup>th</sup> Oct'15 Kohima District

Bulletin No:76/2015

Weather summary of the preceding week			Weather forecast valid upto 11 <sup>th</sup> Oct '15
<ul style="list-style-type: none"> <li>No rain occurred the past week</li> <li>Maximum and minimum temperatures ranged 25<sup>0</sup>C to 26<sup>0</sup>C and 14<sup>0</sup>C to 15<sup>0</sup>C, respectively.</li> <li>Relative humidity varied from 43% to 95%.</li> <li>Wind speed ranged from 1 to 2 kmph</li> </ul>			Probability of light rain the coming week Max temp is expected to be around 24 <sup>0</sup> C - 25 <sup>0</sup> C and min temp is expected to be around 14 <sup>0</sup> C to 15 <sup>0</sup> C. Possibility of mainly cloudy sky the coming week Relative Humidity will range from 52% to 94% Wind speed 2 kmph Wind direction will be mostly easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<b>Field crops</b>			
<i>Terrace/ Wetland paddy</i>	Dough development stage		
<i>Jhum paddy</i>	Harvesting stage		Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
<i>Maize</i>	Harvesting stage	-	Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects
<i>Groundnut</i>	Fruiting stage	-	Proper earthing up should be done.
<b>Horticultural crops</b>			
<i>Citrus</i>	-		Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).
<i>Chilli</i>	Harvesting stage	-	Harvest fully ripe fruits. If its for storage, ensure proper drying.
<i>Ginger</i>	Maturity stage		Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.


	<b><i>Piggery</i></b>	Precaution should be taken against food and Mouth disease
	<b><i>Poultry</i></b>	In case of <b>diarrhoea in poultry</b> , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<b><i>Fishery</i></b>	Farmers are advised to keep the surrounding clean by cutting grasses and shrubs growing on embankment of the pond to remove shelter of predator of fish like snake, otter etc.



## Integrated Agromet Advisory Service Bulletin 7<sup>th</sup> to 11<sup>th</sup> Oct'15 Longleng District

Bulletin No:76/2015

Weather summary of the preceding week			Weather forecast valid upto 11 <sup>th</sup> Oct '15
<ul style="list-style-type: none"> <li>No rain occurred the past week</li> <li>Maximum and minimum temperatures ranged 30<sup>0</sup>C to 32<sup>0</sup>C and 17<sup>0</sup>C to 20<sup>0</sup>C, respectively.</li> <li>Relative humidity varied from 73% to 95%.</li> <li>Wind speed ranged from 1 to 2 kmph</li> </ul>			Probability of light rain the coming week <b>Max temp is expected to be around 28<sup>0</sup>C - 31<sup>0</sup>C and min temp is expected to be around 15<sup>0</sup>C to 17<sup>0</sup>C.</b> <b>Possibility of cloudy sky the coming week</b> <b>Relative Humidity</b> will range from 52% to 96% The <b>wind speed</b> 1 - 2 kmph Wind direction will be mostly southeasterly
Main Crops	Stage	Pest/diseases	Agro-meteorological Advisories
<b>Field crops</b>			
<b>Terrace/Wetland paddy</b>	Dough development stage		
<b>Jhum paddy</b>	Harvesting stage		Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
<b>Maize</b>	Harvesting stage	-	Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects
<b>Groundnut</b>	Fruiting stage	-	Proper earthing up should be done.
<b>Horticultural crops</b>			
<b>Citrus</b>	-		Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).
<b>Chilli</b>	Harvesting stage	-	Harvest fully ripe fruits. If its for storage, ensure proper drying.
<b>Ginger</b>	Maturity stage		Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.

	<b><i>Piggery</i></b>	Precaution should be taken against food and Mouth disease
	<b><i>Poultry</i></b>	In case of <b>diarrhoea in poultry</b> , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<b><i>Fishery</i></b>	Farmers are advised to keep the surrounding clean by cutting grasses and shrubs growing on embankment of the pond to remove shelter of predator of fish like snake, otter etc.

# Integrated Agromet Advisory Service Bulletin 7<sup>th</sup> to 11<sup>th</sup> Oct'15

## Mokokchung District

Bulletin No: 76/2015

Weather summary of the preceding week			Weather forecast valid upto 11 <sup>th</sup> Oct '15
<ul style="list-style-type: none"> <li>No rain occurred the past week</li> <li>Maximum and minimum temperatures ranged 29<sup>0</sup>C to 30<sup>0</sup>C and 16<sup>0</sup>C to 17<sup>0</sup>C, respectively.</li> <li>Relative humidity varied from 51% to 90%.</li> <li>Wind speed ranged from 1 to 2 kmph</li> </ul>			<p>Probability light rain will occur the coming week</p> <p>Max temp is expected to be around 27<sup>0</sup>C - 28<sup>0</sup>C and min temp is expected to be around 15<sup>0</sup>C to 17<sup>0</sup>C.</p> <p>Possibility of mainly cloudy sky the coming week</p> <p>Relative Humidity will range from 65% to 94% and</p> <p>Wind speed 2 kmph</p> <p>Wind direction will be mostly south easterly</p>
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<b>Field crops</b>			
<b>Terrace/ Wetland paddy</b>	Dough development stage		
<b>Jhum paddy</b>	Harvesting stage		Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
<b>Maize</b>	Harvesting stage	-	Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects
<b>Groundnut</b>	Fruiting stage	-	Proper earthing up should be done.
<b>Horticultural crops</b>			
<b>Citrus</b>	-		Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).
<b>Chilli</b>	Harvesting stage	-	Harvest fully ripe fruits. If its for storage, ensure proper drying.
<b>Ginger</b>	Maturity stage		Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.
<b>Piggery</b>	Precaution should be taken against food and Mouth disease		

<b><i>Poultry</i></b>	In case of <b>diarrhoea in poultry</b> , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
<b><i>Fishery</i></b>	Farmers are advised to keep the surrounding clean by cutting grasses and shrubs growing on embankment of the pond to remove shelter of predator of fish like snake, otter etc.






## Integrated Agromet Advisory Service Bulletin from 7<sup>th</sup> to 11<sup>th</sup> Oct'15 Mon District

Bulletin No:76/2015

Weather summary of the preceding week			Weather forecast valid upto 11 <sup>th</sup> Oct '15
<ul style="list-style-type: none"> <li>No rain occurred the past week</li> <li>Maximum and minimum temperatures ranged 29<sup>0</sup>C to 30<sup>0</sup>C and 15<sup>0</sup>C to 16<sup>0</sup>C, respectively.</li> <li>Relative humidity varied from 54% to 90%.</li> <li>Wind speed ranged from 1 to 2 kmph</li> </ul>			<ul style="list-style-type: none"> <li>Probability of light rain will occur the coming week.</li> <li>The max temp is likely to be 27<sup>0</sup>C - 30<sup>0</sup>C and the min temp 14<sup>0</sup>C - 18<sup>0</sup>C.</li> <li>Sky is likely to be mainly cloudy the coming week</li> <li>Relative Humidity is likely to range from 51% to 90%</li> <li>Wind speed may reach upto 1- 2 kmph</li> <li>Wind direction will be mostly south easterly</li> </ul>
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<b>Field crops</b>			
<b>Terrace/ Wetland paddy</b>	Dough development stage		
<b>Jhum paddy</b>	Harvesting stage		Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
<b>Maize</b>	Harvesting stage	-	Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects
<b>Groundnut</b>	Fruiting stage	-	Proper earthing up should be done.
<b>Horticultural crops</b>			
<b>Citrus</b>	-		Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).
<b>Chilli</b>	Harvesting stage	-	Harvest fully ripe fruits. If its for storage, ensure proper drying.
<b>Ginger</b>	Maturity stage		Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.
<b>Piggery</b>	Precaution should be taken against food and Mouth disease		

	<b><i>Poultry</i></b>	In case of <b>diarrhoea in poultry</b> , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<b><i>Fishery</i></b>	Farmers are advised to keep the surrounding clean by cutting grasses and shrubs growing on embankment of the pond to remove shelter of predator of fish like snake, otter etc.






## Integrated Agromet Advisory Service Bulletin 7<sup>th</sup> to 11<sup>th</sup> Oct'15 Peren District

Bulletin No:76/2015

Weather summary of the preceding week			Weather forecast valid upto 11 <sup>th</sup> Oct '15
<ul style="list-style-type: none"> <li>Light rain occurred the past week</li> <li>Maximum and minimum temperatures ranged 32<sup>0</sup>C to 34<sup>0</sup>C and 18<sup>0</sup>C to 19<sup>0</sup>C, respectively.</li> <li>Relative humidity varied from 44% to 99%.</li> <li>Wind speed ranged from 1 to 2 kmph</li> </ul>			Probability of light rain the coming week. <b>Max temp is expected to be around 26<sup>0</sup>C - 31<sup>0</sup>C and min temp is expected to be around 17<sup>0</sup>C to 19<sup>0</sup>C.</b> <b>Possibility of mainly cloudy sky the coming week</b> <b>Relative Humidity</b> will range from 49% to 95% The <b>wind speed</b> 2 kmph Wind direction will be easterly
Main Crops	Stage	Pest/diseases	Agro-meteorological Advisories
<b>Field crops</b>			
<b>Lowland paddy</b>	Dough stage		
<b>Jhum paddy</b>	Harvesting stage		<i>As the grain is to be harvested, the store house should be fumigated; store bins cleaned and should be weather proof, rodent proof and damp proof.</i>
<b>Maize</b>	Harvesting stage	-	<i>For composite and hybrid, Do not wait for stalks and leaves to dry because mostly they remain green</i>
<b>Groundnut</b>	Fruiting stage	-	<i>Proper earthing up should be done.</i>
<b>Horticultural crops</b>			
<b>Citrus</b>	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
<b>Chilli</b>	Harvesting stage	-	<i>Avoid injuries during harvesting.</i>
<b>Ginger</b>	Vegetative stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>

	<b><i>Piggery</i></b>	Avoid stagnant water in and around the sty, maintain proper hygienic condition.
	<b><i>Poultry</i></b>	In case of <b>diarrhoea in poultry</b> , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<b><i>Fishery</i></b>	Chemicals found in or used on a pond's watershed can enter the pond and cause fish kills. Runoff following a heavy rain can transport a variety of unwanted chemicals into the pond including pesticides, herbicides, fertilizers, petroleum products. Avoid areas where the likelihood of chemical runoff is high and avoid using chemicals on the watershed




# Integrated Agromet Advisory Service Bulletin 7<sup>th</sup> to 11<sup>th</sup> Oct'15

## Phek District

Bulletin No:76/2015

CAR RESEARCH COMPLEX FOR NEH REGION  
(Indian Council of Agricultural Research)  
NAGALAND CENTER, JHARNAPANI, MEDZIPHEMA

Weather summary of the preceding week			Weather forecast valid upto 11 <sup>th</sup> Oct '15
<ul style="list-style-type: none"> <li>No rain occurred the past week</li> <li>Maximum and minimum temperatures ranged 27<sup>0</sup>C to 29<sup>0</sup>C and 14<sup>0</sup>C to 16<sup>0</sup>C, respectively.</li> <li>Relative humidity varied from 51% to 98%.</li> <li>Wind speed ranged from 1 to 2 kmph</li> </ul>			<p>Probability there will be light rain the coming week</p> <p>Max temp is expected to be around 25<sup>0</sup>C - 29<sup>0</sup>C and min temp is expected to be around 14<sup>0</sup>C to 17<sup>0</sup>C.</p> <p>Possibility of mainly cloudy sky the coming week</p> <p>Relative Humidity will range from 51% to 95%</p> <p>The wind speed 1-2 kmph</p> <p>Wind direction will blow mostly from easterly</p>
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<b>Field crops</b>			
<b>Terrace/ Wetland paddy</b>	Dough development stage		
<b>Jhum paddy</b>	Harvesting stage		Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
<b>Maize</b>	Harvesting stage	-	Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects
<b>Groundnut</b>	Fruiting stage	-	Proper earthing up should be done.
<b>Horticultural crops</b>			
<b>Citrus</b>	-		Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).
<b>Chilli</b>	Harvesting stage	-	Harvest fully ripe fruits. If its for storage, ensure proper drying.
<b>Ginger</b>	Maturity stage		Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.

	<b><i>Piggery</i></b>	Precaution should be taken against food and Mouth disease
	<b><i>Poultry</i></b>	In case of <b>diarrhoea in poultry</b> , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<b><i>Fishery</i></b>	Farmers are advised to keep the surrounding clean by cutting grasses and shrubs growing on embankment of the pond to remove shelter of predator of fish like snake, otter etc.




## Integrated Agromet Advisory Service Bulletin 7<sup>th</sup> to 11<sup>th</sup> Oct'15

### Tuensang District

Bulletin No:76/2015

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 11<sup>th</sup> Oct '15</i>
<ul style="list-style-type: none"> <li>Light rain occurred the past week</li> <li>Maximum and minimum temperatures ranged 24<sup>0</sup>C to 25<sup>0</sup>C and 9<sup>0</sup>C to 10<sup>0</sup>C, respectively.</li> <li>Relative humidity varied from 56% to 95%.</li> <li>Wind speed ranged from 1 to 2 kmph</li> </ul>			Probability there will be light rain the coming week <b>Max temp is expected to be around 23<sup>0</sup>C - 25<sup>0</sup>C and min temp is expected to be around 9<sup>0</sup>C to 10<sup>0</sup>C.</b> <b>Possibility of mainly cloudy sky the coming week</b> <b>Relative humidity</b> will range from 52% to 93% <b>Wind speed</b> 1-2 kmph Wind direction will be mostly south easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<b>Field crops</b>			
<b>Terrace/ Wetland paddy</b>	Dough development stage		
<b>Jhum paddy</b>	Harvesting stage		Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
<b>Maize</b>	Harvesting stage	-	Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects
<b>Groundnut</b>	Fruiting stage	-	Proper earthing up should be done.
<b>Horticultural crops</b>			
<b>Citrus</b>	-		Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).
<b>Chilli</b>	Harvesting stage	-	Harvest fully ripe fruits. If its for storage, ensure proper drying.
<b>Ginger</b>	Maturity stage		Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.


	<b><i>Piggery</i></b>	Precaution should be taken against food and Mouth disease
	<b><i>Poultry</i></b>	In case of <b>diarrhoea in poultry</b> , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<b><i>Fishery</i></b>	Farmers are advised to keep the surrounding clean by cutting grasses and shrubs growing on embankment of the pond to remove shelter of predator of fish like snake, otter etc.



## Integrated Agromet Advisory Service Bulletin 7<sup>th</sup> to 11<sup>th</sup> Oct'15 Wokha District

Bulletin No:76/2015

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 11<sup>th</sup> Oct '15</i>
<ul style="list-style-type: none"> <li>No rain occurred the past week</li> <li>Maximum and minimum temperatures ranged 28<sup>0</sup>C to 31<sup>0</sup>C and 17<sup>0</sup>C to 19<sup>0</sup>C, respectively.</li> <li>Relative humidity varied from 51% to 95%.</li> <li>Wind speed ranged from 1 to 2 kmph</li> </ul>			Probability light rain will occur the coming week Max temp is expected to be around 29 <sup>0</sup> C - 31 <sup>0</sup> C and min temp is expected to be around 16 <sup>0</sup> C to 19 <sup>0</sup> C. Possibility of mainly cloudy sky the coming week Relative Humidity will range from 55% to 95% The wind speed 1- 2 kmph Wind direction will blow mostly from easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<b>Field crops</b>			
<i>Terrace/ Wetland paddy</i>	Dough development stage		
<i>Jhum paddy</i>	Harvesting stage		Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
<i>Maize</i>	Harvesting stage	-	Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects
<i>Groundnut</i>	Fruiting stage	-	Proper earthing up should be done.
<b>Horticultural crops</b>			
<i>Citrus</i>	-		Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).
<i>Chilli</i>	Harvesting stage	-	Harvest fully ripe fruits. If its for storage, ensure proper drying.
<i>Ginger</i>	Maturity stage		Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.

	<b><i>Piggery</i></b>	Precaution should be taken against food and Mouth disease
	<b><i>Poultry</i></b>	In case of <b>diarrhoea in poultry</b> , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<b><i>Fishery</i></b>	Farmers are advised to keep the surrounding clean by cutting grasses and shrubs growing on embankment of the pond to remove shelter of predator of fish like snake, otter etc.








--	--



## Integrated Agromet Advisory Service Bulletin 7<sup>th</sup> to 11<sup>th</sup> Oct'15 Zunheboto District

Bulletin No:76/2015

Weather summary of the preceding week			Weather forecast valid upto 11 <sup>th</sup> Oct '15
<ul style="list-style-type: none"> <li>No rain occurred the past week</li> <li>Maximum and minimum temperatures ranged 27<sup>0</sup>C to 29<sup>0</sup>C and 14<sup>0</sup>C to 16<sup>0</sup>C, respectively.</li> <li>Relative humidity varied from 51% to 94%.</li> <li>Wind speed ranged from 1 to 2 kmph</li> </ul>			Probability light rain will occur the coming week Max temp is expected to be around 27 <sup>0</sup> C - 29 <sup>0</sup> C and min temp is expected to be around 13 <sup>0</sup> C to 15 <sup>0</sup> C. Possibility of mainly cloudy sky the coming week Relative Humidity will range from 51% to 91% The wind speed 2 kmph Wind direction will blow mostly from easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<b>Field crops</b>			
<b>Terrace/ Wetland paddy</b>	Dough development stage		
<b>Jhum paddy</b>	Harvesting stage		Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
<b>Maize</b>	Harvesting stage	-	Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects
<b>Groundnut</b>	Fruiting stage	-	Proper earthing up should be done.
<b>Horticultural crops</b>			
<b>Citrus</b>	-		Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).
<b>Chilli</b>	Harvesting stage	-	Harvest fully ripe fruits. If its for storage, ensure proper drying.
<b>Ginger</b>	Maturity stage		Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.

	<b><i>Piggery</i></b>	Precaution should be taken against food and Mouth disease
	<b><i>Poultry</i></b>	In case of <b>diarrhoea in poultry</b> , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<b><i>Fishery</i></b>	Farmers are advised to keep the surrounding clean by cutting grasses and shrubs growing on embankment of the pond to remove shelter of predator of fish like snake, otter etc.






# Integrated Agromet Advisory Service Bulletin 7<sup>th</sup> to 11<sup>th</sup> Oct'15 Dimapur District

Bulletin No:76/2015

Weather summary of the preceding week			Weather forecast valid upto 11 <sup>th</sup> Oct '15
<ul style="list-style-type: none"> <li>Light rain occurred the past week</li> <li>Maximum and minimum temperatures ranged 32<sup>0</sup>C to 33<sup>0</sup>C and 19<sup>0</sup>C to 21<sup>0</sup>C, respectively.</li> <li>Relative humidity varied from 48% to 95%.</li> <li>Wind speed ranged from 1 to 2 kmph</li> </ul>			<ul style="list-style-type: none"> <li>Probability of light rain the coming week.</li> <li>The <b>max temp</b> is likely to be 30<sup>0</sup>C - 33<sup>0</sup>C and the <b>min temp</b> 20<sup>0</sup>C - 21<sup>0</sup>C.</li> <li><b>Sky is likely to be mainly cloudy sky the coming week</b></li> <li><b>Relative Humidity</b> is likely to range from 47% to 92%</li> <li><b>Wind speed</b> 2 kmph</li> <li><b>Wind direction</b> will be mostly southeasterly</li> </ul>
Main Crops	Stage	Pest/ diseases	Agro-meteorological Advisories
<b>Field crops</b>			
Lowland paddy	Dough stage		
Jhum paddy	Harvesting stage		Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
Maize	Harvesting stage	-	Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects
Groundnut	Fruiting stage	-	Proper earthing up should be done.
<b>Horticultural crops</b>			
Citrus	-		Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).
Chilli	Harvesting stage	-	Harvest fully ripe fruits. If its for storage, ensure proper drying.
Ginger	Maturity stage		Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.
Piggery	Precaution should be taken against food and Mouth disease		

	<b><i>Poultry</i></b>	In case of <b>diarrhoea in poultry</b> , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<b><i>Fishery</i></b>	Farmers are advised to keep the surrounding clean by cutting grasses and shrubs growing on embankment of the pond to remove shelter of predator of fish like snake, otter etc.



# Integrated Agromet Advisory Service Bulletin 7<sup>th</sup> to 11<sup>th</sup> Oct'15 Kiphire District

Bulletin No:76/2015

Weather summary of the preceding week			Weather forecast valid upto 11 <sup>th</sup> Oct '15
<ul style="list-style-type: none"> <li>No rain occurred the past week</li> <li>Maximum and minimum temperatures ranged 26<sup>0</sup>C to 28<sup>0</sup>C and 13<sup>0</sup>C to 15<sup>0</sup>C, respectively.</li> <li>Relative humidity varied from 57% to 90%.</li> <li>Wind speed ranged from 1 to 2 kmph</li> </ul>			Probability of no rain the coming week Max temp is expected to be around 23 <sup>0</sup> C - 27 <sup>0</sup> C and min temp is expected to be around 11 <sup>0</sup> C to 15 <sup>0</sup> C. Possibility of mainly cloudy sky the coming week Relative Humidity will range from 52% to 95% The wind speed 1- 2 kmph Wind direction will blow mostly from south easterly
Main Crops	Stage	Pest/diseases	Agro-meteorological Advisories
<b>Field crops</b>			
<b>Terrace/ Wetland paddy</b>	Dough development stage		
<b>Jhum paddy</b>	Harvesting stage		Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
<b>Maize</b>	Harvesting stage	-	Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects
<b>Groundnut</b>	Fruiting stage	-	Proper earthing up should be done.
<b>Horticultural crops</b>			
<b>Citrus</b>	-		Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).
<b>Chilli</b>	Harvesting stage	-	Harvest fully ripe fruits. If its for storage, ensure proper drying.
<b>Ginger</b>	Maturity stage		Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.
<b>Piggery</b>	Precaution should be taken against food and Mouth disease		
<b>Poultry</b>	In case of <b>diarrhoea in poultry</b> , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date		





### ***Fishery***

Farmers are advised to keep the surrounding clean by cutting grasses and shrubs growing on embankment of the pond to remove shelter of predator of fish like snake, otter etc.