



भा क अनुष - पु प सं
ICAR - RCNEH

ICAR RESEARCH COMPLEX FOR NEH REGION
(Indian Council of Agricultural Research)
NAGALAND CENTER, JHARNAPANI, MEDZIPHEMA

Integrated Agromet Advisory Service Bulletin 10th to 14th Oct'15

Longleng District

Bulletin No:77/2015

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 14th Oct '15</i>
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Maximum and minimum temperatures ranged 30⁰C to 32⁰C and 17⁰C to 20⁰C, respectively. ▪ Relative humidity varied from 73% to 95%. ▪ Wind speed ranged from 1 to 2 kmph 			Probability of no rain the coming week Max temp is expected to be around 27 ⁰ C-29 ⁰ C and min temp is expected to be around 14 ⁰ C to 15 ⁰ C. Possibility of mainly clear sky the coming week Relative Humidity will range from 49% to 91% The wind speed 1 - 2 kmph Wind direction will be mostly easterly
Main Crops	Stage	Pest/diseases	Agro-meteorological Advisories
Field crops			
<i>Terrace/Wetland paddy</i>	Dough development stage		
<i>Jhum paddy</i>	Harvesting stage		<i>Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds</i>
<i>Maize</i>	Harvesting stage	-	<i>Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects</i>
<i>Groundnut</i>	Fruiting stage	-	<i>Proper earthing up should be done.</i>
Horticultural crops			
<i>Citrus</i>	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
<i>Chilli</i>	Harvesting stage	-	<i>Harvest fully ripe fruits. If its for storage, ensure proper drying.</i>
<i>Ginger</i>	Maturity stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>

	<i>Piggery</i>	Precaution should be taken against food and Mouth disease
	<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<i>Fishery</i>	Farmers are advised to keep the surrounding clean by cutting grasses and shrubs growing on embankment of the pond to remove shelter of predator of fish like snake, otter etc.



Integrated Agromet Advisory Service Bulletin 10th to 14th Oct'15

Mokokchung District

Bulletin No: 77/2015

ICAR RESEARCH COMPLEX FOR NEH REGION
 (Indian Council of Agricultural Research)
 NAGALAND CENTER, JHARNAPANI, MEDZIPHEMA

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 14th Oct '15</i>
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Maximum and minimum temperatures ranged 29^oC to 30^oC and 16^oC to 17^oC, respectively. ▪ Relative humidity varied from 51% to 90%. ▪ Wind speed ranged from 1 to 2 kmph 			Probability no rain the coming week Max temp is expected to be around 27^oC - 29^oC and min temp is expected to be around 15^oC to 16^oC. Possibility of mainly clear sky the coming week Relative Humidity will range from 54% to 94% and Wind speed 2-3 kmph Wind direction will be mostly south easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
<i>Terrace/ Wetland paddy</i>	Dough development stage		
<i>Jhum paddy</i>	Harvesting stage		<i>Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds</i>
<i>Maize</i>	Harvesting stage	-	<i>Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects</i>
<i>Groundnut</i>	Fruiting stage	-	<i>Proper earthing up should be done.</i>
Horticultural crops			
<i>Citrus</i>	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
<i>Chilli</i>	Harvesting stage	-	<i>Harvest fully ripe fruits. If its for storage, ensure proper drying.</i>
<i>Ginger</i>	Maturity stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>
Piggery	Precaution should be taken against food and Mouth disease		

	<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
 भारतीय ICAR	<i>Fishery</i>	Farmers are advised to keep the surrounding clean by cutting grasses and shrubs growing on embankment of the pond to remove shelter of predator of fish like snake, otter etc.



Integrated Agromet Advisory Service Bulletin from 10th to 14th Oct'15 Mon District

Bulletin No:77/2015

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 14th Oct '15</i>
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Maximum and minimum temperatures ranged 29^oC to 30^oC and 15^oC to 16^oC, respectively. ▪ Relative humidity varied from 54% to 90%. ▪ Wind speed ranged from 1 to 2 kmph 			<ul style="list-style-type: none"> ▪ Probability of no rain the coming week. ▪ The max temp is likely to be 26^oC - 28^oC and the min temp 13^oC - 14^oC. ▪ Sky is likely to be mainly clear the coming week ▪ Relative Humidity is likely to range from 47% to 90% ▪ Wind speed may reach upto 2- 3 kmph ▪ Wind direction will be mostly south easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Field crops</i>			
<i>Terrace/ Wetland paddy</i>	Dough development stage		
<i>Jhum paddy</i>	Harvesting stage		<i>Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds</i>
<i>Maize</i>	Harvesting stage	-	<i>Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects</i>
<i>Groundnut</i>	Fruiting stage	-	<i>Proper earthing up should be done.</i>
<i>Horticultural crops</i>			
<i>Citrus</i>	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
<i>Chilli</i>	Harvesting stage	-	<i>Harvest fully ripe fruits. If its for storage, ensure proper drying.</i>
<i>Ginger</i>	Maturity stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>
<i>Piggery</i>	Precaution should be taken against food and Mouth disease		

ICAR RESEARCH COMPLEX FOR NEH REGION
 (Indian Council of Agricultural Research)
 NAGALAND CENTER, JHARNAPANI, MEDZIPHEMA

	<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
 The logo of the Indian Council of Agricultural Research (ICAR) is located in the bottom-left corner of the table. It features a stylized green plant with three leaves inside a circular frame, with the acronym 'ICAR' written in bold black letters below it. हिचडडडु ICAR	<i>Fishery</i>	Farmers are advised to keep the surrounding clean by cutting grasses and shrubs growing on embankment of the pond to remove shelter of predator of fish like snake, otter etc.



Integrated Agromet Advisory Service Bulletin 10th to 14th Oct'15

Peren District

Bulletin No:77/2015

Weather summary of the preceding week

- Light rain occurred the past week
- Maximum and minimum temperatures ranged 32^oC to 34^oC and 18^oC to 19^oC, respectively.
- Relative humidity varied from 44% to 99%.
- Wind speed ranged from 1 to 2 kmph

Weather forecast valid upto 14th Oct '15

Probability of light rain the coming week.
Max temp is expected to be around 27^oC - 28^oC and min temp is expected to be around 15^oC to 16^oC.
Possibility of mainly cloudy sky the coming week
Relative Humidity will range from 47% to 95%
The wind speed 2 kmph
Wind direction will be easterly

Main Crops

Stage

**Pest/
diseases**

Agro-meteorological Advisories

Field crops

Lowland paddy	Dough stage		
Jhum paddy	Harvesting stage		<i>As the grain is to be harvested, the store house should be fumigated; store bins cleaned and should be weather proof, rodent proof and damp proof.</i>
Maize	Harvesting stage	-	<i>For composite and hybrid, Do not wait for stalks and leaves to dry because mostly they remain green</i>
Groundnut	Fruiting stage	-	<i>Proper earthing up should be done.</i>

Horticultural crops

Citrus	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
Chilli	Harvesting stage	-	<i>Avoid injuries during harvesting.</i>
Ginger	Vegetative stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>

ICAR RESEARCH COMPLEX FOR NEH REGION

(Indian Council of Agricultural Research)

NAGALAND CENTER, JHARNAPANI, MEDZIPHEMA

	<i>Piggery</i>	Avoid stagnant water in and around the sty, maintain proper hygienic condition.
 <p>भारतीय ICAR</p>	<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<i>Fishery</i>	Chemicals found in or used on a pond's watershed can enter the pond and cause fish kills. Runoff following a heavy rain can transport a variety of unwanted chemicals into the pond including pesticides, herbicides, fertilizers, petroleum products. Avoid areas where the likelihood of chemical runoff is high and avoid using chemicals on the watershed



Integrated Agromet Advisory Service Bulletin 10th to 14th Oct'15

Phek District

Bulletin No:77/2015

CAR RESEARCH COMPLEX FOR NEH REGION

(Indian Council of Agricultural Research)

NAGALAND CENTER, JHARNAPANI, MEDZIPHEMA

Weather summary of the preceding week			Weather forecast valid upto 14 th Oct '15
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Maximum and minimum temperatures ranged 27^oC to 29^oC and 14^oC to 16^oC, respectively. ▪ Relative humidity varied from 51% to 98%. ▪ Wind speed ranged from 1 to 2 kmph 			<p>Probability there will be light rain the coming week Max temp is expected to be around 25^oC - 28^oC and min temp is expected to be around 12^oC to 15^oC. Possibility of mainly clear sky the coming week Relative Humidity will range from 42% to 95% The wind speed 1-2 kmph Wind direction will blow mostly from easterly</p>
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Terrace/ Wetland paddy	Dough development stage		
Jhum paddy	Harvesting stage		<i>Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds</i>
Maize	Harvesting stage	-	<i>Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects</i>
Groundnut	Fruiting stage	-	<i>Proper earthing up should be done.</i>
Horticultural crops			
Citrus	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
Chilli	Harvesting stage	-	<i>Harvest fully ripe fruits. If its for storage, ensure proper drying.</i>
Ginger	Maturity stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>

	<i>Piggery</i>	Precaution should be taken against food and Mouth disease
	<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
 <p>भारतीय ICAR</p>	<i>Fishery</i>	Farmers are advised to keep the surrounding clean by cutting grasses and shrubs growing on embankment of the pond to remove shelter of predator of fish like snake, otter etc.



Integrated Agromet Advisory Service Bulletin 10th to 14th Oct'15

Tuensang District

Bulletin No:77/2015

ICAR RESEARCH COMPLEX FOR NEH REGION
 (Indian Council of Agricultural Research)
 NAGALAND CENTER, JHARNAPANI, MEDZIPHEMA

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 14th Oct '15</i>
<ul style="list-style-type: none"> ▪ Light rain occurred the past week ▪ Maximum and minimum temperatures ranged 24⁰C to 25⁰C and 9⁰C to 10⁰C, respectively. ▪ Relative humidity varied from 56% to 95%. ▪ Wind speed ranged from 1 to 2 kmph 			Probability there will be light rain the coming week Max temp is expected to be around 20⁰C - 25⁰C and min temp is expected to be around 7⁰C to 11⁰C. Possibility of partly cloudy sky the coming week Relative humidity will range from 56% to 93% Wind speed 2-3 kmph Wind direction will be mostly south easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Field crops</i>			
<i>Terrace/ Wetland paddy</i>	Dough development stage		
<i>Jhum paddy</i>	Harvesting stage		<i>Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds</i>
<i>Maize</i>	Harvesting stage	-	<i>Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects</i>
<i>Groundnut</i>	Fruiting stage	-	<i>Proper earthing up should be done.</i>
<i>Horticultural crops</i>			
<i>Citrus</i>	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
<i>Chilli</i>	Harvesting stage	-	<i>Harvest fully ripe fruits. If its for storage, ensure proper drying.</i>
<i>Ginger</i>	Maturity stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>

	<i>Piggery</i>	Precaution should be taken against food and Mouth disease
	<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<i>Fishery</i>	Farmers are advised to keep the surrounding clean by cutting grasses and shrubs growing on embankment of the pond to remove shelter of predator of fish like snake, otter etc.



Integrated Agromet Advisory Service Bulletin 10th to 14th Oct'15

Wokha District

Bulletin No:77/2015

ICAR RESEARCH COMPLEX FOR NEH REGION
 (Indian Council of Agricultural Research)
 NAGALAND CENTER, JHARNAPANI, MEDZIPHEMA

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 14th Oct '15</i>
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Maximum and minimum temperatures ranged 28^oC to 31^oC and 17^oC to 19^oC, respectively. ▪ Relative humidity varied from 51% to 95%. ▪ Wind speed ranged from 1 to 2 kmph 			Probability light rain will occur the coming week Max temp is expected to be around 27^oC - 30^oC and min temp is expected to be around 15^oC to 16^oC. Possibility of mainly clear sky the coming week Relative Humidity will range from 42% to 95% The wind speed 1- 2 kmph Wind direction will blow mostly from easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Field crops</i>			
<i>Terrace/ Wetland paddy</i>	Dough development stage		
<i>Jhum paddy</i>	Harvesting stage		<i>Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds</i>
<i>Maize</i>	Harvesting stage	-	<i>Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects</i>
<i>Groundnut</i>	Fruiting stage	-	<i>Proper earthing up should be done.</i>
<i>Horticultural crops</i>			
<i>Citrus</i>	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
<i>Chilli</i>	Harvesting stage	-	<i>Harvest fully ripe fruits. If its for storage, ensure proper drying.</i>
<i>Ginger</i>	Maturity stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>

	<i>Piggery</i>	Precaution should be taken against food and Mouth disease
	<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<i>Fishery</i>	Farmers are advised to keep the surrounding clean by cutting grasses and shrubs growing on embankment of the pond to remove shelter of predator of fish like snake, otter etc.





भो कृ अनुप - पु प सं
ICAR - RCNEH

ICAR RESEARCH COMPLEX FOR NEH REGION
(Indian Council of Agricultural Research)
NAGALAND CENTER, JHARNAPANI, MEDZIPHEMA

Integrated Agromet Advisory Service Bulletin 10th to 14th Oct'15 Zunheboto District

Bulletin No:77/2015

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 14th Oct '15</i>
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Maximum and minimum temperatures ranged 27^oC to 29^oC and 14^oC to 16^oC, respectively. ▪ Relative humidity varied from 51% to 94%. ▪ Wind speed ranged from 1 to 2 kmph 			<p>Probability light rain will occur the coming week Max temp is expected to be around 25^oC - 28^oC and min temp is expected to be around 13^oC to 15^oC. Possibility of mainly cloudy sky the coming week Relative Humidity will range from 45% to 91% The wind speed 2 kmph Wind direction will blow mostly from easterly</p>
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Terrace/ Wetland paddy	Dough development stage		
Jhum paddy	Harvesting stage		<i>Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds</i>
Maize	Harvesting stage	-	<i>Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects</i>
Groundnut	Fruiting stage	-	<i>Proper earthing up should be done.</i>
Horticultural crops			
Citrus	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
Chilli	Harvesting stage	-	<i>Harvest fully ripe fruits. If its for storage, ensure proper drying.</i>
Ginger	Maturity stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>

	<i>Piggery</i>	Precaution should be taken against food and Mouth disease
	<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
 The logo of the Indian Council of Agricultural Research (ICAR) is located in the bottom-left corner of the table. It features a stylized green plant with three leaves inside a circular frame, with the text 'भारतीय' and 'ICAR' below it.	<i>Fishery</i>	Farmers are advised to keep the surrounding clean by cutting grasses and shrubs growing on embankment of the pond to remove shelter of predator of fish like snake, otter etc.



Integrated Agromet Advisory Service Bulletin 10th to 14th Oct'15

Dimapur District

Bulletin No:77/2015

Weather summary of the preceding week

- Light rain occurred the past week
- Maximum and minimum temperatures ranged 32^oC to 33^oC and 19^oC to 21^oC, respectively.
- Relative humidity varied from 48% to 95%.
- Wind speed ranged from 1 to 2 kmph

Weather forecast valid upto 14th Oct '15

- Probability of light rain the coming week.
- The **max temp** is likely to be 29^oC - 31^oC and the **min temp** 17^oC - 18^oC.
- **Sky is likely to be mainly clear sky the coming week**
- **Relative Humidity** is likely to range from 43% to 92%
- **Wind speed** 2-3 kmph
- **Wind direction** will be mostly easterly

Main Crops	Stage	Pest/ diseases	Agro-meteorological Advisories
------------	-------	-------------------	--------------------------------

Field crops

Lowland paddy	Dough stage		
Jhum paddy	Harvesting stage		<i>Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds</i>
Maize	Harvesting stage	-	<i>Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects</i>
Groundnut	Fruiting stage	-	<i>Proper earthing up should be done.</i>

Horticultural crops

Citrus	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
Chilli	Harvesting stage	-	<i>Harvest fully ripe fruits. If its for storage, ensure proper drying.</i>
Ginger	Maturity stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>
Piggery	Precaution should be taken against food and Mouth disease		

	<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
 <p>भारतीय ICAR</p>	<i>Fishery</i>	Farmers are advised to keep the surrounding clean by cutting grasses and shrubs growing on embankment of the pond to remove shelter of predator of fish like snake, otter etc.



Integrated Agromet Advisory Service Bulletin 10th to 14th Oct'15

Kiphire District

Bulletin No:77/2015

ICAR RESEARCH COMPLEX FOR NEH REGION
 (Indian Council of Agricultural Research)
 NAGALAND CENTER, JHARNAPANI, MEDZIPHEMA

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 14th Oct '15</i>
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Maximum and minimum temperatures ranged 26^oC to 28^oC and 13^oC to 15^oC, respectively. ▪ Relative humidity varied from 57% to 90%. ▪ Wind speed ranged from 1 to 2 kmph 			Probability of no rain the coming week Max temp is expected to be around 22 ^o C - 26 ^o C and min temp is expected to be around 10 ^o C to 12 ^o C. Possibility of mainly clear sky the coming week Relative Humidity will range from 42% to 95% The wind speed 2- 4 kmph Wind direction will blow mostly from south easterly
Main Crops	Stage	Pest/ diseases	Agro-meteorological Advisories
<i>Field crops</i>			
<i>Terrace/ Wetland paddy</i>	Dough development stage		
<i>Jhum paddy</i>	Harvesting stage		<i>Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds</i>
<i>Maize</i>	Harvesting stage	-	<i>Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects</i>
<i>Groundnut</i>	Fruiting stage	-	<i>Proper earthing up should be done.</i>
<i>Horticultural crops</i>			
<i>Citrus</i>	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
<i>Chilli</i>	Harvesting stage	-	<i>Harvest fully ripe fruits. If its for storage, ensure proper drying.</i>
<i>Ginger</i>	Maturity stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>
<i>Piggery</i>	Precaution should be taken against food and Mouth disease		
<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date		



Fishery

Farmers are advised to keep the surrounding clean by cutting grasses and shrubs growing on embankment of the pond to remove shelter of predator of fish like snake, otter etc.



Integrated Agromet Advisory Service Bulletin 10th to 14th Oct'15 Kohima District

Bulletin No:77/2015

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 14th Oct '15</i>
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Maximum and minimum temperatures ranged 25⁰C to 26⁰C and 14⁰C to 15⁰C, respectively. ▪ Relative humidity varied from 43% to 95%. ▪ Wind speed ranged from 1 to 2 kmph 			Probability of light rain the coming week Max temp is expected to be around 23⁰C - 25⁰C and min temp is expected to be around 13⁰C to 14⁰C. Possibility of mainly cloudy sky the coming week Relative Humidity will range from 48% to 94% Wind speed 2-3 kmph Wind direction will be mostly easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Field crops</i>			
<i>Terrace/ Wetland paddy</i>	Dough development stage		
<i>Jhum paddy</i>	Harvesting stage		<i>Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds</i>
<i>Maize</i>	Harvesting stage	-	<i>Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects</i>
<i>Groundnut</i>	Fruiting stage	-	<i>Proper earthing up should be done.</i>
<i>Horticultural crops</i>			
<i>Citrus</i>	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
<i>Chilli</i>	Harvesting stage	-	<i>Harvest fully ripe fruits. If its for storage, ensure proper drying.</i>
<i>Ginger</i>	Maturity stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>

	<i>Piggery</i>	Precaution should be taken against food and Mouth disease
	<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<i>Fishery</i>	Farmers are advised to keep the surrounding clean by cutting grasses and shrubs growing on embankment of the pond to remove shelter of predator of fish like snake, otter etc.