




Integrated Agromet Advisory Service Bulletin from 14th to 18th Oct'15 Mon District

Bulletin No:78/2015

Weather summary of the preceding week			Weather forecast valid upto 18 th Oct '15
<ul style="list-style-type: none"> No rain occurred the past week Maximum and minimum temperatures ranged 29⁰C to 30⁰C and 15⁰C to 16⁰C, respectively. Relative humidity varied from 54% to 90%. Wind speed ranged from 1 to 2 kmph 			<ul style="list-style-type: none"> Probability of no rain the coming week. The max temp is likely to be 26⁰C - 28⁰C and the min temp 11⁰C - 13⁰C. Sky is likely to be mainly clear the coming week Relative Humidity is likely to range from 47% to 90% Wind speed may reach upto 1- 2 kmph Wind direction will be mostly south southeasterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Terrace/ Wetland paddy	Dough development stage		
Jhum paddy	Harvesting stage		Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
Maize	Harvesting stage	-	Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects
Groundnut	Fruiting stage	-	Proper earthing up should be done.
Horticultural crops			
Citrus	-		Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).
Chilli	Harvesting stage	-	Harvest fully ripe fruits. If its for storage, ensure proper drying.
Ginger	Maturity stage		Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.
Piggery	Precaution should be taken against food and Mouth disease		

	<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<i>Fishery</i>	Farmers are advised to keep the surrounding clean by cutting grasses and shrubs growing on embankment of the pond to remove shelter of predator of fish like snake, otter etc.




Integrated Agromet Advisory Service Bulletin 14th to 18th Oct'15

Peren District

Bulletin No:78/2015

Weather summary of the preceding week			Weather forecast valid upto 18 th Oct '15
<ul style="list-style-type: none"> Light rain occurred the past week Maximum and minimum temperatures ranged 32⁰C to 34⁰C and 18⁰C to 19⁰C, respectively. Relative humidity varied from 44% to 99%. Wind speed ranged from 1 to 2 kmph 			Probability of light rain the coming week. Max temp is expected to be around 27⁰C - 30⁰C and min temp is expected to be around 14⁰C to 15⁰C. Possibility of mainly cloudy sky the coming week Relative Humidity will range from 43% to 95% The wind speed 2 kmph Wind direction will be easterly
Main Crops	Stage	Pest/diseases	Agro-meteorological Advisories
Field crops			
Lowland paddy	Dough stage		
Jhum paddy	Harvesting stage		As the grain is to be harvested, the store house should be fumigated; store bins cleaned and should be weather proof, rodent proof and damp proof.
Maize	Harvesting stage	-	For composite and hybrid, Do not wait for stalks and leaves to dry because mostly they remain green
Groundnut	Fruiting stage	-	Proper earthing up should be done.
Horticultural crops			
Citrus	-		Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).
Chilli	Harvesting stage	-	Avoid injuries during harvesting.
Ginger	Vegetative stage		Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.

	<i>Piggery</i>	Avoid stagnant water in and around the sty, maintain proper hygienic condition.
	<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<i>Fishery</i>	Chemicals found in or used on a pond's watershed can enter the pond and cause fish kills. Runoff following a heavy rain can transport a variety of unwanted chemicals into the pond including pesticides, herbicides, fertilizers, petroleum products. Avoid areas where the likelihood of chemical runoff is high and avoid using chemicals on the watershed




Integrated Agromet Advisory Service Bulletin 14th to 18th Oct'15

Phek District

Bulletin No:78/2015

CAR RESEARCH COMPLEX FOR NEH REGION
(Indian Council of Agricultural Research)
NAGALAND CENTER, JHARNAPANI, MEDZIPHEMA

Weather summary of the preceding week			Weather forecast valid upto 18 th Oct '15
<ul style="list-style-type: none"> No rain occurred the past week Maximum and minimum temperatures ranged 27⁰C to 29⁰C and 14⁰C to 16⁰C, respectively. Relative humidity varied from 51% to 98%. Wind speed ranged from 1 to 2 kmph 			<p>Probability there will be light rain the coming week</p> <p>Max temp is expected to be around 25⁰C - 27⁰C and min temp is expected to be around 11⁰C to 14⁰C.</p> <p>Possibility of mainly clear sky the coming week</p> <p>Relative Humidity will range from 49% to 95%</p> <p>The wind speed 1-2 kmph</p> <p>Wind direction will blow mostly from easterly</p>
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Terrace/ Wetland paddy	Dough development stage		
Jhum paddy	Harvesting stage		Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
Maize	Harvesting stage	-	Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects
Groundnut	Fruiting stage	-	Proper earthing up should be done.
Horticultural crops			
Citrus	-		Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).
Chilli	Harvesting stage	-	Harvest fully ripe fruits. If its for storage, ensure proper drying.
Ginger	Maturity stage		Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.

	<i>Piggery</i>	Precaution should be taken against food and Mouth disease
	<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<i>Fishery</i>	Farmers are advised to keep the surrounding clean by cutting grasses and shrubs growing on embankment of the pond to remove shelter of predator of fish like snake, otter etc.




Integrated Agromet Advisory Service Bulletin 14th to 18th Oct'15

Tuensang District

Bulletin No:78/2015

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 18th Oct '15</i>
<ul style="list-style-type: none"> Light rain occurred the past week Maximum and minimum temperatures ranged 24⁰C to 25⁰C and 9⁰C to 10⁰C, respectively. Relative humidity varied from 56% to 95%. Wind speed ranged from 1 to 2 kmph 			Probability there will be light rain the coming week Max temp is expected to be around 21⁰C - 24⁰C and min temp is expected to be around 5⁰C to 11⁰C. Possibility of partly cloudy sky the coming week Relative humidity will range from 47% to 98% Wind speed 1-2 kmph Wind direction will be mostly easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Terrace/ Wetland paddy	Dough development stage		
Jhum paddy	Harvesting stage		Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
Maize	Harvesting stage	-	Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects
Groundnut	Fruiting stage	-	Proper earthing up should be done.
Horticultural crops			
Citrus	-		Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).
Chilli	Harvesting stage	-	Harvest fully ripe fruits. If its for storage, ensure proper drying.
Ginger	Maturity stage		Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.

	<i>Piggery</i>	Precaution should be taken against food and Mouth disease
	<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<i>Fishery</i>	Farmers are advised to keep the surrounding clean by cutting grasses and shrubs growing on embankment of the pond to remove shelter of predator of fish like snake, otter etc.



Integrated Agromet Advisory Service Bulletin 14th to 18th Oct'15 Wokha District

Bulletin No:78/2015

Weather summary of the preceding week

- No rain occurred the past week
- Maximum and minimum temperatures ranged 28⁰C to 31⁰C and 17⁰C to 19⁰C, respectively.
- Relative humidity varied from 51% to 95%.
- Wind speed ranged from 1 to 2 kmph

Weather forecast valid upto 18th Oct '15

Probability light rain will occur the coming week
Max temp is expected to be around 27⁰C - 29⁰C and min temp is expected to be around 14⁰C to 15⁰C.
Possibility of mainly clear sky the coming week
Relative Humidity will range from 46% to 95%
The wind speed 2- 3 kmph
 Wind direction will blow mostly from easterly

Main Crops

Stage

**Pest/
Diseases**

Agro-meteorological Advisories

Field crops

Terrace/ Wetland paddy	Dough development stage		
Jhum paddy	Harvesting stage		<i>Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds</i>
Maize	Harvesting stage	-	<i>Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects</i>
Groundnut	Fruiting stage	-	<i>Proper earthing up should be done.</i>


Horticultural crops

Citrus	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
Chilli	Harvesting stage	-	<i>Harvest fully ripe fruits. If its for storage, ensure proper drying.</i>
Ginger	Maturity stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>

ICAR RESEARCH COMPLEX FOR NEH REGION

(Indian Council of Agricultural Research)

NAGALAND CENTER, JHARNAPANI, MEDZIPHEMA

	<i>Piggery</i>	Precaution should be taken against food and Mouth disease
	<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<i>Fishery</i>	Farmers are advised to keep the surrounding clean by cutting grasses and shrubs growing on embankment of the pond to remove shelter of predator of fish like snake, otter etc.



--	--




Integrated Agromet Advisory Service Bulletin 14th to 18th Oct'15

Zunheboto District

Bulletin No:78/2015

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 18th Oct '15</i>
<ul style="list-style-type: none"> No rain occurred the past week Maximum and minimum temperatures ranged 27⁰C to 29⁰C and 14⁰C to 16⁰C, respectively. Relative humidity varied from 51% to 94%. Wind speed ranged from 1 to 2 kmph 			Probability light rain will occur the coming week Max temp is expected to be around 25⁰C - 28⁰C and min temp is expected to be around 11⁰C to 12⁰C. Possibility of mainly cloudy sky the coming week Relative Humidity will range from 45% to 91% The wind speed 2 kmph Wind direction will blow mostly from easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Terrace/ Wetland paddy	Dough development stage		
Jhum paddy	Harvesting stage		<i>Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds</i>
Maize	Harvesting stage	-	<i>Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects</i>
Groundnut	Fruiting stage	-	<i>Proper earthing up should be done.</i>
Horticultural crops			
Citrus	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
Chilli	Harvesting stage	-	<i>Harvest fully ripe fruits. If its for storage, ensure proper drying.</i>
Ginger	Maturity stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>


	<i>Piggery</i>	Precaution should be taken against food and Mouth disease
	<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<i>Fishery</i>	Farmers are advised to keep the surrounding clean by cutting grasses and shrubs growing on embankment of the pond to remove shelter of predator of fish like snake, otter etc.



Integrated Agromet Advisory Service Bulletin 14th to 18th Oct'15 Dimapur District

Bulletin No:78/2015

Weather summary of the preceding week			Weather forecast valid upto 18 th Oct '15
<ul style="list-style-type: none"> Light rain occurred the past week Maximum and minimum temperatures ranged 32^oC to 33^oC and 19^oC to 21^oC, respectively. Relative humidity varied from 48% to 95%. Wind speed ranged from 1 to 2 kmph 			<ul style="list-style-type: none"> Probability of light rain the coming week. The max temp is likely to be 29^oC - 30^oC and the min temp 16^oC - 18^oC. Sky is likely to be mainly clear sky the coming week Relative Humidity is likely to range from 48% to 95% Wind speed 2-3 kmph Wind direction will be mostly easterly
Main Crops	Stage	Pest/ diseases	Agro-meteorological Advisories
Field crops			
Lowland paddy	Dough stage		
Jhum paddy	Harvesting stage		Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
Maize	Harvesting stage	-	Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects
Groundnut	Fruiting stage	-	Proper earthing up should be done.
Horticultural crops			
Citrus	-		Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).
Chilli	Harvesting stage	-	Harvest fully ripe fruits. If its for storage, ensure proper drying.
Ginger	Maturity stage		Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.
Piggery	Precaution should be taken against food and Mouth disease		

	<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<i>Fishery</i>	Farmers are advised to keep the surrounding clean by cutting grasses and shrubs growing on embankment of the pond to remove shelter of predator of fish like snake, otter etc.



Integrated Agromet Advisory Service Bulletin 14th to 18th Oct'15 Kiphire District

Bulletin No:78/2015

Weather summary of the preceding week			Weather forecast valid upto 18 th Oct '15
<ul style="list-style-type: none"> No rain occurred the past week Maximum and minimum temperatures ranged 26⁰C to 28⁰C and 13⁰C to 15⁰C, respectively. Relative humidity varied from 57% to 90%. Wind speed ranged from 1 to 2 kmph 			Probability of light rain the coming week Max temp is expected to be around 23 ⁰ C - 25 ⁰ C and min temp is expected to be around 8 ⁰ C to 12 ⁰ C. Possibility of mainly clear sky the coming week Relative Humidity will range from 47% to 95% The wind speed 2- 4 kmph Wind direction will blow mostly from south easterly
Main Crops	Stage	Pest/diseases	Agro-meteorological Advisories
Field crops			
Terrace/ Wetland paddy	Dough development stage		
Jhum paddy	Harvesting stage		Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
Maize	Harvesting stage	-	Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects
Groundnut	Fruiting stage	-	Proper earthing up should be done.
Horticultural crops			
Citrus	-		Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).
Chilli	Harvesting stage	-	Harvest fully ripe fruits. If its for storage, ensure proper drying.
Ginger	Maturity stage		Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.
Piggery	Precaution should be taken against food and Mouth disease		
Poultry	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date		



Fishery


Farmers are advised to keep the surrounding clean by cutting grasses and shrubs growing on embankment of the pond to remove shelter of predator of fish like snake, otter etc.



Integrated Agromet Advisory Service Bulletin 14th to 18th Oct'15 Kohima District

Bulletin No:78/2015

Weather summary of the preceding week			Weather forecast valid upto 18 th Oct '15
<ul style="list-style-type: none"> No rain occurred the past week Maximum and minimum temperatures ranged 25⁰C to 26⁰C and 14⁰C to 15⁰C, respectively. Relative humidity varied from 43% to 95%. Wind speed ranged from 1 to 2 kmph 			Probability of light rain the coming week Max temp is expected to be around 24 ⁰ C - 25 ⁰ C and min temp is expected to be around 13 ⁰ C to 14 ⁰ C. Possibility of mainly clear sky the coming week Relative Humidity will range from 42% to 94% Wind speed 2-3 kmph Wind direction will be mostly easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
<i>Terrace/ Wetland paddy</i>	Dough development stage		
<i>Jhum paddy</i>	Harvesting stage		Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
<i>Maize</i>	Harvesting stage	-	Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects
<i>Groundnut</i>	Fruiting stage	-	Proper earthing up should be done.
Horticultural crops			
<i>Citrus</i>	-		Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).
<i>Chilli</i>	Harvesting stage	-	Harvest fully ripe fruits. If its for storage, ensure proper drying.
<i>Ginger</i>	Maturity stage		Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.


	<i>Piggery</i>	Precaution should be taken against food and Mouth disease
	<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<i>Fishery</i>	Farmers are advised to keep the surrounding clean by cutting grasses and shrubs growing on embankment of the pond to remove shelter of predator of fish like snake, otter etc.



Integrated Agromet Advisory Service Bulletin 14th to 18th Oct'15 Longleng District

Bulletin No:78/2015

Weather summary of the preceding week			Weather forecast valid upto 18 th Oct '15
<ul style="list-style-type: none"> No rain occurred the past week Maximum and minimum temperatures ranged 30⁰C to 32⁰C and 17⁰C to 20⁰C, respectively. Relative humidity varied from 73% to 95%. Wind speed ranged from 1 to 2 kmph 			Probability of no rain the coming week Max temp is expected to be around 27⁰C-28⁰C and min temp is expected to be around 12⁰C to 13⁰C. Possibility of mainly clear sky the coming week Relative Humidity will range from 47% to 91% The wind speed 1 - 2 kmph Wind direction will be mostly easterly
Main Crops	Stage	Pest/ diseases	Agro-meteorological Advisories
Field crops			
Terrace/ Wetland paddy	Dough development stage		
Jhum paddy	Harvesting stage		Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
Maize	Harvesting stage	-	Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects
Groundnut	Fruiting stage	-	Proper earthing up should be done.
Horticultural crops			
Citrus	-		Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).
Chilli	Harvesting stage	-	Harvest fully ripe fruits. If its for storage, ensure proper drying.
Ginger	Maturity stage		Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.

	<i>Piggery</i>	Precaution should be taken against food and Mouth disease
	<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<i>Fishery</i>	Farmers are advised to keep the surrounding clean by cutting grasses and shrubs growing on embankment of the pond to remove shelter of predator of fish like snake, otter etc.

Integrated Agromet Advisory Service Bulletin 14th to 18th Oct'15

Mokokchung District

Bulletin No: 78/2015

Weather summary of the preceding week			Weather forecast valid upto 18 th Oct '15
<ul style="list-style-type: none"> No rain occurred the past week Maximum and minimum temperatures ranged 29⁰C to 30⁰C and 16⁰C to 17⁰C, respectively. Relative humidity varied from 51% to 90%. Wind speed ranged from 1 to 2 kmph 			Probability no rain the coming week Max temp is expected to be around 27 ⁰ C - 29 ⁰ C and min temp is expected to be around 13 ⁰ C to 14 ⁰ C. Possibility of mainly clear sky the coming week Relative Humidity will range from 47% to 95% and Wind speed 1-2 kmph Wind direction will be mostly easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Terrace/ Wetland paddy	Dough development stage		
Jhum paddy	Harvesting stage		Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
Maize	Harvesting stage	-	Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects
Groundnut	Fruiting stage	-	Proper earthing up should be done.
Horticultural crops			
Citrus	-		Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).
Chilli	Harvesting stage	-	Harvest fully ripe fruits. If its for storage, ensure proper drying.
Ginger	Maturity stage		Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.
Piggery	Precaution should be taken against food and Mouth disease		

<i>Poultry</i>	In case of diarrhoea in poultry , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
<i>Fishery</i>	Farmers are advised to keep the surrounding clean by cutting grasses and shrubs growing on embankment of the pond to remove shelter of predator of fish like snake, otter etc.

