


Integrated Agromet Advisory Service Bulletin 24th to 28th Oct'15

Mokokchung District

Bulletin No: 81/2015

Weather summary of the preceding week			Weather forecast valid upto 28 th Oct '15
<ul style="list-style-type: none"> No rain occurred the past week Maximum and minimum temperatures ranged 29^oC to 30^oC and 16^oC to 17^oC, respectively. Relative humidity varied from 51% to 90%. Wind speed ranged from 1 to 2 kmph 			<p>Probability no rain the coming week</p> <p>Max temp is expected to be around 27^oC - 28^oC and min temp is expected to be around 14^oC to 15^oC.</p> <p>Possibility of clear sky the coming week</p> <p>Relative Humidity will range from 40% to 97% and</p> <p>Wind speed 2 kmph</p> <p>Wind direction will be easterly</p>
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<p>Field crops</p> <p>Field preparation should be started for rabi crops like toria, rabi maize</p>			
Terrace/ Wetland paddy	Dough development stage		
Jhum paddy	Harvesting stage		<p>Avoid excessive drying of paddy to avoid breakage of the grains.</p> <p>Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds</p>
Maize	Harvesting stage	-	Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects
Groundnut	Harvesting stage	-	The weather is favorable for harvesting of groundnut. Proper handling should be done while harvesting and storage too.
<p>Horticultural crops</p> <p>Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done. Spread well decomposed cowdung 5 to 6 tones per hectare for the main field and mix it well in the soil by cultivating the field. Keep the field clean by collecting the stubbles and unwanted materials</p>			
Citrus	-		Spray Malathion (2ml/l) over fruit to prevent fruit fly egg laying. If possible apply 650g urea + 1220g SSP + 450g MOP/ tree.
Chilli	Harvesting stage	-	Harvest fully ripe fruits. If its for storage, ensure proper drying.
Ginger	Maturity stage		Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of


				<i>rhizomes.</i>
	<i>Piggery</i>	Proper sanitary measures should be done to control the diseases. Slaughtering of all apparently ailing birds, segregating of in - contact in group of 10 to 15 each; removal of all infective materials such as droppings, residues of poultry cleanliness.		
	<i>Poultry</i>	Farmers are advised to administer proper vaccination schedule to protect their cattle from viral and bacterial diseases and should contact the nearest concerned authorities.		
	<i>Fishery</i>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media		



Integrated Agromet Advisory Service Bulletin from 24th to 28th Oct'15 Mon District

Bulletin No:81/2015

Weather summary of the preceding week			Weather forecast valid upto 28 th Oct '15
<ul style="list-style-type: none"> No rain occurred the past week Maximum and minimum temperatures ranged 29^oC to 30^oC and 15^oC to 16^oC, respectively. Relative humidity varied from 54% to 90%. Wind speed ranged from 1 to 2 kmph 			<ul style="list-style-type: none"> Probability of no rain the coming week. The max temp is likely to be 27^oC - 28^oC and the min temp 14^oC - 15^oC. Sky is likely to be clear the coming week Relative Humidity is likely to range from 41% to 97% Wind speed may reach upto 0- 2 kmph Wind direction will be easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops Field preparation should be started for rabi crops like toria, rabi maize			
Terrace/ Wetland paddy	Dough development stage		
Jhum paddy	Harvesting stage		Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
Maize	Harvesting stage	-	Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects
Groundnut	Harvesting stage	-	The weather is favorable for harvesting of groundnut. Proper handling should be done while harvesting and storage too.
Horticultural crops Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done. Spread well decomposed cowdung 5 to 6 tones per hectare for the main field and mix it well in the soil by cultivating the field. Keep the field clean by collecting the stubbles and unwanted materials			
Citrus	-		Spray Malathion (2ml/l) over fruit to prevent fruit fly egg laying. If possible apply 650g urea + 1220g SSP + 450g MOP/ tree.
Chilli	Harvesting stage	-	Harvest fully ripe fruits. If its for storage, ensure proper drying.

	<i>Ginger</i>	Maturity stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>
	<i>Poultry</i>	Proper sanitary measures should be done to control the diseases. Slaughtering of all apparently ailing birds, segregating of in - contact in group of 10 to 15 each; removal of all infective materials such as droppings, residues of poultry cleanliness.		
	<i>Piggery</i>	Farmers are advised to administer proper vaccination schedule to protect their cattle from viral and bacterial diseases and should contact the nearest concerned authorities.		
	<i>Fishery</i>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media		




Integrated Agromet Advisory Service Bulletin 24th to 28th Oct'15

Peren District

Bulletin No:81/2015

Weather summary of the preceding week			Weather forecast valid upto 28 th Oct '15
<ul style="list-style-type: none"> Light rain occurred the past week Maximum and minimum temperatures ranged 32⁰C to 34⁰C and 18⁰C to 19⁰C, respectively. Relative humidity varied from 44% to 99%. Wind speed ranged from 1 to 2 kmph 			Probability of no rain the coming week. Max temp is expected to be around 28⁰C - 29⁰C and min temp is expected to be around 13⁰C to 14⁰C. Possibility of clear sky the coming week Relative Humidity will range from 40% to 96% The wind speed 4 kmph Wind direction will be mostly northeasterly
Main Crops	Stage	Pest/diseases	Agro-meteorological Advisories
Field crops			
Field preparation should be started for rabi crops like toria, rabi maize			
Lowland paddy	Dough development stage		
Jhum paddy	Harvesting stage		Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
Maize	Harvesting stage	-	Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects
Groundnut	Harvesting stage	-	The weather is favorable for harvesting of groundnut. Proper handling should be done while harvesting and storage too.
Horticultural crops			
Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done. Spread well decomposed cowdung 5 to 6 tones per hectare for the main field and mix it well in the soil by cultivating the field. Keep the field clean by collecting the stubbles and unwanted materials			
Citrus	-		Spray Malathion (2ml/l) over fruit to prevent fruit fly egg laying. If possible apply 650g urea + 1220g SSP + 450g MOP/ tree.
Chilli	Harvesting stage	-	Harvest fully ripe fruits. If its for storage, ensure proper drying.

	<i>Ginger</i>	Maturity stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>
	<i>Piggery</i>	Proper sanitary measures should be done to control the diseases. Slaughtering of all apparently ailing birds, segregating of in - contact in group of 10 to 15 each; removal of all infective materials such as droppings, residues of poultry cleanliness.		
	<i>Poultry</i>	Farmers are advised to administer proper vaccination schedule to protect their cattle from viral and bacterial diseases and should contact the nearest concerned authorities.		
	<i>Fishery</i>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media		



Integrated Agromet Advisory Service Bulletin 24th to 28th Oct'15

Phek District

Bulletin No:81/2015

Weather summary of the preceding week

- No rain occurred the past week
- Maximum and minimum temperatures ranged 27⁰C to 29⁰C and 14⁰C to 16⁰C, respectively.
- Relative humidity varied from 51% to 98%.
- Wind speed ranged from 1 to 2 kmph

Weather forecast valid upto 28th Oct '15

Probability there will be no rain the coming week
Max temp is expected to be around 26⁰C - 27⁰C and min temp is expected to be around 13⁰C to 14⁰C.
Possibility of clear sky the coming week
Relative Humidity will range from 39% to 97%
The wind speed 2 kmph
Wind direction will blow mostly from easterly

Main Crops

Stage

Pest/ Diseases

Agro-meteorological Advisories

Field crops

Field preparation should be started for rabi crops like toria, rabi maize

Terrace/ Wetland paddy	Dough development stage		
Jhum paddy	Harvesting stage		<i>Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds</i>
Maize	Harvesting stage	-	<i>Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects</i>
Groundnut	Harvesting stage	-	<i>The weather is favorable for harvesting of groundnut. Proper handling should be done while harvesting and storage too.</i>

Horticultural crops


Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done. Spread well decomposed cowdung 5 to 6 tones per hectare for the main field and mix it well in the soil by cultivating the field. Keep the field clean by collecting the stubbles and unwanted materials

Citrus	-		<i>Spray Malathion (2ml/l) over fruit to prevent fruit fly egg laying. If possible apply 650g urea + 1220g SSP + 450g MOP/ tree.</i>
---------------	---	--	--

CAR RESEARCH COMPLEX FOR NEH REGION

(Indian Council of Agricultural Research)

NAGALAND CENTER, JHARNAPANI, MEDZIPHEMA

	Chilli	Harvesting stage	-	<i>Harvest fully ripe fruits. If its for storage, ensure proper drying.</i>
	Ginger	Maturity stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>
	Piggery	Proper sanitary measures should be done to control the diseases. Slaughtering of all apparently ailing birds, segregating of in - contact in group of 10 to 15 each; removal of all infective materials such as droppings, residues of poultry cleanliness.		
	Poultry	Farmers are advised to administer proper vaccination schedule to protect their cattle from viral and bacterial diseases and should contact the nearest concerned authorities.		
	Fishery	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media		




Integrated Agromet Advisory Service Bulletin 24th to 28th Oct'15

Tuensang District

Bulletin No:81/2015

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 28th Oct '15</i>
<ul style="list-style-type: none"> Light rain occurred the past week Maximum and minimum temperatures ranged 24⁰C to 25⁰C and 9⁰C to 10⁰C, respectively. Relative humidity varied from 56% to 95%. Wind speed ranged from 1 to 2 kmph 			Probability there will be no rain the coming week Max temp is expected to be around 23⁰C - 24⁰C and min temp is expected to be around 9⁰C to 10⁰C. Possibility of clear sky the coming week Relative humidity will range from 35% to 98% Wind speed 2 kmph Wind direction will be mostly easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops Field preparation should be started for rabi crops like toria, rabi maize			
Terrace/ Wetland paddy	Dough development stage		
Jhum paddy	Harvesting stage		<i>Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds</i>
Maize	Harvesting stage	-	<i>Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects</i>
Groundnut	Harvesting stage	-	<i>The weather is favorable for harvesting of groundnut. Proper handling should be done while harvesting and storage too.</i>
Horticultural crops Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done. Spread well decomposed cowdung 5 to 6 tones per hectare for the main field and mix it well in the soil by cultivating the field. Keep the field clean by collecting the stubbles and unwanted materials			
Citrus	-		<i>Spray Malathion (2ml/l) over fruit to prevent fruit fly egg laying. If possible apply 650g urea + 1220g SSP + 450g MOP/ tree.</i>
Chilli	Harvesting stage	-	<i>Harvest fully ripe fruits. If its for storage, ensure proper drying.</i>
Ginger	Maturity stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to</i>


				<i>break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>
	<i>Piggery</i>	Proper sanitary measures should be done to control the diseases. Slaughtering of all apparently ailing birds, segregating of in - contact in group of 10 to 15 each; removal of all infective materials such as droppings, residues of poultry cleanliness.		
	<i>Poultry</i>	Farmers are advised to administer proper vaccination schedule to protect their cattle from viral and bacterial diseases and should contact the nearest concerned authorities.		
	<i>Fishery</i>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media		



Integrated Agromet Advisory Service Bulletin 24th to 28th Oct'15 Wokha District

Bulletin No:81/2015

Weather summary of the preceding week			Weather forecast valid upto 28 th Oct '15
<ul style="list-style-type: none"> No rain occurred the past week Maximum and minimum temperatures ranged 28⁰C to 31⁰C and 17⁰C to 19⁰C, respectively. Relative humidity varied from 51% to 95%. Wind speed ranged from 1 to 2 kmph 			<p>Probability no rain will occur the coming week Max temp is expected to be around 27⁰C - 28⁰C and min temp is expected to be around 13⁰C to 14⁰C. Possibility of clear sky the coming week Relative Humidity will range from 40% to 96% The wind speed 2- 4 kmph Wind direction will blow from easterly</p>
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops Field preparation should be started for rabi crops like toria, rabi maize			
Terrace/ Wetland paddy	Dough development stage		
Jhum paddy	Harvesting stage		<i>Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds</i>
Maize	Harvesting stage	-	<i>Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects</i>
Groundnut	Harvesting stage	-	<i>The weather is favorable for harvesting of groundnut. Proper handling should be done while harvesting and storage too.</i>
Horticultural crops Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done. Spread well decomposed cowdung 5 to 6 tones per hectare for the main field and mix it well in the soil by cultivating the field. Keep the field clean by collecting the stubbles and unwanted materials			
Citrus	-		<i>Spray Malathion (2ml/l) over fruit to prevent fruit fly egg laying. If possible apply 650g urea + 1220g SSP + 450g MOP/ tree.</i>
Chilli	Harvesting stage	-	<i>Harvest fully ripe fruits. If its for storage, ensure proper drying.</i>

	<i>Ginger</i>	Maturity stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>
	<i>Piggery</i>	Proper sanitary measures should be done to control the diseases. Slaughtering of all apparently ailing birds, segregating of in - contact in group of 10 to 15 each; removal of all infective materials such as droppings, residues of poultry cleanliness.		
	<i>Poultry</i>	Farmers are advised to administer proper vaccination schedule to protect their cattle from viral and bacterial diseases and should contact the nearest concerned authorities.		
	<i>Fishery</i>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media		




--	--



Integrated Agromet Advisory Service Bulletin 24th to 28th Oct'15 Zunheboto District

Bulletin No:81/2015

Weather summary of the preceding week			Weather forecast valid upto 28 th Oct '15
<ul style="list-style-type: none"> No rain occurred the past week Maximum and minimum temperatures ranged 27⁰C to 29⁰C and 14⁰C to 16⁰C, respectively. Relative humidity varied from 51% to 94%. Wind speed ranged from 1 to 2 kmph 			Probability no rain the coming week Max temp is expected to be around 26 ⁰ C - 27 ⁰ C and min temp is expected to be around 12 ⁰ C to 13 ⁰ C. Possibility of clear sky the coming week Relative Humidity will range from 39% to 97% The wind speed 2 kmph Wind direction will blow from easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops Field preparation should be started for rabi crops like toria, rabi maize			
Terrace/ Wetland paddy	Dough development stage		
Jhum paddy	Harvesting stage		Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
Maize	Harvesting stage	-	Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects
Groundnut	Harvesting stage	-	The weather is favorable for harvesting of groundnut. Proper handling should be done while harvesting and storage too.
Horticultural crops Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done. Spread well decomposed cowdung 5 to 6 tones per hectare for the main field and mix it well in the soil by cultivating the field. Keep the field clean by collecting the stubbles and unwanted materials			
Citrus	-		Spray Malathion (2ml/l) over fruit to prevent fruit fly egg laying. If possible apply 650g urea + 1220g SSP + 450g MOP/ tree.
Chilli	Harvesting stage	-	Harvest fully ripe fruits. If its for storage, ensure proper drying.


	<i>Ginger</i>	Maturity stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>
	<i>Piggery</i>	Proper sanitary measures should be done to control the diseases. Slaughtering of all apparently ailing birds, segregating of in - contact in group of 10 to 15 each; removal of all infective materials such as droppings, residues of poultry cleanliness.		
	<i>Poultry</i>	Farmers are advised to administer proper vaccination schedule to protect their cattle from viral and bacterial diseases and should contact the nearest concerned authorities.		
	<i>Fishery</i>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media		



Integrated Agromet Advisory Service Bulletin 24th to 28th Oct'15 Dimapur District

Bulletin No:81/2015

Weather summary of the preceding week			Weather forecast valid upto 28 th Oct '15
<ul style="list-style-type: none"> Light rain occurred the past week Maximum and minimum temperatures ranged 32⁰C to 33⁰C and 19⁰C to 21⁰C, respectively. Relative humidity varied from 48% to 95%. Wind speed ranged from 1 to 2 kmph 			<ul style="list-style-type: none"> Probability of no rain the coming week. The max temp is likely to be 29⁰C - 30⁰C and the min temp 19⁰C - 20⁰C. Sky is likely to be clear the coming week Relative Humidity is likely to range from 40% to 96% Wind speed 2-3 kmph Wind direction will be mostly easterly
Main Crops	Stage	Pest/ diseases	Agro-meteorological Advisories
Field crops Field preparation should be started for rabi crops like toria, rabi maize			
Lowland paddy	Dough development stage		
Jhum paddy	Harvesting stage		Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
Maize	Harvesting stage	-	Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects
Groundnut	Harvesting stage	-	The weather is favorable for harvesting of groundnut. Proper handling should be done while harvesting and storage too.
Horticultural crops Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done. Spread well decomposed cowdung 5 to 6 tones per hectare for the main field and mix it well in the soil by cultivating the field. Keep the field clean by collecting the stubbles and unwanted materials			
Citrus	-		Spray Malathion (2ml/l) over fruit to prevent fruit fly egg laying. If possible apply 650g urea + 1220g SSP + 450g MOP/ tree.
Chilli	Harvesting stage	-	Harvest fully ripe fruits. If its for storage, ensure proper drying.
Ginger	Maturity stage		Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the

				<i>rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>
	<i>Piggery</i>	Proper sanitary measures should be done to control the diseases. Slaughtering of all apparently ailing birds, segregating of in - contact in group of 10 to 15 each; removal of all infective materials such as droppings, residues of poultry cleanliness.		
	<i>Poultry</i>	Farmers are advised to administer proper vaccination schedule to protect their cattle from viral and bacterial diseases and should contact the nearest concerned authorities.		
 भारतीय ICAR	<i>Fishery</i>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media		




Integrated Agromet Advisory Service Bulletin 24th to 28th Oct'15

Kiphire District

Bulletin No:81/2015

Weather summary of the preceding week			Weather forecast valid upto 28 th Oct '15
<ul style="list-style-type: none"> No rain occurred the past week Maximum and minimum temperatures ranged 26⁰C to 28⁰C and 13⁰C to 15⁰C, respectively. Relative humidity varied from 57% to 90%. Wind speed ranged from 1 to 2 kmph 			Probability of no rain the coming week Max temp is expected to be around 24 ⁰ C - 25 ⁰ C and min temp is expected to be around 14 ⁰ C to 15 ⁰ C. Possibility of clear sky the coming week Relative Humidity will range from 37% to 96% The wind speed 2 kmph Wind direction will blow mostly from easterly
Main Crops	Stage	Pest/diseases	Agro-meteorological Advisories
Field crops			
Field preparation should be started for rabi crops like toria, rabi maize			
Terrace/Wetland paddy	Dough development stage		
Jhum paddy	Harvesting stage		Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
Maize	Harvesting stage	-	Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects
Groundnut	Harvesting stage	-	The weather is favorable for harvesting of groundnut. Proper handling should be done while harvesting and storage too.
Horticultural crops			
Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done. Spread well decomposed cowdung 5 to 6 tones per hectare for the main field and mix it well in the soil by cultivating the field. Keep the field clean by collecting the stubbles and unwanted materials			
Citrus	-		Spray Malathion (2ml/l) over fruit to prevent fruit fly egg laying. If possible apply 650g urea + 1220g SSP + 450g MOP/ tree.
Chilli	Harvesting stage	-	Harvest fully ripe fruits. If its for storage, ensure proper drying.
Ginger	Maturity stage		Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.


	<i>Piggery</i>	Proper sanitary measures should be done to control the diseases. Slaughtering of all apparently ailing birds, segregating of in - contact in group of 10 to 15 each; removal of all infective materials such as droppings, residues of poultry cleanliness.
	<i>Poultry</i>	Farmers are advised to administer proper vaccination schedule to protect their cattle from viral and bacterial diseases and should contact the nearest concerned authorities.
	<i>Fishery</i>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media



Integrated Agromet Advisory Service Bulletin 24th to 28th Oct'15 Kohima District

Bulletin No:81/2015

Weather summary of the preceding week			Weather forecast valid upto 28 th Oct '15
<ul style="list-style-type: none"> No rain occurred the past week Maximum and minimum temperatures ranged 25⁰C to 26⁰C and 14⁰C to 15⁰C, respectively. Relative humidity varied from 43% to 95%. Wind speed ranged from 1 to 2 kmph 			Probability of no rain the coming week Max temp is expected to be around 27 ⁰ C - 28 ⁰ C and min temp is expected to be around 11 ⁰ C to 12 ⁰ C. Possibility of clear sky the coming week Relative Humidity will range from 36% to 96% Wind speed 4 kmph Wind direction will be mostly northeasterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops			
Field preparation should be started for rabi crops like toria, rabi maize			
Terrace/ Wetland paddy	Dough development stage		
Jhum paddy	Harvesting stage		Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
Maize	Harvesting stage	-	Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects
Groundnut	Harvesting stage	-	The weather is favorable for harvesting of groundnut. Proper handling should be done while harvesting and storage too.
Horticultural crops			
Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done. Spread well decomposed cowdung 5 to 6 tones per hectare for the main field and mix it well in the soil by cultivating the field. Keep the field clean by collecting the stubbles and unwanted materials			
Citrus	-		Spray Malathion (2ml/l) over fruit to prevent fruit fly egg laying. If possible apply 650g urea + 1220g SSP + 450g MOP/ tree.
Chilli	Harvesting stage	-	Harvest fully ripe fruits. If its for storage, ensure proper drying.


	<i>Ginger</i>	Maturity stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>
	<i>Piggery</i>	Proper sanitary measures should be done to control the diseases. Slaughtering of all apparently ailing birds, segregating of in - contact in group of 10 to 15 each; removal of all infective materials such as droppings, residues of poultry cleanliness.		
	<i>Poultry</i>	Farmers are advised to administer proper vaccination schedule to protect their cattle from viral and bacterial diseases and should contact the nearest concerned authorities.		
	<i>Fishery</i>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media		



Integrated Agromet Advisory Service Bulletin 24th to 28th Oct'15 Longleng District

Bulletin No:81/2015

Weather summary of the preceding week			Weather forecast valid upto 28 th Oct '15
<ul style="list-style-type: none"> No rain occurred the past week Maximum and minimum temperatures ranged 30⁰C to 32⁰C and 17⁰C to 20⁰C, respectively. Relative humidity varied from 73% to 95%. Wind speed ranged from 1 to 2 kmph 			Probability of no rain the coming week Max temp is expected to be around 27⁰C-28⁰C and min temp is expected to be around 13⁰C to 14⁰C. Possibility of clear sky the coming week Relative Humidity will range from 39% to 97% The wind speed 0 - 2 kmph Wind direction will be easterly
Main Crops	Stage	Pest/diseases	Agro-meteorological Advisories
Field crops Field preparation should be started for rabi crops like toria, rabi maize			
Terrace/Wetland paddy	Dough development stage		
Jhum paddy	Harvesting stage		Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
Maize	Harvesting stage	-	Harvesting should not be delayed as it leads to many problems. It is exposed to attack by termites, rodents and domestic animals and can be infested by insects
Groundnut	Harvesting stage	-	The weather is favorable for harvesting of groundnut. Proper handling should be done while harvesting and storage too.
Horticultural crops Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done. Spread well decomposed cowdung 5 to 6 tones per hectare for the main field and mix it well in the soil by cultivating the field. Keep the field clean by collecting the stubbles and unwanted materials			
Citrus	-		Spray Malathion (2ml/l) over fruit to prevent fruit fly egg laying. If possible apply 650g urea + 1220g SSP + 450g MOP/ tree.
Chilli	Harvesting stage	-	Harvest fully ripe fruits. If its for storage, ensure proper drying.
Ginger	Maturity stage		Soil around the plants should be worked with the help of khurpi . It

				<i>helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>
	Piggery	Proper sanitary measures should be done to control the diseases. Slaughtering of all apparently ailing birds, segregating of in - contact in group of 10 to 15 each; removal of all infective materials such as droppings, residues of poultry cleanliness.		
	Poultry	Farmers are advised to administer proper vaccination schedule to protect their cattle from viral and bacterial diseases and should contact the nearest concerned authorities.		
	Fishery	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media		