




# Integrated Agromet Advisory Service Bulletin 25<sup>th</sup> to 29<sup>th</sup> Sept '15

## Kohima District

Bulletin No:73/2015

Weather summary of the preceding week			Weather forecast valid upto 29 <sup>th</sup> Sept '15
			Probability of medium rain on 25 <sup>th</sup> Sept Max temp is expected to be around 23 <sup>0</sup> C - 30 <sup>0</sup> C and min temp is expected to be around 15 <sup>0</sup> C to 16 <sup>0</sup> C. Possibility of mainly cloudy sky the coming week Relative Humidity will range from 41% to 98% Wind speed 2 kmph Wind direction will be mostly south westerly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<b>Field crops</b>			
<b>Terrace/ Wetland paddy</b>	Milk development stage	Gundi bug	<i>They feed on all parts of the plant. Proper water management and sanitation practices should be done.</i>
<b>Jhum paddy</b>	Harvesting stage		<i>As the grain is to be harvested, the store house should be fumigated; store bins cleaned and should be weather proof, rodent proof and damp proof.</i>
<b>Maize</b>	Harvesting stage	-	<i>For composite and hybrid, Do not wait for stalks and leaves to dry because mostly they remain green</i>
<b>Groundnut</b>	Fruiting stage	-	<i>Proper earthing up should be done.</i>
<b>Horticultural crops</b>			
<b>Citrus</b>	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
<b>Chilli</b>	Harvesting stage	-	<i>Avoid injuries during harvesting.</i>
<b>Ginger</b>	Vegetative stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>
<b>Piggery</b>	Avoid stagnant water in and around the sty, maintain proper hygienic condition.		

	<b><i>Poultry</i></b>	In case of <b>diarrhoea in poultry</b> , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<b><i>Fishery</i></b>	<p>Chemicals found in or used on a pond's watershed can enter the pond and cause fish kills. Runoff following a heavy rain can transport a variety of unwanted chemicals into the pond including pesticides, herbicides, fertilizers, petroleum products.</p> <p>Avoid areas where the likelihood of chemical runoff is high and avoid using chemicals on the watershed</p>




## Integrated Agromet Advisory Service Bulletin 25<sup>th</sup> to 29<sup>th</sup> Sept '15

### Longleng District

Bulletin No:73/2015

Weather summary of the preceding week			Weather forecast valid upto 29 <sup>th</sup> Sept '15
			Probability of light rain will occur the coming week Max temp is expected to be around 26 <sup>0</sup> C - 30 <sup>0</sup> C and min temp is expected to be around 17 <sup>0</sup> C to 19 <sup>0</sup> C. <b>Possibility of cloudy sky the coming week</b> Relative Humidity will range from 53% to 98% The wind speed 1 - 2 kmph Wind direction will be mostly easterly
Main Crops	Stage	Pest/diseases	Agro-meteorological Advisories
<b>Field crops</b>			
<b>Terrace/ Wetland paddy</b>	Milk development stage	Gundi bug	<i>They feed on all parts of the plant. Proper water management and sanitation practices should be done.</i>
<b>Jhum paddy</b>	Harvesting stage		<i>As the grain is to be harvested, the store house should be fumigated; store bins cleaned and should be weather proof, rodent proof and damp proof.</i>
<b>Maize</b>	Harvesting stage	-	<i>For composite and hybrid, Do not wait for stalks and leaves to dry because mostly they remain green</i>
<b>Groundnut</b>	Fruiting stage	-	<i>Proper earthing up should be done.</i>
<b>Horticultural crops</b>			
<b>Citrus</b>	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
<b>Chilli</b>	Harvesting stage	-	<i>Avoid injuries during harvesting.</i>
<b>Ginger</b>	Vegetative stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>
<b>Piggery</b>	Avoid stagnant water in and around the sty, maintain proper hygienic condition.		

	<b><i>Poultry</i></b>	In case of <b>diarrhoea in poultry</b> , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<b><i>Fishery</i></b>	<p>Chemicals found in or used on a pond's watershed can enter the pond and cause fish kills. Runoff following a heavy rain can transport a variety of unwanted chemicals into the pond including pesticides, herbicides, fertilizers, petroleum products.</p> <p>Avoid areas where the likelihood of chemical runoff is high and avoid using chemicals on the watershed</p>



# Integrated Agromet Advisory Service Bulletin 25<sup>th</sup> to 29<sup>th</sup> Sept '15

## Mokokchung District

Bulletin No: 73/2015

Weather summary of the preceding week			Weather forecast valid upto 29 <sup>th</sup> Sept '15
			Probability light rain will occur the coming week Max temp is expected to be around 26 <sup>0</sup> C - 29 <sup>0</sup> C and min temp is expected to be around 17 <sup>0</sup> C to 19 <sup>0</sup> C. Possibility of mainly cloudy sky the coming week Relative Humidity will range from 55% to 91% and Wind speed 2 kmph Wind direction will be mostly southerly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<b>Field crops</b>			
<b>Terrace/ Wetland paddy</b>	Milk development stage	Gundi bug	They feed on all parts of the plant. Proper water management and sanitation practices should be done.
<b>Jhum paddy</b>	Harvesting stage		As the grain is to be harvested, the store house should be fumigated; store bins cleaned and should be weather proof, rodent proof and damp proof.
<b>Maize</b>	Harvesting stage	-	For composite and hybrid, Do not wait for stalks and leaves to dry because mostly they remain green
<b>Groundnut</b>	Fruiting stage	-	Proper earthing up should be done.
<b>Horticultural crops</b>			
<b>Citrus</b>	-		Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).
<b>Chilli</b>	Harvesting stage	-	Avoid injuries during harvesting.
<b>Ginger</b>	Vegetative stage		Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.
<b>Piggery</b>	Avoid stagnant water in and around the sty, maintain proper hygienic condition.		
<b>Poultry</b>	In case of <b>diarrhoea in poultry</b> , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date		



### ***Fishery***

Chemicals found in or used on a pond's watershed can enter the pond and cause fish kills. Runoff following a heavy rain can transport a variety of unwanted chemicals into the pond including pesticides, herbicides, fertilizers, petroleum products.


Avoid areas where the likelihood of chemical runoff is high and avoid using chemicals on the watershed



# Integrated Agromet Advisory Service Bulletin from 25<sup>th</sup> to 29<sup>th</sup> Sept '15 Mon District

Bulletin No:73/2015

Weather summary of the preceding week			Weather forecast valid upto 29 <sup>th</sup> Sept '15
			<ul style="list-style-type: none"> <li>▪ Probability of light rain will occur the coming week.</li> <li>▪ The max temp is likely to be 26<sup>0</sup>C - 29<sup>0</sup>C and the min temp 16<sup>0</sup>C - 18<sup>0</sup>C.</li> <li>▪ Sky is likely to be mainly cloudy the coming week</li> <li>▪ Relative Humidity is likely to range from 56% to 92%</li> <li>▪ Wind speed may reach upto 1- 2 kmph</li> <li>▪ Wind direction will be mostly southwesterly</li> </ul>
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<b>Field crops</b>			
<b>Terrace/ Wetland paddy</b>	Milk development stage	Gundi bug	<i>They feed on all parts of the plant. Proper water management and sanitation practices should be done.</i>
<b>Jhum paddy</b>	Harvesting stage		<i>As the grain is to be harvested, the store house should be fumigated; store bins cleaned and should be weather proof, rodent proof and damp proof.</i>
<b>Maize</b>	Harvesting stage	-	<i>For composite and hybrid, Do not wait for stalks and leaves to dry because mostly they remain green</i>
<b>Groundnut</b>	Fruiting stage	-	<i>Proper earthing up should be done.</i>
<b>Horticultural crops</b>			
<b>Citrus</b>	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
<b>Chilli</b>	Harvesting stage	-	<i>Avoid injuries during harvesting.</i>
<b>Ginger</b>	Vegetative stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>
<b>Piggery</b>	Avoid stagnant water in and around the sty, maintain proper hygienic condition.		

	<b><i>Poultry</i></b>	In case of <b>diarrhoea in poultry</b> , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<b><i>Fishery</i></b>	<p>Chemicals found in or used on a pond's watershed can enter the pond and cause fish kills. Runoff following a heavy rain can transport a variety of unwanted chemicals into the pond including pesticides, herbicides, fertilizers, petroleum products.</p> <p>Avoid areas where the likelihood of chemical runoff is high and avoid using chemicals on the watershed</p>





# Integrated Agromet Advisory Service Bulletin 25<sup>th</sup> to 29<sup>th</sup> Sept '15

## Peren District

Bulletin No:73/2015

### Weather summary of the preceding week

### Weather forecast valid upto 29<sup>th</sup> Sept '15

Probability of medium rain on 25<sup>th</sup> Sept.  
 Max temp is expected to be around 25<sup>0</sup>C - 30<sup>0</sup>C and min temp is expected to be around 17<sup>0</sup>C to 19<sup>0</sup>C.  
 Possibility of mainly cloudy sky the coming week  
 Relative Humidity will range from 43% to 98%  
 The wind speed 2 kmph  
 Wind direction will be southerly

### Main Crops

### Stage

### Pest/ diseases


### Agro-meteorological Advisories

### Field crops

<b>Lowland paddy</b>	Milk development stage	Gundi bug	<i>They feed on all parts of the plant. Proper water management and sanitation practices should be done.</i>
<b>Jhum paddy</b>	Harvesting stage		<i>As the grain is to be harvested, the store house should be fumigated; store bins cleaned and should be weather proof, rodent proof and damp proof.</i>
<b>Maize</b>	Harvesting stage	-	<i>For composite and hybrid, Do not wait for stalks and leaves to dry because mostly they remain green</i>
<b>Groundnut</b>	Fruiting stage	-	<i>Proper earthing up should be done.</i>

### Horticultural crops

<b>Citrus</b>	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
<b>Chilli</b>	Harvesting stage	-	<i>Avoid injuries during harvesting.</i>
<b>Ginger</b>	Vegetative stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>

	<b><i>Piggery</i></b>	Avoid stagnant water in and around the sty, maintain proper hygienic condition.
	<b><i>Poultry</i></b>	In case of <b>diarrhoea in poultry</b> , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<b><i>Fishery</i></b>	Chemicals found in or used on a pond's watershed can enter the pond and cause fish kills. Runoff following a heavy rain can transport a variety of unwanted chemicals into the pond including pesticides, herbicides, fertilizers, petroleum products. Avoid areas where the likelihood of chemical runoff is high and avoid using chemicals on the watershed




# Integrated Agromet Advisory Service Bulletin 25<sup>th</sup> to 29<sup>th</sup> Sept '15

## Phek District

Bulletin No:73/2015

CAR RESEARCH COMPLEX FOR NEH REGION  
(Indian Council of Agricultural Research)  
NAGALAND CENTER, JHARNAPANI, MEDZIPHEMA

Weather summary of the preceding week			Weather forecast valid upto 29 <sup>th</sup> Sept '15
			<p>Probability there will be light rain the coming week</p> <p>Max temp is expected to be around 25<sup>0</sup>C - 28<sup>0</sup>C and min temp is expected to be around 15<sup>0</sup>C to 18<sup>0</sup>C.</p> <p>Possibility of mainly cloudy sky the coming week</p> <p>Relative Humidity will range from 54% to 95%</p> <p>The wind speed 0-2 kmph</p> <p>Wind direction will blow mostly from southerly</p>
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<b>Field crops</b>			
<b>Terrace/ Wetland paddy</b>	Milk development stage	Gundi bug	<i>They feed on all parts of the plant. Proper water management and sanitation practices should be done.</i>
<b>Jhum paddy</b>	Harvesting stage		<i>As the grain is to be harvested, the store house should be fumigated; store bins cleaned and should be weather proof, rodent proof and damp proof.</i>
<b>Maize</b>	Harvesting stage	-	<i>For composite and hybrid, Do not wait for stalks and leaves to dry because mostly they remain green</i>
<b>Groundnut</b>	Fruiting stage	-	<i>Proper earthing up should be done.</i>
<b>Horticultural crops</b>			
<b>Citrus</b>	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
<b>Chilli</b>	Harvesting stage	-	<i>Avoid injuries during harvesting.</i>
<b>Ginger</b>	Vegetative stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>

	<b><i>Piggery</i></b>	Avoid stagnant water in and around the sty, maintain proper hygienic condition.
	<b><i>Poultry</i></b>	In case of <b>diarrhoea in poultry</b> , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<b><i>Fishery</i></b>	Chemicals found in or used on a pond's watershed can enter the pond and cause fish kills. Runoff following a heavy rain can transport a variety of unwanted chemicals into the pond including pesticides, herbicides, fertilizers, petroleum products. Avoid areas where the likelihood of chemical runoff is high and avoid using chemicals on the watershed




## Integrated Agromet Advisory Service Bulletin 25<sup>th</sup> to 29<sup>th</sup> Sept '15

### Tuensang District

Bulletin No:73/2015

Weather summary of the preceding week			Weather forecast valid upto 29 <sup>th</sup> Sept '15
			Probability there will be light rain the coming week Max temp is expected to be around 23 <sup>0</sup> C - 24 <sup>0</sup> C and min temp is expected to be around 12 <sup>0</sup> C to 13 <sup>0</sup> C. Possibility of mainly cloudy sky the coming week Relative humidity will range from 57% to 93% Wind speed 2 kmph Wind direction will be mostly south easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<b>Field crops</b>			
<b>Terrace/ Wetland paddy</b>	Milk development stage	Gundi bug	<i>They feed on all parts of the plant. Proper water management and sanitation practices should be done.</i>
<b>Jhum paddy</b>	Harvesting stage		<i>As the grain is to be harvested, the store house should be fumigated; store bins cleaned and should be weather proof, rodent proof and damp proof.</i>
<b>Maize</b>	Harvesting stage	-	<i>For composite and hybrid, Do not wait for stalks and leaves to dry because mostly they remain green</i>
<b>Groundnut</b>	Fruiting stage	-	<i>Proper earthing up should be done.</i>
<b>Horticultural crops</b>			
<b>Citrus</b>	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
<b>Chilli</b>	Harvesting stage	-	<i>Avoid injuries during harvesting.</i>
<b>Ginger</b>	Vegetative stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>
<b>Piggery</b>	Avoid stagnant water in and around the sty, maintain proper hygienic condition.		


	<b><i>Poultry</i></b>	In case of <b>diarrhoea in poultry</b> , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<b><i>Fishery</i></b>	<p>Chemicals found in or used on a pond's watershed can enter the pond and cause fish kills. Runoff following a heavy rain can transport a variety of unwanted chemicals into the pond including pesticides, herbicides, fertilizers, petroleum products.</p> <p>Avoid areas where the likelihood of chemical runoff is high and avoid using chemicals on the watershed</p>



## Integrated Agromet Advisory Service Bulletin 25<sup>th</sup> to 29<sup>th</sup> Sept '15 Wokha District

Bulletin No:73/2015

Weather summary of the preceding week			Weather forecast valid upto 29 <sup>th</sup> Sept '15
			<p>Probability light to medium rain will occur the coming week  Max temp is expected to be around 27<sup>0</sup>C - 30<sup>0</sup>C and min temp is expected to be around 17<sup>0</sup>C to 20<sup>0</sup>C.  Possibility of mainly cloudy sky the coming week  Relative Humidity will range from 54% to 98%  The wind speed 2 kmph  Wind direction will blow mostly from southerly</p>
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<b>Field crops</b>			
<b>Terrace/ Wetland paddy</b>	Milk development stage	Gundi bug	<i>They feed on all parts of the plant. Proper water management and sanitation practices should be done.</i>
<b>Jhum paddy</b>	Harvesting stage		<i>As the grain is to be harvested, the store house should be fumigated; store bins cleaned and should be weather proof, rodent proof and damp proof.</i>
<b>Maize</b>	Harvesting stage	-	<i>For composite and hybrid, Do not wait for stalks and leaves to dry because mostly they remain green</i>
<b>Groundnut</b>	Fruiting stage	-	<i>Proper earthing up should be done.</i>
<b>Horticultural crops</b>			
<b>Citrus</b>	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
<b>Chilli</b>	Harvesting stage	-	<i>Avoid injuries during harvesting.</i>
<b>Ginger</b>	Vegetative stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>

	<b><i>Piggery</i></b>	Avoid stagnant water in and around the sty, maintain proper hygienic condition.
	<b><i>Poultry</i></b>	In case of <b>diarrhoea in poultry</b> , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<b><i>Fishery</i></b>	Chemicals found in or used on a pond's watershed can enter the pond and cause fish kills. Runoff following a heavy rain can transport a variety of unwanted chemicals into the pond including pesticides, herbicides, fertilizers, petroleum products. Avoid areas where the likelihood of chemical runoff is high and avoid using chemicals on the watershed








--	--



## Integrated Agromet Advisory Service Bulletin 25<sup>th</sup> to 29<sup>th</sup> Sept '15 Zunheboto District

Bulletin No:73/2015

Weather summary of the preceding week			Weather forecast valid upto 29 <sup>th</sup> Sept '15
			<p>Probability light rain will occur the coming week  Max temp is expected to be around 25<sup>0</sup>C - 28<sup>0</sup>C and min temp is expected to be around 16<sup>0</sup>C to 18<sup>0</sup>C.  Possibility of mainly cloudy sky the coming week  Relative Humidity will range from 58% to 91%  The wind speed 2 kmph  Wind direction will blow mostly from easterly</p>
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<b>Field crops</b>			
<b>Terrace/ Wetland paddy</b>	Milk development stage	Gundi bug	<i>They feed on all parts of the plant. Proper water management and sanitation practices should be done.</i>
<b>Jhum paddy</b>	Harvesting stage		<i>As the grain is to be harvested, the store house should be fumigated; store bins cleaned and should be weather proof, rodent proof and damp proof.</i>
<b>Maize</b>	Harvesting stage	-	<i>For composite and hybrid, Do not wait for stalks and leaves to dry because mostly they remain green</i>
<b>Groundnut</b>	Fruiting stage	-	<i>Proper earthing up should be done.</i>
<b>Horticultural crops</b>			
<b>Citrus</b>	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
<b>Chilli</b>	Harvesting stage	-	<i>Avoid injuries during harvesting.</i>
<b>Ginger</b>	Vegetative stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>
<b>Piggery</b>	Avoid stagnant water in and around the sty, maintain proper hygienic condition.		

	<b><i>Poultry</i></b>	In case of <b>diarrhoea in poultry</b> , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date
	<b><i>Fishery</i></b>	<p>Chemicals found in or used on a pond's watershed can enter the pond and cause fish kills. Runoff following a heavy rain can transport a variety of unwanted chemicals into the pond including pesticides, herbicides, fertilizers, petroleum products.</p> <p>Avoid areas where the likelihood of chemical runoff is high and avoid using chemicals on the watershed</p>





## Integrated Agromet Advisory Service Bulletin 25<sup>th</sup> to 29<sup>th</sup> Sept '15

### Dimapur District

Bulletin No:73/2015

Weather summary of the preceding week			Weather forecast valid upto 29 <sup>th</sup> Sept '15
			<ul style="list-style-type: none"> <li>Probability of medium rain on 25<sup>th</sup> Sept.</li> <li>The <b>max temp</b> is likely to be 27<sup>0</sup>C - 33<sup>0</sup>C and the <b>min temp</b> 18<sup>0</sup>C - 20<sup>0</sup>C.</li> <li><b>Sky is likely to be mainly cloudy sky the coming week</b></li> <li><b>Relative Humidity</b> is likely to range from 48% to 95%</li> <li><b>Wind speed</b> 2 kmph</li> <li><b>Wind direction</b> will be mostly easterly</li> </ul>
Main Crops	Stage	Pest/ diseases	Agro-meteorological Advisories
<b>Field crops</b>			
<b>Lowland paddy</b>	Milk development stage	Gundi bug	<i>They feed on all parts of the plant. Proper water management and sanitation practices should be done.</i>
<b>Jhum paddy</b>	Harvesting stage		<i>As the grain is to be harvested, the store house should be fumigated; store bins cleaned and should be weather proof, rodent proof and damp proof.</i>
<b>Maize</b>	Harvesting stage	-	<i>For composite and hybrid, Do not wait for stalks and leaves to dry because mostly they remain green</i>
<b>Groundnut</b>	Fruiting stage	-	<i>Proper earthing up should be done.</i>
<b>Horticultural crops</b>			
<b>Citrus</b>	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
<b>Chilli</b>	Harvesting stage	-	<i>Avoid injuries during harvesting.</i>
<b>Ginger</b>	Vegetative stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>
<b>Piggery</b>	Avoid stagnant water in and around the sty, maintain proper hygienic condition.		
<b>Poultry</b>	In case of <b>diarrhoea in poultry</b> , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date		

***Fishery***

Chemicals found in or used on a pond's watershed can enter the pond and cause fish kills. Runoff following a heavy rain can transport a variety of unwanted chemicals into the pond including pesticides, herbicides, fertilizers, petroleum products.

Avoid areas where the likelihood of chemical runoff is high and avoid using chemicals on the watershed



# Integrated Agromet Advisory Service Bulletin 25<sup>th</sup> to 29<sup>th</sup> Sept '15

## Kiphire District

Bulletin No:73/2015

Weather summary of the preceding week			Weather forecast valid upto 29 <sup>th</sup> Sept '15
			Probability of light rain on 25 <sup>th</sup> Sept. Max temp is expected to be around 24 <sup>0</sup> C - 26 <sup>0</sup> C and min temp is expected to be around 14 <sup>0</sup> C to 15 <sup>0</sup> C. Possibility of mainly cloudy sky the coming week Relative Humidity will range from 54% to 98% The wind speed 1- 2 kmph Wind direction will blow mostly from south easterly
Main Crops	Stage	Pest/ diseases	Agro-meteorological Advisories
<b>Field crops</b>			
<b>Terrace/ Wetland paddy</b>	Milk development stage	Gundi bug	<i>They feed on all parts of the plant. Proper water management and sanitation practices should be done.</i>
<b>Jhum paddy</b>	Harvesting stage		<i>As the grain is to be harvested, the store house should be fumigated; store bins cleaned and should be weather proof, rodent proof and damp proof.</i>
<b>Maize</b>	Harvesting stage	-	<i>For composite and hybrid, Do not wait for stalks and leaves to dry because mostly they remain green</i>
<b>Groundnut</b>	Fruiting stage	-	<i>Proper earthing up should be done.</i>
<b>Horticultural crops</b>			
<b>Citrus</b>	-		<i>Application of bordeaux paste on the tree trunk should be continued till Sept (up to 60cm height from ground level).</i>
<b>Chilli</b>	Harvesting stage	-	<i>Avoid injuries during harvesting.</i>
<b>Ginger</b>	Vegetative stage		<i>Soil around the plants should be worked with the help of khurpi . It helps to break the fibrous roots and thereby supports new growth. Soil near the rhizomes becomes loose and friable and helps in proper development of rhizomes.</i>
<b>Piggery</b>	Avoid stagnant water in and around the sty, maintain proper hygienic condition.		
<b>Poultry</b>	In case of <b>diarrhoea in poultry</b> , Always ensure that your birds have fresh clean drinking water, have fresh good quality feed, that has been stored correctly and is in date		





### ***Fishery***

Chemicals found in or used on a pond's watershed can enter the pond and cause fish kills. Runoff following a heavy rain can transport a variety of unwanted chemicals into the pond including pesticides, herbicides, fertilizers, petroleum products.

Avoid areas where the likelihood of chemical runoff is high and avoid using chemicals on the watershed