




Integrated Agromet Advisory Service Bulletin from 26th to 30th Nov'16 Dimapur District

Bulletin No:94/2016

Weather summary of the preceding week			Weather forecast valid upto 30 th Nov'16
<ul style="list-style-type: none"> No rain occurred the past week Maximum and minimum temperatures ranged 26⁰C to 28⁰C and 11⁰C to 14⁰C, respectively. Relative humidity varied from 37% to 95%. Wind speed ranged from 3 to 2 kmph 			<ul style="list-style-type: none"> Probability of no rain the coming week. Max temp is likely to be 26⁰C - 27⁰C and the min temp 12⁰C to 13⁰C Sky is likely to be clear the coming week Relative Humidity is likely to range from 41% to 83%. Wind speed may reach upto 3- 4 kmph Wind direction will be mostly easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles. With prevalence of continued dry weather, need based irrigation is advisable to growing/sown (rabi) crops and vegetables			
TRC/WRC paddy	Harvesting stage		Dry the paddy grains properly and regularly before storing to reduce storage insect pests and fungus. Paddy grains can be mixed with neem leaves at the time of storage to reduce storage pests.
Toria	Sowing	-	Quality seeds should be used. M-27, TS-36 and TS-38 are some of the varieties recommended for our region
Rabi maize	Seedling stage		Provide mulching to conserve moisture
Horticultural crop Keep the field clean by collecting the stubbles and unwanted materials.			
Citrus		Fruit fly	Spray malathion (2ml/l) over fruit to prevent fruit fly egg laying.
Vegetables	Germination stage		Thinning should be done and gap filling should be carried out
Pea	Germinating stage	Powdery mildew	Sowing should be avoided by the end of Nov and Dec. White powdery patches form on leaves and pods and dry weather favors the spread of the disease. Seeds should be treated with hot water.
Poultry	Protect poultry birds from cold winds by hanging curtain/clothing from outside the poultry shed. Use of electric bulbs for heat is advisable.		


	<i>Piggery</i>	Depending on number of pigs farmers should clean pens naturally twice in a week. Farmers should be aware about the symptoms of diseases. Like no interest in food, diarrhea, eye discharge, excessive coughing, hernia, dry skin and irregular spots on skin, excessively long hair, back bone showing etc.
	<i>Fishery</i>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media



Integrated Agromet Advisory Service Bulletin from 26th to 30th Nov'16 Kiphire District

Bulletin No:94/2016

Weather summary of the preceding week			Weather forecast valid upto 30 th Nov'16
<ul style="list-style-type: none"> No rain occurred the past week Maximum and minimum temperatures ranged 22°C to 24°C and 10°C to 12°C, respectively. Relative humidity varied from 26% to 95%. Wind speed ranged from 1 to 2 kmph 			<ul style="list-style-type: none"> Probability of no rain the coming week. Max temp is likely to be 27°C - 28°C and the min temp 13°C to 14°C Sky is likely to be clear the coming week Relative Humidity is likely to range from 30% to 95%. Wind speed may reach upto 1- 2 kmph Wind direction will be mostly southeasterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<p align="center">Field crops</p> <p>Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles. With prevalence of continued dry weather, need based irrigation is advisable to growing/sown (rabi) crops and vegetables</p>			
TRC/WRC paddy	Harvesting stage		Dry the paddy grains properly and regularly before storing to reduce storage insect pests and fungus. Paddy grains can be mixed with neem leaves at the time of storage to reduce storage pests.
Toria	Sowing	-	Quality seeds should be used. M-27, TS-36 and TS-38 are some of the varieties recommended for our region
Rabi maize	Seedling stage		Provide mulching to conserve moisture
<p align="center">Horticultural crop</p> <p>Keep the field clean by collecting the stubbles and unwanted materials.</p>			
Citrus		Fruit fly	Spray malathion (2ml/l) over fruit to prevent fruit fly egg laying.
Vegetables	Germination stage		Thinning should be done and gap filling should be carried out
Pea	Germinating stage	Powdery mildew	Sowing should be avoided by the end of Nov and Dec. White powdery patches form on leaves and pods and dry weather favors the spread of the disease. Seeds should be treated with hot water.
Poultry	Protect poultry birds from cold winds by hanging curtain/clothing from outside the poultry shed. Use of electric bulbs for heat is advisable.		


	<i>Piggery</i>	Depending on number of pigs farmers should clean pens naturally twice in a week. Farmers should be aware about the symptoms of diseases. Like no interest in food, diarrhea, eye discharge, excessive coughing, hernia, dry skin and irregular spots on skin, excessively long hair, back bone showing etc.
	<i>Fishery</i>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media



Integrated Agromet Advisory Service Bulletin from 26th to 30th Nov'16 Kohima District

Bulletin No:94/2016

Weather summary of the preceding week			Weather forecast valid upto 30 th Nov'16
<ul style="list-style-type: none"> No rain occurred the past week Maximum and minimum temperatures ranged 25^oC to 26^oC and 8^oC to 10^oC, respectively. Relative humidity varied from 33% to 94%. Wind speed ranged from 1 to 2 kmph 			<ul style="list-style-type: none"> Probability of no rain the coming week. Max temp is likely to be 25^oC - 26^oC and the min temp 11^oC to 12^oC Sky is likely to be clear the coming week Relative Humidity is likely to range from 35% to 95%. Wind speed may reach upto 3- 4 kmph Wind direction will be mostly easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<p align="center">Field crops</p> <p>Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles. With prevalence of continued dry weather, need based irrigation is advisable to growing/sown (rabi) crops and vegetables</p>			
TRC/WRC paddy	Harvesting stage		<i>Dry the paddy grains properly and regularly before storing to reduce storage insect pests and fungus. Paddy grains can be mixed with neem leaves at the time of storage to reduce storage pests.</i>
Toria	Sowing	-	<i>Quality seeds should be used. M-27, TS-36 and TS-38 are some of the varieties recommended for our region</i>
Rabi maize	Seedling stage		<i>Provide mulching to conserve moisture</i>
<p align="center">Horticultural crop</p> <p>Keep the field clean by collecting the stubbles and unwanted materials.</p>			
Citrus		Fruit fly	<i>Spray malathion (2ml/l) over fruit to prevent fruit fly egg laying.</i>
Vegetables	Germination stage		<i>Thinning should be done and gap filling should be carried out</i>
Pea	Germinating stage	Powdery mildew	<i>Sowing should be avoided by the end of Nov and Dec. White powdery patches form on leaves and pods and dry weather favors the spread of the disease. Seeds should be treated with hot water.</i>
Poultry	Protect poultry birds from cold winds by hanging curtain/clothing from outside the poultry shed. Use of		


		electric bulbs for heat is advisable.
	<i>Piggery</i>	Depending on number of pigs farmers should clean pens naturally twice in a week. Farmers should be aware about the symptoms of diseases. Like no interest in food, diarrhea, eye discharge, excessive coughing, hernia, dry skin and irregular spots on skin, excessively long hair, back bone showing etc.
	<i>Fishery</i>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media



Integrated Agromet Advisory Service Bulletin from 26th to 30th Nov'16 Longleng District

Bulletin No:94/2016

Weather summary of the preceding week			Weather forecast valid upto 30 th Nov'16
<ul style="list-style-type: none"> No rain occurred the past week Maximum and minimum temperatures ranged 11⁰C to 26⁰C and 11⁰C to 13⁰C, respectively. Relative humidity varied from 28% to 95%. Wind speed ranged from 1 to 2 kmph 			<ul style="list-style-type: none"> Probability of no rain the coming week. Max temp is likely to be 27⁰C - 29⁰C and the min temp 12⁰C to 13⁰C Sky is likely to be clear the coming week Relative Humidity is likely to range from 34% to 95%. Wind speed may reach upto 1- 2 kmph Wind direction will be mostly easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<p align="center">Field crops</p> <p>Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles. With prevalence of continued dry weather, need based irrigation is advisable to growing/sown (rabi) crops and vegetables</p>			
TRC/WRC paddy	Harvesting stage		Dry the paddy grains properly and regularly before storing to reduce storage insect pests and fungus. Paddy grains can be mixed with neem leaves at the time of storage to reduce storage pests.
Toria	Sowing	-	Quality seeds should be used. M-27, TS-36 and TS-38 are some of the varieties recommended for our region
Rabi maize	Seedling stage		Provide mulching to conserve moisture
<p align="center">Horticultural crop</p> <p>Keep the field clean by collecting the stubbles and unwanted materials.</p>			
Citrus		Fruit fly	Spray malathion (2ml/l) over fruit to prevent fruit fly egg laying.
Vegetables	Germination stage		Thinning should be done and gap filling should be carried out
Pea	Germinating stage	Powdery mildew	Sowing should be avoided by the end of Nov and Dec. White powdery patches form on leaves and pods and dry weather favors the spread of the disease. Seeds should be treated with hot water.
Poultry	Protect poultry birds from cold winds by hanging curtain/clothing from outside the poultry shed. Use of electric bulbs for heat is advisable.		

	Piggery	Depending on number of pigs farmers should clean pens naturally twice in a week. Farmers should be aware about the symptoms of diseases. Like no interest in food, diarrhea, eye discharge, excessive coughing, hernia, dry skin and irregular spots on skin, excessively long hair, back bone showing etc.
	Fishery	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media

Integrated Agromet Advisory Service Bulletin from 26th to 30th Nov'16

Mokokchung District

Bulletin No: 94/2016

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 30th Nov'16</i>
<ul style="list-style-type: none"> No rain occurred the past week Maximum and minimum temperatures ranged 24⁰C to 26⁰C and 11⁰C to 12⁰C, respectively. Relative humidity varied from 30% to 95%. Wind speed ranged from 1 to 2 kmph 			<ul style="list-style-type: none"> Probability of no rain the coming week. Max temp is likely to be 27⁰C - 28⁰C and the min temp 13⁰C to 14⁰C Sky is likely to be clear the coming week Relative Humidity is likely to range from 34% to 90%. Wind speed may reach upto 1- 2 kmph Wind direction will be mostly southeasterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<p align="center">Field crops</p> <p>Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles. With prevalence of continued dry weather, need based irrigation is advisable to growing/sown (rabi) crops and vegetables</p>			
TRC/WRC paddy	Harvesting stage		<i>Dry the paddy grains properly and regularly before storing to reduce storage insect pests and fungus. Paddy grains can be mixed with neem leaves at the time of storage to reduce storage pests.</i>
Toria	Sowing	-	<i>Quality seeds should be used. M-27, TS-36 and TS-38 are some of the varieties recommended for our region</i>
Rabi maize	Seedling stage		<i>Provide mulching to conserve moisture</i>
<p align="center">Horticultural crop</p> <p>Keep the field clean by collecting the stubbles and unwanted materials.</p>			
Citrus		Fruit fly	<i>Spray malathion (2ml/l) over fruit to prevent fruit fly egg laying.</i>
Vegetables	Germination stage		<i>Thinning should be done and gap filling should be carried out</i>
Pea	Germinating stage	Powdery mildew	<i>Sowing should be avoided by the end of Nov and Dec. White powdery patches form on leaves and pods and dry weather favors the spread of the disease. Seeds should be treated with hot water.</i>
Poultry	Protect poultry birds from cold winds by hanging curtain/clothing from outside the poultry shed. Use of electric bulbs for heat is advisable.		

<i>Piggery</i>	Depending on number of pigs farmers should clean pens naturally twice in a week. Farmers should be aware about the symptoms of diseases. Like no interest in food, diarrhea, eye discharge, excessive coughing, hernia, dry skin and irregular spots on skin, excessively long hair, back bone showing etc.
<i>Fishery</i>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media






Integrated Agromet Advisory Service Bulletin from 26th to 30th Nov'16 Mon District

Bulletin No:94/2016

Weather summary of the preceding week			Weather forecast valid upto 30 th Nov'16
<ul style="list-style-type: none"> No rain occurred the past week Maximum and minimum temperatures ranged 24⁰C to 25⁰C and 13⁰C to 14⁰C, respectively. Relative humidity varied from 26% to 92% Wind speed ranged from 1 to 2 kmph 			<ul style="list-style-type: none"> Probability of no rain the coming week. Max temp is likely to be 27⁰C - 28⁰C and the min temp 13⁰C to 14⁰C Sky is likely to be clear the coming week Relative Humidity is likely to range from 33% to 94%. Wind speed may reach upto 1- 2 kmph Wind direction will be mostly southeasterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<p align="center">Field crops</p> <p>Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles. With prevalence of continued dry weather, need based irrigation is advisable to growing/sown (rabi) crops and vegetables</p>			
TRC/WRC paddy	Harvesting stage		<i>Dry the paddy grains properly and regularly before storing to reduce storage insect pests and fungus. Paddy grains can be mixed with neem leaves at the time of storage to reduce storage pests.</i>
Toria	Sowing	-	<i>Quality seeds should be used. M-27, TS-36 and TS-38 are some of the varieties recommended for our region</i>
Rabi maize	Seedling stage		<i>Provide mulching to conserve moisture</i>
<p align="center">Horticultural crop</p> <p>Keep the field clean by collecting the stubbles and unwanted materials.</p>			
Citrus		Fruit fly	<i>Spray malathion (2ml/l) over fruit to prevent fruit fly egg laying.</i>
Vegetables	Germination stage		<i>Thinning should be done and gap filling should be carried out</i>
Pea	Germinating stage	Powdery mildew	<i>Sowing should be avoided by the end of Nov and Dec. White powdery patches form on leaves and pods and dry weather favors the spread of the disease. Seeds should be treated with hot water.</i>
Poultry	Protect poultry birds from cold winds by hanging curtain/clothing from outside the poultry shed. Use of electric bulbs for heat is advisable.		

	Piggery	Depending on number of pigs farmers should clean pens naturally twice in a week. Farmers should be aware about the symptoms of diseases. Like no interest in food, diarrhea, eye discharge, excessive coughing, hernia, dry skin and irregular spots on skin, excessively long hair, back bone showing etc.
	Fishery	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media



Integrated Agromet Advisory Service Bulletin from 26th to 30th Nov'16

Peren District

Bulletin No:94/2016

Weather summary of the preceding week			Weather forecast valid upto 30 th Nov'16
<ul style="list-style-type: none"> No rain occurred the past week Maximum and minimum temperatures ranged 26⁰C to 27⁰C and 10⁰C to 13⁰C, respectively. Relative humidity varied from 34% to 95%. Wind speed ranged from 1 to 2 kmph 			<ul style="list-style-type: none"> Probability of no rain the coming week. Max temp is likely to be 26⁰C - 28⁰C and the min temp 13⁰C to 14⁰C Sky is likely to be clear the coming week Relative Humidity is likely to range from 36% to 90%. Wind speed may reach upto 3- 4 kmph Wind direction will be mostly easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles. With prevalence of continued dry weather, need based irrigation is advisable to growing/sown (rabi) crops and vegetables			
TRC/WRC paddy	Harvesting stage		Dry the paddy grains properly and regularly before storing to reduce storage insect pests and fungus. Paddy grains can be mixed with neem leaves at the time of storage to reduce storage pests.
Toria	Sowing	-	Quality seeds should be used. M-27, TS-36 and TS-38 are some of the varieties recommended for our region
Rabi maize	Seedling stage		Provide mulching to conserve moisture
Horticultural crop Keep the field clean by collecting the stubbles and unwanted materials.			
Citrus		Fruit fly	Spray malathion (2ml/l) over fruit to prevent fruit fly egg laying.
Vegetables	Germination stage		Thinning should be done and gap filling should be carried out
Pea	Germinating stage	Powdery mildew	Sowing should be avoided by the end of Nov and Dec. White powdery patches form on leaves and pods and dry weather favors the spread of the disease. Seeds should be treated with hot water.
Poultry	Protect poultry birds from cold winds by hanging curtain/clothing from outside the poultry shed. Use of electric bulbs for heat is advisable.		




<i>Piggery</i>	Depending on number of pigs farmers should clean pens naturally twice in a week. Farmers should be aware about the symptoms of diseases. Like no interest in food, diarrhea, eye discharge, excessive coughing, hernia, dry skin and irregular spots on skin, excessively long hair, back bone showing etc.
<i>Fishery</i>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media



Integrated Agromet Advisory Service Bulletin from 26th to 30th Nov'16 Phek District

Bulletin No:94/2016

Weather summary of the preceding week			Weather forecast valid upto 30 th Nov'16
<ul style="list-style-type: none"> No rain occurred the past week Maximum and minimum temperatures ranged 24⁰C to 25⁰C and 13⁰C to 14⁰C, respectively. Relative humidity varied from 30% to 95%. Wind speed ranged from 1 to 2 kmph 			<ul style="list-style-type: none"> Probability of no rain the coming week. Max temp is likely to be 25⁰C - 26⁰C and the min temp 13⁰C to 14⁰C Sky is likely to be clear the coming week Relative Humidity is likely to range from 34% to 92%. Wind speed may reach upto 2- 4 kmph Wind direction will be mostly south easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<p align="center">Field crops</p> <p>Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles. With prevalence of continued dry weather, need based irrigation is advisable to growing/sown (rabi) crops and vegetables</p>			
TRC/WRC paddy	Harvesting stage		<i>Dry the paddy grains properly and regularly before storing to reduce storage insect pests and fungus. Paddy grains can be mixed with neem leaves at the time of storage to reduce storage pests.</i>
Toria	Sowing	-	<i>Quality seeds should be used. M-27, TS-36 and TS-38 are some of the varieties recommended for our region</i>
Rabi maize	Seedling stage		<i>Provide mulching to conserve moisture</i>
<p align="center">Horticultural crop</p> <p>Keep the field clean by collecting the stubbles and unwanted materials.</p>			
Citrus		Fruit fly	<i>Spray malathion (2ml/l) over fruit to prevent fruit fly egg laying.</i>
Vegetables	Germination stage		<i>Thinning should be done and gap filling should be carried out</i>
Pea	Germinating stage	Powdery mildew	<i>Sowing should be avoided by the end of Nov and Dec. White powdery patches form on leaves and pods and dry weather favors the spread of the disease. Seeds should be treated with hot water.</i>


	<i>Poultry</i>	Protect poultry birds from cold winds by hanging curtain/clothing from outside the poultry shed. Use of electric bulbs for heat is advisable.
	<i>Piggery</i>	Depending on number of pigs farmers should clean pens naturally twice in a week. Farmers should be aware about the symptoms of diseases. Like no interest in food, diarrhea, eye discharge, excessive coughing, hernia, dry skin and irregular spots on skin, excessively long hair, back bone showing etc.
 भारतीय ICAR	<i>Fishery</i>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media



Integrated Agromet Advisory Service Bulletin from 26th to 30th Nov'16 Tuensang District

Bulletin No:94/2016

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 30th Nov'16</i>
<ul style="list-style-type: none"> No rain occurred the past week Maximum and minimum temperatures ranged 21⁰C to 23⁰C and 9⁰C to 10⁰C, respectively. Relative humidity varied from 25% to 94%. Wind speed ranged from 1 to 2 kmph 			<ul style="list-style-type: none"> Probability of no rain the coming week. Max temp is likely to be 23⁰C - 25⁰C and the min temp 12⁰C to 14⁰C Sky is likely to be clear the coming week Relative Humidity is likely to range from 28% to 95%. Wind speed may reach upto 1- 2 kmph Wind direction will be mostly southeasterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<p align="center">Field crops</p> <p>Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles. With prevalence of continued dry weather, need based irrigation is advisable to growing/sown (rabi) crops and vegetables</p>			
TRC/WRC paddy	Harvesting stage		<i>Dry the paddy grains properly and regularly before storing to reduce storage insect pests and fungus. Paddy grains can be mixed with neem leaves at the time of storage to reduce storage pests.</i>
Toria	Sowing	-	<i>Quality seeds should be used. M-27, TS-36 and TS-38 are some of the varieties recommended for our region</i>
Rabi maize	Seedling stage		<i>Provide mulching to conserve moisture</i>
<p align="center">Horticultural crop</p> <p>Keep the field clean by collecting the stubbles and unwanted materials.</p>			
Citrus		Fruit fly	<i>Spray malathion (2ml/l) over fruit to prevent fruit fly egg laying.</i>
Vegetables	Germination stage		<i>Thinning should be done and gap filling should be carried out</i>
Pea	Germinating stage	Powdery mildew	<i>Sowing should be avoided by the end of Nov and Dec. White powdery patches form on leaves and pods and dry weather favors the spread of the disease. Seeds should be treated with hot water.</i>
Poultry	Protect poultry birds from cold winds by hanging curtain/clothing from outside the poultry shed. Use of electric bulbs for heat is advisable.		


	<i>Piggery</i>	Depending on number of pigs farmers should clean pens naturally twice in a week. Farmers should be aware about the symptoms of diseases. Like no interest in food, diarrhea, eye discharge, excessive coughing, hernia, dry skin and irregular spots on skin, excessively long hair, back bone showing etc.
 भारतीय ICAR	<i>Fishery</i>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media



Integrated Agromet Advisory Service Bulletin from 26th to 30th Nov'16 Wokha District

Bulletin No:94/2016

Weather summary of the preceding week			Weather forecast valid upto 30 th Nov'16
<ul style="list-style-type: none"> No rain occurred the past week Maximum and minimum temperatures ranged 25⁰C to 26⁰C and 8⁰C to 11⁰C, respectively. Relative humidity varied from 33% to 95%. Wind speed ranged from 1 to 3 kmph 			<ul style="list-style-type: none"> Probability of no rain the coming week. Max temp is likely to be 26⁰C - 27⁰C and the min temp 13⁰C to 14⁰C Sky is likely to be clear the coming week Relative Humidity is likely to range from 38% to 84%. Wind speed may reach upto 3- 4 kmph Wind direction will be mostly southeasterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles. With prevalence of continued dry weather, need based irrigation is advisable to growing/sown (rabi) crops and vegetables			
TRC/WRC paddy	Harvesting stage		<i>Dry the paddy grains properly and regularly before storing to reduce storage insect pests and fungus. Paddy grains can be mixed with neem leaves at the time of storage to reduce storage pests.</i>
Toria	Sowing	-	<i>Quality seeds should be used. M-27, TS-36 and TS-38 are some of the varieties recommended for our region</i>
Rabi maize	Seedling stage		<i>Provide mulching to conserve moisture</i>
Horticultural crop Keep the field clean by collecting the stubbles and unwanted materials.			
Citrus		Fruit fly	<i>Spray malathion (2ml/l) over fruit to prevent fruit fly egg laying.</i>
Vegetables	Germination stage		<i>Thinning should be done and gap filling should be carried out</i>
Pea	Germinating stage	Powdery mildew	<i>Sowing should be avoided by the end of Nov and Dec. White powdery patches form on leaves and pods and dry weather favors the spread of the disease. Seeds should be treated with hot water.</i>

	<i>Poultry</i>	Protect poultry birds from cold winds by hanging curtain/clothing from outside the poultry shed. Use of electric bulbs for heat is advisable.
	<i>Piggery</i>	Depending on number of pigs farmers should clean pens naturally twice in a week. Farmers should be aware about the symptoms of diseases. Like no interest in food, diarrhea, eye discharge, excessive coughing, hernia, dry skin and irregular spots on skin, excessively long hair, back bone showing etc.
	<i>Fishery</i>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media




--	--



Integrated Agromet Advisory Service Bulletin from 26th to 30th Nov'16 Zunheboto District

Bulletin No:94/2016

Weather summary of the preceding week			Weather forecast valid upto 30 th Nov'16
<ul style="list-style-type: none"> No rain occurred the past week Maximum and minimum temperatures ranged 24⁰C to 25⁰C and 13⁰C to 14⁰C, respectively. Relative humidity varied from 29% to 95%. Wind speed ranged from 1 to 2 kmph 			<ul style="list-style-type: none"> Probability of no rain the coming week. Max temp is likely to be 26⁰C - 27⁰C and the min temp 13⁰C to 16⁰C Sky is likely to be clear the coming week Relative Humidity is likely to range from 31% to 90%. Wind speed may reach upto 2- 4 kmph Wind direction will be mostly southeasterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Field crops Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles. With prevalence of continued dry weather, need based irrigation is advisable to growing/sown (rabi) crops and vegetables			
TRC/WRC paddy	Harvesting stage		Dry the paddy grains properly and regularly before storing to reduce storage insect pests and fungus. Paddy grains can be mixed with neem leaves at the time of storage to reduce storage pests.
Toria	Sowing	-	Quality seeds should be used. M-27, TS-36 and TS-38 are some of the varieties recommended for our region
Rabi maize	Seedling stage		Provide mulching to conserve moisture
Horticultural crop Keep the field clean by collecting the stubbles and unwanted materials.			
Citrus		Fruit fly	Spray malathion (2ml/l) over fruit to prevent fruit fly egg laying.
Vegetables	Germination stage		Thinning should be done and gap filling should be carried out
Pea	Germinating stage	Powdery mildew	Sowing should be avoided by the end of Nov and Dec. White powdery patches form on leaves and pods and dry weather favors the spread of the disease. Seeds should be treated with hot water.
Poultry	Protect poultry birds from cold winds by hanging curtain/clothing from outside the poultry shed. Use of electric bulbs for heat is advisable.		

	Piggery	Depending on number of pigs farmers should clean pens naturally twice in a week. Farmers should be aware about the symptoms of diseases. Like no interest in food, diarrhea, eye discharge, excessive coughing, hernia, dry skin and irregular spots on skin, excessively long hair, back bone showing etc.
	Fishery	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media

