




Integrated Agromet Advisory Service Bulletin from 29th Oct to 2nd Nov'16 Zunheboto District

Bulletin No:86/2016

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 2nd Nov'16</i>
<ul style="list-style-type: none"> Light rain occurred the past week Maximum and minimum temperatures ranged 29⁰C to 31⁰C and 18⁰C to 20⁰C , respectively. Relative humidity varied from 39% to 93%. Wind speed ranged from 1 to 2 kmph 			<ul style="list-style-type: none"> Probability of Light rain the coming week. Max temp is likely to be 29⁰C - 30⁰C and the min temp 18⁰C to 20⁰C Sky is likely to be mainly cloudy the coming week Relative Humidity is likely to range from 42% to 95%. Wind speed may reach upto 1-2 kmph Wind direction will be mostly easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<p style="text-align: center;">Field crops</p> <p style="text-align: center;">Undertake timely harvest of kharif crops with safe storage of produce. Seed treatment with fungicides (@ 2 to 3 g/kg seed) prevents seed borne diseases, promotes better germination and crop stand leading to higher productivity in all rabi crops</p>			
TRC/WRC paddy	Ripening stage	-	<i>Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds</i>
Toria	Land preparation	-	<i>Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region</i>
Rabi maize	Land preparation		<i>Land preparation should be started for rabi maize. It can be grown in all types of soils having adequate provision of drainage. Field should be ploughed properly so as to expose the pupae of red hairy caterpillar. Sow certified seeds from a reliable source to prevent seed rot and seedling blight. If one wants to use his own seed, seed treatment is a must. Some of the varieties recommended for our region are HQPM-1, Allrounder and RCM-76.</i>
<p style="text-align: center;">Horticultural crop</p> <p style="text-align: center;">Spread well decomposed cowdung 5 to 6 tones per hectare for the main field and mix it well in the soil by cultivating the field. Keep the field clean by collecting the stubbles and unwanted materials.</p>			
Citrus		Leaf miner	<i>Proper management should be done for leaf miner. Monitor pest regularly, manage ants and whitewash of the trunk should be done.</i>


	Vegetables			<p><i>Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done.</i></p> <p><i>Suitable varieties for our region:</i></p> <ol style="list-style-type: none"> 1. <i>Cabbage – Green express (Hybrid), Rareball (Hybrid), Pride of India (OP).</i> 2. <i>Broccoli – Pushpa, princess.</i> 3. <i>Tomato – Pusa ruby, Arka vikash, Punjab swari.</i> 4. <i>Knolkhol – White Vienna.</i> 5. <i>Radish – Japanese white, Pusa chetki.</i> 6. <i>Pea – Arkel</i>
	Poultry	<p>Prevailing high humidity may cause ammonia fumes which irritate the respiratory tract and eyes of the poultry. Good ventilation provides a comfortable environment without draft. Farmers are advised to provide clean drinking water round the clock and water troughs/ feeders should be regularly cleaned. Antibiotics or other stress medications may be added to water if desired.</p>		
	Piggery	<p>Routine vaccination programme is recommended and once any disease is diagnosed treat immediately with antibiotics.</p>		
 <p>भारतीय ICAR</p>	Fishery	<p>Organic manure/ cattle manure may be applied @ 2 ton per ha to maintain the natural productivity of the ponds. To avoid losses, avoid application of manure at maximum water depth.</p>		



Integrated Agromet Advisory Service Bulletin from 29th Oct to 2nd Nov'16 Dimapur District

Bulletin No:86/2016

Weather summary of the preceding week			Weather forecast valid upto 2 nd Nov'16
<ul style="list-style-type: none"> Light rain occurred the past week Maximum and minimum temperatures ranged 29⁰C to 30⁰C and 19⁰C to 21⁰C, respectively. Relative humidity varied from 42% to 95%. Wind speed ranged from 1 to 2 kmph 			<ul style="list-style-type: none"> Probability of light rain the coming week. Max temp is likely to be 28⁰C - 29⁰C and the min temp 18⁰C to 20⁰C Sky is likely to be mainly cloudy the coming week Relative Humidity is likely to range from 43% to 95%. Wind speed may reach upto 1-2 kmph Wind direction will be mostly easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<p align="center">Field crops</p> <p align="center">Undertake timely harvest of kharif crops with safe storage of produce. Seed treatment with fungicides (@ 2 to 3 g/kg seed) prevents seed borne diseases, promotes better germination and crop stand leading to higher productivity in all rabi crops</p>			
TRC/WRC paddy	Ripening stage	-	Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
Toria	Land preparation	-	Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region
Rabi maize	Land preparation		Land preparation should be started for rabi maize. It can be grown in all types of soils having adequate provision of drainage. Field should be ploughed properly so as to expose the pupae of red hairy caterpillar. Sow certified seeds from a reliable source to prevent seed rot and seedling blight. If one wants to use his own seed, seed treatment is a must. Some of the varieties recommended for our region are HQPM-1, Allrounder and RCM-76.
<p align="center">Horticultural crop</p> <p align="center">Spread well decomposed cowdung 5 to 6 tones per hectare for the main field and mix it well in the soil by cultivating the field. Keep the field clean by collecting the stubbles and unwanted materials.</p>			
Citrus		Leaf miner	Proper management should be done for leaf miner. Monitor pest regularly, manage ants and whitewash of the trunk should be done.


	Vegetables			<p><i>Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done.</i></p> <p><i>Suitable varieties for our region:</i></p> <ol style="list-style-type: none"> 1. <i>Cabbage – Green express (Hybrid), Rareball (Hybrid), Pride of India (OP).</i> 2. <i>Broccoli – Pushpa, princess.</i> 3. <i>Tomato – Pusa ruby, Arka vikash, Punjab swari.</i> 4. <i>Knolkhol – White Vienna.</i> 5. <i>Radish – Japanese white, Pusa chetki.</i> 6. <i>Pea – Arkel</i>
	Poultry	<p>Prevailing high humidity may cause ammonia fumes which irritate the respiratory tract and eyes of the poultry. Good ventilation provides a comfortable environment without draft. Farmers are advised to provide clean drinking water round the clock and water troughs/ feeders should be regularly cleaned. Antibiotics or other stress medications may be added to water if desired.</p>		
	Piggery	<p>Routine vaccination programme is recommended and once any disease is diagnosed treat immediately with antibiotics.</p>		
	Fishery	<p>Organic manure/ cattle manure may be applied @ 2 ton per ha to maintain the natural productivity of the ponds. To avoid losses, avoid application of manure at maximum water depth.</p>		



Integrated Agromet Advisory Service Bulletin from 29th Oct to 2nd Nov'16 Kiphire District

Bulletin No:86/2016

Weather summary of the preceding week			Weather forecast valid upto 2 nd Nov'16
<ul style="list-style-type: none"> Light rain occurred the past week Maximum and minimum temperatures ranged 29^oC to 31^oC and 17^oC to 20^oC, respectively. Relative humidity varied from 38% to 93%. Wind speed ranged from 1 to 2 kmph 			<ul style="list-style-type: none"> Probability of light rain the coming week. Max temp is likely to be 28^oC - 30^oC and the min temp 19^oC to 21^oC Sky is likely to be mainly cloudy the coming week Relative Humidity is likely to range from 43% to 93%. Wind speed may reach upto 1-2 kmph Wind direction will be mostly southeasterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<p align="center">Field crops</p> <p align="center">Undertake timely harvest of kharif crops with safe storage of produce. Seed treatment with fungicides (@ 2 to 3 g/kg seed) prevents seed borne diseases, promotes better germination and crop stand leading to higher productivity in all rabi crops</p>			
TRC/WRC paddy	Ripening stage	-	Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
Toria	Land preparation	-	Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region
Rabi maize	Land preparation		Land preparation should be started for rabi maize. It can be grown in all types of soils having adequate provision of drainage. Field should be ploughed properly so as to expose the pupae of red hairy caterpillar. Sow certified seeds from a reliable source to prevent seed rot and seedling blight. If one wants to use his own seed, seed treatment is a must. Some of the varieties recommended for our region are HQPM-1, Allrounder and RCM-76.
<p align="center">Horticultural crop</p> <p align="center">Spread well decomposed cowdung 5 to 6 tones per hectare for the main field and mix it well in the soil by cultivating the field. Keep the field clean by collecting the stubbles and unwanted materials.</p>			
Citrus		Leaf miner	Proper management should be done for leaf miner. Monitor pest regularly, manage ants and whitewash of the trunk should be done.
Vegetables			Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done.


				<p><i>Suitable varieties for our region:</i></p> <ol style="list-style-type: none"> 1. <i>Cabbage – Green express (Hybrid), Rareball (Hybrid), Pride of India (OP).</i> 2. <i>Broccoli – Pushpa, princess.</i> 3. <i>Tomato – Pusa ruby, Arka vikash, Punjab swari.</i> 4. <i>Knolkhol – White Vienna.</i> 5. <i>Radish – Japanese white, Pusa chetki.</i> 6. <i>Pea – Arkel</i>
	<i>Poultry</i>	<p>Prevailing high humidity may cause ammonia fumes which irritate the respiratory tract and eyes of the poultry. Good ventilation provides a comfortable environment without draft. Farmers are advised to provide clean drinking water round the clock and water troughs/ feeders should be regularly cleaned. Antibiotics or other stress medications may be added to water if desired.</p>		
	<i>Piggery</i>	<p>Routine vaccination programme is recommended and once any disease is diagnosed treat immediately with antibiotics.</p>		
	<i>Fishery</i>	<p>Organic manure/ cattle manure may be applied @ 2 ton per ha to maintain the natural productivity of the ponds. To avoid losses, avoid application of manure at maximum water depth.</p>		



Integrated Agromet Advisory Service Bulletin from 29th Oct to 2nd Nov'16 Kohima District

Bulletin No:86/2016

Weather summary of the preceding week			Weather forecast valid upto 2 nd Nov'16
<ul style="list-style-type: none"> Light rain occurred the past week Maximum and minimum temperatures ranged 27^oC to 28^oC and 17^oC to 18^oC, respectively. Relative humidity varied from 36% to 94%. Wind speed ranged from 1 to 2 kmph 			<ul style="list-style-type: none"> Probability of light rain the coming week. Max temp is likely to be 28^oC - 29^oC and the min temp 19^oC to 20^oC Sky is likely to be mainly cloudy the coming week Relative Humidity is likely to range from 45% to 95%. Wind speed may reach upto 2-3 kmph Wind direction will be mostly easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<p align="center">Field crops</p> <p align="center">Undertake timely harvest of kharif crops with safe storage of produce. Seed treatment with fungicides (@ 2 to 3 g/kg seed) prevents seed borne diseases, promotes better germination and crop stand leading to higher productivity in all rabi crops</p>			
TRC/WRC paddy	Ripening stage	-	Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
Toria	Land preparation	-	Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region
Rabi maize	Land preparation		Land preparation should be started for rabi maize. It can be grown in all types of soils having adequate provision of drainage. Field should be ploughed properly so as to expose the pupae of red hairy caterpillar. Sow certified seeds from a reliable source to prevent seed rot and seedling blight. If one wants to use his own seed, seed treatment is a must. Some of the varieties recommended for our region are HQPM-1, Allrounder and RCM-76.
<p align="center">Horticultural crop</p> <p align="center">Spread well decomposed cowdung 5 to 6 tones per hectare for the main field and mix it well in the soil by cultivating the field. Keep the field clean by collecting the stubbles and unwanted materials.</p>			
Citrus		Leaf miner	Proper management should be done for leaf miner. Monitor pest regularly, manage ants and whitewash of the trunk should be done.


	Vegetables			<p>Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done.</p> <p>Suitable varieties for our region:</p> <ol style="list-style-type: none"> 1. Cabbage – Green express (Hybrid), Rareball (Hybrid), Pride of India (OP). 2. Broccoli – Pushpa, princess. 3. Tomato – Pusa ruby, Arka vikash, Punjab swari. 4. Knolkhol – White Vienna. 5. Radish – Japanese white, Pusa chetki. 6. Pea – Arkel
	Poultry	Prevailing high humidity may cause ammonia fumes which irritate the respiratory tract and eyes of the poultry. Good ventilation provides a comfortable environment without draft. Farmers are advised to provide clean drinking water round the clock and water troughs/ feeders should be regularly cleaned. Antibiotics or other stress medications may be added to water if desired.		
	Piggery	Routine vaccination programme is recommended and once any disease is diagnosed treat immediately with antibiotics.		
	Fishery	Organic manure/ cattle manure may be applied @ 2 ton per ha to maintain the natural productivity of the ponds. To avoid losses, avoid application of manure at maximum water depth.		



Integrated Agromet Advisory Service Bulletin from 29th Oct to 2nd Nov'16 Longleng District

Bulletin No:86/2016

Weather summary of the preceding week			Weather forecast valid upto 2 nd Nov'16
<ul style="list-style-type: none"> No rain occurred the past week Maximum and minimum temperatures ranged 28⁰C to 31⁰C and 18⁰C to 20⁰C, respectively. Relative humidity varied from 37% to 95%. Wind speed ranged from 2 to 3 kmph 			<ul style="list-style-type: none"> Probability of light rain the coming week. Max temp is likely to be 27⁰C - 28⁰C and the min temp 17⁰C to 19⁰C Sky is likely to be mainly cloudy the coming week Relative Humidity is likely to range from 39% to 95%. Wind speed may reach upto 1- 2 kmph Wind direction will be mostly easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<p>Field crops</p> <p>Undertake timely harvest of kharif crops with safe storage of produce. Seed treatment with fungicides (@ 2 to 3 g/kg seed) prevents seed borne diseases, promotes better germination and crop stand leading to higher productivity in all rabi crops</p>			
TRC/WRC paddy	Ripening stage	-	Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
Toria	Land preparation	-	Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region
Rabi maize	Land preparation		Land preparation should be started for rabi maize. It can be grown in all types of soils having adequate provision of drainage. Field should be ploughed properly so as to expose the pupae of red hairy caterpillar. Sow certified seeds from a reliable source to prevent seed rot and seedling blight. If one wants to use his own seed, seed treatment is a must. Some of the varieties recommended for our region are HQPM-1, Allrounder and RCM-76.
<p>Horticultural crop</p> <p>Spread well decomposed cowdung 5 to 6 tones per hectare for the main field and mix it well in the soil by cultivating the field. Keep the field clean by collecting the stubbles and unwanted materials.</p>			
Citrus		Leaf miner	Proper management should be done for leaf miner. Monitor pest regularly, manage ants and whitewash of the trunk should


				<i>be done.</i>
	Vegetables			<p><i>Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done.</i></p> <p><i>Suitable varieties for our region:</i></p> <ol style="list-style-type: none"> 1. <i>Cabbage – Green express (Hybrid), Rareball (Hybrid), Pride of India (OP).</i> 2. <i>Broccoli – Pushpa, princess.</i> 3. <i>Tomato – Pusa ruby, Arka vikash, Punjab swari.</i> 4. <i>Knolkhol – White Vienna.</i> 5. <i>Radish – Japanese white, Pusa chetki.</i> 6. <i>Pea – Arkel</i>
	Poultry	Prevailing high humidity may cause ammonia fumes which irritate the respiratory tract and eyes of the poultry. Good ventilation provides a comfortable environment without draft. Farmers are advised to provide clean drinking water round the clock and water troughs/ feeders should be regularly cleaned. Antibiotics or other stress medications may be added to water if desired.		
	Piggery	Routine vaccination programme is recommended and once any disease is diagnosed treat immediately with antibiotics.		
	Fishery	Organic manure/ cattle manure may be applied @ 2 ton per ha to maintain the natural productivity of the ponds. To avoid losses, avoid application of manure at maximum water depth.		

Integrated Agromet Advisory Service Bulletin from 29th Oct to 2nd Nov'16

Mokokchung District

Bulletin No: 86/2016

Weather summary of the preceding week			Weather forecast valid upto 2 nd Nov'16
<ul style="list-style-type: none"> Light rain occurred the past week Maximum and minimum temperatures ranged 28⁰C to 30⁰C and 19⁰C to 20⁰C, respectively. Relative humidity varied from 38% to 94%. Wind speed ranged from 1 to 2 kmph 			<ul style="list-style-type: none"> Probability of light rain the coming week. Max temp is likely to be 29⁰C - 30⁰C and the min temp 18⁰C to 19⁰C Sky is likely to be mainly cloudy the coming week Relative Humidity is likely to range from 41% to 95%. Wind speed may reach upto 1-2 kmph Wind direction will be mostly easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<p align="center">Field crops</p> <p align="center">Undertake timely harvest of kharif crops with safe storage of produce.</p> <p align="center">Seed treatment with fungicides (@ 2 to 3 g/kg seed) prevents seed borne diseases, promotes better germination and crop stand leading to higher productivity in all rabi crops</p>			
TRC/WRC paddy	Ripening stage	-	Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
Toria	Land preparation	-	Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region
Rabi maize	Land preparation		Land preparation should be started for rabi maize. It can be grown in all types of soils having adequate provision of drainage. Field should be ploughed properly so as to expose the pupae of red hairy caterpillar. Sow certified seeds from a reliable source to prevent seed rot and seedling blight. If one wants to use his own seed, seed treatment is a must. Some of the varieties recommended for our region are HQPM-1, Allrounder and RCM-76.
<p align="center">Horticultural crop</p> <p align="center">Spread well decomposed cowdung 5 to 6 tones per hectare for the main field and mix it well in the soil by cultivating the field. Keep the field clean by collecting the stubbles and unwanted materials.</p>			
Citrus		Leaf miner	Proper management should be done for leaf miner. Monitor pest regularly, manage ants and whitewash of the trunk should be done.
Vegetables			Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be


			<p>done.</p> <p><i>Suitable varieties for our region:</i></p> <ol style="list-style-type: none"> 1. <i>Cabbage – Green express (Hybrid), Rareball (Hybrid), Pride of India (OP).</i> 2. <i>Broccoli – Pushpa, princess.</i> 3. <i>Tomato – Pusa ruby, Arka vikash, Punjab swari.</i> 4. <i>Knolkhol – White Vienna.</i> 5. <i>Radish – Japanese white, Pusa chetki.</i> 6. <i>Pea – Arkel</i>
	<i>Poultry</i>	<p>Prevailing high humidity may cause ammonia fumes which irritate the respiratory tract and eyes of the poultry. Good ventilation provides a comfortable environment without draft. Farmers are advised to provide clean drinking water round the clock and water troughs/ feeders should be regularly cleaned. Antibiotics or other stress medications may be added to water if desired.</p>	
	<i>Piggery</i>	<p>Routine vaccination programme is recommended and once any disease is diagnosed treat immediately with antibiotics.</p>	
	<i>Fishery</i>	<p>Organic manure/ cattle manure may be applied @ 2 ton per ha to maintain the natural productivity of the ponds. To avoid losses, avoid application of manure at maximum water depth.</p>	



Integrated Agromet Advisory Service Bulletin from 29th Oct to 2nd Nov'16 Mon District

Bulletin No:86/2016

Weather summary of the preceding week			Weather forecast valid upto 2 nd Nov'16
<ul style="list-style-type: none"> Light rain occurred the past week Maximum and minimum temperatures ranged 29^oC to 31^oC and 18^oC to 19^oC, respectively. Relative humidity varied from 36% to 92% Wind speed ranged from 1 to 2 kmph 			<ul style="list-style-type: none"> Probability of light rain the coming week. Max temp is likely to be 29^oC - 30^oC and the min temp 19^oC to 20^oC Sky is likely to be mainly cloudy the coming week Relative Humidity is likely to range from 39% to 94%. Wind speed may reach upto 1- 2 kmph Wind direction will be mostly easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<p>Field crops</p> <p>Undertake timely harvest of kharif crops with safe storage of produce.</p> <p>Seed treatment with fungicides (@ 2 to 3 g/kg seed) prevents seed borne diseases, promotes better germination and crop stand leading to higher productivity in all rabi crops</p>			
TRC/WRC paddy	Ripening stage		Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
Toria	Land preparation	-	Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region
Rabi maize	Land preparation		Land preparation should be started for rabi maize. It can be grown in all types of soils having adequate provision of drainage. Field should be ploughed properly so as to expose the pupae of red hairy caterpillar. Sow certified seeds from a reliable source to prevent seed rot and seedling blight. If one wants to use his own seed, seed treatment is a must. Some of the varieties recommended for our region are HQPM-1, Allrounder and RCM-76.
<p>Horticultural crop</p> <p>Spread well decomposed cowdung 5 to 6 tones per hectare for the main field and mix it well in the soil by cultivating the field. Keep the field clean by collecting the stubbles and unwanted materials.</p>			
Citrus		Leaf miner	Proper management should be done for leaf miner. Monitor pest regularly, manage ants and whitewash of the trunk should be done.


	Vegetables			<p><i>Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done.</i></p> <p><i>Suitable varieties for our region:</i></p> <ol style="list-style-type: none"> 1. <i>Cabbage – Green express (Hybrid), Rareball (Hybrid), Pride of India (OP).</i> 2. <i>Broccoli – Pushpa, princess.</i> 3. <i>Tomato – Pusa ruby, Arka vikash, Punjab swari.</i> 4. <i>Knolkhol – White Vienna.</i> 5. <i>Radish – Japanese white, Pusa chetki.</i> 6. <i>Pea – Arkel</i>
	Poultry	Proper sanitary measures should be done to control the diseases. Slaughtering of all apparently ailing birds, segregating of in - contact in group of 10 to 15 each; removal of all infective materials such as droppings, residues of poultry cleanliness.		
	Piggery	It is vital that piglets should be nurse soon after birth since colostrum gives them some protection against infection the first few weeks.		
	Fishery	Pond water should not just maintain at a certain depth; its quality must also be kept high to ensure optimal growth of the culture organism.		



Integrated Agromet Advisory Service Bulletin from 29th Oct to 2nd Nov'16 Peren District

Bulletin No:86/2016

Weather summary of the preceding week			Weather forecast valid upto 2 nd Nov'16
<ul style="list-style-type: none"> Light rain occurred the past week Maximum and minimum temperatures ranged 28^oC to 30^oC and 19^oC to 20^oC, respectively. Relative humidity varied from 37% to 94%. Wind speed ranged from 1 to 2 kmph 			<ul style="list-style-type: none"> Probability of light rain the coming week. Max temp is likely to be 28^oC - 30^oC and the min temp 19^oC to 20^oC Sky is likely to be mainly cloudy the coming week Relative Humidity is likely to range from 44% to 95%. Wind speed may reach upto 2-4 kmph Wind direction will be mostly easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<p align="center">Field crops</p> <p align="center">Undertake timely harvest of kharif crops with safe storage of produce. Seed treatment with fungicides (@ 2 to 3 g/kg seed) prevents seed borne diseases, promotes better germination and crop stand leading to higher productivity in all rabi crops</p>			
TRC/WRC paddy	Ripening stage	-	<i>Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds</i>
Toria	Land preparation	-	<i>Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region</i>
Rabi maize	Land preparation		<i>Land preparation should be started for rabi maize. It can be grown in all types of soils having adequate provision of drainage. Field should be ploughed properly so as to expose the pupae of red hairy caterpillar. Sow certified seeds from a reliable source to prevent seed rot and seedling blight. If one wants to use his own seed, seed treatment is a must. Some of the varieties recommended for our region are HQPM-1, Allrounder and RCM-76.</i>
<p align="center">Horticultural crop</p> <p align="center">Spread well decomposed cowdung 5 to 6 tones per hectare for the main field and mix it well in the soil by cultivating the field. Keep the field clean by collecting the stubbles and unwanted materials.</p>			
Citrus		Leaf miner	<i>Proper management should be done for leaf miner. Monitor pest regularly, manage ants and whitewash of the trunk should be done.</i>


	Vegetables			<p><i>Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done.</i></p> <p><i>Suitable varieties for our region:</i></p> <ol style="list-style-type: none"> 1. Cabbage – Green express (Hybrid), Rareball (Hybrid), Pride of India (OP). 2. Broccoli – Pushpa, princess. 3. Tomato – Pusa ruby, Arka vikash, Punjab swari. 4. Knolkhol – White Vienna. 5. Radish – Japanese white, Pusa chetki. 6. Pea – Arkel
	Poultry	<p>Prevailing high humidity may cause ammonia fumes which irritate the respiratory tract and eyes of the poultry. Good ventilation provides a comfortable environment without draft. Farmers are advised to provide clean drinking water round the clock and water troughs/ feeders should be regularly cleaned. Antibiotics or other stress medications may be added to water if desired.</p>		
	Piggery	<p>Routine vaccination programme is recommended and once any disease is diagnosed treat immediately with antibiotics.</p>		
	Fishery	<p>Organic manure/ cattle manure may be applied @ 2 ton per ha to maintain the natural productivity of the ponds. To avoid losses, avoid application of manure at maximum water depth.</p>		



Integrated Agromet Advisory Service Bulletin from 29th Oct to 2nd Nov'16 Phek District

Bulletin No:86/2016

Weather summary of the preceding week			Weather forecast valid upto 2 nd Nov'16
<ul style="list-style-type: none"> Light rain occurred the past week Maximum and minimum temperatures ranged 29⁰C to 31⁰C and 18⁰C to 20⁰C, respectively. Relative humidity varied from 39% to 95%. Wind speed ranged from 1 to 2 kmph 			<ul style="list-style-type: none"> Probability of light rain the coming week. Max temp is likely to be 29⁰C - 30⁰C and the min temp 18⁰C to 20⁰C Sky is likely to be mainly cloudy the coming week Relative Humidity is likely to range from 44% to 91%. Wind speed may reach upto 2-3 kmph Wind direction will be mostly easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<p align="center">Field crops</p> <p align="center">Undertake timely harvest of kharif crops with safe storage of produce. Seed treatment with fungicides (@ 2 to 3 g/kg seed) prevents seed borne diseases, promotes better germination and crop stand leading to higher productivity in all rabi crops</p>			
TRC/WRC paddy	Ripening stage	-	<i>Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds</i>
Toria	Land preparation	-	<i>Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region</i>
Rabi maize	Land preparation		<i>Land preparation should be started for rabi maize. It can be grown in all types of soils having adequate provision of drainage. Field should be ploughed properly so as to expose the pupae of red hairy caterpillar. Sow certified seeds from a reliable source to prevent seed rot and seedling blight. If one wants to use his own seed, seed treatment is a must. Some of the varieties recommended for our region are HQPM-1, Allrounder and RCM-76.</i>
<p align="center">Horticultural crop</p> <p align="center">Spread well decomposed cowdung 5 to 6 tones per hectare for the main field and mix it well in the soil by cultivating the field. Keep the field clean by collecting the stubbles and unwanted materials.</p>			
Citrus		Leaf miner	<i>Proper management should be done for leaf miner. Monitor pest regularly, manage ants and whitewash of the trunk should be done.</i>


	Vegetables			<p><i>Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done.</i></p> <p><i>Suitable varieties for our region:</i></p> <ol style="list-style-type: none"> 1. Cabbage – Green express (Hybrid), Rareball (Hybrid), Pride of India (OP). 2. Broccoli – Pushpa, princess. 3. Tomato – Pusa ruby, Arka vikash, Punjab swari. 4. Knolkhol – White Vienna. 5. Radish – Japanese white, Pusa chetki. 6. Pea – Arkel
	Poultry	Prevailing high humidity may cause ammonia fumes which irritate the respiratory tract and eyes of the poultry. Good ventilation provides a comfortable environment without draft. Farmers are advised to provide clean drinking water round the clock and water troughs/ feeders should be regularly cleaned. Antibiotics or other stress medications may be added to water if desired.		
	Piggery	Routine vaccination programme is recommended and once any disease is diagnosed treat immediately with antibiotics.		
	Fishery	Organic manure/ cattle manure may be applied @ 2 ton per ha to maintain the natural productivity of the ponds. To avoid losses, avoid application of manure at maximum water depth.		



Integrated Agromet Advisory Service Bulletin from 29th Oct to 2nd Nov'16 Tuensang District

Bulletin No:86/2016

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 2nd Nov'16</i>
<ul style="list-style-type: none"> Light rain occurred the past week Maximum and minimum temperatures ranged 28⁰C to 30⁰C and 16⁰C to 18⁰C, respectively. Relative humidity varied from 40% to 92%. Wind speed ranged from 1 to 2 kmph 			<ul style="list-style-type: none"> Probability of light rain the coming week. Max temp is likely to be 28⁰C - 30⁰C and the min temp 19⁰C to 20⁰C Sky is likely to be mainly cloudy the coming week Relative Humidity is likely to range from 42% to 92%. Wind speed may reach upto 1-2 kmph Wind direction will be mostly south easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<p style="text-align: center;">Field crops</p> <p style="text-align: center;">Undertake timely harvest of kharif crops with safe storage of produce. Seed treatment with fungicides (@ 2 to 3 g/kg seed) prevents seed borne diseases, promotes better germination and crop stand leading to higher productivity in all rabi crops</p>			
TRC/WRC paddy	Ripening stage	-	<i>Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds</i>
Toria	Land preparation	-	<i>Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region</i>
Rabi maize	Land preparation		<i>Land preparation should be started for rabi maize. It can be grown in all types of soils having adequate provision of drainage. Field should be ploughed properly so as to expose the pupae of red hairy caterpillar. Sow certified seeds from a reliable source to prevent seed rot and seedling blight. If one wants to use his own seed, seed treatment is a must. Some of the varieties recommended for our region are HQPM-1, Allrounder and RCM-76.</i>
<p style="text-align: center;">Horticultural crop</p> <p style="text-align: center;">Spread well decomposed cowdung 5 to 6 tones per hectare for the main field and mix it well in the soil by cultivating the field. Keep the field clean by collecting the stubbles and unwanted materials.</p>			
Citrus		Leaf miner	<i>Proper management should be done for leaf miner. Monitor pest regularly, manage ants and whitewash of the trunk should be done.</i>


	Vegetables		<p>Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done.</p> <p>Suitable varieties for our region:</p> <ol style="list-style-type: none"> 1. Cabbage – Green express (Hybrid), Rareball (Hybrid), Pride of India (OP). 2. Broccoli – Pushpa, princess. 3. Tomato – Pusa ruby, Arka vikash, Punjab swari. 4. Knolkhol – White Vienna. 5. Radish – Japanese white, Pusa chetki. 6. Pea – Arkel
	Poultry	<p>Prevailing high humidity may cause ammonia fumes which irritate the respiratory tract and eyes of the poultry. Good ventilation provides a comfortable environment without draft. Farmers are advised to provide clean drinking water round the clock and water troughs/ feeders should be regularly cleaned. Antibiotics or other stress medications may be added to water if desired.</p>	
	Piggery	<p>Routine vaccination programme is recommended and once any disease is diagnosed treat immediately with antibiotics.</p>	
	Fishery	<p>Organic manure/ cattle manure may be applied @ 2 ton per ha to maintain the natural productivity of the ponds. To avoid losses, avoid application of manure at maximum water depth.</p>	



Integrated Agromet Advisory Service Bulletin from 29th Oct to 2nd Nov'16 Wokha District

Bulletin No:86/2016

Weather summary of the preceding week			Weather forecast valid upto 2 nd Nov'16
<ul style="list-style-type: none"> No rain occurred the past week Maximum and minimum temperatures ranged 28⁰C to 31⁰C and 19⁰C to 20⁰C, respectively. Relative humidity varied from 40% to 91%. Wind speed ranged from 1 to 2 kmph 			<ul style="list-style-type: none"> Probability of light rain the coming week. Max temp is likely to be 27⁰C - 29⁰C and the min temp 17⁰C to 18⁰C Sky is likely to be mainly cloudy the coming week Relative Humidity is likely to range from 43% to 95%. Wind speed may reach upto 1-2 kmph Wind direction will be mostly south easterly
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<p align="center">Field crops</p> <p align="center">Undertake timely harvest of kharif crops with safe storage of produce. Seed treatment with fungicides (@ 2 to 3 g/kg seed) prevents seed borne diseases, promotes better germination and crop stand leading to higher productivity in all rabi crops</p>			
TRC/WRC paddy	Ripening stage	-	<i>Avoid excessive drying of paddy to avoid breakage of the grains. Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds</i>
Toria	Land preparation	-	<i>Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region</i>
Rabi maize	Land preparation		<i>Land preparation should be started for rabi maize. It can be grown in all types of soils having adequate provision of drainage. Field should be ploughed properly so as to expose the pupae of red hairy caterpillar. Sow certified seeds from a reliable source to prevent seed rot and seedling blight. If one wants to use his own seed, seed treatment is a must. Some of the varieties recommended for our region are HQPM-1, Allrounder and RCM-76.</i>
<p align="center">Horticultural crop</p> <p align="center">Spread well decomposed cowdung 5 to 6 tones per hectare for the main field and mix it well in the soil by cultivating the field. Keep the field clean by collecting the stubbles and unwanted materials.</p>			
Citrus		Leaf miner	<i>Proper management should be done for leaf miner. Monitor pest regularly, manage ants and whitewash of the trunk should</i>

				<i>be done.</i>
	Vegetables			<p><i>Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done.</i></p> <p><i>Suitable varieties for our region:</i></p> <ol style="list-style-type: none"> <i>1. Cabbage – Green express (Hybrid), Rareball (Hybrid), Pride of India (OP).</i> <i>2. Broccoli – Pushpa, princess.</i> <i>3. Tomato – Pusa ruby, Arka vikash, Punjab swari.</i> <i>4. Knolkhol – White Vienna.</i> <i>5. Radish – Japanese white, Pusa chetki.</i> <i>6. Pea – Arkel</i>
	Poultry	Prevailing high humidity may cause ammonia fumes which irritate the respiratory tract and eyes of the poultry. Good ventilation provides a comfortable environment without draft. Farmers are advised to provide clean drinking water round the clock and water troughs/ feeders should be regularly cleaned. Antibiotics or other stress medications may be added to water if desired.		
	Piggery	Routine vaccination programme is recommended and once any disease is diagnosed treat immediately with antibiotics.		
	Fishery	Organic manure/ cattle manure may be applied @ 2 ton per ha to maintain the natural productivity of the ponds. To avoid losses, avoid application of manure at maximum water depth.		



--	--