




# Integrated Agromet Advisory Service Bulletin from 5<sup>th</sup> to 9<sup>th</sup> Oct'16 Dimapur District

Bulletin No:80/2016

Weather summary of the preceding week			Weather forecast valid upto 9 <sup>th</sup> Oct'16
<ul style="list-style-type: none"> <li>light rain occurred the past week</li> <li>Maximum and minimum temperatures ranged 27<sup>0</sup>C to 30<sup>0</sup>C and 19<sup>0</sup>C to 21<sup>0</sup>C, respectively.</li> <li>Relative humidity varied from 49% to 93%.</li> <li>Wind speed ranged from 1 to 2 kmph</li> </ul>			<ul style="list-style-type: none"> <li>Probability of light rain the coming week.</li> <li><b>Max temp</b> is likely to be 33<sup>0</sup>C - 34<sup>0</sup>C and the <b>min temp</b> 24<sup>0</sup>C to 25<sup>0</sup>C</li> <li><b>Sky is likely to be mainly cloudy</b> the coming week</li> <li><b>Relative Humidity</b> is likely to range from 56% to 90%.</li> <li><b>Wind speed</b> may reach upto 1-2 kmph</li> <li><b>Wind direction</b> will be mostly southeasterly</li> </ul>
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<p align="center"><b>Field crops</b></p> <p>Timely field sanitation should be done for all the crops. Always use freshly prepared neem seed kernel extract. Apply pesticides only when required. Don't apply more than the recommended dose of the pesticide</p>			
<b>Jhum paddy</b>	Harvesting stage		<i>During sunny days harvest the early and early medium duration varieties when 80% of the grains matured in a panicle. Harvest, process, dry and pack different paddy varieties separately without mixing for better price for their produce</i>
<b>TRC/WRC paddy</b>	Milk development stage	Brown spot	<i>Destroy the infested residue, stubbles, grass and weeds present in the bunds of the field as they act as a source for the infection</i>
		Bacterial blight	<i>If the infestation is more, spray Streptocycline 2gm/5lt of water.</i>
		Blast	<i>For rice blast management, destroy the infested residue, stubbles, grass and weeds present in the bunds of the field as they act as a source for the infection.</i>
<b>Soybean</b>	Harvesting stage		<i>If you're storing soybeans for next season, screen them prior to loading them into the bin</i>
<b>Toria</b>	Land preparation	-	<i>Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region</i>
<b>Rabi maize</b>	Land preparation		<i>Land preparation should be started for rabi maize. It can be grown in all types of soils having adequate provision of drainage. Field should be ploughed properly so as to expose the pupae of red hairy caterpillar. Sow certified seeds from a reliable source to prevent seed rot and seedling blight. If one wants to use his own seed, seed treatment is a must. Some of the varieties recommended for our region are HQPM-1,</i>

				<i>Allrounder and RCM-76.</i>
	<b><i>Horticultural crops</i></b>	Harvesting stage	-	<i>To avoid loss because of the shattering of pods, the crop should be harvested before it is dead ripe.</i>
	<b><i>Citrus</i></b>			
	<b><i>Vegetables</i></b>			<i>Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done.</i> <i>Suitable varieties for our region:</i> <ol style="list-style-type: none"> <li><i>1. Cabbage – Green express (Hybrid), Rareball (Hybrid), Pride of India (OP).</i></li> <li><i>2. Broccoli – Pushpa, princess.</i></li> <li><i>3. Tomato – Pusa ruby, Arka vikash, Punjab swari.</i></li> <li><i>4. Knolkhol – White Vienna.</i></li> <li><i>5. Radish – Japanese white, Pusa chetki.</i></li> <li><i>6. Pea – Arkel</i></li> </ol>
	<b><i>Poultry</i></b>	Good ventilation in the chicken shed is must, because chickens naturally produce a lot of moisture in their breath and droppings. Poor ventilation inside their house may create an ammonia build-up, which will affect the		


		poultry's eyes and respiratory system and makes them sick.		
	<b>Piggery</b>	A clean, sanitary environment provides the best prevention for internal and external parasite which can be serious problems. For diseases that can be prevented by vaccination, a veterinarian should be contacted to provide such services routinely.		
	<b>Fishery</b>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media		
	<b>Fishery</b>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media		



# Integrated Agromet Advisory Service Bulletin from 5<sup>th</sup> to 9<sup>th</sup> Oct'16 Kiphire District

Bulletin No:80/2016

Weather summary of the preceding week			Weather forecast valid upto 9 <sup>th</sup> Oct'16
<ul style="list-style-type: none"> <li>Light rain occurred the past week</li> <li>Maximum and minimum temperatures ranged 26<sup>0</sup>C to 30<sup>0</sup>C and 19<sup>0</sup>C to 22<sup>0</sup>C, respectively.</li> <li>Relative humidity varied from 56% to 93%.</li> <li>Wind speed ranged from 1 to 2 kmph</li> </ul>			<ul style="list-style-type: none"> <li>Probability of light rain the coming week.</li> <li><b>Max temp</b> is likely to be 27<sup>0</sup>C - 29<sup>0</sup>C and the <b>min temp</b> 20<sup>0</sup>C to 22<sup>0</sup>C</li> <li><b>Sky is likely to be mainly cloudy</b> the coming week</li> <li><b>Relative Humidity</b> is likely to range from 55% to 90%.</li> <li><b>Wind speed</b> may reach upto 2-4 kmph</li> <li><b>Wind direction</b> will be mostly southeasterly</li> </ul>
Main Crops	Stage	Pest/Diseases	Agro-meteorological Advisories
<p align="center"><b>Field crops</b></p> <p>Timely field sanitation should be done for all the crops. Always use freshly prepared neem seed kernel extract. Apply pesticides only when required. Don't apply more than the recommended dose of the pesticide</p>			
<b>Jhum paddy</b>	Harvesting stage		<i>During sunny days harvest the early and early medium duration varieties when 80% of the grains matured in a panicle. Harvest, process, dry and pack different paddy varieties separately without mixing for better price for their produce</i>
<b>TRC/WRC paddy</b>	Milk development stage	Brown spot	<i>Destroy the infested residue, stubbles, grass and weeds present in the bunds of the field as they act as a source for the infection</i>
		Bacterial blight	<i>If the infestation is more, spray Streptocycline 2gm/5lt of water.</i>
		Blast	<i>For rice blast management, destroy the infested residue, stubbles, grass and weeds present in the bunds of the field as they act as a source for the infection.</i>
<b>Soybean</b>	Harvesting stage		<i>If you're storing soybeans for next season, screen them prior to loading them into the bin</i>
<b>Toria</b>	Land preparation	-	<i>Field preparation should be started for <b>toria</b> and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region</i>
<b>Rabi maize</b>	Land preparation		<p><i>Land preparation should be started for rabi maize. It can be grown in all types of soils having adequate provision of drainage. Field should be ploughed properly so as to expose the pupae of red hairy caterpillar. Sow certified seeds from a reliable source to prevent seed rot and seedling blight. If one wants to use his own seed, seed treatment is a must.</i></p> <p><i>Some of the varieties recommended for our region are HQPM-1, Allrounder and RCM-76.</i></p>
<b>Horticultural crops</b>			


	<b>Citrus</b>		Trunk borer	<i>To kill the trunk borer grubs, clean the bored holes of the infested trunk with iron wire and insert a cotton swab soaked in petrol or kerosene and plug with mud.</i>
	<b>Vegetables</b>			<i>Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done.</i>  <i>Suitable varieties for our region:</i> <ol style="list-style-type: none"> <li>1. <i>Cabbage – Green express (Hybrid), Rareball (Hybrid), Pride of India (OP).</i></li> <li>2. <i>Broccoli – Pushpa, princess.</i></li> <li>3. <i>Tomato – Pusa ruby, Arka vikash, Punjab swari.</i></li> <li>4. <i>Knolkhol – White Vienna.</i></li> <li>5. <i>Radish – Japanese white, Pusa chetki.</i></li> <li>6. <i>Pea – Arkel</i></li> </ol>
	<b>Poultry</b>	Good ventilation in the chicken shed is must, because chickens naturally produce a lot of moisture in their breath and droppings. Poor ventilation inside their house may create an ammonia build-up, which will affect the poultry's eyes and respiratory system and makes them sick.		
	<b>Piggery</b>	A clean, sanitary environment provides the best prevention for internal and external parasite which can be serious problems. For diseases that can be prevented by vaccination, a veterinarian should be contacted to provide such services routinely.		
	<b>Fishery</b>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media		



## Integrated Agromet Advisory Service Bulletin from 5<sup>th</sup> to 9<sup>th</sup> Oct'16 Kohima District

Bulletin No:80/2016

Weather summary of the preceding week			Weather forecast valid upto 9 <sup>th</sup> Oct'16
<ul style="list-style-type: none"> <li>Medium rain occurred the past week</li> <li>Maximum and minimum temperatures ranged 26<sup>o</sup>C to 27<sup>o</sup>C and 18<sup>o</sup>C to 19<sup>o</sup>C, respectively.</li> <li>Relative humidity varied from 58% to 94%.</li> <li>Wind speed ranged from 1 to 2 kmph</li> </ul>			<ul style="list-style-type: none"> <li>Probability of medium rain the coming week.</li> <li><b>Max temp</b> is likely to be 26<sup>o</sup>C - 27<sup>o</sup>C and the <b>min temp</b> 20<sup>o</sup>C to 22<sup>o</sup>C</li> <li><b>Sky is likely to be mainly cloudy</b> the coming week</li> <li><b>Relative Humidity</b> is likely to range from 60% to 90%.</li> <li><b>Wind speed</b> may reach upto 1-2 kmph</li> <li><b>Wind direction</b> will be mostly easterly</li> </ul>
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<p><b>Field crops</b></p> <p>Timely field sanitation should be done for all the crops. Always use freshly prepared neem seed kernel extract. Apply pesticides only when required. Don't apply more than the recommended dose of the pesticide</p>			
<b>Jhum paddy</b>	Harvesting stage		<i>During sunny days harvest the early and early medium duration varieties when 80% of the grains matured in a panicle. Harvest, process, dry and pack different paddy varieties separately without mixing for better price for their produce</i>
<b>TRC/WRC paddy</b>	Milk development stage	Brown spot	<i>Destroy the infested residue, stubbles, grass and weeds present in the bunds of the field as they act as a source for the infection</i>
		Bacterial blight	<i>If the infestation is more, spray Streptocycline 2gm/5lt of water.</i>
		Blast	<i>For rice blast management, destroy the infested residue, stubbles, grass and weeds present in the bunds of the field as they act as a source for the infection.</i>
<b>Soybean</b>	Harvesting stage		<i>If you're storing soybeans for next season, screen them prior to loading them into the bin</i>
<b>Toria</b>	Land preparation	-	<i>Field preparation should be started for <b>toria</b> and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region</i>
<b>Rabi maize</b>	Land preparation		<i>Land preparation should be started for rabi maize. It can be grown in all types of soils having adequate provision of drainage. Field should be ploughed properly so as to expose the pupae of red hairy caterpillar. Sow certified seeds from a reliable source to prevent seed rot and seedling blight. If one wants to use his own seed, seed treatment is a must. Some of the varieties recommended for our region are HQPM-1,</i>

				<i>Allrounder and RCM-76.</i>
	<b><i>Horticultural crops</i></b>			
	<b><i>Citrus</i></b>		Trunk borer	<i>To kill the trunk borer grubs, clean the bored holes of the infested trunk with iron wire and insert a cotton swab soaked in petrol or kerosene and plug with mud.</i>
	<b><i>Vegetables</i></b>			<i>Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done.</i> <i>Suitable varieties for our region:</i> <ol style="list-style-type: none"> <li><i>1. Cabbage – Green express (Hybrid), Rareball (Hybrid), Pride of India (OP).</i></li> <li><i>2. Broccoli – Pushpa, princess.</i></li> <li><i>3. Tomato – Pusa ruby, Arka vikash, Punjab swari.</i></li> <li><i>4. Knolkhol – White Vienna.</i></li> <li><i>5. Radish – Japanese white, Pusa chetki.</i></li> <li><i>6. Pea – Arkel</i></li> </ol>
	<b><i>Poultry</i></b>	Good ventilation in the chicken shed is must, because chickens naturally produce a lot of moisture in their breath and droppings. Poor ventilation inside their house may create an ammonia build-up, which will affect the poultry's eyes and respiratory system and makes them sick.		
	<b><i>Piggery</i></b>	A clean, sanitary environment provides the best prevention for internal and external parasite which can be serious problems. For diseases that can be prevented by vaccination, a veterinarian should be contacted to provide such services routinely.		
	<b><i>Fishery</i></b>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media		






## Integrated Agromet Advisory Service Bulletin from 5<sup>th</sup> to 9<sup>th</sup> Oct'16 Longleng District

Bulletin No:80/2016

Weather summary of the preceding week			Weather forecast valid upto 9 <sup>th</sup> Oct'16
<ul style="list-style-type: none"> <li>Light rain occurred the past week</li> <li>Maximum and minimum temperatures ranged 26<sup>0</sup>C to 28<sup>0</sup>C and 18<sup>0</sup>C to 20<sup>0</sup>C, respectively.</li> <li>Relative humidity varied from 53% to 95%.</li> <li>Wind speed ranged from 2 to 3 kmph</li> </ul>			<ul style="list-style-type: none"> <li>Probability of light rain the coming week.</li> <li><b>Max temp</b> is likely to be 28<sup>0</sup>C - 29<sup>0</sup>C and the <b>min temp</b> 19<sup>0</sup>C to 20<sup>0</sup>C</li> <li><b>Sky is likely to be mainly cloudy</b> the coming week</li> <li><b>Relative Humidity</b> is likely to range from 55% to 84%.</li> <li><b>Wind speed</b> may reach upto 1- 2 kmph</li> <li><b>Wind direction</b> will be mostly southeasterly</li> </ul>
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<p><b>Field crops</b></p> <p>Timely field sanitation should be done for all the crops. Always use freshly prepared neem seed kernel extract. Apply pesticides only when required. Don't apply more than the recommended dose of the pesticide</p>			
<b>Jhum paddy</b>	Harvesting stage		During sunny days harvest the early and early medium duration varieties when 80% of the grains matured in a panicle. Harvest, process, dry and pack different paddy varieties separately without mixing for better price for their produce
<b>TRC/WRC paddy</b>	Milk development stage	Brown spot	Destroy the infested residue, stubbles, grass and weeds present in the bunds of the field as they act as a source for the infection
		Bacterial blight	If the infestation is more, spray Streptocycline 2gm/5lt of water.
		Blast	For rice blast management, destroy the infested residue, stubbles, grass and weeds present in the bunds of the field as they act as a source for the infection.
<b>Soybean</b>	Harvesting stage		If you're storing soybeans for next season, screen them prior to loading them into the bin
<b>Toria</b>	Land preparation	-	Field preparation should be started for <b>toria</b> and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region
<b>Rabi maize</b>	Land preparation		Land preparation should be started for rabi maize. It can be grown in all types of soils having adequate provision of drainage. Field should be ploughed properly so as to expose the pupae of red hairy caterpillar. Sow certified seeds from a reliable source to prevent seed rot and seedling blight. If one wants to use his own seed, seed



				<p><i>treatment is a must.</i></p> <p><i>Some of the varieties recommended for our region are HQPM-1, Allrounder and RCM-76.</i></p>
	<b>Horticultural crops</b>			
	<b>Citrus</b>		Trunk borer	<p><i>To kill the trunk borer grubs, clean the bored holes of the infested trunk with iron wire and insert a cotton swab soaked in petrol or kerosene and plug with mud.</i></p>
	<b>Vegetables</b>			<p><i>Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done.</i></p> <p><i>Suitable varieties for our region:</i></p> <ol style="list-style-type: none"> <li><i>1. Cabbage – Green express (Hybrid), Rareball (Hybrid), Pride of India (OP).</i></li> <li><i>2. Broccoli – Pushpa, princess.</i></li> <li><i>3. Tomato – Pusa ruby, Arka vikash, Punjab swari.</i></li> <li><i>4. Knolkhol – White Vienna.</i></li> <li><i>5. Radish – Japanese white, Pusa chetki.</i></li> <li><i>6. Pea – Arkel</i></li> </ol>
	<b>Poultry</b>	<p>Good ventilation in the chicken shed is must, because chickens naturally produce a lot of moisture in their breath and droppings. Poor ventilation inside their house may create an ammonia build-up, which will affect the poultry's eyes and respiratory system and makes them sick.</p>		
	<b>Piggery</b>	<p>A clean, sanitary environment provides the best prevention for internal and external parasite which can be serious problems. For diseases that can be prevented by vaccination, a veterinarian should be contacted to provide such services routinely.</p>		
	<b>Fishery</b>	<p>Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media</p>		

# Integrated Agromet Advisory Service Bulletin from 5<sup>th</sup> to 9<sup>th</sup> Oct'16

## Mokokchung District

Bulletin No: 80/2016

Weather summary of the preceding week			Weather forecast valid upto 9 <sup>th</sup> Oct'16
<ul style="list-style-type: none"> <li>Light rain occurred the past week</li> <li>Maximum and minimum temperatures ranged 29<sup>0</sup>C to 30<sup>0</sup>C and 16<sup>0</sup>C to 17<sup>0</sup>C, respectively.</li> <li>Relative humidity varied from 60% to 94%.</li> <li>Wind speed ranged from 1 to 2 kmph</li> </ul>			<ul style="list-style-type: none"> <li>Probability of light rain the coming week.</li> <li><b>Max temp</b> is likely to be 27<sup>0</sup>C - 29<sup>0</sup>C and the <b>min temp</b> 22<sup>0</sup>C to 23<sup>0</sup>C</li> <li><b>Sky is likely to be mainly cloudy</b> the coming week</li> <li><b>Relative Humidity</b> is likely to range from 44% to 90%.</li> <li><b>Wind speed</b> may reach upto 1-2 kmph</li> <li><b>Wind direction</b> will be mostly southeasterly</li> </ul>
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<p><b>Field crops</b></p> <p>Timely field sanitation should be done for all the crops. Always use freshly prepared neem seed kernel extract. Apply pesticides only when required. Don't apply more than the recommended dose of the pesticide</p>			
<b>Jhum paddy</b>	Harvesting stage		<i>During sunny days harvest the early and early medium duration varieties when 80% of the grains matured in a panicle. Harvest, process, dry and pack different paddy varieties separately without mixing for better price for their produce</i>
<b>TRC/WRC paddy</b>	Milk development stage	Brown spot	<i>Destroy the infested residue, stubbles, grass and weeds present in the bunds of the field as they act as a source for the infection</i>
		Bacterial blight	<i>If the infestation is more, spray Streptocycline 2gm/5lt of water.</i>
		Blast	<i>For rice blast management, destroy the infested residue, stubbles, grass and weeds present in the bunds of the field as they act as a source for the infection.</i>
<b>Soybean</b>	Harvesting stage		<i>If you're storing soybeans for next season, screen them prior to loading them into the bin</i>
<b>Toria</b>	Land preparation	-	<i>Field preparation should be started for <b>toria</b> and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region</i>
<b>Rabi maize</b>	Land preparation		<i>Land preparation should be started for rabi maize. It can be grown in all types of soils having adequate provision of drainage. Field should be ploughed properly so as to expose the pupae of red hairy caterpillar. Sow certified seeds from a reliable source to prevent seed rot and seedling blight. If one wants to use his own seed, seed treatment is a must. Some of the varieties recommended for our region are HQPM-1, Allrounder and RCM-76.</i>

<b>Horticultural crops</b>			
<b>Citrus</b>		Trunk borer	<i>To kill the trunk borer grubs, clean the bored holes of the infested trunk with iron wire and insert a cotton swab soaked in petrol or kerosene and plug with mud.</i>
<b>Vegetables</b>			<p><i>Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done.</i></p> <p><i>Suitable varieties for our region:</i></p> <ol style="list-style-type: none"> <li><i>1. Cabbage – Green express (Hybrid), Rareball (Hybrid), Pride of India (OP).</i></li> <li><i>2. Broccoli – Pushpa, princess.</i></li> <li><i>3. Tomato – Pusa ruby, Arka vikash, Punjab swari.</i></li> <li><i>4. Knolkhol – White Vienna.</i></li> <li><i>5. Radish – Japanese white, Pusa chetki.</i></li> <li><i>6. Pea – Arkel</i></li> </ol>
<b>Poultry</b>	Good ventilation in the chicken shed is must, because chickens naturally produce a lot of moisture in their breath and droppings. Poor ventilation inside their house may create an ammonia build-up, which will affect the poultry's eyes and respiratory system and makes them sick.		
<b>Piggery</b>	A clean, sanitary environment provides the best prevention for internal and external parasite which can be serious problems. For diseases that can be prevented by vaccination, a veterinarian should be contacted to provide such services routinely.		
<b>Fishery</b>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media		





# Integrated Agromet Advisory Service Bulletin from 5<sup>th</sup> to 9<sup>th</sup> Oct'16 Mon District

Bulletin No:80/2016

## Weather summary of the preceding week

- Light rain occurred the past week
- Maximum and minimum temperatures ranged 26<sup>0</sup>C to 32<sup>0</sup>C and 15<sup>0</sup>C to 19<sup>0</sup>C, respectively.
- Relative humidity varied from 54% to 95%
- Wind speed ranged from 1 to 2 kmph

## Weather forecast valid upto 9<sup>th</sup> Oct'16

- Probability of light rain the coming week.
- **Max temp** is likely to be 28<sup>0</sup>C - 30<sup>0</sup>C and the **min temp** 18<sup>0</sup>C to 21<sup>0</sup>C
- **Sky is likely to be partly cloudy** the coming week
- **Relative Humidity** is likely to range from 45% to 84%.
- **Wind speed** may reach upto 1- 2 kmph
- **Wind direction** will be mostly southeasterly

## Main Crops

## Stage


## Pest/ Diseases

## Agro-meteorological Advisories

## Field crops

Timely field sanitation should be done for all the crops. Always use freshly prepared neem seed kernel extract. Apply pesticides only when required. Don't apply more than the recommended dose of the pesticide

<b>Jhum paddy</b>	Harvesting stage		<i>During sunny days harvest the early and early medium duration varieties when 80% of the grains matured in a panicle. Harvest, process, dry and pack different paddy varieties separately without mixing for better price for their produce</i>
<b>TRC/WRC paddy</b>	Dough development stage	Brown spot	<i>Destroy the infested residue, stubbles, grass and weeds present in the bunds of the field as they act as a source for the infection</i>
		Bacterial blight	<i>If the infestation is more, spray Streptocycline 2gm/5lt of water.</i>
		Blast	<i>For rice blast management, destroy the infested residue, stubbles, grass and weeds present in the bunds of the field as they act as a source for the infection.</i>
<b>Soybean</b>	Harvesting stage		<i>If you're storing soybeans for next season, screen them prior to loading them into the bin</i>
<b>Toria</b>	Land preparation	-	<i>Field preparation should be started for <b>toria</b> and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region</i>
<b>Rabi maize</b>	Land preparation		<i>Land preparation should be started for rabi maize. It can be grown in all types of soils having adequate provision of drainage. Field should be ploughed properly so as to expose the pupae of red hairy caterpillar. Sow certified seeds from a reliable source to prevent seed rot and seedling blight. If one wants to use his own seed, seed treatment is a must.</i>

				<i>Some of the varieties recommended for our region are HQPM-1, Allrounder and RCM-76.</i>
	<p align="center"><b>Horticultural crop</b></p> <p>Spread well decomposed cowdung 5 to 6 tones per hectare for the main field and mix it well in the soil by cultivating the field. Keep the field clean by collecting the stubbles and unwanted materials.</p>			
	<b>Citrus</b>		Trunk borer	<i>To kill the trunk borer grubs, clean the bored holes of the infested trunk with iron wire and insert a cotton swab soaked in petrol or kerosene and plug with mud.</i>
	<b>Vegetables</b>			<p><i>Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done.</i></p> <p><i>Suitable varieties for our region:</i></p> <ol style="list-style-type: none"> <li><i>1. Cabbage – Green express (Hybrid), Rareball (Hybrid), Pride of India (OP).</i></li> <li><i>2. Broccoli – Pushpa, princess.</i></li> <li><i>3. Tomato – Pusa ruby, Arka vikash, Punjab swari.</i></li> <li><i>4. Knolkhol – White Vienna.</i></li> <li><i>5. Radish – Japanese white, Pusa chetki.</i></li> <li><i>6. Pea – Arkel</i></li> </ol>
	<b>Poultry</b>	Good ventilation in the chicken shed is must, because chickens naturally produce a lot of moisture in their breath and droppings. Poor ventilation inside their house may create an ammonia build-up, which will affect the poultry's eyes and respiratory system and makes them sick.		
	<b>Piggery</b>	A clean, sanitary environment provides the best prevention for internal and external parasite which can be serious problems. For diseases that can be prevented by vaccination, a veterinarian should be contacted to provide such services routinely.		
	<b>Fishery</b>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media		



# Integrated Agromet Advisory Service Bulletin from 5<sup>th</sup> to 9<sup>th</sup> Oct'16

## Peren District

Bulletin No:80/2016

### Weather summary of the preceding week

- Light rain occurred the past week
- Maximum and minimum temperatures ranged 24<sup>0</sup>C to 27<sup>0</sup>C and 16<sup>0</sup>C to 20<sup>0</sup>C, respectively.
- Relative humidity varied from 48% to 94%.
- Wind speed ranged from 1 to 2 kmph

### Weather forecast valid upto 9<sup>th</sup> Oct'16

- Probability of light rain the coming week.
- **Max temp** is likely to be 26<sup>0</sup>C - 28<sup>0</sup>C and the **min temp** 20<sup>0</sup>C to 21<sup>0</sup>C
- **Sky is likely to be mainly cloudy** the coming week
- **Relative Humidity** is likely to range from 63% to 90%.
- **Wind speed** may reach upto 1-2 kmph
- **Wind direction** will be mostly easterly

Main Crops

Stage

Pest/  
Diseases



Agro-meteorological Advisories

### Field crops

Timely field sanitation should be done for all the crops. Always use freshly prepared neem seed kernel extract. Apply pesticides only when required. Don't apply more than the recommended dose of the pesticide

<b>Jhum paddy</b>	Harvesting stage		<i>During sunny days harvest the early and early medium duration varieties when 80% of the grains matured in a panicle. Harvest, process, dry and pack different paddy varieties separately without mixing for better price for their produce</i>
<b>TRC/WRC paddy</b>	Milk development stage	Brown spot	<i>Destroy the infested residue, stubbles, grass and weeds present in the bunds of the field as they act as a source for the infection</i>
		Bacterial blight	<i>If the infestation is more, spray Streptocycline 2gm/5lt of water.</i>
		Blast	<i>For rice blast management, destroy the infested residue, stubbles, grass and weeds present in the bunds of the field as they act as a source for the infection.</i>
<b>Soybean</b>	Harvesting stage		<i>If you're storing soybeans for next season, screen them prior to loading them into the bin</i>
<b>Toria</b>	Land preparation	-	<i>Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region</i>
<b>Rabi maize</b>	Land preparation		<i>Land preparation should be started for rabi maize. It can be grown in all types of soils having adequate provision of drainage. Field should be ploughed properly so as to expose the pupae of red hairy caterpillar. Sow certified seeds from a reliable source to prevent seed rot and seedling blight. If one wants to use his own seed, seed treatment is a must.</i>



				Some of the varieties recommended for our region are HQPM-1, Allrounder and RCM-76.
	<b>Horticultural crops</b>			
	<b>Citrus</b>		Trunk borer	To kill the trunk borer grubs, clean the bored holes of the infested trunk with iron wire and insert a cotton swab soaked in petrol or kerosene and plug with mud.
	<b>Vegetables</b>			Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done. Suitable varieties for our region: 1. Cabbage – Green express (Hybrid), Rareball (Hybrid), Pride of India (OP). 2. Broccoli – Pushpa, princess. 3. Tomato – Pusa ruby, Arka vikash, Punjab swari. 4. Knolkhol – White Vienna. 5. Radish – Japanese white, Pusa chetki. 6. Pea – Arkel
	<b>Poultry</b>	Good ventilation in the chicken shed is must, because chickens naturally produce a lot of moisture in their breath and droppings. Poor ventilation inside their house may create an ammonia build-up, which will affect the poultry's eyes and respiratory system and makes them sick.		
	<b>Piggery</b>	A clean, sanitary environment provides the best prevention for internal and external parasite which can be serious problems. For diseases that can be prevented by vaccination, a veterinarian should be contacted to provide such services routinely.		
	<b>Fishery</b>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media		





# Integrated Agromet Advisory Service Bulletin from 5<sup>th</sup> to 9<sup>th</sup> Oct'16 Phek District

Bulletin No:80/2016

## Weather summary of the preceding week

- Light rain occurred the past week
- Maximum and minimum temperatures ranged 26<sup>0</sup>C to 28<sup>0</sup>C and 17<sup>0</sup>C to 19<sup>0</sup>C, respectively.
- Relative humidity varied from 53% to 95%.
- Wind speed ranged from 1 to 2 kmph

## Weather forecast valid upto 9<sup>th</sup> Oct'16

- Probability of light rain the coming week.
- **Max temp** is likely to be 28<sup>0</sup>C - 30<sup>0</sup>C and the **min temp** 19<sup>0</sup>C to 21<sup>0</sup>C
- **Sky is likely to be mainly cloudy** the coming week
- **Relative Humidity** is likely to range from 42% to 80%.
- **Wind speed** may reach upto 1-2 kmph
- **Wind direction** will be mostly southeasterly

**Main  
Crops**

**Stage**


**Pest/  
Diseases**

**Agro-meteorological Advisories**

## Field crops

Timely field sanitation should be done for all the crops. Always use freshly prepared neem seed kernel extract. Apply pesticides only when required. Don't apply more than the recommended dose of the pesticide

<b>Jhum paddy</b>	Harvesting stage		<i>During sunny days harvest the early and early medium duration varieties when 80% of the grains matured in a panicle. Harvest, process, dry and pack different paddy varieties separately without mixing for better price for their produce</i>
<b>TRC/WRC paddy</b>	Milk development stage	Brown spot	<i>Destroy the infested residue, stubbles, grass and weeds present in the bunds of the field as they act as a source for the infection</i>
		Bacterial blight	<i>If the infestation is more, spray Streptocycline 2gm/5lt of water.</i>
		Blast	<i>For rice blast management, destroy the infested residue, stubbles, grass and weeds present in the bunds of the field as they act as a source for the infection.</i>
<b>Soybean</b>	Harvesting stage		<i>If you're storing soybeans for next season, screen them prior to loading them into the bin</i>
<b>Toria</b>	Land preparation	-	<i>Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region</i>
<b>Rabi maize</b>	Land preparation		<i>Land preparation should be started for rabi maize. It can be grown in all types of soils having adequate provision of drainage. Field should be ploughed properly so as to expose the pupae of red hairy caterpillar. Sow certified seeds from a reliable source to prevent seed rot and seedling blight. If one wants to use his own seed,</i>


				<p><i>seed treatment is a must.</i></p> <p><i>Some of the varieties recommended for our region are HQPM-1, Allrounder and RCM-76.</i></p>
	<b>Horticultural crops</b>			
	<b>Citrus</b>		Trunk borer	<p><i>To kill the trunk borer grubs, clean the bored holes of the infested trunk with iron wire and insert a cotton swab soaked in petrol or kerosene and plug with mud.</i></p>
	<b>Vegetables</b>			<p><i>Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done.</i></p> <p><i>Suitable varieties for our region:</i></p> <ol style="list-style-type: none"> <li><i>1. Cabbage – Green express (Hybrid), Rareball (Hybrid), Pride of India (OP).</i></li> <li><i>2. Broccoli – Pushpa, princess.</i></li> <li><i>3. Tomato – Pusa ruby, Arka vikash, Punjab swari.</i></li> <li><i>4. Knolkhol – White Vienna.</i></li> <li><i>5. Radish – Japanese white, Pusa chetki.</i></li> <li><i>6. Pea – Arkel</i></li> </ol>
	<b>Poultry</b>	<p>Good ventilation in the chicken shed is must, because chickens naturally produce a lot of moisture in their breath and droppings. Poor ventilation inside their house may create an ammonia build-up, which will affect the poultry's eyes and respiratory system and makes them sick.</p>		
	<b>Piggery</b>	<p>A clean, sanitary environment provides the best prevention for internal and external parasite which can be serious problems. For diseases that can be prevented by vaccination, a veterinarian should be contacted to provide such services routinely.</p>		
	<b>Fishery</b>	<p>Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media</p>		



# Integrated Agromet Advisory Service Bulletin from 5<sup>th</sup> to 9<sup>th</sup> Oct'16 Tuensang District

Bulletin No:80/2016

Weather summary of the preceding week			Weather forecast valid upto 9 <sup>th</sup> Oct'16
<ul style="list-style-type: none"> <li>Light rain occurred the past week</li> <li>Maximum and minimum temperatures ranged 27<sup>0</sup>C to 30<sup>0</sup>C and 18<sup>0</sup>C to 23<sup>0</sup>C, respectively.</li> <li>Relative humidity varied from 64% to 92%.</li> <li>Wind speed ranged from 1 to 2 kmph</li> </ul>			<ul style="list-style-type: none"> <li>Probability of light rain the coming week.</li> <li><b>Max temp</b> is likely to be 28<sup>0</sup>C - 30<sup>0</sup>C and the <b>min temp</b> 19<sup>0</sup>C to 21<sup>0</sup>C</li> <li><b>Sky is likely to be mainly cloudy</b> the coming week</li> <li><b>Relative Humidity</b> is likely to range from 40% to 90%.</li> <li><b>Wind speed</b> may reach upto 1-2 kmph</li> <li><b>Wind direction</b> will be mostly southeasterly</li> </ul>
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<p><b>Field crops</b></p> <p>Timely field sanitation should be done for all the crops. Always use freshly prepared neem seed kernel extract. Apply pesticides only when required. Don't apply more than the recommended dose of the pesticide</p>			
<b>Jhum paddy</b>	Harvesting stage		<i>During sunny days harvest the early and early medium duration varieties when 80% of the grains matured in a panicle. Harvest, process, dry and pack different paddy varieties separately without mixing for better price for their produce</i>
<b>TRC/WRC paddy</b>	Milk development stage	Brown spot	<i>Destroy the infested residue, stubbles, grass and weeds present in the bunds of the field as they act as a source for the infection</i>
		Bacterial blight	<i>If the infestation is more, spray Streptocycline 2gm/5lt of water.</i>
		Blast	<i>For rice blast management, destroy the infested residue, stubbles, grass and weeds present in the bunds of the field as they act as a source for the infection.</i>
<b>Soybean</b>	Harvesting stage		<i>If you're storing soybeans for next season, screen them prior to loading them into the bin</i>
<b>Toria</b>	Land preparation	-	<i>Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region</i>
<b>Rabi maize</b>	Land preparation		<i>Land preparation should be started for rabi maize. It can be grown in all types of soils having adequate provision of drainage. Field should be ploughed properly so as to expose the pupae of red hairy caterpillar. Sow certified seeds from a reliable source to prevent seed rot and seedling blight. If one wants to use his own seed, seed treatment is a must. Some of the varieties recommended for our region are HQPM-1,</i>


				Allrounder and RCM-76.
	<b>Horticultural crops</b>			
	<b>Citrus</b>		Trunk borer	To kill the trunk borer grubs, clean the bored holes of the infested trunk with iron wire and insert a cotton swab soaked in petrol or kerosene and plug with mud.
	<b>Vegetables</b>			<p>Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done.</p> <p>Suitable varieties for our region:</p> <ol style="list-style-type: none"> <li>1. Cabbage – Green express (Hybrid), Rareball (Hybrid), Pride of India (OP).</li> <li>2. Broccoli – Pushpa, princess.</li> <li>3. Tomato – Pusa ruby, Arka vikash, Punjab swari.</li> <li>4. Knolkhol – White Vienna.</li> <li>5. Radish – Japanese white, Pusa chetki.</li> <li>6. Pea – Arkel</li> </ol>
	<b>Poultry</b>	Good ventilation in the chicken shed is must, because chickens naturally produce a lot of moisture in their breath and droppings. Poor ventilation inside their house may create an ammonia build-up, which will affect the poultry's eyes and respiratory system and makes them sick.		
	<b>Piggery</b>	A clean, sanitary environment provides the best prevention for internal and external parasite which can be serious problems. For diseases that can be prevented by vaccination, a veterinarian should be contacted to provide such services routinely.		
	<b>Fishery</b>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media		



## Integrated Agromet Advisory Service Bulletin from 5<sup>th</sup> to 9<sup>th</sup> Oct'16 Wokha District

Bulletin No:80/2016

Weather summary of the preceding week			Weather forecast valid upto 9 <sup>th</sup> Oct'16
<ul style="list-style-type: none"> <li>Light rain occurred the past week</li> <li>Maximum and minimum temperatures ranged 27<sup>0</sup>C to 28<sup>0</sup>C and 18<sup>0</sup>C to 22<sup>0</sup>C, respectively.</li> <li>Relative humidity varied from 51% to 91%.</li> <li>Wind speed ranged from 1 to 2 kmph</li> </ul>			<ul style="list-style-type: none"> <li>Probability of light rain the coming week.</li> <li><b>Max temp</b> is likely to be 26<sup>0</sup>C - 28<sup>0</sup>C and the <b>min temp</b> 20<sup>0</sup>C to 21<sup>0</sup>C</li> <li><b>Sky is likely to be mainly cloudy</b> the coming week</li> <li><b>Relative Humidity</b> is likely to range from 45% to 91%.</li> <li><b>Wind speed</b> may reach upto 1-3 kmph</li> <li><b>Wind direction</b> will be mostly southeasterly</li> </ul>
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<p><b>Field crops</b></p> <p>Timely field sanitation should be done for all the crops. Always use freshly prepared neem seed kernel extract. Apply pesticides only when required. Don't apply more than the recommended dose of the pesticide</p>			
<b>Jhum paddy</b>	Harvesting stage		<i>During sunny days harvest the early and early medium duration varieties when 80% of the grains matured in a panicle. Harvest, process, dry and pack different paddy varieties separately without mixing for better price for their produce</i>
<b>TRC/WRC paddy</b>	Milk development stage	Brown spot	<i>Destroy the infested residue, stubbles, grass and weeds present in the bunds of the field as they act as a source for the infection</i>
		Bacterial blight	<i>If the infestation is more, spray Streptocycline 2gm/5lt of water.</i>
		Blast	<i>For rice blast management, destroy the infested residue, stubbles, grass and weeds present in the bunds of the field as they act as a source for the infection.</i>
<b>Soybean</b>	Harvesting stage		<i>If you're storing soybeans for next season, screen them prior to loading them into the bin</i>
<b>Toria</b>	Land preparation	-	<i>Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region</i>
<b>Rabi maize</b>	Land preparation		<i>Land preparation should be started for rabi maize. It can be grown in all types of soils having adequate provision of drainage. Field should be ploughed properly so as to expose the pupae of red hairy caterpillar. Sow certified seeds from a reliable source to prevent seed rot and seedling blight. If one wants to use his own seed, seed treatment is a must.</i>

				<i>Some of the varieties recommended for our region are HQPM-1, Allrounder and RCM-76.</i>
	<b><i>Horticultural crops</i></b>			
	<b><i>Citrus</i></b>		Trunk borer	<i>To kill the trunk borer grubs, clean the bored holes of the infested trunk with iron wire and insert a cotton swab soaked in petrol or kerosene and plug with mud.</i>
	<b><i>Vegetables</i></b>			<i>Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done.</i> <i>Suitable varieties for our region:</i> <ol style="list-style-type: none"> <li><i>1. Cabbage – Green express (Hybrid), Rareball (Hybrid), Pride of India (OP).</i></li> <li><i>2. Broccoli – Pushpa, princess.</i></li> <li><i>3. Tomato – Pusa ruby, Arka vikash, Punjab swari.</i></li> <li><i>4. Knolkhol – White Vienna.</i></li> <li><i>5. Radish – Japanese white, Pusa chetki.</i></li> <li><i>6. Pea – Arkel</i></li> </ol>
	<b><i>Poultry</i></b>	Good ventilation in the chicken shed is must, because chickens naturally produce a lot of moisture in their breath and droppings. Poor ventilation inside their house may create an ammonia build-up, which will affect the poultry's eyes and respiratory system and makes them sick.		
	<b><i>Piggery</i></b>	A clean, sanitary environment provides the best prevention for internal and external parasite which can be serious problems. For diseases that can be prevented by vaccination, a veterinarian should be contacted to provide such services routinely.		
	<b><i>Fishery</i></b>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media		








--	--



## Integrated Agromet Advisory Service Bulletin from 5<sup>th</sup> to 9<sup>th</sup> Oct'16 Zunheboto District

Bulletin No:80/2016

Weather summary of the preceding week			Weather forecast valid upto 9 <sup>th</sup> Oct'16
<ul style="list-style-type: none"> <li>Light rain occurred the past week</li> <li>Maximum and minimum temperatures ranged 25<sup>0</sup>C to 30<sup>0</sup>C and 17<sup>0</sup>C to 21<sup>0</sup>C , respectively.</li> <li>Relative humidity varied from 53% to 93%.</li> <li>Wind speed ranged from 1 to 2 kmph</li> </ul>			<ul style="list-style-type: none"> <li>Probability of light rain the coming week.</li> <li><b>Max temp</b> is likely to be 29<sup>0</sup>C - 30<sup>0</sup>C and the <b>min temp</b> 21<sup>0</sup>C to 22<sup>0</sup>C</li> <li><b>Sky is likely to be mainly cloudy</b> the coming week</li> <li><b>Relative Humidity</b> is likely to range from 45% to 95%.</li> <li><b>Wind speed</b> may reach upto 1-2 kmph</li> <li><b>Wind direction</b> will be mostly southeasterly</li> </ul>
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<p align="center"><b>Field crops</b></p> <p>Timely field sanitation should be done for all the crops. Always use freshly prepared neem seed kernel extract. Apply pesticides only when required. Don't apply more than the recommended dose of the pesticide</p>			
<b>Jhum paddy</b>	Harvesting stage		<i>During sunny days harvest the early and early medium duration varieties when 80% of the grains matured in a panicle. Harvest, process, dry and pack different paddy varieties separately without mixing for better price for their produce</i>
<b>TRC/WRC paddy</b>	Milk development stage	Brown spot	<i>Destroy the infested residue, stubbles, grass and weeds present in the bunds of the field as they act as a source for the infection</i>
		Bacterial blight	<i>If the infestation is more, spray Streptocycline 2gm/5lt of water.</i>
		Blast	<i>For rice blast management, destroy the infested residue, stubbles, grass and weeds present in the bunds of the field as they act as a source for the infection.</i>
<b>Soybean</b>	Harvesting stage		<i>If you're storing soybeans for next season, screen them prior to loading them into the bin</i>
<b>Toria</b>	Land preparation	-	<i>Field preparation should be started for <b>toria</b> and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region</i>
<b>Rabi maize</b>	Land preparation		<i>Land preparation should be started for rabi maize. It can be grown in all types of soils having adequate provision of drainage. Field should be ploughed properly so as to expose the pupae of red hairy caterpillar. Sow certified seeds from a reliable source to prevent seed rot and seedling blight. If one wants to use his own seed, seed treatment is a must.</i>

				<i>Some of the varieties recommended for our region are HQPM-1, Allrounder and RCM-76.</i>
	<b><i>Horticultural crops</i></b>			
	<b><i>Citrus</i></b>		Trunk borer	<i>To kill the trunk borer grubs, clean the bored holes of the infested trunk with iron wire and insert a cotton swab soaked in petrol or kerosene and plug with mud.</i>
	<b><i>Vegetables</i></b>			<i>Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done.</i> <i>Suitable varieties for our region:</i> <ol style="list-style-type: none"> <li><i>1. Cabbage – Green express (Hybrid), Rareball (Hybrid), Pride of India (OP).</i></li> <li><i>2. Broccoli – Pushpa, princess.</i></li> <li><i>3. Tomato – Pusa ruby, Arka vikash, Punjab swari.</i></li> <li><i>4. Knolkhol – White Vienna.</i></li> <li><i>5. Radish – Japanese white, Pusa chetki.</i></li> <li><i>6. Pea – Arkel</i></li> </ol>
	<b><i>Poultry</i></b>	Good ventilation in the chicken shed is must, because chickens naturally produce a lot of moisture in their breath and droppings. Poor ventilation inside their house may create an ammonia build-up, which will affect the poultry's eyes and respiratory system and makes them sick.		
	<b><i>Piggery</i></b>	A clean, sanitary environment provides the best prevention for internal and external parasite which can be serious problems. For diseases that can be prevented by vaccination, a veterinarian should be contacted to provide such services routinely.		
	<b><i>Fishery</i></b>	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface. If there are too many weeds, clean your pond. Take the advice of experts and information from mass media		

