




## Integrated Agromet Advisory Service Bulletin from 21<sup>st</sup> to 25<sup>th</sup> June'17

### Longleng District

Bulletin No:47/2017

Weather summary of the preceding week			Weather forecast valid upto 25 <sup>th</sup> June'17
<ul style="list-style-type: none"> <li>Moderate rain occurred the past week</li> <li>Maximum and minimum temperatures ranged 30<sup>o</sup>C to 34<sup>o</sup>C and 21<sup>o</sup>C to 24<sup>o</sup>C, respectively.</li> <li>Relative humidity varied from 47% to 93%.</li> <li>Wind speed ranged from 1 to 2 kmph</li> </ul>			<ul style="list-style-type: none"> <li>Probability of light rain the coming week.</li> <li><b>Max temp</b> is likely to be 30<sup>o</sup>C - 31<sup>o</sup>C and the <b>min temp</b> 24<sup>o</sup>C to 25<sup>o</sup>C</li> <li><b>Sky is likely to be mainly cloudy</b> the coming week</li> <li><b>Relative Humidity</b> is likely to range from 64% to 95%.</li> <li><b>Wind speed</b> may reach upto 1- 2 kmph</li> <li><b>Wind direction</b> will be mostly southerly</li> </ul>
Field crops			
Proper drainage should be maintained in the field.			
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Jhum paddy	Tillering stage	Blast	If the infection is severe, spray tricyclazole 0.1% at 12-15 days interval
TRC/WRC paddy	Nursery		Prepare the nursery by plowing at least twice and harrow at least once. Level the soil surface and put in drainage lines across the field. Broadcast seed in the nursery evenly, over the water covered soil surface Seed treatment should be done
Kharif maize	Vegetative stage	Stem borer	The prevalent weather is favourable for the development of stem borer. Removed the infested plant parts.
Green gram	Flowering stage		Good drainage may be given to manage soil borne root diseases
Horticultural crop			
Citrus	Flowering to fruiting stage	Mealy bug	<ul style="list-style-type: none"> <li>In citrus orchard, destroy the ant colonies (black ants) as they act as the carriers of mealy bug. The affected plants become pale, wilt and may die also. In heavily attacked plants, citrus fruits do not set.</li> <li>Timely weeding should be carried out</li> </ul>
Okra	Flowering to fruiting stage	Yellow vein mosaic	<ul style="list-style-type: none"> <li>Remove and burn the infected plants from the field</li> <li>Remove the unwanted weed host from the field</li> <li>If the infection is severe, spray imidacloprid at 1ml/lit of water</li> </ul>
		Spotted bollworm	<ul style="list-style-type: none"> <li>In the present weather condition, spotted bollworm will be seen. The infested fruits will have a varying number of holes.</li> <li>Remove regularly the attacked fruits and bury deep in the soil.</li> </ul>

	<b>Cucurbits</b>	Vegetative stage	Powdery mildew	<i>Remove the infected leaves</i>
			Stink bug	<i>Maintain proper sanitation as they are mostly found in weedy areas.</i>
				<ul style="list-style-type: none"><li>➤ <i>For trailing, dried twigs can be spread on the ground. Weeding should be done whenever necessary.</i></li></ul>
	<b>Brinjal</b>	Vegetative stage		<ul style="list-style-type: none"><li>➤ <i>Timely weeding and intercultural operation should be done.</i></li></ul>
			Wilt	<ul style="list-style-type: none"><li>➤ <i>Remove and destroy the affected plant</i></li><li>➤ <i>Proper drainage should be maintained in the field</i></li></ul>
	<b>Naga king chilli</b>	Vegetative stage	viral disease	<ul style="list-style-type: none"><li>➤ <i>Timely weeding should be done</i></li><li>➤ <i>Virus infected plants should be removed or uprooted and buried into the soil</i></li><li>➤ <i>Daily monitoring should be done for insect pest and need based neem oil should be spray</i></li></ul>
	<b>Fruit trees</b>			<i>Maintain proper sanitation in and around the vicinity of the tree</i>
	<b>Ginger / turmeric</b>	Vegetative stage	-	<i>Timely earthing up and weeding/hoeing should be done. Mulch the plant with dry leaves and straw</i>
	<b>Livestock</b>			
	Provide clean and cool drinking water to the animals			
<b>Poultry</b>	<ul style="list-style-type: none"><li>➤ Protect the birds from rain</li><li>➤ Keep the litter material dry</li><li>➤ Give electrolyte in drinking water to birds of 1-6 weeks.</li></ul>			
<b>Piggery</b>	<ul style="list-style-type: none"><li>➤ Vaccinate against swine fever</li><li>➤ Maintain hygiene in pig house to protect from diarrhea</li><li>➤ Deworm the animal at 3 month interval</li><li>➤ Do AI, if animal comes into heat</li></ul>			
<b>Fisheries</b>				
		<ul style="list-style-type: none"><li>➤ Apply lime as per recommended dose depending on the pH</li><li>➤ After 15 days of lime application, apply cow dung and wait till the pond water turns green.</li><li>➤ Put some aquatic plant in one corner of the pond for sheltering stocked fish during high temperature.</li></ul>		

## SCIENTIFIC EXPERT COMMITTEE

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



# Integrated Agromet Advisory Service Bulletin from 21<sup>st</sup> to 25<sup>th</sup> June'17

## Mokokchung District

Bulletin No:47/2017

### Weather summary of the preceding week

- Moderate rain occurred the past week
- Maximum and minimum temperatures ranged 30<sup>0</sup>C to 32<sup>0</sup>C and 20<sup>0</sup>C to 23<sup>0</sup>C, respectively.
- Relative humidity varied from 57% to 96%.
- Wind speed ranged from 1 to 2 kmph

### Weather forecast valid upto 25<sup>th</sup> June'17

- Probability of light rain the coming week.
- **Max temp** is likely to be 26<sup>0</sup>C - 27<sup>0</sup>C and the **min temp** 19<sup>0</sup>C to 21<sup>0</sup>C
- **Sky is likely to be mainly cloudy** the coming week
- **Relative Humidity** is likely to range from 63% to 96%.
- **Wind speed** may reach upto 1- 2 kmph
- **Wind direction** will be southerly

### Field crops

Proper drainage should be maintained in the field.

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum paddy</i>	Tillering stage	Blast	<i>If the infection is severe, spray tricyclazole 0.1% at 12-15 days interval</i>
<i>TRC/WRC paddy</i>	Nursery		<i>Prepare the nursery by plowing at least twice and harrow at least once. Level the soil surface and put in drainage lines across the field. Broadcast seed in the nursery evenly, over the water covered soil surface Seed treatment should be done</i>
<i>Kharif maize</i>	Vegetative stage	Stem borer	<i>The prevalent weather is favourable for the development of stem borer. Removed the infested plant parts.</i>
<i>Green gram</i>	Flowering stage		<i>Good drainage may be given to manage soil borne root diseases</i>

### Horticultural crop

<i>Citrus</i>	Flowering to fruiting stage	Mealy bug	<ul style="list-style-type: none"> <li>➤ <i>In citrus orchard, destroy the ant colonies (black ants) as they act as the carriers of mealy bug. The affected plants become pale, wilt and may die also. In heavily attacked plants, citrus fruits do not set.</i></li> <li>➤ <i>Timely weeding should be carried out</i></li> </ul>
<i>Okra</i>	Flowering to fruiting stage	Yellow vein mosaic	<ul style="list-style-type: none"> <li>➤ <i>Remove and burn the infected plants from the field</i></li> <li>➤ <i>Remove the unwanted weed host from the field</i></li> <li>➤ <i>If the infection is severe, spray imidacloprid at 1ml/lit of water</i></li> </ul>
		Spotted bollworm	<ul style="list-style-type: none"> <li>➤ <i>In the present weather condition, spotted bollworm will be seen. The infested fruits will have a varying number of holes.</i></li> <li>➤ <i>Remove regularly the attacked fruits and bury deep in the soil.</i></li> </ul>
<i>Cucurbits</i>	Vegetative stage	Powdery	<i>Remove the infected leaves</i>

		mildew	
		Stink bug	<i>Maintain proper sanitation as they are mostly found in weedy areas.</i>
			➤ <i>For trailing, dried twigs can be spread on the ground. Weeding should be done whenever necessary.</i>
<b>Brinjal</b>	Vegetative stage		➤ <i>Timely weeding and intercultural operation should be done.</i>
		Wilt	➤ <i>Remove and destroy the affected plant</i> ➤ <i>Proper drainage should be maintained in the field</i>
<b>Naga king chilli</b>	Vegetative stage	viral disease	➤ <i>Timely weeding should be done</i> ➤ <i>Virus infected plants should be removed or uprooted and buried into the soil</i> ➤ <i>Daily monitoring should be done for insect pest and need based neem oil should be spray</i>
<b>Fruit trees</b>			<i>Maintain proper sanitation in and around the vicinity of the tree</i>
<b>Ginger / turmeric</b>	Vegetative stage	-	<i>Timely earthing up and weeding/hoeing should be done. Mulch the plant with dry leaves and straw</i>
<b>Livestock</b>			
Provide clean and cool drinking water to the animals			
<b>Poultry</b>	➤ Protect the birds from rain ➤ Keep the litter material dry ➤ Give electrolyte in drinking water to birds of 1-6 weeks.		
<b>Piggery</b>	➤ Vaccinate against swine fever ➤ Maintain hygiene in pig house to protect from diarrhea ➤ Deworm the animal at 3 month interval ➤ Do AI, if animal comes into heat		
	<b>Fisheries</b>		
	➤ Apply lime as per recommended dose depending on the pH ➤ After 15 days of lime application, apply cow dung and wait till the pond water turns green. ➤ Put some aquatic plant in one corner of the pond for sheltering stocked fish during high temperature.		



## SCIENTIFIC EXPERT COMMITTEE


Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



## Integrated Agromet Advisory Service Bulletin from 21<sup>st</sup> to 25<sup>th</sup> June'17 Mon District

Bulletin No:47/2017

Weather summary of the preceding week			Weather forecast valid upto 25 <sup>th</sup> June'17
<ul style="list-style-type: none"> <li>Moderate rain occurred the past week</li> <li>Maximum and minimum temperatures ranged 30<sup>o</sup>C to 34<sup>o</sup>C and 20<sup>o</sup>C to 23<sup>o</sup>C, respectively.</li> <li>Relative humidity varied from 57% to 92%</li> <li>Wind speed ranged from 1 to 2 kmph</li> </ul>			<ul style="list-style-type: none"> <li>Probability of light rain the coming week.</li> <li><b>Max temp</b> is likely to be 25<sup>o</sup>C - 27<sup>o</sup>C and the <b>min temp</b> 19<sup>o</sup>C to 20<sup>o</sup>C</li> <li><b>Sky is likely to be mainly cloudy</b> the coming week</li> <li><b>Relative Humidity</b> is likely to range from 62% to 95%.</li> <li><b>Wind speed</b> may reach upto 1- 2 kmph</li> <li><b>Wind direction</b> will be mostly southwesterly</li> </ul>
Field crops			
Proper drainage should be maintained in the field.			
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum paddy</i>	Tillering stage	Blast	<i>If the infection is severe, spray tricyclazole 0.1% at 12-15 days interval</i>
<i>TRC/WRC paddy</i>	Nursery		<i>Prepare the nursery by plowing at least twice and harrow at least once. Level the soil surface and put in drainage lines across the field. Broadcast seed in the nursery evenly, over the water covered soil surface Seed treatment should be done</i>
<i>Kharif maize</i>	Vegetative stage	Stem borer	<i>The prevalent weather is favourable for the development of stem borer. Removed the infested plant parts.</i>
<i>Green gram</i>	Flowering stage		<i>Good drainage may be given to manage soil borne root diseases</i>
Horticultural crop			
<i>Citrus</i>	Flowering to fruiting stage	Mealy bug	<ul style="list-style-type: none"> <li>➤ <i>In citrus orchard, destroy the ant colonies (black ants) as they act as the carriers of mealy bug. The affected plants become pale, wilt and may die also. In heavily attacked plants, citrus fruits do not set.</i></li> <li>➤ <i>Timely weeding should be carried out</i></li> </ul>
<i>Okra</i>	Flowering to fruiting stage	Yellow vein mosaic	<ul style="list-style-type: none"> <li>➤ <i>Remove and burn the infected plants from the field</i></li> <li>➤ <i>Remove the unwanted weed host from the field</i></li> <li>➤ <i>If the infection is severe, spray imidacloprid at 1ml/lit of water</i></li> </ul>
		Spotted bollworm	<ul style="list-style-type: none"> <li>➤ <i>In the present weather condition, spotted bollworm will be seen. The infested fruits will have a varying number of holes.</i></li> <li>➤ <i>Remove regularly the attacked fruits and bury deep in the soil.</i></li> </ul>
<i>Cucurbits</i>	Vegetative stage	Powdery mildew	<i>Remove the infected leaves</i>

			Stink bug	Maintain proper sanitation as they are mostly found in weedy areas.
				➤ For trailing, dried twigs can be spread on the ground. Weeding should be done whenever necessary.
	Brinjal	Vegetative stage		➤ Timely weeding and intercultural operation should be done.
			Wilt	➤ Remove and destroy the affected plant ➤ Proper drainage should be maintained in the field
	Naga king chilli	Vegetative stage	viral disease	➤ Timely weeding should be done ➤ Virus infected plants should be removed or uprooted and buried into the soil ➤ Daily monitoring should be done for insect pest and need based neem oil should be spray
	Fruit trees			Maintain proper sanitation in and around the vicinity of the tree
	Ginger / turmeric	Vegetative stage	-	Timely earthing up and weeding/hoeing should be done. Mulch the plant with dry leaves and straw
	<b>Livestock</b> Provide clean and cool drinking water to the animals			
	Poultry	➤ Protect the birds from rain ➤ Keep the litter material dry ➤ Give electrolyte in drinking water to birds of 1-6 weeks. ➤ Good management and sanitation are the best ways to avoid infectious disease in poultry.		
	Piggery	➤ Vaccinate against swine fever ➤ Maintain hygiene in pig house to protect from diarrhea ➤ Deworm the animal at 3 month interval ➤ Do AI, if animal comes into heat		
<b>Fisheries</b>				
		➤ Apply lime as per recommended dose depending on the pH ➤ After 15 days of lime application, apply cow dung and wait till the pond water turns green. ➤ Put some aquatic plant in one corner of the pond for sheltering stocked fish during high temperature ➤ Avoid excessive manuring and feeding which can lead to oxygen depletion in cloudy weather, hence both should be avoided.		



## SCIENTIFIC EXPERT COMMITTEE

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com




# Integrated Agromet Advisory Service Bulletin from 21<sup>st</sup> to 25<sup>th</sup> June'17

## Peren District

Bulletin No:47/2017

Weather summary of the preceding week			Weather forecast valid upto 25 <sup>th</sup> June'17
<ul style="list-style-type: none"> <li>Moderate rain occurred the past week</li> <li>Maximum and minimum temperatures ranged 31<sup>0</sup>C to 33<sup>0</sup>C and 21<sup>0</sup>C to 25<sup>0</sup>C, respectively.</li> <li>Relative humidity varied from 45% to 95%.</li> <li>Wind speed ranged from 1 to 2 kmph</li> </ul>			<ul style="list-style-type: none"> <li>Probability of moderate rain the coming week.</li> <li><b>Max temp</b> is likely to be 29<sup>0</sup>C - 31<sup>0</sup>C and the <b>min temp</b> 23<sup>0</sup>C to 24<sup>0</sup>C</li> <li><b>Sky is likely to be mainly cloudy</b> the coming week</li> <li><b>Relative Humidity</b> is likely to range from 54% to 96%.</li> <li><b>Wind speed</b> may reach upto 1- 2 kmph</li> <li><b>Wind direction</b> will be mostly southerly</li> </ul>
Field crops			
Proper drainage should be maintained in the field.			
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Jhum paddy	Tillering stage	Blast	If the infection is severe, spray tricyclazole 0.1% at 12-15 days interval
TRC/WRC paddy	Nursery		Prepare the nursery by plowing at least twice and harrow at least once. Level the soil surface and put in drainage lines across the field. Broadcast seed in the nursery evenly, over the water covered soil surface Seed treatment should be done
Kharif maize	Vegetative stage	Stem borer	The prevalent weather is favourable for the development of stem borer. Removed the infested plant parts.
Green gram	Flowering stage		Good drainage may be given to manage soil borne root diseases
Horticultural crop			
Citrus	Flowering to fruiting stage	Mealy bug	<ul style="list-style-type: none"> <li>In citrus orchard, destroy the ant colonies (black ants) as they act as the carriers of mealy bug. The affected plants become pale, wilt and may die also. In heavily attacked plants, citrus fruits do not set.</li> <li>Timely weeding should be carried out</li> </ul>
Okra	Flowering to fruiting stage	Yellow vein mosaic	<ul style="list-style-type: none"> <li>Remove and burn the infected plants from the field</li> <li>Remove the unwanted weed host from the field</li> <li>If the infection is severe, spray imidacloprid at 1ml/lit of water</li> </ul>
		Spotted	<ul style="list-style-type: none"> <li>In the present weather condition, spotted bollworm will be seen. The infested fruits will have a varying number of holes.</li> </ul>

			bollworm	➤ <i>Remove regularly the attacked fruits and bury deep in the soil.</i>
	<b>Cucurbits</b>	Vegetative stage	Powdery mildew	<i>Remove the infected leaves</i>
			Stink bug	<i>Maintain proper sanitation as they are mostly found in weedy areas.</i>
				➤ <i>For trailing, dried twigs can be spread on the ground. Weeding should be done whenever necessary.</i>
	<b>Brinjal</b>	Vegetative stage		➤ <i>Timely weeding and intercultural operation should be done.</i>
			Wilt	➤ <i>Remove and destroy the affected plant</i> ➤ <i>Proper drainage should be maintained in the field</i>
	<b>Naga king chilli</b>	Vegetative stage	viral disease	➤ <i>Timely weeding should be done</i> ➤ <i>Virus infected plants should be removed or uprooted and buried into the soil</i> ➤ <i>Daily monitoring should be done for insect pest and need based neem oil should be spray</i>
	<b>Fruit trees</b>			<i>Maintain proper sanitation in and around the vicinity of the tree</i>
	<b>Ginger / turmeric</b>	Vegetative stage	-	<i>Timely earthing up and weeding/hoeing should be done. Mulch the plant with dry leaves and straw</i>
	<b>Livestock</b>			
Provide clean and cool drinking water to the animals				
	<b>Poultry</b>	➤ Protect the birds from rain ➤ Keep the litter material dry ➤ Give electrolyte in drinking water to birds of 1-6 weeks.		
	<b>Piggery</b>	➤ Vaccinate against swine fever ➤ Maintain hygiene in pig house to protect from diarrhea ➤ Deworm the animal at 3 month interval ➤ Do AI, if animal comes into heat		
		<b>Fisheries</b>		
		➤ Apply lime as per recommended dose depending on the pH ➤ After 15 days of lime application, apply cow dung and wait till the pond water turns green. ➤ Put some aquatic plant in one corner of the pond for sheltering stocked fish during high temperature.		

## SCIENTIFIC EXPERT COMMITTEE

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



# Integrated Agromet Advisory Service Bulletin from 21<sup>st</sup> to 25<sup>th</sup> June'17 Phek District

Bulletin No:47/2017

Weather summary of the preceding week			Weather forecast valid upto 25 <sup>th</sup> June'17
<ul style="list-style-type: none"> <li>Moderate rain occurred the past week</li> <li>Maximum and minimum temperatures ranged 28<sup>0</sup>C to 31<sup>0</sup>C and 20<sup>0</sup>C to 22<sup>0</sup>C, respectively.</li> <li>Relative humidity varied from 45% to 95%.</li> <li>Wind speed ranged from 1 to 2 kmph</li> </ul>			<ul style="list-style-type: none"> <li>Probability of light rain the coming week.</li> <li><b>Max temp</b> is likely to be 24<sup>0</sup>C - 25<sup>0</sup>C and the <b>min temp</b> 19<sup>0</sup>C to 21<sup>0</sup>C</li> <li><b>Sky is likely to be mainly cloudy</b> the coming week</li> <li><b>Relative Humidity</b> is likely to range from 60% to 95%.</li> <li><b>Wind speed</b> may reach upto 1- 2 kmph</li> <li><b>Wind direction</b> will be mostly southwesterly</li> </ul>
Field crops			
Proper drainage should be maintained in the field.			
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Jhum paddy	Tillering stage	Blast	If the infection is severe, spray tricyclazole 0.1% at 12-15 days interval
TRC/WRC paddy	Nursery		Prepare the nursery by plowing at least twice and harrow at least once. Level the soil surface and put in drainage lines across the field. Broadcast seed in the nursery evenly, over the water covered soil surface Seed treatment should be done
Kharif maize	Vegetative stage	Stem borer	The prevalent weather is favourable for the development of stem borer. Removed the infested plant parts.
Green gram	Flowering stage		Good drainage may be given to manage soil borne root diseases
Horticultural crop			
Citrus	Flowering to fruiting stage	Mealy bug	<ul style="list-style-type: none"> <li>In citrus orchard, destroy the ant colonies (black ants) as they act as the carriers of mealy bug. The affected plants become pale, wilt and may die also. In heavily attacked plants, citrus fruits do not set.</li> <li>Timely weeding should be carried out</li> </ul>
Okra	Flowering to fruiting stage	Yellow vein mosaic	<ul style="list-style-type: none"> <li>Remove and burn the infected plants from the field</li> <li>Remove the unwanted weed host from the field</li> <li>If the infection is severe, spray imidacloprid at 1ml/lit of water</li> </ul>

			Spotted bollworm	<ul style="list-style-type: none"> <li>➤ In the present weather condition, spotted bollworm will be seen. The infested fruits will have a varying number of holes.</li> <li>➤ Remove regularly the attacked fruits and bury deep in the soil.</li> </ul>
	<b>Cucurbits</b>	Vegetative stage	Powdery mildew	Remove the infected leaves
			Stink bug	Maintain proper sanitation as they are mostly found in weedy areas.
				<ul style="list-style-type: none"> <li>➤ For trailing, dried twigs can be spread on the ground. Weeding should be done whenever necessary.</li> </ul>
	<b>Brinjal</b>	Vegetative stage		<ul style="list-style-type: none"> <li>➤ Timely weeding and intercultural operation should be done.</li> </ul>
			Wilt	<ul style="list-style-type: none"> <li>➤ Remove and destroy the affected plant</li> <li>➤ Proper drainage should be maintained in the field</li> </ul>
	<b>Naga king chilli</b>	Vegetative stage	viral disease	<ul style="list-style-type: none"> <li>➤ Timely weeding should be done</li> <li>➤ Virus infected plants should be removed or uprooted and buried into the soil</li> <li>➤ Daily monitoring should be done for insect pest and need based neem oil should be spray</li> </ul>
	<b>Fruit trees</b>			Maintain proper sanitation in and around the vicinity of the tree
	<b>Ginger / turmeric</b>	Vegetative stage	-	Timely earthing up and weeding/hoeing should be done. Mulch the plant with dry leaves and straw
	<p style="text-align: center;"><b>Livestock</b></p> <p style="text-align: center;">Provide clean and cool drinking water to the animals</p>			
	<b>Poultry</b>	<ul style="list-style-type: none"> <li>➤ Protect the birds from rain</li> <li>➤ Keep the litter material dry</li> <li>➤ Give electrolyte in drinking water to birds of 1-6 weeks.</li> </ul>		
	<b>Piggery</b>	<ul style="list-style-type: none"> <li>➤ Vaccinate against swine fever</li> <li>➤ Maintain hygiene in pig house to protect from diarrhea</li> <li>➤ Deworm the animal at 3 month interval</li> <li>➤ Do AI, if animal comes into heat</li> </ul>		
		<b>Fisheries</b>		



- Apply lime as per recommended dose depending on the pH
- After 15 days of lime application, apply cow dung and wait till the pond water turns green.
- Put some aquatic plant in one corner of the pond for sheltering stocked fish during high temperature.

## SCIENTIFIC EXPERT COMMITTEE

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



# Integrated Agromet Advisory Service Bulletin from 21<sup>st</sup> to 25<sup>th</sup> June'17 Tuensang District

Bulletin No:47/2017

## Weather summary of the preceding week

- Moderate rain occurred the past week
- Maximum and minimum temperatures ranged 30<sup>0</sup>C to 32<sup>0</sup>C and 21<sup>0</sup>C to 22<sup>0</sup>C, respectively.
- Relative humidity varied from 51% to 95%.
- Wind speed ranged from 1 to 2 kmph

## Weather forecast valid upto 25<sup>th</sup> June'17

- Probability of light rain the coming week.
- **Max temp** is likely to be 26<sup>0</sup>C - 27<sup>0</sup>C and the **min temp** 19<sup>0</sup>C to 21<sup>0</sup>C
- **Sky is likely to be mainly cloudy** the coming week
- **Relative Humidity** is likely to range from 65% to 96%.
- **Wind speed** may reach upto 1 - 2 kmph
- **Wind direction** will be mostly southwesterly

## Field crops


Proper drainage should be maintained in the field.

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum paddy</i>	Tillering stage	Blast	<i>If the infection is severe, spray tricyclazole 0.1% at 12-15 days interval</i>
<i>TRC/WRC paddy</i>	Nursery		<i>Prepare the nursery by plowing at least twice and harrow at least once. Level the soil surface and put in drainage lines across the field. Broadcast seed in the nursery evenly, over the water covered soil surface Seed treatment should be done</i>
<i>Kharif maize</i>	Vegetative stage	Stem borer	<i>The prevalent weather is favourable for the development of stem borer. Removed the infested plant parts.</i>
<i>Green gram</i>	Flowering stage		<i>Good drainage may be given to manage soil borne root diseases</i>

## Horticultural crop

<i>Citrus</i>	Flowering to fruiting stage	Mealy bug	<ul style="list-style-type: none"> <li>➤ <i>In citrus orchard, destroy the ant colonies (black ants) as they act as the carriers of mealy bug. The affected plants become pale, wilt and may die also. In heavily attacked plants, citrus fruits do not set.</i></li> <li>➤ <i>Timely weeding should be carried out</i></li> </ul>
<i>Okra</i>	Flowering to fruiting stage	Yellow vein mosaic	<ul style="list-style-type: none"> <li>➤ <i>Remove and burn the infected plants from the field</i></li> <li>➤ <i>Remove the unwanted weed host from the field</i></li> <li>➤ <i>If the infection is severe, spray imidacloprid at 1ml/lit of water</i></li> </ul>
		Spotted bollworm	<ul style="list-style-type: none"> <li>➤ <i>In the present weather condition, spotted bollworm will be seen. The infested fruits will have a varying number of holes.</i></li> <li>➤ <i>Remove regularly the attacked fruits and bury deep in the soil.</i></li> </ul>
<i>Cucurbits</i>	Vegetative stage	Powdery	<i>Remove the infected leaves</i>



			mildew	
			Stink bug	<i>Maintain proper sanitation as they are mostly found in weedy areas.</i>
				<ul style="list-style-type: none"> <li>➤ <i>For trailing, dried twigs can be spread on the ground. Weeding should be done whenever necessary.</i></li> </ul>
	<b>Brinjal</b>	Vegetative stage		<ul style="list-style-type: none"> <li>➤ <i>Timely weeding and intercultural operation should be done.</i></li> </ul>
			Wilt	<ul style="list-style-type: none"> <li>➤ <i>Remove and destroy the affected plant</i></li> <li>➤ <i>Proper drainage should be maintained in the field</i></li> </ul>
	<b>Naga king chilli</b>	Vegetative stage	viral disease	<ul style="list-style-type: none"> <li>➤ <i>Timely weeding should be done</i></li> <li>➤ <i>Virus infected plants should be removed or uprooted and buried into the soil</i></li> <li>➤ <i>Daily monitoring should be done for insect pest and need based neem oil should be spray</i></li> </ul>
	<b>Fruit trees</b>			<i>Maintain proper sanitation in and around the vicinity of the tree</i>
	<b>Ginger / turmeric</b>	Vegetative stage	-	<i>Timely earthing up and weeding/hoeing should be done. Mulch the plant with dry leaves and straw</i>
	<p style="text-align: center;"><b>Livestock</b></p> <p style="text-align: center;">Provide clean and cool drinking water to the animals</p>			
	<b>Poultry</b>	<ul style="list-style-type: none"> <li>➤ <i>Protect the birds from rain</i></li> <li>➤ <i>Keep the litter material dry</i></li> <li>➤ <i>Give electrolyte in drinking water to birds of 1-6 weeks.</i></li> </ul>		
	<b>Piggery</b>	<ul style="list-style-type: none"> <li>➤ <i>Vaccinate against swine fever</i></li> <li>➤ <i>Maintain hygiene in pig house to protect from diarrhea</i></li> <li>➤ <i>Deworm the animal at 3 month interval</i></li> <li>➤ <i>Do AI, if animal comes into heat</i></li> </ul>		
		<b>Fisheries</b>		
		<ul style="list-style-type: none"> <li>➤ <i>Apply lime as per recommended dose depending on the pH</i></li> <li>➤ <i>After 15 days of lime application, apply cow dung and wait till the pond water turns green.</i></li> <li>➤ <i>Put some aquatic plant in one corner of the pond for sheltering stocked fish during high temperature.</i></li> </ul>		

## SCIENTIFIC EXPERT COMMITTEE


Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



## Integrated Agromet Advisory Service Bulletin from 21<sup>st</sup> to 25<sup>th</sup> June'17 Wokha District

Bulletin No:47/2017

Weather summary of the preceding week			Weather forecast valid upto 25 <sup>th</sup> June'17
<ul style="list-style-type: none"> <li>Moderate rain occurred the past week</li> <li>Maximum and minimum temperatures ranged 28<sup>0</sup>C to 31<sup>0</sup>C and 19<sup>0</sup>C to 22<sup>0</sup>C, respectively.</li> <li>Relative humidity varied from 47% to 94%.</li> <li>Wind speed ranged from 2 to 4 kmph</li> </ul>			<ul style="list-style-type: none"> <li>Probability of light rain the coming week.</li> <li><b>Max temp</b> is likely to be 24<sup>0</sup>C - 25<sup>0</sup>C and the <b>min temp</b> 21<sup>0</sup>C to 22<sup>0</sup>C</li> <li><b>Sky is likely to be mainly cloudy</b> the coming week</li> <li><b>Relative Humidity</b> is likely to range from 58% to 95%.</li> <li><b>Wind speed</b> may reach upto 1- 2 kmph</li> <li><b>Wind direction</b> will be mostly southerly</li> </ul>
Field crops			
Proper drainage should be maintained in the field.			
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Jhum paddy	Tillering stage	Blast	If the infection is severe, spray tricyclazole 0.1% at 12-15 days interval
TRC/WRC paddy	Nursery		Prepare the nursery by plowing at least twice and harrow at least once. Level the soil surface and put in drainage lines across the field. Broadcast seed in the nursery evenly, over the water covered soil surface. Seed treatment should be done
Kharif maize	Vegetative stage	Stem borer	The prevalent weather is favourable for the development of stem borer. Removed the infested plant parts.
Green gram	Flowering stage		Good drainage may be given to manage soil borne root diseases
Horticultural crop			
Citrus	Flowering to fruiting stage	Mealy bug	<ul style="list-style-type: none"> <li>In citrus orchard, destroy the ant colonies (black ants) as they act as the carriers of mealy bug. The affected plants become pale, wilt and may die also. In heavily attacked plants, citrus fruits do not set.</li> <li>Timely weeding should be carried out</li> </ul>
Okra	Flowering to fruiting stage	Yellow vein mosaic	<ul style="list-style-type: none"> <li>Remove and burn the infected plants from the field</li> <li>Remove the unwanted weed host from the field</li> <li>If the infection is severe, spray imidacloprid at 1ml/lit of water</li> </ul>
		Spotted bollworm	<ul style="list-style-type: none"> <li>In the present weather condition, spotted bollworm will be seen. The infested fruits will have a varying number of holes. Remove regularly the attacked fruits and bury deep in the soil.</li> </ul>

	<b>Cucurbits</b>	Vegetative stage	Powdery mildew	<i>Remove the infected leaves</i>
			Stink bug	<i>Maintain proper sanitation as they are mostly found in weedy areas.</i>
				➤ <i>For trailing, dried twigs can be spread on the ground. Weeding should be done whenever necessary.</i>
	<b>Brinjal</b>	Vegetative stage		➤ <i>Timely weeding and intercultural operation should be done.</i>
			Wilt	➤ <i>Remove and destroy the affected plant</i> ➤ <i>Proper drainage should be maintained in the field</i>
	<b>Naga king chilli</b>	Vegetative stage	viral disease	➤ <i>Timely weeding should be done</i> ➤ <i>Virus infected plants should be removed or uprooted and buried into the soil</i> ➤ <i>Daily monitoring should be done for insect pest and need based neem oil should be spray</i>
	<b>Fruit trees</b>			<i>Maintain proper sanitation in and around the vicinity of the tree</i>
	<b>Ginger / turmeric</b>	Vegetative stage	-	<i>Timely earthing up and weeding/hoeing should be done. Mulch the plant with dry leaves and straw</i>
	<b>Livestock</b> Provide clean and cool drinking water to the animals			
	<b>Poultry</b>	➤ Protect the birds from rain ➤ Keep the litter material dry ➤ Give electrolyte in drinking water to birds of 1-6 weeks.		
	<b>Piggery</b>	➤ Vaccinate against swine fever ➤ Maintain hygiene in pig house to protect from diarrhea ➤ Deworm the animal at 3 month interval ➤ Do AI, if animal comes into heat		
		<b>Fisheries</b>		
		➤ Apply lime as per recommended dose depending on the pH ➤ After 15 days of lime application, apply cow dung and wait till the pond water turns green. ➤ Put some aquatic plant in one corner of the pond for sheltering stocked fish during high temperature.		

## SCIENTIFIC EXPERT COMMITTEE

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com





--	--



## Integrated Agromet Advisory Service Bulletin from 21<sup>st</sup> to 25<sup>th</sup> June'17 Zunheboto District

Bulletin No:47/2017

### Weather summary of the preceding week

- Moderate rain occurred the past week
- Maximum and minimum temperatures ranged 30<sup>0</sup>C to 32<sup>0</sup>C and 21<sup>0</sup>C to 23<sup>0</sup>C, respectively.
- Relative humidity varied from 49% to 95%.
- Wind speed ranged from 1 to 2 kmph

### Weather forecast valid upto 25<sup>th</sup> June'17

- Probability of light rain the coming week.
- **Max temp** is likely to be 26<sup>0</sup>C - 27<sup>0</sup>C and the **min temp** 18<sup>0</sup>C to 20<sup>0</sup>C
- **Sky is likely to be mainly cloudy** the coming week
- **Relative Humidity** is likely to range from 61% to 95%.
- **Wind speed** may reach upto 1- 2 kmph
- **Wind direction** will be mostly southerly

### Field crops


Proper drainage should be maintained in the field.

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum paddy</i>	Tillering stage	Blast	<i>If the infection is severe, spray tricyclazole 0.1% at 12-15 days interval</i>
<i>TRC/WRC paddy</i>	Nursery		<i>Prepare the nursery by plowing at least twice and harrow at least once. Level the soil surface and put in drainage lines across the field. Broadcast seed in the nursery evenly, over the water covered soil surface Seed treatment should be done</i>
<i>Kharif maize</i>	Vegetative stage	Stem borer	<i>The prevalent weather is favourable for the development of stem borer. Removed the infested plant parts.</i>
<i>Green gram</i>	Flowering stage		<i>Good drainage may be given to manage soil borne root diseases</i>

### Horticultural crop

<i>Citrus</i>	Flowering to fruiting stage	Mealy bug	<ul style="list-style-type: none"> <li>➤ <i>In citrus orchard, destroy the ant colonies (black ants) as they act as the carriers of mealy bug. The affected plants become pale, wilt and may die also. In heavily attacked plants, citrus fruits do not set.</i></li> <li>➤ <i>Timely weeding should be carried out</i></li> </ul>
<i>Okra</i>	Flowering to fruiting stage	Yellow vein mosaic	<ul style="list-style-type: none"> <li>➤ <i>Remove and burn the infected plants from the field</i></li> <li>➤ <i>Remove the unwanted weed host from the field</i></li> <li>➤ <i>If the infection is severe, spray imidacloprid at 1ml/lit of water</i></li> </ul>
		Spotted	<ul style="list-style-type: none"> <li>➤ <i>In the present weather condition, spotted bollworm will be seen. The</i></li> </ul>



			bollworm	<i>infested fruits will have a varying number of holes. Remove regularly the attacked fruits and bury deep in the soil.</i>
	<b>Cucurbits</b>	Vegetative stage	Powdery mildew	<i>Remove the infected leaves</i>
			Stink bug	<i>Maintain proper sanitation as they are mostly found in weedy areas.</i>
				<ul style="list-style-type: none"><li>➤ <i>For trailing, dried twigs can be spread on the ground. Weeding should be done whenever necessary.</i></li></ul>
	<b>Brinjal</b>	Vegetative stage		<ul style="list-style-type: none"><li>➤ <i>Timely weeding and intercultural operation should be done.</i></li></ul>
			Wilt	<ul style="list-style-type: none"><li>➤ <i>Remove and destroy the affected plant</i></li><li>➤ <i>Proper drainage should be maintained in the field</i></li></ul>
	<b>Naga king chilli</b>	Vegetative stage	viral disease	<ul style="list-style-type: none"><li>➤ <i>Timely weeding should be done</i></li><li>➤ <i>Virus infected plants should be removed or uprooted and buried into the soil</i></li><li>➤ <i>Daily monitoring should be done for insect pest and need based neem oil should be spray</i></li></ul>
	<b>Fruit trees</b>			<i>Maintain proper sanitation in and around the vicinity of the tree</i>
	<b>Ginger / turmeric</b>	Vegetative stage	-	<i>Timely earthing up and weeding/hoeing should be done. Mulch the plant with dry leaves and straw</i>
	<b>Livestock</b> Provide clean and cool drinking water to the animals			
	<b>Poultry</b>	<ul style="list-style-type: none"><li>➤ Protect the birds from rain</li><li>➤ Keep the litter material dry</li><li>➤ Give electrolyte in drinking water to birds of 1-6 weeks.</li></ul>		
	<b>Piggery</b>	<ul style="list-style-type: none"><li>➤ Vaccinate against swine fever</li><li>➤ Maintain hygiene in pig house to protect from diarrhea</li><li>➤ Deworm the animal at 3 month interval</li><li>➤ Do AI, if animal comes into heat</li></ul>		
	<b>Fisheries</b>			
		<ul style="list-style-type: none"><li>➤ Apply lime as per recommended dose depending on the pH</li><li>➤ After 15 days of lime application, apply cow dung and wait till the pond water turns green.</li><li>➤ Put some aquatic plant in one corner of the pond for sheltering stocked fish during high temperature.</li></ul>		

## SCIENTIFIC EXPERT COMMITTEE

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com





# Integrated Agromet Advisory Service Bulletin from 21<sup>st</sup> to 25<sup>th</sup> June'17

## Dimapur District

Bulletin No:47/2017

### Weather summary of the preceding week

- Moderate rain occurred the past week
- Maximum and minimum temperatures ranged 27<sup>0</sup>C to 35<sup>0</sup>C and 23<sup>0</sup>C to 25<sup>0</sup>C, respectively.
- Relative humidity varied from 56% to 96%.
- Wind speed ranged from 1 to 2 kmph

### Weather forecast valid upto 25<sup>th</sup> June'17

- Probability of moderate rain the coming week.
- **Max temp** is likely to be 32<sup>0</sup>C - 34<sup>0</sup>C and the **min temp** 23<sup>0</sup>C to 25<sup>0</sup>C
- **Sky is likely to be mainly cloudy** the coming week
- **Relative Humidity** is likely to range from 51% to 93%.
- **Wind speed** may reach upto 1- 2 kmph
- **Wind direction** will be mostly southeasterly

### Field crops


Proper drainage should be maintained in the field.

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum paddy</i>	Tillering stage	Blast	<i>If the infection is severe, spray tricyclazole 0.1% at 12-15 days interval</i>
<i>TRC/WRC paddy</i>	Nursery		<i>Prepare the nursery by plowing at least twice and harrow at least once. Level the soil surface and put in drainage lines across the field. Broadcast seed in the nursery evenly, over the water covered soil surface Seed treatment should be done</i>
<i>Kharif maize</i>	Vegetative stage	Stem borer	<i>The prevalent weather is favourable for the development of stem borer. Removed the infested plant parts.</i>
<i>Green gram</i>	Flowering stage		<i>Good drainage may be given to manage soil borne root diseases</i>

### Horticultural crop

<i>Citrus</i>	Flowering to fruiting stage	Mealy bug	<ul style="list-style-type: none"> <li>➤ <i>In citrus orchard, destroy the ant colonies (black ants) as they act as the carriers of mealy bug. The affected plants become pale, wilt and may die also. In heavily attacked plants, citrus fruits do not set.</i></li> <li>➤ <i>Timely weeding should be carried out</i></li> </ul>
<i>Okra</i>	Flowering to fruiting stage	Yellow vein mosaic	<ul style="list-style-type: none"> <li>➤ <i>Remove and burn the infected plants from the field</i></li> <li>➤ <i>Remove the unwanted weed host from the field</i></li> <li>➤ <i>If the infection is severe, spray imidacloprid at 1ml/lit of water</i></li> </ul>
		Spotted bollworm	<ul style="list-style-type: none"> <li>➤ <i>In the present weather condition, spotted bollworm will be seen. The infested fruits will have a varying number of holes. Remove regularly the attacked fruits and bury deep in the soil.</i></li> </ul>
<i>Cucurbits</i>	Vegetative stage	Powdery	<i>Remove the infected leaves</i>

			mildew	
			Stink bug	➤ <i>Maintain proper sanitation as they are mostly found in weedy areas.</i>
				➤ <i>For trailing, dried twigs can be spread on the ground. Weeding should be done whenever necessary.</i>
	<b>Brinjal</b>	Vegetative stage		➤ <i>Timely weeding and intercultural operation should be done.</i>
			Wilt	➤ <i>Remove and destroy the affected plant</i> ➤ <i>Proper drainage should be maintained in the field</i>
	<b>Naga king chilli</b>	Vegetative stage	viral disease	➤ <i>Timely weeding should be done</i> ➤ <i>Virus infected plants should be removed or uprooted and buried into the soil</i> ➤ <i>Daily monitoring should be done for insect pest and need based neem oil should be spray</i>
	<b>Fruit trees</b>			<i>Maintain proper sanitation in and around the vicinity of the tree</i>
	<b>Ginger / turmeric</b>	Vegetative stage	-	<i>Timely earthing up and weeding/hoeing should be done. Mulch the plant with dry leaves and straw</i>
	<b><i>Livestock</i></b>			
	Provide clean and cool drinking water to the animals			
	<b>Poultry</b>	➤ Protect the birds from rain ➤ Keep the litter material dry ➤ Give electrolyte in drinking water to birds of 1-6 weeks. ➤ Good management and sanitation are the best ways to avoid infectious disease in poultry.		
<b>Piggery</b>	➤ Vaccinate against swine fever ➤ Maintain hygiene in pig house to protect from diarrhea ➤ Deworm the animal at 3 month interval ➤ Do AI, if animal comes into heat			
<b><i>Fisheries</i></b>				
	➤ Apply lime as per recommended dose depending on the pH ➤ After 15 days of lime application, apply cow dung and wait till the pond water turns green. ➤ Put some aquatic plant in one corner of the pond for sheltering stocked fish during high temperature ➤ Avoid excessive manuring and feeding which can lead to oxygen depletion in cloudy weather, hence both should be avoided.			



## SCIENTIFIC EXPERT COMMITTEE


Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



# Integrated Agromet Advisory Service Bulletin from 21<sup>st</sup> to 25<sup>th</sup> June'17 Kiphire District

Bulletin No:47/2017

Weather summary of the preceding week			Weather forecast valid upto 25 <sup>th</sup> June'17
<ul style="list-style-type: none"> <li>Moderate rain occurred the past week</li> <li>Maximum and minimum temperatures ranged 30<sup>o</sup>C to 31<sup>o</sup>C and 21<sup>o</sup>C to 24<sup>o</sup>C, respectively.</li> <li>Relative humidity varied from 52% to 95%.</li> <li>Wind speed ranged from 1 to 2 kmph</li> </ul>			<ul style="list-style-type: none"> <li>Probability of light rain the coming week.</li> <li><b>Max temp</b> is likely to be 30<sup>o</sup>C - 31<sup>o</sup>C and the <b>min temp</b> 23<sup>o</sup>C to 24<sup>o</sup>C</li> <li><b>Sky is likely to be mainly cloudy</b> the coming week</li> <li><b>Relative Humidity</b> is likely to range from 64% to 96%.</li> <li><b>Wind speed</b> may reach upto 1- 2 kmph</li> <li><b>Wind direction</b> will be mostly westerly</li> </ul>
Field crops			
Proper drainage should be maintained in the field.			
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum paddy</i>	Tillering stage	Blast	<i>If the infection is severe, spray tricyclazole 0.1% at 12-15 days interval</i>
<i>TRC/WRC paddy</i>	Nursery		<i>Prepare the nursery by plowing at least twice and harrow at least once. Level the soil surface and put in drainage lines across the field. Broadcast seed in the nursery evenly, over the water covered soil surface Seed treatment should be done</i>
<i>Kharif maize</i>	Vegetative stage	Stem borer	<i>The prevalent weather is favourable for the development of stem borer. Removed the infested plant parts.</i>
<i>Green gram</i>	Flowering stage		<i>Good drainage may be given to manage soil borne root diseases</i>
Horticultural crop			
<i>Citrus</i>	Flowering to fruiting stage	Mealy bug	<ul style="list-style-type: none"> <li>➤ <i>In citrus orchard, destroy the ant colonies (black ants) as they act as the carriers of mealy bug. The affected plants become pale, wilt and may die also. In heavily attacked plants, citrus fruits do not set.</i></li> <li>➤ <i>Timely weeding should be carried out</i></li> </ul>
<i>Okra</i>	Flowering to fruiting stage	Yellow vein mosaic	<ul style="list-style-type: none"> <li>➤ <i>Remove and burn the infected plants from the field</i></li> <li>➤ <i>Remove the unwanted weed host from the field</i></li> <li>➤ <i>If the infection is severe, spray imidacloprid at 1ml/lit of water</i></li> </ul>
		Spotted bollworm	<ul style="list-style-type: none"> <li>➤ <i>In the present weather condition, spotted bollworm will be seen. The infested fruits will have a varying number of holes.</i></li> <li><i>Remove regularly the attacked fruits and bury deep in the soil.</i></li> </ul>
<i>Cucurbits</i>	Vegetative stage	Powdery mildew	<i>Remove the infected leaves</i>

			Stink bug	Maintain proper sanitation as they are mostly found in weedy areas.
				<ul style="list-style-type: none"> <li>➤ For trailing, dried twigs can be spread on the ground. Weeding should be done whenever necessary.</li> </ul>
	<b>Brinjal</b>	Vegetative stage		<ul style="list-style-type: none"> <li>➤ Timely weeding and intercultural operation should be done.</li> </ul>
			Wilt	<ul style="list-style-type: none"> <li>➤ Remove and destroy the affected plant</li> <li>➤ Proper drainage should be maintained in the field</li> </ul>
	<b>Naga king chilli</b>	Vegetative stage	viral disease	<ul style="list-style-type: none"> <li>➤ Timely weeding should be done</li> <li>➤ Virus infected plants should be removed or uprooted and buried into the soil</li> <li>➤ Daily monitoring should be done for insect pest and need based neem oil should be spray</li> </ul>
	<b>Fruit trees</b>			Maintain proper sanitation in and around the vicinity of the tree
	<b>Ginger / turmeric</b>	Vegetative stage	-	Timely earthing up and weeding/hoeing should be done. Mulch the plant with dry leaves and straw
	<p style="text-align: center;"><b>Livestock</b></p> <p style="text-align: center;">Provide clean and cool drinking water to the animals</p>			
	<b>Poultry</b>	<ul style="list-style-type: none"> <li>➤ Protect the birds from rain</li> <li>➤ Keep the litter material dry</li> <li>➤ Give electrolyte in drinking water to birds of 1-6 weeks.</li> </ul>		
	<b>Piggery</b>	<ul style="list-style-type: none"> <li>➤ Vaccinate against swine fever</li> <li>➤ Maintain hygiene in pig house to protect from diarrhea</li> <li>➤ Deworm the animal at 3 month interval</li> <li>➤ Do AI, if animal comes into heat</li> </ul>		
		<b>Fisheries</b>		
		<ul style="list-style-type: none"> <li>➤ Apply lime as per recommended dose depending on the pH</li> <li>➤ After 15 days of lime application, apply cow dung and wait till the pond water turns green.</li> <li>➤ Put some aquatic plant in one corner of the pond for sheltering stocked fish during high temperature.</li> </ul>		



## SCIENTIFIC EXPERT COMMITTEE

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



## Integrated Agromet Advisory Service Bulletin from 21<sup>st</sup> to 25<sup>th</sup> June'17 Kohima District

Bulletin No:47/2017

Weather summary of the preceding week			Weather forecast valid upto 25 <sup>th</sup> June'17
<ul style="list-style-type: none"> <li>Moderate rain occurred the past week</li> <li>Maximum and minimum temperatures ranged 24<sup>o</sup>C to 28<sup>o</sup>C and 16<sup>o</sup>C to 17<sup>o</sup>C, respectively.</li> <li>Relative humidity varied from 55% to 98%.</li> <li>Wind speed ranged from 1 to 2 kmph</li> </ul>			<ul style="list-style-type: none"> <li>Probability of moderate rain the coming week.</li> <li><b>Max temp</b> is likely to be 24<sup>o</sup>C - 25<sup>o</sup>C and the <b>min temp</b> 15<sup>o</sup>C to 16<sup>o</sup>C</li> <li><b>Sky is likely to be mainly cloudy</b> the coming week</li> <li><b>Relative Humidity</b> is likely to range from 54% to 97%.</li> <li><b>Wind speed</b> may reach upto 1- 2 kmph</li> <li><b>Wind direction</b> will be southwesterly</li> </ul>
Field crops			
Proper drainage should be maintained in the field.			
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum paddy</i>	Tillering stage	Blast	<i>If the infection is severe, spray tricyclazole 0.1% at 12-15 days interval</i>
<i>TRC/WRC paddy</i>	Nursery		<i>Prepare the nursery by plowing at least twice and harrow at least once. Level the soil surface and put in drainage lines across the field. Broadcast seed in the nursery evenly, over the water covered soil surface Seed treatment should be done</i>
<i>Kharif maize</i>	Vegetative stage	Stem borer	<i>The prevalent weather is favourable for the development of stem borer. Removed the infested plant parts.</i>
<i>Green gram</i>	Flowering stage		<i>Good drainage may be given to manage soil borne root diseases</i>
Horticultural crop			
<i>Citrus</i>	Flowering to fruiting stage	Mealy bug	<ul style="list-style-type: none"> <li>➤ <i>In citrus orchard, destroy the ant colonies (black ants) as they act as the carriers of mealy bug. The affected plants become pale, wilt and may die also. In heavily attacked plants, citrus fruits do not set.</i></li> <li>➤ <i>Timely weeding should be carried out</i></li> </ul>
<i>Okra</i>	Flowering to fruiting stage	Yellow vein mosaic	<ul style="list-style-type: none"> <li>➤ <i>Remove and burn the infected plants from the field</i></li> <li>➤ <i>Remove the unwanted weed host from the field</i></li> <li>➤ <i>If the infection is severe, spray imidacloprid at 1ml/lit of water</i></li> </ul>
		Spotted	<ul style="list-style-type: none"> <li>➤ <i>In the present weather condition, spotted bollworm will be seen. The infested fruits will have a varying number of holes.</i></li> </ul>

			bollworm	➤ Remove regularly the attacked fruits and bury deep in the soil.
	<b>Cucurbits</b>	Vegetative stage	Powdery mildew	Remove the infected leaves
			Stink bug	Maintain proper sanitation as they are mostly found in weedy areas.
				➤ For trailing, dried twigs can be spread on the ground. Weeding should be done whenever necessary.
	<b>Brinjal</b>	Vegetative stage		➤ Timely weeding and intercultural operation should be done.
			Wilt	➤ Remove and destroy the affected plant Proper drainage should be maintained in the field
	<b>Naga king chilli</b>	Vegetative stage	viral disease	➤ Timely weeding should be done ➤ Virus infected plants should be removed or uprooted and buried into the soil ➤ Daily monitoring should be done for insect pest and need based neem oil should be spray
	<b>Fruit trees</b>			Maintain proper sanitation in and around the vicinity of the tree
	<b>Ginger / turmeric</b>	Vegetative stage	-	Timely earthing up and weeding/hoeing should be done. Mulch the plant with dry leaves and straw
	<b>Livestock</b> Provide clean and cool drinking water to the animals			
	<b>Poultry</b>	➤ Protect the birds from rain ➤ Keep the litter material dry ➤ Give electrolyte in drinking water to birds of 1-6 weeks.		
	<b>Piggery</b>	➤ Vaccinate against swine fever ➤ Maintain hygiene in pig house to protect from diarrhea ➤ Deworm the animal at 3 month interval ➤ Do AI, if animal comes into heat		
		<b>Fisheries</b>		
		➤ Apply lime as per recommended dose depending on the pH ➤ After 15 days of lime application, apply cow dung and wait till the pond water turns green. ➤ Put some aquatic plant in one corner of the pond for sheltering stocked fish during high temperature.		



## SCIENTIFIC EXPERT COMMITTEE

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com