



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 01.01.19	<b>District: Dimapur</b>	Bulletin No: 01/2019
Weather of preceding week		Weather forecast valid upto 06.01.2019
<ul> <li>No rain occurred the past week</li> </ul>		<ul> <li>Probability of no rain.</li> </ul>
• Max and min temperatures ranged 21 <sup>o</sup> C to	$22^{\circ}C$ and $10^{\circ}C$ to	• Max temp is likely to be 22 <sup>o</sup> C - 23 <sup>o</sup> C and min temp 8 <sup>o</sup> C to 9 <sup>o</sup> C
$12^{0}$ C, respectively.		Sky is likely to be clear the coming week
<ul> <li>Relative humidity varied from 45% to 90%.</li> </ul>		• <b>Relative Humidity</b> is likely to range from 38% to 72%.
• Wind speed ranged from 3 to 4 kmph		Wind speed may reach upto 4-5 kmph
		Wind may blow from south easterly direction
		Weekly cumulative Rainfall: 0mm

#### WEATHER FORECAST BASED AGROMET ADVISORIES

General Advisories:

•				
Main crop	Growth	<b>Pest/Disease</b>	Weather based Agro-Advisories	
	Stage			
TRC/WRC	Harvesting to		Keeping weather in view, harvesting and threshing should	
paddy	threshing		be postpone	
• •	stage			
Toria/Mustard	Flowering	-	Daily monitoring should be done for pest and disease	
	stage		infestation	
Rabi maize	Tasseling		Light hoeing should be done when necessary for better	
	stage		control of weeds	
Assam lemon			Weeding should be done frequently. Daily monitoring	
			should be done for aphid	
Cabbage	Pre cupping		Constant monitoring of leaf feeders in cole crop is advised	
_	stage			
Broccoli	Head		Shallow hoeing should be done to remove weeds, a light	
	initiation		earthing-up at final hoeing is beneficial.	
	stage			
Radish, carrot	Harvesting		Timely harvesting should be carried out	
D. K.	stage			
Poultry	-		ovide vitamin complex. Get ready to provide warmth by use	
		of incandescent bulbs to protect them from cold winds by use of plastic sheets, jute sacks and		
	plastic sacks			
Piggery	Protect the animals from cold draft using gunny bags and other locally available materials			
	Provide protein rich ration and ad libitum clean drinking water			
Fishery	If the pond has minimum 1m depth water, apply lime @ 50kg/bigha along with raw cow dung			
	@ 125 kg/bigha, Urea 3.5 kg/bigha, SSP 2.7 kg /bigha.			
	If water depth i	s less than 1m harv	est 50% of the stocked fish	





Date: 01.01.19	District: Kiphire	Bulletin No: 01/2019
Weather of preceding week		Weather forecast valid upto 06.01.2019
• No rain occurred the past week		<ul> <li>Probability of no rain.</li> </ul>
• Max and min temperatures ranged 23 <sup>o</sup> C to 24 <sup>o</sup> C and 7 <sup>o</sup> C to 8 <sup>o</sup> C,		• Max temp is likely to be 20 <sup>o</sup> C - 21 <sup>o</sup> C and min temp 9 <sup>o</sup> C to
respectively.		$10^{0}$ C
Relative humidity varied from 29% to 90%.		Sky is likely to be clear the coming week
• Wind speed ranged from 2 to 4 kmph		Relative Humidity is likely to range from 22% to 71%.
		Wind speed may reach upto 2-4 kmph
		Wind may blow from easterly direction
		Weekly cumulative Rainfall: 0mm

	WEATHER FORECAST BASED AGROMET ADVISORIES			
General Advisor	General Advisories:			
Main crop	Growth	<b>Pest/Disease</b>	Weather based Agro-Advisories	
	Stage			
TRC/WRC	Threshing		Keeping weather in view, threshing should be carried out	
paddy	stage		this week	
Toria/Mustard	Flowering	Aphids	In the present weather proper monitoring should be done	
	stage		for aphids	
Rabi maize	Tasseling		Light hoeing should be done when necessary for better	
	stage		control of weeds	
Cabbage	Pre Cupping	Diamond back	In the present weather daily monitoring should be done for	
	stage	moth	cabbage caterpillar.	
Radish, turnip,	Maturing		As roots have a tendency to protrude above soil surface,	
Carrot	stage		earthing up should be done immediately after weeding.	
Pea, Rajmash	Vegetative		Keeping weather in view, weeding should be done as and	
	stage		when necessary	
Tomato	Vegetative		To protect the fruits from bruises and soil borne diseases,	
	stage		make arrangement for staking and trellis of the plant	
Poultry	Chickens show	ring flu like sympto	ms should be treated with antibiotic Vendox, N-dox powder.	
	For prevention it should be given for three days and for treatment give for 5 to 7 days.			
Piggery	Provide high energy feeds like maize + wheat bran during cold days			
Fisherry				
Fishery	As far as possible do not disturb the pond. Prune the tree branches along the pond embankment			
	and allow enough sunlight to fall on the pond water surface			





Date: 01.01.19	District: Kohim	na	Bulletin No: 01/2019
Weather of preceding week		Weather	forecast valid upto 06.01.2019
• No rain occurred the past week		<ul> <li>Probability of no rain.</li> </ul>	
• Max and min temperatures ranged $22^{\circ}$ C to $23^{\circ}$ C and $6^{\circ}$ C to		• Max temp is likely to	be $16^{\circ}$ C - $18^{\circ}$ C and <b>min temp</b> $8^{\circ}$ C to $9^{\circ}$ C
8°C, respectively.		Sky is likely to be clear	
<ul> <li>Relative humidity varied from 33% to 92%.</li> </ul>		• Relative Humidity is 1	likely to range from 42% to 80%.
• Wind speed ranged from 4 to 5 kmph		• Wind speed may reach	1 upto 4-5 kmph
		• Wind may blow from	south easterly direction
		Weekly cumulative Rain	fall: 0 mm

	WEATHER FORECAST BASED AGROMET ADVISORIES			
General Advisor	General Advisories:			
Main crop	Growth	Pest/Disease	Weather based Agro-Advisories	
	Stage			
TRC/WRC	Threshing		Keeping weather in view, threshing should be carried out	
paddy	stage			
Toria/Mustard	Flowering stage	Aphids	In the present weather proper monitoring should be done for aphids	
Rabi maize	Silking stage		<i>Light hoeing should be done when necessary for better control of weeds</i>	
Pea	Vegetative stage		Proper staking up should be done	
Tomato	Vegetative stage		Spray 5% neem seed kernel extract to kill early stages larvae. Uproot if any virus infected plants are there	
Khasi mandarin			For good yield, maintain proper sanitation in the orchard	
Winter	Vegetative		Proper cultural practices should be carried out	
vegetables	stage			
Poultry	Chickens and brooding chicks should be provided warmth by use of incandescent bulbs and protection from cold winds by use of plastic sheets, jute sacks and plastic sacks.			
Piggery	Provide high energy feeds like maize + wheat bran during cold days. Always disinfect the animal shed regularly.			
Fishery	If the water lev	el is very low, pa	artially harvest some of the stocked fishes	





Date: 01.01.19	District: Longleng	Bulletin No: 01/2019
Weather of preceding week		Weather forecast valid upto 06.01.2019
• No rain occurred the past week		<ul> <li>Probability of no rain.</li> </ul>
• Max and min temperatures ranged 22°C to 23°C and 8°C to 9°C,		• Max temp is likely to be 20 <sup>o</sup> C - 21 <sup>o</sup> C and min temp 9 <sup>o</sup> C to
respectively.		$10^{0}$ C
Relative humidity varied from 45% to 84%.		Sky is likely to be clear the coming week
• Wind speed ranged from 3 to 4 kmph		• <b>Relative Humidity</b> is likely to range from 27% to 70%.
		• Wind speed may reach upto 3-4 kmph
		Wind may blow from south easterly direction
		Weekly cumulative Rainfall: 0mm

	WEATHER FORECAST BASED AGROMET ADVISORIES			
General Advisorie	s:			
Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories	
TRC/WRC paddy	Storage		As the weather will be clear with no rain grains should be dried properly before storing to bring down moisture level below 12%.	
Toria/Mustard	Flowering stage	Aphids	In the present weather proper monitoring should be done for aphids	
Rabi maize	Silking stage		Light hoeing should be done when necessary for better control of weeds	
Khasi Mandarin	Fruiting to harvesting stage		Mulching should be done with dry grasses near the tree base to conserve soil moisture	
Rajmash	Podding stage		Timely monitoring should be done	
Cabbage	Cupping stage	Diamond back moth	In the present weather daily monitoring should be done for cabbage caterpillar.	
Tomato	Vegetative stage		To protect the fruits from bruises and soil borne diseases, make arrangement for staking and trellis of the plant	
Poultry			Chicks and brooding chicks should be provided warmth by use of incandescent bulbs and protection from cold winds by use of plastic sheets, jute sacks and plastic sacks	
Piggery			As the days are getting cooler, provide high energy feeds like maize + wheat bran during cold days	
Fishery			As far as possible do not disturb the pond. Prune the tree branches along the pond embankment and allow enough sunlight to fall on the pond water surface	





Date: 01.01.19	District: Mokokchur	Ig	Bulletin No: 01/2019
Weather of preceding week		Weather f	orecast valid upto 06.01.2019
No rain occurred the past week		<ul> <li>Probability of no rain</li> </ul>	
• Max and min temperatures ranged 20 <sup>°</sup> C to	$24^{\circ}$ C and $6^{\circ}$ C to $8^{\circ}$ C,	• Max temp is likely	to be $21^{\circ}$ C - $22^{\circ}$ C and <b>min temp</b> $9^{\circ}$ C to
respectively.		$10^{0}$ C	
• Relative humidity varied from 40% to 80%.		Sky is likely to be cl	ear the coming week
• Wind speed ranged from 1 to 2 kmph		<ul> <li>Relative Humidity i</li> </ul>	s likely to range from 29% to 76%.
		Wind speed may rea	ch upto 2- 3 kmph
		• Wind may blow from	m south easterly direction
		Weekly cumulative Ra	infall: 0mm

	WEATHER FORECAST BASED AGROMET ADVISORIES			
General Advisor	ries:			
Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories	
TRC/WRC	Storage		As the weather will be clear with no rain grains should be	
paddy			dried properly before storing to bring down moisture level	
			below 12%.	
Toria/Mustard	Flowering	Aphids	In the present weather proper monitoring should be done	
	stage		for aphids	
Rabi maize	Silking stage		Light hoeing should be done when necessary for better	
			control of weeds	
Khasi	Fruiting to		Mulching should be done with dry grasses near the tree	
Mandarin	harvesting		base to conserve soil moisture	
	stage			
Rajmash	Podding stage		Timely monitoring should be done	
Cabbage	Cupping stage	Diamond back	In the present weather daily monitoring should be done for	
		moth	cabbage caterpillar.	
Tomato	Vegetative		To protect the fruits from bruises and soil borne diseases,	
D L	stage		make arrangement for staking and trellis of the plant	
Poultry			Chicks and brooding chicks should be provided warmth by	
			<i>use of incandescent bulbs and protection from cold winds</i> <i>by use of plastic sheets, jute sacks and plastic sacks</i>	
Piggery			As the days are getting cooler, provide high energy feeds	
1 18801 9			like maize + wheat bran during cold days	
Fishery			As far as possible do not disturb the pond. Prune the tree	
,			branches along the pond embankment and allow enough	
			sunlight to fall on the pond water surface	
			sumplie to juit on the pond water surjuce	





Date: 01.01.19	District: Mon	Bulletin No: 01/2019
Weather of preceding week		Weather forecast valid upto 06.01.2019
• No rain occurred the past week		<ul> <li>Probability of no rain.</li> </ul>
• Max and min temperatures ranged $22^{\circ}$ C to $22^{\circ}$ C and $5^{\circ}$ C to		• Max temp is likely to be $18^{\circ}$ C - $20^{\circ}$ C and min temp $7^{\circ}$ C to $8^{\circ}$ C
$6^{0}$ C, respectively.		Sky is likely to be clear the coming week
<ul> <li>Relative humidity varied from 47% to 95%.</li> </ul>		• <b>Relative Humidity</b> is likely to range from 28% to 78%.
• Wind speed ranged from 4 to 5 kmph		• Wind speed may reach upto 3- 4 kmph
		Wind may blow from south easterly direction
		Weekly cumulative Rainfall: 0 mm

WEATHER FORECAST BASED AGROMET ADVISORIES General Advisories: Timely intercultural operations in vegetables are advised			
Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
TRC/WRC paddy	Storage		Before storing of rice grains in storage, grains should be dried properly to bring down moisture level below 12%. Godowns should be thoroughly cleaned to destroy any insect-pests hiding in the creeks and crevices and dried before storing grains.
Potato	Vegetative stage		Try to avoid any stress to the crop as much as possible. Collect and destroy disease infected and insect infested plant parts
Toria/Mustard	Flowering stage	-	Daily monitoring should be done for pest and disease infestation
Pea	Vegetative stage		Proper staking should be done
Rabi maize	Tasseling stage		Plants should be earthed up after every weeding.
Radish	Harvesting stage		Timely harvesting should be carried out
Rajmash	Podding stage		In the present weather condition, proper mulching should be done. Collect and destroy disease infected and insect infested plant parts if there is any
Poultry			Chicks and brooding chicks should be provided warmth by use of incandescent bulbs and protection from cold winds by use of plastic sheets, jute sacks and plastic sacks Provide vitamin complex during sudden drop of temperature
Piggery			Get ready with provision of warmth for piglets during cold days in the upcoming winter with incandescent bulbs and wind breaks made from sack or plastic sheets
Fishery			If the water level is very low, partially harvest some of the stocked fishes





Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 01.01.19	9 District: Peren		Bulletin No: 01/2019
Weather of preceding week		Weather	forecast valid upto 06.01.2019
• No rain occurred the past week		<ul> <li>Probability of no rain</li> </ul>	
• Max and min temperatures ranged 20°C to 23°C and 8°C to		• Max temp is likely to	be $18^{\circ}$ C - $23^{\circ}$ C and <b>min temp</b> $5^{\circ}$ C to $6^{\circ}$ C
$10^{\circ}$ C, respectively.		Sky is likely to be clear	ar the coming week
Relative humidity varied from 41% to 94%.		Relative Humidity is	likely to range from 35% to 38%.
• Wind speed ranged from 3 to 4 kmph		Wind speed may reach	h upto 3- 4 kmph
		• Wind may blow from	south easterly direction
		Weekly cumulative Rain	ifall: 0mm

#### WEATHER FORECAST BASED AGROMET ADVISORIES

**General Advisories:** 

•	1		
Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
TRC/WRC	Harvesting to		Keeping weather in view, harvesting and threshing should
paddy	threshing		be postpone
	stage		
Toria/Mustard	Flowering	-	Daily monitoring should be done for pest and disease
	stage		infestation
Rabi maize	Tasseling		Light hoeing should be done when necessary for better
	stage		control of weeds
Assam lemon			Weeding should be done frequently. Daily monitoring
			should be done for aphid
Cabbage	Pre cupping		Constant monitoring of leaf feeders in cole crop is advised
	stage		
Broccoli	Head		Shallow hoeing should be done to remove weeds, a light
	initiation		earthing-up at final hoeing is beneficial.
	stage		
Radish, carrot	Harvesting		Timely harvesting should be carried out
	stage		
Poultry	As the temperature is dropping, provide vitamin complex. Get ready to provide warmth by use		
	of incandescent bulbs to protect them from cold winds by use of plastic sheets, jute sacks and		
	plastic sacks		
Piggery	Protect the animals from cold draft using gunny bags and other locally available materials		
	Provide protein rich ration and ad libitum clean drinking water		
Fishery	If the pond has minimum 1m depth water, apply lime @ 50kg/bigha along with raw cow dung		
	@ 125 kg/bigha, Urea 3.5 kg/bigha, SSP 2.7 kg /bigha.		
	If water depth is less than 1m harvest 50% of the stocked fish		





Date: 01.01.19	District: Phek	Bulletin No: 01/2019	
Weather of preceding week		Weather forecast valid upto 06.01.2019	
• No rain occurred the past week		<ul> <li>Probability of no rain.</li> </ul>	
• Max and min temperatures ranged 22 <sup>o</sup> C to 24 <sup>o</sup> C and 6 <sup>o</sup> C to		• Max temp is likely to be $18^{\circ}$ C - $20^{\circ}$ C and min temp $7^{\circ}$ C to $8^{\circ}$ C	
9 <sup>°</sup> C, respectively.		• Sky is likely to be clear the coming week	
Relative humidity varied from 32% to 94%.		• <b>Relative Humidity</b> is likely to range from 28% to 80%.	
• Wind speed ranged from 2 to 3 kmph		Wind speed may reach upto 2- 3 kmph	
		<ul> <li>Wind may blow mostly south easterly direction</li> </ul>	
		Weekly cumulative Rainfall: 0mm	

WEATHER FORECAST BASED AGROMET ADVISORIES General Advisories:				
		Ge		
Main crop	Growth	<b>Pest/Disease</b>	Weather based Agro-Advisories	
	Stage			
TRC/WRC	Threshing		Keeping weather in view, threshing should be carried out	
paddy	stage			
Toria/Mustard	Flowering	Aphids	In the present weather proper monitoring should be done for	
	stage		aphids	
Rabi maize	Silking stage		Light hoeing should be done when necessary for better control	
			of weeds	
Pea	Vegetative		Proper staking up should be done	
	stage			
Tomato	Vegetative		Spray 5% neem seed kernel extract to kill early stages larvae.	
771 • 1 •	stage		Uproot if any virus infected plants are there	
Khasi mandarin			For good yield, maintain proper sanitation in the orchard	
117.	Manadationa			
Winter	Vegetative		Proper cultural practices should be carried out	
vegetables	stage			
Poultry	Chickens and brooding chicks should be provided warmth by use of incandescent bulbs and			
	protection from cold winds by use of plastic sheets, jute sacks and plastic sacks.			
Piggery	Provide high energy feeds like maize + wheat bran during cold days. Always disinfect the			
	animal shed regularly.			
Fishery	If the water level is very low, partially harvest some of the stocked fishes			





Date: 01.01.19	District: Tuensang	Bulletin No: 01/2019
Weather of preceding week		Weather forecast valid upto 06.01.2019
• No rain occurred the past week		<ul> <li>Probability of no rain.</li> </ul>
• Max and min temperatures ranged $22^{\circ}$ C to $24^{\circ}$ C and $4^{\circ}$ C to $9^{\circ}$ C,		• Max temp is likely to be $16^{\circ}$ C - $18^{\circ}$ C and min temp $8^{\circ}$ C to
respectively.		9 <sup>0</sup> C
<ul> <li>Relative humidity varied from 30% to 70%.</li> </ul>		• Sky is likely to be clear the coming week
Wind speed ranged from 2 to 4 kmph		• <b>Relative Humidity</b> is likely to range from 24% to 80%.
		<ul> <li>Wind speed may reach upto 2- 3 kmph</li> </ul>
		<ul> <li>Wind may blow from south easterly direction</li> </ul>
		Weekly cumulative Rainfall: 0mm

WEATHER FORECAST BASED AGROMET ADVISORIES General Advisories:			
Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
TRC/WRC	Threshing		Keeping weather in view, threshing should be carried out
paddy	stage		this week
Toria/Mustard	Flowering	Aphids	In the present weather proper monitoring should be done
	stage		for aphids
Rabi maize	Tasseling		Light hoeing should be done when necessary for better
	stage		control of weeds
Cabbage	Pre Cupping	Diamond back	In the present weather daily monitoring should be done for
	stage	moth	cabbage caterpillar.
Radish, turnip,	Maturing		As roots have a tendency to protrude above soil surface,
Carrot	stage		earthing up should be done immediately after weeding.
Pea, Rajmash	Vegetative		Keeping weather in view, weeding should be done as and
	stage		when necessary
Tomato	Vegetative		To protect the fruits from bruises and soil borne diseases,
	stage		make arrangement for staking and trellis of the plant
Poultry	Chickens showing flu like symptoms should be treated with antibiotic Vendox, N-dox powder.		
	For prevention it should be given for three days and for treatment give for 5 to 7 days.		
Piggery	Provide high energy feeds like maize + wheat bran during cold days		
Fishery	As far as possible do not disturb the pond. Prune the tree branches along the pond embankment		
	and allow enough sunlight to fall on the pond water surface		
		igh sunnght to fall C	n ne pone water surface





Date: 01.01.19	District: Wokha	]	Bulletin No: 01/2019
Weather of preceding week		Weather for	recast valid upto 06.01.2019
• No rain occurred the past week		<ul> <li>Probability of no rain.</li> </ul>	
• Max and min temperatures ranged $22^{\circ}$ C to $24^{\circ}$ C and $10^{\circ}$ C to $11^{\circ}$ C		• Max temp is likely t	o be $18^{\circ}$ C - $20^{\circ}$ C and <b>min temp</b> $7^{\circ}$ C to
respectively.		$8^{0}C$	
Relative humidity varied from 60% to 95%.		Sky is likely to be clear	ar the coming week
Wind speed ranged from 3 to 4 kmph		Relative Humidity is	likely to range from 35% to 78%.
		Wind speed may reac	h upto 3- 4 kmph
		Wind may blow from	n easterly direction
		Weekly cumulative Rain	ıfall: Omm

WEATHER FORECAST BASED AGROMET ADVISORIES				
General Advisories:				
Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories	
TRC/WRC paddy	Storage		As the weather will be clear with no rain grains should be dried properly before storing to bring down moisture level below 12%.	
Toria/Mustard	Flowering stage	Aphids	In the present weather proper monitoring should be done for aphids	
Rabi maize	Silking stage		Light hoeing should be done when necessary for better control of weeds	
Khasi Mandarin	Fruiting to harvesting stage		Mulching should be done with dry grasses near the tree base to conserve soil moisture	
Rajmash	Podding stage		Timely monitoring should be done	
Cabbage	Cupping stage	Diamond back moth	In the present weather daily monitoring should be done for cabbage caterpillar.	
Tomato	Vegetative stage		To protect the fruits from bruises and soil borne diseases, make arrangement for staking and trellis of the plant	
Poultry			Chicks and brooding chicks should be provided warmth by use of incandescent bulbs and protection from cold winds by use of plastic sheets, jute sacks and plastic sacks	
Piggery			As the days are getting cooler, provide high energy feeds like maize + wheat bran during cold days	
Fishery			As far as possible do not disturb the pond. Prune the tree branches along the pond embankment and allow enough sunlight to fall on the pond water surface	





Date: 01.01.19	<b>District: Zunheboto</b>	Bulletin No: 01/2019
Weather of preceding week		Weather forecast valid upto 06.01.2019
• No rain occurred the past week		<ul> <li>Probability of no rain .</li> </ul>
• Max and min temperatures ranged $22^{\circ}$ C to $23^{\circ}$ C and $7^{\circ}$ C to $8^{\circ}$ C,		• Max temp is likely to be $18^{\circ}$ C - $20^{\circ}$ C and min temp $7^{\circ}$ C to
respectively.		$8^{0}C$
Relative humidity varied from 42% to 90%.		Sky is likely to be clear the coming week
• Wind speed ranged from 2 to 3 kmph		• <b>Relative Humidity</b> is likely to range from 34% to 80%.
		Wind speed may reach upto 3-4 kmph
		Wind may blow from south easterly direction
		Weekly cumulative Rainfall: 0mm

WEATHER FORECAST BASED AGROMET ADVISORIES					
<b>General Advisori</b>	General Advisories:				
Main crop	Growth	Pest/Disease	Weather based Agro-Advisories		
TRC/WRC paddy	Storage		As the weather will be clear with no rain grains should be dried properly before storing to bring down moisture level below 12%.		
Toria/Mustard	Flowering stage	Aphids	In the present weather proper monitoring should be done for aphids		
Rabi maize	Silking stage		Light hoeing should be done when necessary for better control of weeds		
Khasi Mandarin	Fruiting to harvesting stage		Mulching should be done with dry grasses near the tree base to conserve soil moisture		
Rajmash	Podding stage		Timely monitoring should be done		
Cabbage	Cupping stage	Diamond back moth	In the present weather daily monitoring should be done for cabbage caterpillar.		
Tomato	Vegetative stage		To protect the fruits from bruises and soil borne diseases, make arrangement for staking and trellis of the plant		
Poultry			Chicks and brooding chicks should be provided warmth by use of incandescent bulbs and protection from cold winds by use of plastic sheets, jute sacks and plastic sacks		
Piggery			As the days are getting cooler, provide high energy feeds like maize + wheat bran during cold days		
Fishery			As far as possible do not disturb the pond. Prune the tree branches along the pond embankment and allow enough sunlight to fall on the pond water surface		