



“GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland

Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in



Date: 22.01.19		District: Kiphire		Bulletin No: 07/2019	
Weather of preceding week			Weather forecast valid upto 27.01.2019		
<ul style="list-style-type: none">No rain occurred the past weekMax and min temperatures ranged 24⁰C to 25⁰C and 11⁰C to 12⁰C, respectively.Relative humidity varied from 23% to 87%.Wind speed ranged from 2 to 4 kmph			<ul style="list-style-type: none">Probability of no rain.Max temp is likely to be 23⁰C - 24⁰C and min temp 9⁰C to 10⁰CSky is likely to be clear the coming weekRelative Humidity is likely to range from 23% to 80%.Wind speed may reach upto 2-4 kmphWind may blow from easterly direction		
			Weekly cumulative Rainfall: 0mm		
WEATHER FORECAST BASED AGROMET ADVISORIES					
General Advisories: In view of falling both maximum and minimum temperature, Farmers are advised for light irrigation at evening time to protect the crop from probable frost.					
Main crop		Growth Stage	Pest/Disease	Weather based Agro-Advisories	
TRC/WRC paddy		Storage		Storage rooms should be physically rodent and bird proof, if possible, place sticky traps in the drying and storage areas for rats.	
Toria/Mustard		Maturing stage		In the present weather, as soon as the pods turn yellowish-brown, harvest the crop in order to avoid losses. Cleaned seed must be dried in the sun for four to five days	
Cabbage		Cupping stage		Daily monitoring should be done for pest and diseases	
Radish, turnip, Carrot		Harvesting stage		In the present weather, timely harvesting should be done	
Pea, Rajmash		Flowering stage		Keeping weather in view, weeding should be done as and when necessary	
Tomato		Flowering stage		To protect the fruits from bruises and soil borne diseases, make arrangement for staking and trellis of the plant	
Poultry		Protect the birds from cold draft by covering the exposed windward areas using gunny bags and other locally available materials Chickens showing flu like symptoms should be treated with antibiotic Vendox, N-dox powder. For prevention it should be given for three days and for treatment give for 5 to 7 days			
Piggery		Protect the animals from cold draft using gunny bags and other locally available materials Provide high energy feeds like maize + wheat bran during cold days			
Fishery		As the water temperature of your pond falls, the fish will require less food. Never over feed your fish. Excess food will pollute the pond water, so only feed amounts that will be completely consumed by your fish within 2 minutes			



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland

Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in



Date: 22.01.19	District: Kohima	Bulletin No: 07/2019
Weather of preceding week		Weather forecast valid upto 27.01.2019
<ul style="list-style-type: none">No rain occurred the past weekMax and min temperatures ranged 20⁰C to 21⁰C and 6⁰C to 7⁰C, respectively.Relative humidity varied from 37% to 92%.Wind speed ranged from 4 to 5 kmph		<ul style="list-style-type: none">Probability of no rain.Max temp is likely to be 20⁰C - 21⁰C and min temp 6⁰C to 8⁰CSky is likely to be clear the coming weekRelative Humidity is likely to range from 33% to 80%.Wind speed may reach upto 4-5 kmphWind may blow from south easterly direction Weekly cumulative Rainfall: 0 mm

WEATHER FORECAST BASED AGROMET ADVISORIES

General Advisories: In view of falling both maximum and minimum temperature, Farmers are advised for light irrigation at evening time to protect the crop from probable frost.

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
TRC/WRC paddy	Storage		Storage rooms should be physically rodent and bird proof, if possible, place sticky traps in the drying and storage areas for rats.
Toria/Mustard	Podding stage	Aphids	In the present weather proper monitoring should be done for aphids
Pea	Podding stage		Daily inspect the field frequently for mature pods as all the pods do not mature at the same time.
Tomato	Flowering stage		Spray 5% neem seed kernel extract to kill early stages larvae. Uproot if any virus infected plants are there
Khasi mandarin			If harvesting is over, water sprout, diseases and dried or dead twigs should be removed and cut end pasted with Bordeaux paste
Winter vegetables	Harvesting stage		Timely harvesting should be carried out
Poultry	Protect the birds from cold draft by covering the exposed windward areas using gunny bags and other locally available materials Deworm the birds once every 2 months Sick birds showing respiratory distress, drooping of head or any other illness should be promptly segregated and the entire flock be treated against the illness.		
Piggery	Sick or injured animals should be attended to immediately, preferably move to a warm comfortable environment. Medications should be used only on the advice of a veterinarian, following recommended treatment		
Fishery	As the water temperature of your pond falls, the fish will require less food. Never over feed your fish. Excess food will pollute the pond water, so only feed amounts that will be completely consumed by your fish within 2 minutes		



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 22.01.19	District: Longleng	Bulletin No: 07/2019
Weather of preceding week		Weather forecast valid upto 27.01.2019
<ul style="list-style-type: none"> No rain occurred the past week Max and min temperatures ranged 24⁰C to 25⁰C and 11⁰C to 12⁰C, respectively. Relative humidity varied from 27% to 87%. Wind speed ranged from 3 to 4 kmph 		<ul style="list-style-type: none"> Probability of no rain. Max temp is likely to be 23⁰C - 24⁰C and min temp 9⁰C to 10⁰C Sky is likely to be clear the coming week Relative Humidity is likely to range from 27% to 80%. Wind speed may reach upto 4-5 kmph Wind may blow from south easterly direction
		Weekly cumulative Rainfall: 0mm

WEATHER FORECAST BASED AGROMET ADVISORIES

General Advisories: In view of falling both maximum and minimum temperature, Farmers are advised for light irrigation at evening time to protect the crop from probable frost.

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
TRC/WRC paddy	Storage		Keep storage areas, crevices, wooden pallets etc clean by spraying with the recommended insecticide to disinfect insect breeding places. Place sticky traps in the drying and storage areas for rats
Toria/Mustard	Podding stage		Monitor the fields closely to detect any insect problems
Khasi Mandarin	Harvesting stage		If harvesting is over, water sprout, diseases and dried or dead twigs should be removed and cut end pasted with Bordeaux paste
Rajmash	Harvesting stage		Harvesting should be done timely
Cabbage	Maturing stage		The crop is ready for harvest and should be done timely as it may split if not harvested in time.
Tomato	Fruiting to harvesting stage		To protect the fruits from bruises and soil borne diseases, make arrangement for staking and trellis of the plant
Poultry			Seal all the cracks and crevices in the house Side curtains can be used during severe winds and draft thus sealing the house from cold draft
Piggery			Sick or injured animals should be attended to immediately, preferably move to a warm comfortable environment. Medications should be used only on the advice of a veterinarian, following recommended treatment
Fishery			As far as possible do not disturb the pond. Prune the tree branches along the pond embankment and allow enough sunlight to fall on the pond water surface



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland

Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in



Date: 22.01.19	District: Mokokchung	Bulletin No: 07/2019
Weather of preceding week		Weather forecast valid upto 27.01.2019
<ul style="list-style-type: none"> No rain occurred the past week Max and min temperatures ranged 24⁰C to 25⁰C and 11⁰C to 12⁰C, respectively. Relative humidity varied from 26% to 87%. Wind speed ranged from 2 to 3 kmph 		<ul style="list-style-type: none"> Probability of no rain. Max temp is likely to be 23⁰C - 24⁰C and min temp 8⁰C to 9⁰C Sky is likely to be clear the coming week Relative Humidity is likely to range from 25% to 80%. Wind speed may reach upto 2- 4 kmph Wind may blow from south easterly direction
		Weekly cumulative Rainfall: 0mm

WEATHER FORECAST BASED AGROMET ADVISORIES

General Advisories: In view of falling both maximum and minimum temperature, Farmers are advised for light irrigation at evening time to protect the crop from probable frost.

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
TRC/WRC paddy	Storage		Keep storage areas, crevices, wooden pallets etc clean by spraying with the recommended insecticide to disinfect insect breeding places. Place sticky traps in the drying and storage areas for rats
Toria/Mustard	Podding stage		Monitor the fields closely to detect any insect problems
Khasi Mandarin	Harvesting stage		If harvesting is over, water sprout, diseases and dried or dead twigs should be removed and cut end pasted with Bordeaux paste
Rajmash	Harvesting stage		Harvesting should be done timely
Cabbage	Maturing stage		The crop is ready for harvest and should be done timely as it may split if not harvested in time.
Tomato	Fruiting to harvesting stage		To protect the fruits from bruises and soil borne diseases, make arrangement for staking and trellis of the plant
Poultry			Seal all the cracks and crevices in the house Side curtains can be used during severe winds and draft thus sealing the house from cold draft
Piggery			Sick or injured animals should be attended to immediately, preferably move to a warm comfortable environment. Medications should be used only on the advice of a veterinarian, following recommended treatment
Fishery			As far as possible do not disturb the pond. Prune the tree branches along the pond embankment and allow enough sunlight to fall on the pond water surface



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 22.01.19	District: Mon	Bulletin No: 07/2019
Weather of preceding week		Weather forecast valid upto 27.01.2019
<ul style="list-style-type: none"> No rain occurred the past week Max and min temperatures ranged 22⁰C to 23⁰C and 8⁰C to 10⁰C, respectively. Relative humidity varied from 29% to 85%. Wind speed ranged from 3 to 4 kmph 		<ul style="list-style-type: none"> Probability of no rain. Max temp is likely to be 22⁰C - 23⁰C and min temp 7⁰C to 8⁰C Sky is likely to be clear the coming week Relative Humidity is likely to range from 32% to 80%. Wind speed may reach upto 4- 5 kmph Wind may blow from south easterly direction Weekly cumulative Rainfall: 0 mm

WEATHER FORECAST BASED AGROMET ADVISORIES
General Advisories: In view of falling both maximum and minimum temperature, Farmers are advised for light irrigation at evening time to protect the crop from probable frost.

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
<i>TRC/WRC paddy</i>	Storage		<i>Inspect the stored seeds once a week for signs of insect infestation.</i>
<i>Potato</i>	Vegetative stage		<i>Try to avoid any stress to the crop as much as possible. Collect and destroy disease infected and insect infested plant parts</i>
<i>Toria/Mustard</i>	Podding stage	-	<i>Left over weeds before shading of seeds may be removed to reduce the weed seed bank/spread.</i>
<i>Pea</i>	Fruiting stage		<i>Weather is favourable for picking of pods.</i>
<i>Winter vegetables</i>	Harvesting stage		<i>Timely harvesting should be carried out</i>
<i>Rajmash</i>	Podding stage		<i>In the present weather condition, proper mulching should be done. Collect and destroy disease infected and insect infested plant parts if there is any</i>
<i>Poultry</i>			Provide vitamin complex during sudden drop of temperature Chickens showing flu like symptoms should be treated with antibiotic Vendox, N-dox powder. For prevention it should be given for three days and for treatment give for 5 to 7 days
<i>Piggery</i>			Protect the animals from cold draft using gunny bags and other locally available materials Provide high energy feeds like maize + wheat bran during cold days
<i>Fishery</i>			As the water temperature of your pond falls, the fish will require less food. Never over feed your fish. Excess food will pollute the pond water, so only feed amounts that will be completely consumed by your fish within 2 minutes



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 22.01.19	District: Peren	Bulletin No: 07/2019
Weather of preceding week		Weather forecast valid upto 27.01.2019
<ul style="list-style-type: none"> No rain occurred the past week Max and min temperatures ranged 22^oC to 23^oC and 8^oC to 10^oC, respectively. Relative humidity varied from 32% to 85%. Wind speed ranged from 3 to 4 kmph 		<ul style="list-style-type: none"> Probability of no rain Max temp is likely to be 22^oC - 23^oC and min temp 8^oC to 9^oC Sky is likely to be clear the coming week Relative Humidity is likely to range from 30% to 80%. Wind speed may reach upto 3- 4 kmph Wind may blow from south easterly direction
		Weekly cumulative Rainfall: 0mm

WEATHER FORECAST BASED AGROMET ADVISORIES

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
TRC/WRC paddy	Storage		Keep storage areas, crevices, wooden pallets etc clean by spraying with the recommended insecticide to disinfect insect breeding places. Place sticky traps in the drying and storage areas for rats
Toria/Mustard	Podding stage	-	Left over weeds before shading of seeds may be removed to reduce the weed seed bank/spread.
Assam lemon			Weeding should be done frequently. Daily monitoring should be done for aphid
Cabbage	Maturing stage		The crop is ready for harvest and should be done timely as it may split if not harvested in time.
Winter vegetables	Harvesting stage		Timely harvesting should be carried out
Poultry	Protect the birds from cold draft by covering the exposed windward areas using gunny bags and other locally available materials Deworm the birds once every 2 months Sick birds showing respiratory distress, drooping of head or any other illness should be promptly segregated and the entire flock be treated against the illness.		
Piggery	Sick or injured animals should be attended to immediately, preferably move to a warm comfortable environment. Medications should be used only on the advice of a veterinarian, following recommended treatment		
Fishery	As the water temperature of your pond falls, the fish will require less food. Never over feed your fish. Excess food will pollute the pond water, so only feed amounts that will be completely consumed by your fish within 2 minutes		



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland

Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in



Date: 22.01.19	District: Phek	Bulletin No: 07/2019
Weather of preceding week		Weather forecast valid upto 27.01.2019
<ul style="list-style-type: none">No rain occurred the past weekMax and min temperatures ranged 20⁰C to 21⁰C and 6⁰C to 7⁰C, respectively.Relative humidity varied from 28% to 90%.Wind speed ranged from 2 to 3 kmph		<ul style="list-style-type: none">Probability of no rain.Max temp is likely to be 20⁰C - 22⁰C and min temp 7⁰C to 8⁰CSky is likely to be clear the coming weekRelative Humidity is likely to range from 23% to 80%.Wind speed may reach upto 2- 3 kmphWind may blow mostly south easterly direction Weekly cumulative Rainfall: 0mm

WEATHER FORECAST BASED AGROMET ADVISORIES

General Advisories: In view of falling both maximum and minimum temperature, Farmers are advised for light irrigation at evening time to protect the crop from probable frost.

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
TRC/WRC paddy	Storage		Storage rooms should be physically rodent and bird proof, if possible, place sticky traps in the drying and storage areas for rats.
Toria/Mustard	Podding stage	Aphids	In the present weather proper monitoring should be done for aphids
Pea	Podding stage		Daily inspect the field frequently for mature pods as all the pods do not mature at the same time.
Tomato	Flowering stage		Spray 5% neem seed kernel extract to kill early stages larvae. Uproot if any virus infected plants are there
Khasi mandarin			If harvesting is over, water sprout, diseases and dried or dead twigs should be removed and cut end pasted with Bordeaux paste
Winter vegetables	Harvesting stage		Timely harvesting should be carried out
Poultry	Protect the birds from cold draft by covering the exposed windward areas using gunny bags and other locally available materials Deworm the birds once every 2 months Sick birds showing respiratory distress, drooping of head or any other illness should be promptly segregated and the entire flock be treated against the illness.		
Piggery	Sick or injured animals should be attended to immediately, preferably move to a warm comfortable environment. Medications should be used only on the advice of a veterinarian, following recommended treatment		
Fishery	As the water temperature of your pond falls, the fish will require less food. Never over feed your fish. Excess food will pollute the pond water, so only feed amounts that will be completely consumed by your fish within 2 minutes		



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 22.01.19	District: Tuensang	Bulletin No: 07/2019
Weather of preceding week		Weather forecast valid upto 27.01.2019
<ul style="list-style-type: none"> No rain occurred the past week Max and min temperatures ranged 18⁰C to 19⁰C and 3⁰C to 4⁰C, respectively. Relative humidity varied from 22% to 90%. Wind speed ranged from 2 to 3 kmph 		<ul style="list-style-type: none"> Probability of no rain. Max temp is likely to be 18⁰C - 19⁰C and min temp 7⁰C to 9⁰C Sky is likely to be clear the coming week Relative Humidity is likely to range from 20% to 89%. Wind speed may reach upto 2- 3 kmph Wind may blow from south easterly direction Weekly cumulative Rainfall: 0mm

WEATHER FORECAST BASED AGROMET ADVISORIES

General Advisories: In view of falling both maximum and minimum temperature, Farmers are advised for light irrigation at evening time to protect the crop from probable frost.

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
TRC/WRC paddy	Storage		<i>Storage rooms should be physically rodent and bird proof, if possible, place sticky traps in the drying and storage areas for rats.</i>
Toria/Mustard	Maturing stage		<i>In the present weather, as soon as the pods turn yellowish-brown, harvest the crop in order to avoid losses. Cleaned seed must be dried in the sun for four to five days</i>
Cabbage	Cupping stage		<i>Daily monitoring should be done for pest and diseases</i>
Radish, turnip, Carrot	Harvesting stage		<i>In the present weather, timely harvesting should be done</i>
Pea, Rajmash	Flowering stage		<i>Keeping weather in view, weeding should be done as and when necessary</i>
Tomato	Flowering stage		<i>To protect the fruits from bruises and soil borne diseases, make arrangement for staking and trellis of the plant</i>
Poultry	Protect the birds from cold draft by covering the exposed windward areas using gunny bags and other locally available materials Chickens showing flu like symptoms should be treated with antibiotic Vendox, N-dox powder. For prevention it should be given for three days and for treatment give for 5 to 7 days		
Piggery	Protect the animals from cold draft using gunny bags and other locally available materials Provide high energy feeds like maize + wheat bran during cold days		
Fishery	As the water temperature of your pond falls, the fish will require less food. Never over feed your fish. Excess food will pollute the pond water, so only feed amounts that will be completely consumed by your fish within 2 minutes		



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland

Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in



Date: 22.01.19	District: Wokha	Bulletin No: 07/2019
Weather of preceding week		Weather forecast valid upto 27.01.2019
<ul style="list-style-type: none"> No rain occurred the past week Max and min temperatures ranged 22^oC to 23^oC and 8^oC to 10^oC respectively. Relative humidity varied from 34% to 95%. Wind speed ranged from 3 to 4 kmph 		<ul style="list-style-type: none"> Probability of no rain . Max temp is likely to be 22^oC - 24^oC and min temp 8^oC to 9^oC Sky is likely to be clear the coming week Relative Humidity is likely to range from 32% to 85%. Wind speed may reach upto 3- 4 kmph Wind may blow from easterly direction Weekly cumulative Rainfall: 0mm

WEATHER FORECAST BASED AGROMET ADVISORIES

General Advisories: In view of falling both maximum and minimum temperature, Farmers are advised for light irrigation at evening time to protect the crop from probable frost.

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
TRC/WRC paddy	Storage		Keep storage areas, crevices, wooden pallets etc clean by spraying with the recommended insecticide to disinfect insect breeding places. Place sticky traps in the drying and storage areas for rats
Toria/Mustard	Podding stage		Monitor the fields closely to detect any insect problems
Khasi Mandarin	Harvesting stage		If harvesting is over, water sprout, diseases and dried or dead twigs should be removed and cut end pasted with Bordeaux paste
Rajmash	Harvesting stage		Harvesting should be done timely
Cabbage	Maturing stage		The crop is ready for harvest and should be done timely as it may split if not harvested in time.
Tomato	Fruiting to harvesting stage		To protect the fruits from bruises and soil borne diseases, make arrangement for staking and trellis of the plant
Poultry			Seal all the cracks and crevices in the house Side curtains can be used during severe winds and draft thus sealing the house from cold draft
Piggery			Sick or injured animals should be attended to immediately, preferably move to a warm comfortable environment. Medications should be used only on the advice of a veterinarian, following recommended treatment
Fishery			As far as possible do not disturb the pond. Prune the tree branches along the pond embankment and allow enough sunlight to fall on the pond water surface



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland

Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in



Date: 22.01.19	District: Zunheboto	Bulletin No: 07/2019
Weather of preceding week		Weather forecast valid upto 27.01.2019
<ul style="list-style-type: none"> No rain occurred the past week Max and min temperatures ranged 21⁰C to 22⁰C and 8⁰C to 10⁰C, respectively. Relative humidity varied from 27% to 87%. Wind speed ranged from 2 to 3 kmph 		<ul style="list-style-type: none"> Probability of no rain. Max temp is likely to be 22⁰C - 23⁰C and min temp 8⁰C to 9⁰C Sky is likely to be clear the coming week Relative Humidity is likely to range from 29% to 80%. Wind speed may reach upto 3-4 kmph Wind may blow from south easterly direction
		Weekly cumulative Rainfall: 0mm

WEATHER FORECAST BASED AGROMET ADVISORIES

General Advisories: In view of falling both maximum and minimum temperature, Farmers are advised for light irrigation at evening time to protect the crop from probable frost.

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
TRC/WRC paddy	Storage		Keep storage areas, crevices, wooden pallets etc clean by spraying with the recommended insecticide to disinfect insect breeding places. Place sticky traps in the drying and storage areas for rats
Toria/Mustard	Podding stage		Monitor the fields closely to detect any insect problems
Khasi Mandarin	Harvesting stage		If harvesting is over, water sprout, diseases and dried or dead twigs should be removed and cut end pasted with Bordeaux paste
Rajmash	Harvesting stage		Harvesting should be done timely
Cabbage	Maturing stage		The crop is ready for harvest and should be done timely as it may split if not harvested in time.
Tomato	Fruiting to harvesting stage		To protect the fruits from bruises and soil borne diseases, make arrangement for staking and trellis of the plant
Poultry			Seal all the cracks and crevices in the house Side curtains can be used during severe winds and draft thus sealing the house from cold draft
Piggery			Sick or injured animals should be attended to immediately, preferably move to a warm comfortable environment. Medications should be used only on the advice of a veterinarian, following recommended treatment
Fishery			As far as possible do not disturb the pond. Prune the tree branches along the pond embankment and allow enough sunlight to fall on the pond water surface



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 22.01.19		District: Dimapur		Bulletin No: 07/2019	
Weather of preceding week			Weather forecast valid upto 27.01.2019		
<ul style="list-style-type: none">▪ No rain occurred the past week▪ Max and min temperatures ranged 21⁰C to 26⁰C and 8⁰C to 10⁰C, respectively.▪ Relative humidity varied from 34% to 67%.▪ Wind speed ranged from 4 to 6 kmph			<ul style="list-style-type: none">▪ Probability of no rain.▪ Max temp is likely to be 28⁰C - 29⁰C and min temp 11⁰C to 12⁰C▪ Sky is likely to be clear the coming week▪ Relative Humidity is likely to range from 31% to 78%.▪ Wind speed may reach upto 4-6 kmph▪ Wind may blow from south easterly direction Weekly cumulative Rainfall: 0mm		
WEATHER FORECAST BASED AGROMET ADVISORIES					
General Advisories					
Main crop		Growth Stage	Pest/Disease	Weather based Agro-Advisories	
TRC/WRC paddy		Storage		Keep storage areas, crevices, wooden pallets etc clean by spraying with the recommended insecticide to disinfect insect breeding places. Place sticky traps in the drying and storage areas for rats	
Toria/Mustard		Podding stage	-	Left over weeds before shading of seeds may be removed to reduce the weed seed bank/spread.	
Assam lemon				Weeding should be done frequently. Daily monitoring should be done for aphid	
Cabbage		Maturing stage		The crop is ready for harvest and should be done timely as it may split if not harvested in time.	
Winter vegetables		Harvesting stage		Timely harvesting should be carried out	
Poultry		Oil or fat can be added to feed and energy levels in the feed can be raised to meet the higher energy need of the bird. Provide warm water to the birds Chickens showing flu like symptoms should be treated with antibiotic Vendox, N-dox powder. For prevention it should be given for three days and for treatment give for 5 to 7 days.			
Piggery		Protect the animals from cold draft using gunny bags and other locally available materials Provide high energy feeds like maize + wheat bran during cold days			
Fishery		As the water temperature of your pond falls, the fish will require less food. Never over feed your fish. Excess food will pollute the pond water, so only feed amounts that will be completely consumed by your fish within 2 minutes			