



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 25.01.19		District: Dimapur		Bulletin No: 08/2019	
Weather of preceding week			Weather forecast valid upto 30.01.2019		
<ul style="list-style-type: none">▪ No rain occurred the past week▪ Max and min temperatures ranged 21⁰C to 26⁰C and 8⁰C to 10⁰C, respectively.▪ Relative humidity varied from 34% to 67%.▪ Wind speed ranged from 4 to 6 kmph			<ul style="list-style-type: none">▪ Probability of light rain on 29th or 30th.▪ Max temp is likely to be 26⁰C - 27⁰C and min temp 8⁰C to 11⁰C▪ Sky is likely to be partly cloudy the coming week▪ Relative Humidity is likely to range from 34% to 85%.▪ Wind speed may reach upto 4-6 kmph▪ Wind may blow from south easterly direction		
WEATHER FORECAST BASED AGROMET ADVISORIES					
General Advisories					
Main crop		Growth Stage	Pest/Disease	Weather based Agro-Advisories	
TRC/WRC paddy		Storage		Keep storage areas, crevices, wooden pallets etc clean by spraying with the recommended insecticide to disinfect insect breeding places. Place sticky traps in the drying and storage areas for rats	
Toria/Mustard		Podding stage	Leaf feeders	In the present weather spray plant product extract like neem till 28 th Left over weeds before shading of seeds may be removed to reduce the weed seed bank/spread.	
Assam lemon/Citrus				In the present weather, water sprout, diseases and dried/dead twigs of mandarin, lime should be removed and cut end pasted with Bordeaux paste	
Cabbage		Maturing to harvesting stage		The crop is ready for harvest and should be done timely as it may split if not harvested in time and increased incidence of field disease After harvesting, remove the entire stem and root system from the soil to prevent disease buildup	
Cucurbitaceous vegetable				Prepare seedling of Cucurbitaceous vegetable in polythene bag for getting early produce .	
Poultry		Oil or fat can be added to feed and energy levels in the feed can be raised to meet the higher energy need of the bird. Provide warm water to the birds Chickens showing flu like symptoms should be treated with antibiotic Vendox, N-dox powder. For prevention it should be given for three days and for treatment give for 5 to 7 days.			
Piggery		Protect the animals from cold draft using gunny bags and other locally available materials Provide high energy feeds like maize + wheat bran during cold days			
Fishery		As the water temperature of your pond falls, the fish will require less food. Never over feed your fish. Excess food will pollute the pond water, so only feed amounts that will be completely consumed by your fish within 2 minutes			
Livestock		For maintaining milk production in milch animals, they should be protected from low temperature.			



“GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland

Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in



Date: 25.01.19	District: Kiphire	Bulletin No: 08/2019
Weather of preceding week		Weather forecast valid upto 30.01.2019
<ul style="list-style-type: none"> No rain occurred the past week Max and min temperatures ranged 24⁰C to 25⁰C and 11⁰C to 12⁰C, respectively. Relative humidity varied from 23% to 87%. Wind speed ranged from 2 to 4 kmph 		<ul style="list-style-type: none"> Probability of light rain on 29th or 30th. Max temp is likely to be 22⁰C - 23⁰C and min temp 10⁰C to 11⁰C Sky is likely to be partly cloudy the coming week Relative Humidity is likely to range from 27% to 85%. Wind speed may reach upto 2-4 kmph Wind may blow from south easterly direction
		Weekly cumulative Rainfall: 5mm

WEATHER FORECAST BASED AGROMET ADVISORIES

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
TRC/WRC paddy	Storage		Storage rooms should be physically rodent and bird proof, if possible, place sticky traps in the drying and storage areas for rats.
Toria/Mustard	Maturing stage		In the present weather, as soon as the pods turn yellowish-brown, harvest the crop in order to avoid losses but it should be done before 29 th . Cleaned seed must be dried in the sun for four to five days
Cabbage	Cupping stage		Daily monitoring should be done for pest and diseases
Cucurbitaceous vegetable			Prepare seedling of Cucurbitaceous vegetable in polythene bag for getting early produce.
Pea, Rajmash	Flowering to podding stage		Weather is favourable for picking of podstill 28 th
Khasi Mandarin	Harvesting stage		In the present weather, water sprout, diseases and dried/dead twigs of mandarin should be removed after harvesting and cut end pasted with Bordeaux paste
Poultry	Artificial heating through lighting of electric bulb at poultry house during night time may protect the birds from fall of night temperature and reduce the incidence of mortality. Chickens showing flu like symptoms should be treated with antibiotic Vendox, N-dox powder. For prevention it should be given for three days and for treatment give for 5 to 7 days		
Piggery	Protect the animals from cold draft using gunny bags and other locally available materials. Provide high energy feeds like maize + wheat bran during cold days		
Fishery	As the water temperature of your pond falls, the fish will require less food. Never over feed your fish. Excess food will pollute the pond water, so only feed amounts that will be completely consumed by your fish within 2 minutes		



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 25.01.19	District: Kohima	Bulletin No: 08/2019
Weather of preceding week		Weather forecast valid upto 30.01.2019
<ul style="list-style-type: none"> No rain occurred the past week Max and min temperatures ranged 20⁰C to 21⁰C and 6⁰C to 7⁰C, respectively. Relative humidity varied from 37% to 92%. Wind speed ranged from 4 to 5 kmph 		<ul style="list-style-type: none"> Probability of light rain on 29th or 30th Max temp is likely to be 21⁰C - 22⁰C and min temp 9⁰C to 10⁰C Sky is likely to be partly cloudy the coming week Relative Humidity is likely to range from 34% to 90%. Wind speed may reach upto 4-5 kmph Wind may blow from south easterly direction Weekly cumulative Rainfall: 10 mm

WEATHER FORECAST BASED AGROMET ADVISORIES

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
TRC/WRC paddy	Storage		Storage rooms should be physically rodent and bird proof, if possible, place sticky traps in the drying and storage areas for rats.
Toria/Mustard	Podding to maturing stage	-	Left over weeds before shading of seeds may be removed to reduce the weed seed bank/spread. Timely harvesting should be done to avoid losses due to shattering
Pea	Podding stage		Daily inspect the field frequently for mature pods as all the pods do not mature at the same time.
Cucurbitaceous vegetable			Prepare seedling of Cucurbitaceous vegetable in polythene bag for getting early produce .
Khasi mandarin			If harvesting is over, water sprout, diseases and dried or dead twigs should be removed and cut end pasted with Bordeaux paste
Tomato	Fruiting to harvesting stage		To protect the fruits from bruises and soil borne diseases, make arrangement for staking and trellis of the plant
Poultry	Oil or fat can be added to feed and energy levels in the feed can be raised to meet the higher energy need of the bird. Provide warm water to the birds Sick birds showing respiratory distress, drooping of head or any other illness should be promptly segregated and the entire flock be treated against the illness.		
Piggery	Protect the animals from cold draft using gunny bags and other locally available materials Provide high energy feeds like maize + wheat bran during cold days		
Fishery	As the water temperature of your pond falls, the fish will require less food. Never over feed your fish. Excess food will pollute the pond water, so only feed amounts that will be completely consumed by your fish within 2 minutes		
Livestock	For maintaining milk production in milch animals, they should be protected from low temperature.		



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 25.01.19	District: Longleng	Bulletin No: 08/2019
Weather of preceding week		Weather forecast valid upto 30.01.2019
<ul style="list-style-type: none"> No rain occurred the past week Max and min temperatures ranged 24⁰C to 25⁰C and 11⁰C to 12⁰C, respectively. Relative humidity varied from 27% to 87%. Wind speed ranged from 3 to 4 kmph 		<ul style="list-style-type: none"> Probability of no rain. Max temp is likely to be 22⁰C - 23⁰C and min temp 10⁰C to 11⁰C Sky is likely to be partly cloudy the coming week Relative Humidity is likely to range from 28% to 90%. Wind speed may reach upto 4-5 kmph Wind may blow from south easterly direction
		Weekly cumulative Rainfall: 0mm

WEATHER FORECAST BASED AGROMET ADVISORIES

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
TRC/WRC paddy	Storage		Keep storage areas, crevices, wooden pallets etc clean by spraying with the recommended insecticide to disinfect insect breeding places. Place sticky traps in the drying and storage areas for rats
Toria/Mustard	Podding stage		Monitor the fields closely to detect any insect problems
Khasi Mandarin	Harvesting stage		In the present weather, water sprout, diseases and dried/dead twigs of mandarin should be removed after harvesting and cut end pasted with Bordeaux paste
Rajmash	Harvesting stage		Harvesting should be done timely
Cabbage	Maturing to harvesting stage		The crop is ready for harvest and should be done timely as it may split if not harvested in time and increased incidence of field disease After harvesting, remove the entire stem and root system from the soil to prevent disease buildup
Tomato	Fruiting to harvesting stage		To protect the fruits from bruises and soil borne diseases, make arrangement for staking and trellis of the plant
Poultry			Chickens showing flu like symptoms should be treated with antibiotic Vendox, N-dox powder. For prevention it should be given for three days and for treatment give for 5 to 7 days
Piggery			Put up windbreaks and protection from excess cold Look out for skin conditions and treat if necessary Wash the sties with potassium permanganate solution a week atleast Deworm the pig with suitable anthelminthis FMD and classical swine fever vaccination is to be done
Fishery			As far as possible do not disturb the pond. Prune the tree branches along the pond embankment and allow enough sunlight to fall on the pond water surface



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 25.01.19	District: Mokokchung	Bulletin No: 08/2019
Weather of preceding week		Weather forecast valid upto 30.01.2019
<ul style="list-style-type: none"> No rain occurred the past week Max and min temperatures ranged 24⁰C to 25⁰C and 11⁰C to 12⁰C, respectively. Relative humidity varied from 26% to 87%. Wind speed ranged from 2 to 3 kmph 		<ul style="list-style-type: none"> Probability of light rain on 29th or 30th. Max temp is likely to be 22⁰C - 23⁰C and min temp 10⁰C to 11⁰C Sky is likely to be partly cloudy the coming week Relative Humidity is likely to range from 25% to 80%. Wind speed may reach upto 2- 4 kmph Wind may blow from south easterly direction
		Weekly cumulative Rainfall: 5mm

WEATHER FORECAST BASED AGROMET ADVISORIES

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
TRC/WRC paddy	Storage		Keep storage areas, crevices, wooden pallets etc clean by spraying with the recommended insecticide to disinfect insect breeding places. Place sticky traps in the drying and storage areas for rats.
Toria/Mustard	Podding stage		Monitor the fields closely to detect any insect problems
Khasi Mandarin	Harvesting stage		In the present weather, water sprout, diseases and dried/dead twigs of mandarin should be removed after harvesting and cut end pasted with Bordeaux paste
Rajmash	Harvesting stage		Harvesting should be done timely and try to do it before 29 th
Cabbage	Maturing to harvesting stage		The crop is ready for harvest and should be done timely as it may split if not harvested in time and increased incidence of field disease After harvesting, remove the entire stem and root system from the soil to prevent disease buildup
Tomato	Fruiting to harvesting stage		To protect the fruits from bruises and soil borne diseases, make arrangement for staking and trellis of the plant
Poultry			Chickens showing flu like symptoms should be treated with antibiotic Vendox, N-dox powder. For prevention it should be given for three days and for treatment give for 5 to 7 days
Piggery			Put up windbreaks and protection from excess cold Look out for skin conditions and treat if necessary Wash the sties with potassium permanganate solution a week atleast Deworm the pig with suitable anthelminthis FMD and classical swine fever vaccination is to be done
Fishery			As far as possible do not disturb the pond. Prune the tree branches along the pond embankment and allow enough sunlight to fall on the pond water surface



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 25.01.19	District: Mon	Bulletin No: 08/2019
Weather of preceding week		Weather forecast valid upto 30.01.2019
<ul style="list-style-type: none"> No rain occurred the past week Max and min temperatures ranged 22°C to 23°C and 8°C to 10°C, respectively. Relative humidity varied from 29% to 85%. Wind speed ranged from 3 to 4 kmph 		<ul style="list-style-type: none"> Probability of light rain on 29th or 30th. Max temp is likely to be 21°C - 22°C and min temp 9°C to 10°C Sky is likely to be mainly cloudy the coming week Relative Humidity is likely to range from 32% to 90%. Wind speed may reach upto 4- 5 kmph Wind may blow from south easterly direction
		Weekly cumulative Rainfall: 8 mm

WEATHER FORECAST BASED AGROMET ADVISORIES			
Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
TRC/WRC paddy	Storage		Inspect the stored seeds once a week for signs of insect infestation.
Potato	Tuber initiation stage		Fungal diseases like blight in potato may occur. Mancozeb (Indofil M-45) @ 2.5gm per litter of water may be sprayed in the potato field at least twice at an interval of 10 days. As precautionary measures filed must be inspected twice in a day
Toria/Mustard	Podding to maturing stage	-	Left over weeds before shading of seeds may be removed to reduce the weed seed bank/spread. Timely harvesting should be done to avoid losses due to shattering, try to finish it before 29 th Jan
Pea	Fruiting stage		Weather is favourable for picking of pods till 28 th .
Cucurbitaceous vegetable			Prepare seedling of Cucurbitaceous vegetable in polythene bag for getting early produce.
Rajmash	Podding stage		In the present weather condition, proper mulching should be done. Collect and destroy disease infected and insect infested plant parts if there is any
Poultry			Artificial heating through lighting of electric bulb at poultry house during night time may protect the birds from fall of night temperature and reduce the incidence of mortality. Chickens showing flu like symptoms should be treated with antibiotic Vendox, N-dox powder. For prevention it should be given for three days and for treatment give for 5 to 7 days
Piggery			Protect the animals from cold draft using gunny bags and other locally available materials Provide high energy feeds like maize + wheat bran during cold days
Fishery			As the water temperature of your pond falls, the fish will require less food. Never over feed your fish. Excess food will pollute the pond water, so only feed amounts that will be completely consumed by your fish within 2 minutes



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 25.01.19		District: Peren	Bulletin No: 08/2019
Weather of preceding week		Weather forecast valid upto 30.01.2019	
<ul style="list-style-type: none">▪ No rain occurred the past week▪ Max and min temperatures ranged 22⁰C to 23⁰C and 8⁰C to 10⁰C, respectively.▪ Relative humidity varied from 32% to 85%.▪ Wind speed ranged from 3 to 4 kmph		<ul style="list-style-type: none">▪ Probability of no rain▪ Max temp is likely to be 21⁰C - 22⁰C and min temp 9⁰C to 10⁰C▪ Sky is likely to be clear the coming week▪ Relative Humidity is likely to range from 31% to 80%.▪ Wind speed may reach upto 3- 4 kmph▪ Wind may blow from south easterly direction Weekly cumulative Rainfall: 0mm	
WEATHER FORECAST BASED AGROMET ADVISORIES			
Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
TRC/WRC paddy	Storage		Keep storage areas, crevices, wooden pallets etc clean by spraying with the recommended insecticide to disinfect insect breeding places. Place sticky traps in the drying and storage areas for rats
Toria/Mustard	Podding stage	Leaf feeders	In the present weather spray plant product extract like neem till 28 th Left over weeds before shading of seeds may be removed to reduce the weed seed bank/spread.
Assam lemon/Citrus			In the present weather, water sprout, diseases and dried/dead twigs of mandarin, lime should be removed and cut end pasted with Bordeaux paste
Cabbage	Maturing to harvesting stage		The crop is ready for harvest and should be done timely as it may split if not harvested in time and increased incidence of field disease After harvesting, remove the entire stem and root system from the soil to prevent disease buildup
Cucurbitaceous vegetable			Prepare seedling of Cucurbitaceous vegetable in polythene bag for getting early produce .
Poultry	Oil or fat can be added to feed and energy levels in the feed can be raised to meet the higher energy need of the bird. Provide warm water to the birds Chickens showing flu like symptoms should be treated with antibiotic Vendox, N-dox powder. For prevention it should be given for three days and for treatment give for 5 to 7 days.		
Piggery	Protect the animals from cold draft using gunny bags and other locally available materials Provide high energy feeds like maize + wheat bran during cold days		
Fishery	As the water temperature of your pond falls, the fish will require less food. Never over feed your fish. Excess food will pollute the pond water, so only feed amounts that will be completely consumed by your fish within 2 minutes		
Livestock	For maintaining milk production in milch animals, they should be protected from low temperature.		



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland

Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in



Date: 25.01.19	District: Phek	Bulletin No: 08/2019
Weather of preceding week		Weather forecast valid upto 30.01.2019
<ul style="list-style-type: none"> No rain occurred the past week Max and min temperatures ranged 20⁰C to 21⁰C and 6⁰C to 7⁰C, respectively. Relative humidity varied from 28% to 90%. Wind speed ranged from 2 to 3 kmph 		<ul style="list-style-type: none"> Probability of light rain on 29th or 30th. Max temp is likely to be 19⁰C - 20⁰C and min temp 6⁰C to 7⁰C Sky is likely to be partly cloudy the coming week Relative Humidity is likely to range from 31% to 89%. Wind speed may reach upto 2- 3 kmph Wind may blow mostly south easterly direction
		Weekly cumulative Rainfall: 13mm

WEATHER FORECAST BASED AGROMET ADVISORIES

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
TRC/WRC paddy	Storage		Storage rooms should be physically rodent and bird proof, if possible, place sticky traps in the drying and storage areas for rats.
Toria/Mustard	Podding to maturing stage	-	Left over weeds before shading of seeds may be removed to reduce the weed seed bank/spread. Timely harvesting should be done to avoid losses due to shattering
Pea	Podding stage		Daily inspect the field frequently for mature pods as all the pods do not mature at the same time.
Cucurbitaceous vegetable			Prepare seedling of Cucurbitaceous vegetable in polythene bag for getting early produce.
Khasi mandarin			If harvesting is over, water sprout, diseases and dried or dead twigs should be removed and cut end pasted with Bordeaux paste
Tomato	Fruiting to harvesting stage		To protect the fruits from bruises and soil borne diseases, make arrangement for staking and trellis of the plant
Poultry	Oil or fat can be added to feed and energy levels in the feed can be raised to meet the higher energy need of the bird. Provide warm water to the birds Sick birds showing respiratory distress, drooping of head or any other illness should be promptly segregated and the entire flock be treated against the illness.		
Piggery	Protect the animals from cold draft using gunny bags and other locally available materials Provide high energy feeds like maize + wheat bran during cold days		
Fishery	As the water temperature of your pond falls, the fish will require less food. Never over feed your fish. Excess food will pollute the pond water, so only feed amounts that will be completely consumed by your fish within 2 minutes		
Livestock	For maintaining milk production in milch animals, they should be protected from low temperature.		



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland

Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in



Date: 25.01.19	District: Tuensang	Bulletin No: 08/2019
Weather of preceding week		Weather forecast valid upto 30.01.2019
<ul style="list-style-type: none"> No rain occurred the past week Max and min temperatures ranged 18⁰C to 19⁰C and 3⁰C to 4⁰C, respectively. Relative humidity varied from 22% to 90%. Wind speed ranged from 2 to 3 kmph 		<ul style="list-style-type: none"> Probability of light rain on 29th or 30th. Max temp is likely to be 19⁰C - 20⁰C and min temp 6⁰C to 8⁰C Sky is likely to be partly cloudy the coming week Relative Humidity is likely to range from 24% to 85%. Wind speed may reach upto 2- 3 kmph Wind may blow from south easterly direction
		Weekly cumulative Rainfall: 4mm

WEATHER FORECAST BASED AGROMET ADVISORIES

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
TRC/WRC paddy	Storage		Storage rooms should be physically rodent and bird proof, if possible, place sticky traps in the drying and storage areas for rats.
Toria/Mustard	Maturing stage		In the present weather, as soon as the pods turn yellowish-brown, harvest the crop in order to avoid losses but it should be done before 29 th . Cleaned seed must be dried in the sun for four to five days
Cabbage	Cupping stage		Daily monitoring should be done for pest and diseases
Cucurbitaceous vegetable			Prepare seedling of Cucurbitaceous vegetable in polythene bag for getting early produce.
Pea, Rajmash	Flowering to podding stage		Weather is favourable for picking of podstill 28 th
Khasi Mandarin	Harvesting stage		In the present weather, water sprout, diseases and dried/dead twigs of mandarin should be removed after harvesting and cut end pasted with Bordeaux paste
Poultry	Artificial heating through lighting of electric bulb at poultry house during night time may protect the birds from fall of night temperature and reduce the incidence of mortality Chickens showing flu like symptoms should be treated with antibiotic Vendox, N-dox powder. For prevention it should be given for three days and for treatment give for 5 to 7 days		
Piggery	Protect the animals from cold draft using gunny bags and other locally available materials Provide high energy feeds like maize + wheat bran during cold days		
Fishery	As the water temperature of your pond falls, the fish will require less food. Never over feed your fish. Excess food will pollute the pond water, so only feed amounts that will be completely consumed by your fish within 2 minutes		



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 25.01.19		District: Wokha		Bulletin No: 08/2019	
Weather of preceding week			Weather forecast valid upto 30.01.2019		
<ul style="list-style-type: none">No rain occurred the past weekMax and min temperatures ranged 22⁰C to 23⁰C and 8⁰C to 10⁰C respectively.Relative humidity varied from 34% to 95%.Wind speed ranged from 3 to 4 kmph			<ul style="list-style-type: none">Probability of light rain on 29th or 30th.Max temp is likely to be 21⁰C - 22⁰C and min temp 9⁰C to 10⁰CSky is likely to be partly cloudy the coming weekRelative Humidity is likely to range from 32% to 90%.Wind speed may reach upto 3- 4 kmphWind may blow from south easterly direction		
			Weekly cumulative Rainfall: 7mm		
WEATHER FORECAST BASED AGROMET ADVISORIES					
Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories		
TRC/WRC paddy	Storage		Keep storage areas, crevices, wooden pallets etc clean by spraying with the recommended insecticide to disinfect insect breeding places. Place sticky traps in the drying and storage areas for rats		
Toria/Mustard	Podding stage		Monitor the fields closely to detect any insect problems		
Khasi Mandarin	Harvesting stage		In the present weather, water sprout, diseases and dried/dead twigs of mandarin should be removed after harvesting and cut end pasted with Bordeaux paste		
Rajmash	Harvesting stage		Harvesting should be done timely		
Cabbage	Maturing to harvesting stage		The crop is ready for harvest and should be done timely as it may split if not harvested in time and increased incidence of field disease After harvesting, remove the entire stem and root system from the soil to prevent disease buildup		
Tomato	Fruiting to harvesting stage		To protect the fruits from bruises and soil borne diseases, make arrangement for staking and trellis of the plant		
Poultry			Chickens showing flu like symptoms should be treated with antibiotic Vendox, N-dox powder. For prevention it should be given for three days and for treatment give for 5 to 7 days		
Piggery			Put up windbreaks and protection from excess cold Look out for skin conditions and treat if necessary Wash the sties with potassium permanganate solution a week atleast Deworm the pig with suitable anthelminthis FMD and classical swine fever vaccination is to be done		
Fishery			As far as possible do not disturb the pond. Prune the tree branches along the pond embankment and allow enough sunlight to fall on the pond water surface		



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland

Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in



Date: 25.01.19	District: Zunheboto	Bulletin No: 08/2019
Weather of preceding week		Weather forecast valid upto 30.01.2019
<ul style="list-style-type: none"> No rain occurred the past week Max and min temperatures ranged 21⁰C to 22⁰C and 8⁰C to 10⁰C, respectively. Relative humidity varied from 27% to 87%. Wind speed ranged from 2 to 3 kmph 		<ul style="list-style-type: none"> Probability of light rain on 29th or 30th. Max temp is likely to be 21⁰C - 22⁰C and min temp 9⁰C to 10⁰C Sky is likely to be partly cloudy the coming week Relative Humidity is likely to range from 27% to 90%. Wind speed may reach upto 3-4 kmph Wind may blow from south easterly direction
		Weekly cumulative Rainfall: 5mm

WEATHER FORECAST BASED AGROMET ADVISORIES

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
TRC/WRC paddy	Storage		Keep storage areas, crevices, wooden pallets etc clean by spraying with the recommended insecticide to disinfect insect breeding places. Place sticky traps in the drying and storage areas for rats
Toria/Mustard	Podding stage		Monitor the fields closely to detect any insect problems
Khasi Mandarin	Harvesting stage		In the present weather, water sprout, diseases and dried/dead twigs of mandarin should be removed after harvesting and cut end pasted with Bordeaux paste
Rajmash	Harvesting stage		Harvesting should be done timely
Cabbage	Maturing to harvesting stage		The crop is ready for harvest and should be done timely as it may split if not harvested in time and increased incidence of field disease After harvesting, remove the entire stem and root system from the soil to prevent disease buildup
Tomato	Fruiting to harvesting stage		To protect the fruits from bruises and soil borne diseases, make arrangement for staking and trellis of the plant
Poultry			Chickens showing flu like symptoms should be treated with antibiotic Vendox, N-dox powder. For prevention it should be given for three days and for treatment give for 5 to 7 days
Piggery			Put up windbreaks and protection from excess cold Look out for skin conditions and treat if necessary Wash the sties with potassium permanganate solution a week atleast Deworm the pig with suitable anthelminthis FMD and classical swine fever vaccination is to be done
Fishery			As far as possible do not disturb the pond. Prune the tree branches along the pond embankment and allow enough sunlight to fall on the pond water surface

