



**GRAMIN KRISHI MAUSAM SEWA**  
**ICAR RESEARCH COMPLEX FOR NEH REGION**  
**Nagaland Center, Medziphema-797106**  
**Dimapur, Nagaland**



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 26.03.19		District: Dimapur		Bulletin No: 25/2019	
Weather of preceding week			Weather forecast valid upto 31.03.2019		
<ul style="list-style-type: none"><li>▪ Light rain occurred the past week</li><li>▪ Max and min temperatures ranged 29<sup>0</sup>C to 31<sup>0</sup>C and 17<sup>0</sup>C to 19<sup>0</sup>C, respectively.</li><li>▪ Relative humidity varied from 27% to 80%.</li><li>▪ Wind speed ranged from 5 to 6 kmph</li></ul>			<ul style="list-style-type: none"><li>▪ Probability of light rain</li><li>▪ <b>Max temp</b> is likely to be 28<sup>0</sup>C - 31<sup>0</sup>C and <b>min temp</b> 16<sup>0</sup>C to 17<sup>0</sup>C</li><li>▪ <b>Sky is likely to be mainly clear</b> the coming week</li><li>▪ <b>Relative Humidity</b> is likely to range from 24% to 81%.</li><li>▪ <b>Wind speed</b> may reach upto 4-6 kmph</li><li>▪ <b>Wind may blow from south easterly direction</b></li></ul> <b>Weekly cumulative Rainfall:</b> 1mm		
WEATHER FORECAST BASED AGROMET ADVISORIES					
General Advisories: Short duration pulse crops and green manuring crops like dhaincha, sunnhemp etc should be planted as pre-kharif crop in fallow paddy lowland/ Terrace Rice field to enhance the soil fertility					
Main crop		Growth Stage	Pest/Disease	Weather based Agro-Advisories	
TRC/WRC paddy		Storage		Keep storage areas, crevices, wooden pallets etc clean by spraying with the recommended insecticide to disinfect insect breeding places. Place sticky traps in the drying and storage areas for rats	
Green gram		Sowing		Sowing should be carried out in the present weather	
Naga king chilli		Nursery preparation	Damping off	Before sowing of seeds, seed treatment should be done with Thiram @2gm/litre of water. Avoid over watering and close spacing to avoid damping off	
Assam Lemon				Propagation through cuttings or seeds should be started before the onset of hot weather for citrus.	
Okra		Sowing	Hoppers and white flies	While germinating, install yellow sticky trap for monitoring white flies and hoppers	
Cucurbitaceous vegetable		Germination	Fruit fly	Apply neem cake @ 250kg/ha immediately after germination	
Poultry		For day old chicks provide cool water and electrolytes on their arrival to farm before offering feed to avoid dehydration after transportation Supply of plenty of clean and cool water must be ensured during summer months. Provide Electral Energy (1-2 gm/liter) in water during hot hours.			
Piggery		As temperature and humidity rises, pay extra attention to the movement of animals. Provide adequate water Prevent overcrowding			
Fishery		As most of the pond is dry, take up renovation works like repairing and cleaning of bottom of the pond			



**“GRAMIN KRISHI MAUSAM SEWA**  
**ICAR RESEARCH COMPLEX FOR NEH REGION**  
**Nagaland Center, Medziphema-797106**  
**Dimapur, Nagaland**



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 26.03.19		District: Kiphire		Bulletin No: 25/2019	
Weather of preceding week			Weather forecast valid upto 31.03.2019		
<ul style="list-style-type: none"><li>▪ Light rain occurred the past week</li><li>▪ Max and min temperatures ranged 25<sup>0</sup>C to 30<sup>0</sup>C and 15<sup>0</sup>C to 18<sup>0</sup>C, respectively.</li><li>▪ Relative humidity varied from 24% to 73%.</li><li>▪ Wind speed ranged from 3 to 4 kmph</li></ul>			<ul style="list-style-type: none"><li>▪ Probability of light rain</li><li>▪ <b>Max temp</b> is likely to be 28<sup>0</sup>C -30<sup>0</sup>C and <b>min temp</b> 13<sup>0</sup>C to 14<sup>0</sup>C</li><li>▪ <b>Sky is likely to be mainly clear</b> the coming week</li><li>▪ <b>Relative Humidity</b> is likely to range from 23% to 79%.</li><li>▪ <b>Wind speed</b> may reach upto 3-4 kmph</li><li>▪ <b>Wind may blow from southerly direction</b></li></ul> <b>Weekly cumulative Rainfall:</b> 12 mm		
WEATHER FORECAST BASED AGROMET ADVISORIES					
General Recommendations: In the present weather, start sowing of summer crops					
Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories		
Jhum paddy	Sowing	Blast	To manage blast, use high quality, disease free seeds for sowing, Destroy the infested residue, stubbles, grass and weeds present around the field. They act as a source for the infection		
Potato	Harvesting stage		Regular monitoring of pest and disease is advised Crop rotation should be carried out to reduce soil borne pests and diseases		
Rajmash	Vegetative stage		In the present weather, Gap filling should be carried out		
Cucurbitaceous vegetable	Germination	Fruit fly	Apply neem cake @ 250kg/ha immediately after germination		
Brinjal	Nursery	Damping off	To manage damping off, drench the beds with bavistin@15-20 gm/litres of water		
King chilli	Nursery	Damping off	In the present weather, watering should be postponed as there is probability of light rain the coming days too. For damping off, drenching should be done with Thiram@2gm/litre of water at 10 – 15 days interval Start preparation of the main field, use well rotten FYM or vermin compost		
Poultry	For day old chicks provide cool water and electrolytes on their arrival to farm before offering feed to avoid dehydration after transportation Supply of plenty of clean and cool water must be ensured during summer months. Provide Electral Energy (1-2 gm/liter) in water during hot hours				
Piggery	As temperature and humidity rises, pay extra attention to the movement of animals. Provide adequate water Prevent overcrowding				
Fishery	As most of the pond is dry, take up renovation works like repairing and cleaning of bottom of the pond				



**GRAMIN KRISHI MAUSAM SEWA**  
**ICAR RESEARCH COMPLEX FOR NEH REGION**  
**Nagaland Center, Medziphema-797106**  
**Dimapur, Nagaland**

Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in



Date: 26.03.19	District: Kohima	Bulletin No: 25/2019
<b>Weather of preceding week</b>		<b>Weather forecast valid upto 31.03.2019</b>
<ul style="list-style-type: none"> <li>Light rain occurred the past week</li> <li>Max and min temperatures ranged 29<sup>0</sup>C to 32<sup>0</sup>C and 15<sup>0</sup>C to 18<sup>0</sup>C, respectively.</li> <li>Relative humidity varied from 24% to 82%.</li> <li>Wind speed ranged from 5 to 6 kmph</li> </ul>		<ul style="list-style-type: none"> <li>Probability of light rain</li> <li><b>Max temp</b> is likely to be 25<sup>0</sup>C - 27<sup>0</sup>C and <b>min temp</b> 11<sup>0</sup>C to 12<sup>0</sup>C</li> <li><b>Sky is likely to be mainly clear</b> the coming week</li> <li><b>Relative Humidity</b> is likely to range from 32% to 83%.</li> <li><b>Wind speed</b> may reach upto 2-7 kmph</li> <li><b>Wind may blow from south easterly direction</b></li> <li><b>Weekly cumulative Rainfall:</b> 22 mm</li> </ul>

**WEATHER FORECAST BASED AGROMET ADVISORIES**

**General Advisories:** Short duration pulse crops and green manuring crops like dhaincha, sunnhemp etc should be planted as pre-kharif crop in fallow paddy lowland/ Terrace Rice field to enhance the soil fertility

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
TRC/WRC paddy	Storage		Place sticky traps in the drying and storage areas for rats
Potato	Vegetative stage	Early blight, leaf spot	Regular monitoring of disease should be done. Avoid cultivation of solanaceous crops like tomato, brinjal, chillies etc. nearby potato fields Fields should be kept neat and clean, free from weeds
Maize	Land preparation	White grubs	Deep ploughing should be done to expose the resting stages of pest. Collect and destroy the white grubs
Cucurbitaceous vegetable	Germination	Fruit fly	Apply neem cake @ 250kg/ha immediately after germination
Khasi mandarin			Prepare basin around the plants followed by mulching with local grasses.
Summer crops			Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. and certified seeds should be used to produce good yield.
Poultry	For day old chicks provide cool water and electrolytes on their arrival to farm before offering feed to avoid dehydration after transportation Supply of plenty of clean and cool water must be ensured during summer months. Provide Electral Energy (1-2 gm/liter) in water during hot hours.		
Piggery	As temperature and humidity rises, pay extra attention to the movement of animals. Provide adequate water Prevent overcrowding		
Fishery	As most of the pond is dry, take up renovation works like repairing and cleaning of bottom of the pond		



**GRAMIN KRISHI MAUSAM SEWA**  
**ICAR RESEARCH COMPLEX FOR NEH REGION**  
**Nagaland Center, Medziphema-797106**  
**Dimapur, Nagaland**



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

<b>Date: 26.03.19</b>	<b>District: Longleng</b>	<b>Bulletin No: 25/2019</b>
<b>Weather of preceding week</b>		<b>Weather forecast valid upto 31.03.2019</b>
<ul style="list-style-type: none"> <li>Light rain occurred the past week</li> <li>Max and min temperatures ranged 27<sup>0</sup>C to 31<sup>0</sup>C and 17<sup>0</sup>C to 20<sup>0</sup>C, respectively.</li> <li>Relative humidity varied from 32% to 81%.</li> <li>Wind speed ranged from 2 to 4 kmph</li> </ul>		<ul style="list-style-type: none"> <li>Probability of light rain</li> <li><b>Max temp</b> is likely to be 28<sup>0</sup>C-30<sup>0</sup>C and <b>min temp</b> 13<sup>0</sup>C to 15<sup>0</sup>C</li> <li><b>Sky is likely to be mainly clear</b> the coming week</li> <li><b>Relative Humidity</b> is likely to range from 23% to 74%.</li> <li><b>Wind speed</b> may reach upto 3-4 kmph</li> <li><b>Wind may blow from south easterly direction</b></li> </ul>
		<b>Weekly cumulative Rainfall:</b> 3mm

**WEATHER FORECAST BASED AGROMET ADVISORIES**

**General Recommendations:** Short duration pulse crops and green manuring crops like dhaincha, sunnhemp etc should be planted as pre-kharif crop in fallow paddy lowland/ Terrace Rice field to enhance the soil fertility

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
<i>Jhum paddy</i>	Sowing	Blast	To manage blast, use high quality, disease free seeds for sowing, Destroy the infested residue, stubbles, grass and weeds present around the field. They act as a source for the infection
<b>Potato</b>	Vegetative stage	Early blight, leaf spot	Regular monitoring of disease should be done. Avoid cultivation of solanaceous crops like tomato, brinjal, chillies etc. nearby potato fields Fields should be kept neat and clean, free from weeds
<i>Maize</i>	Land preparation	White grubs	Deep ploughing should be done to expose the resting stages of pest. Collect and destroy the white grubs
<i>Brinjal</i>	Nursery		Prepare the nursery bed, the soil should be fine and seeds should be sown thinly in rows 10cm apart and 0.5cm deep. Mulch with dry straw or grass whenever it is necessary.
<i>Summer crops</i>			Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. and certified seeds should be use to produce good yield.
<i>Citrus</i>	Propagation	-	Propagation through cuttings or seeds should be started before the onset of hot weather for citrus. For young trees to encourage good growth, a weed free area should be maintained atleast 4 feet diameter around the young trees.
<i>Poultry</i>			For day old chicks provide cool water and electrolytes on their arrival to farm before offering feed to avoid dehydration after transportation
<i>Piggery</i>			Vaccinate all newly born piglets against <b>swine fever</b> at the age of 3 months followed by annual booster dose. Keep floor dry to avoid skin diseases.
<i>Fishery</i>			As most of the pond is dry, take up renovation works like repairing and cleaning of bottom of the pond





**GRAMIN KRISHI MAUSAM SEWA**  
**ICAR RESEARCH COMPLEX FOR NEH REGION**  
**Nagaland Center, Medziphema-797106**  
**Dimapur, Nagaland**

Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in



Date: 26.03.19		District: Mokokchung		Bulletin No: 25/2019	
Weather of preceding week			Weather forecast valid upto 31.03.2019		
<ul style="list-style-type: none"><li>▪ Light rain occurred the past week</li><li>▪ Max and min temperatures ranged 25<sup>0</sup>C to 30<sup>0</sup>C and 16<sup>0</sup>C to 18<sup>0</sup>C, respectively.</li><li>▪ Relative humidity varied from 27% to 90%.</li><li>▪ Wind speed ranged from 4 to 5 kmph</li></ul>			<ul style="list-style-type: none"><li>▪ Probability of light rain</li><li>▪ <b>Max temp</b> is likely to be 28<sup>0</sup>C-30<sup>0</sup>C and <b>min temp</b> 13<sup>0</sup>C to 14<sup>0</sup>C</li><li>▪ <b>Sky is likely to be mainly clear</b> the coming week</li><li>▪ <b>Relative Humidity</b> is likely to range from 31% to 74%.</li><li>▪ <b>Wind speed</b> may reach upto 3-6 kmph</li><li>▪ <b>Wind may blow from south easterly direction</b></li></ul> <b>Weekly cumulative Rainfall:</b> 24 mm		
WEATHER FORECAST BASED AGROMET ADVISORIES					
<b>General Recommendations:</b> Short duration pulse crops and green manuring crops like dhaincha, sunnhemp etc should be planted as pre-kharif crop in fallow paddy lowland/ Terrace Rice field to enhance the soil fertility					
Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories		
Jhum paddy	Sowing	Blast	To manage blast, use high quality, disease free seeds for sowing, Destroy the infested residue, stubbles, grass and weeds present around the field. They act as a source for the infection		
Potato	Vegetative stage	Early blight, leaf spot	Regular monitoring of disease should be done. Avoid cultivation of solanaceous crops like tomato, brinjal, chillies etc. nearby potato fields Fields should be kept neat and clean, free from weeds		
Maize	Land preparation	White grubs	Deep ploughing should be done to expose the resting stages of pest. Collect and destroy the white grubs		
Brinjal	Nursery		Prepare the nursery bed, the soil should be fine and seeds should be sown thinly in rows 10cm apart and 0.5cm deep. Mulch with dry straw or grass whenever it is necessary.		
Summer crops			Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. and certified seeds should be use to produce good yield.		
Citrus	Propagation	-	Propagation through cuttings or seeds should be started before the onset of hot weather for citrus. For young trees to encourage good growth, a weed free area should be maintained atleast 4 feet diameter around the young trees.		
Poultry			Supply of plenty of clean and cool water must be ensured during summer months. Provide Electral Energy (1-2 gm/liter) in water during hot hours.		
Piggery			As temperature and humidity rises, pay extra attention to the movement of animals.		
Fishery			As most of the pond is dry, take up renovation works like repairing and cleaning of bottom of the pond		





**GRAMIN KRISHI MAUSAM SEWA**  
**ICAR RESEARCH COMPLEX FOR NEH REGION**  
**Nagaland Center, Medziphema-797106**  
**Dimapur, Nagaland**



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 26.03.19	District: Mon	Bulletin No: 25/2019
Weather of preceding week		Weather forecast valid upto 31.03.2019
<ul style="list-style-type: none"> <li>Light rain occurred the past week</li> <li>Max and min temperatures ranged 27<sup>0</sup>C to 30<sup>0</sup>C and 15<sup>0</sup>C to 17<sup>0</sup>C, respectively.</li> <li>Relative humidity varied from 33% to 89%.</li> <li>Wind speed ranged from 4 to 5 kmph</li> </ul>		<ul style="list-style-type: none"> <li>Probability of light rain</li> <li><b>Max temp</b> is likely to be 28<sup>0</sup>C - 30<sup>0</sup>C and <b>min temp</b> 13<sup>0</sup>C to 14<sup>0</sup>C</li> <li><b>Sky is likely to be mainly clear</b> the coming week</li> <li><b>Relative Humidity</b> is likely to range from 32% to 73%.</li> <li><b>Wind speed</b> may reach upto 3-7 kmph</li> <li><b>Wind may blow from south easterly direction</b></li> </ul>
		<b>Weekly cumulative Rainfall:</b> 14mm

**General Recommendations:** Short duration pulse crops and green manuring crops like dhaincha, sunnhemp etc should be planted as pre-kharif crop in fallow paddy lowland field to enhance the soil fertility

**WEATHER FORECAST BASED AGROMET ADVISORIES**

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
<i>Jhum paddy</i>	Sowing	Blast	To manage blast, use high quality, disease free seeds for sowing, Destroy the infested residue, stubbles, grass and weeds present around the field. They act as a source for the infection
<i>TRC/WRC paddy</i>	Storage		Inspect the stored seeds once a week for signs of insect infestation. Place sticky traps in the drying and storage areas for rats
<i>Potato</i>	Vegetative stage	Early blight, leaf spot	Regular monitoring of disease should be done. Avoid cultivation of solanaceous crops like tomato, brinjal, chillies etc. nearby potato fields Fields should be kept neat and clean, free from weeds
<i>Maize</i>	Land preparation	White grubs	Deep ploughing should be done to expose the resting stages of pest. Collect and destroy the white grubs
<i>Cucurbitaceous vegetable</i>	Germination	Fruit fly	Apply neem cake @ 250kg/ha immediately after germination
<i>Citrus</i>		Leaf miner	Citrus leafminer moths are attracted to the new flush of citrus trees. Avoid pruning live branches more than once a year, so that the cycles of flushing are uniform and short. Once the leaves harden, the pest will not be able to mine the leaves. Do not prune off leaves damaged by citrus leaf miner since undamaged areas of leaves continue to produce food for the tree.
<i>Poultry</i>			Supply of plenty of clean and cool water must be ensured during summer months. Provide Electral Energy (1-2 gm/liter) in water during hot hours.
<i>Piggery</i>			As temperature and humidity rises, pay extra attention to the movement of animals. Provide adequate water Prevent overcrowding
<i>Fishery</i>			As most of the pond is dry, take up renovation works like repairing and cleaning of bottom of the pond



**GRAMIN KRISHI MAUSAM SEWA**  
**ICAR RESEARCH COMPLEX FOR NEH REGION**  
**Nagaland Center, Medziphema-797106**  
**Dimapur, Nagaland**



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 26.03.19		District: Peren		Bulletin No: 25/2019	
Weather of preceding week			Weather forecast valid upto 31.03.2019		
<ul style="list-style-type: none"><li>▪ Light rain occurred the past week</li><li>▪ Max and min temperatures ranged 29<sup>0</sup>C to 32<sup>0</sup>C and 15<sup>0</sup>C to 19<sup>0</sup>C, respectively.</li><li>▪ Relative humidity varied from 23% to 90%.</li><li>▪ Wind speed ranged from 4 to 5 kmph</li></ul>			<ul style="list-style-type: none"><li>▪ Probability of light rain</li><li>▪ <b>Max temp</b> is likely to be 28<sup>0</sup>C - 30<sup>0</sup>C and <b>min temp</b> 13<sup>0</sup>C to 14<sup>0</sup>C</li><li>▪ <b>Sky is likely to be mainly clear</b> the coming week</li><li>▪ <b>Relative Humidity</b> is likely to range from 20% to 89%.</li><li>▪ <b>Wind speed</b> may reach upto 2-3 kmph</li><li>▪ <b>Wind may blow from mostly southeasterly direction</b></li></ul> <b>Weekly cumulative Rainfall:</b> 12 mm		
<b>General Advisories:</b> Short duration pulse crops and green manuring crops like dhaincha, sunnhemp etc should be planted as pre-kharif crop in fallow paddy lowland/ Terrace Rice field to enhance the soil fertility					
Main crop		Growth Stage	Pest/Disease	Weather based Agro-Advisories	
TRC/WRC paddy		Storage		Keep storage areas, crevices, wooden pallets etc clean by spraying with the recommended insecticide to disinfect insect breeding places. Place sticky traps in the drying and storage areas for rats	
Green gram		Sowing		Sowing should be carried out in the present weather	
Naga king chilli		Nursery preparation	Damping off	Before sowing of seeds, seed treatment should be done with Thiram @2gm/litre of water. Avoid over watering and close spacing to avoid damping off	
Assam Lemon				Propagation through cuttings or seeds should be started before the onset of hot weather for citrus.	
Okra		Sowing	Hoppers and white flies	While germinating, install yellow sticky trap for monitoring white flies and hoppers	
Cucurbitaceous vegetable		Germination	Fruit fly	Apply neem cake @ 250kg/ha immediately after germination	
Poultry		Vaccination of chicks against Ranikhet disease on 5 days old or 5 – 7 days once, one drop in the eyes or nose During transportation of chicks, provide electrolytes and vitamin C and vitamin E rich syrup in drinking water.			
Piggery		As temperature and humidity rises, pay extra attention to the movement of animals. Provide adequate water Prevent overcrowding			
Fishery		As most of the pond is dry, take up renovation works like repairing and cleaning of bottom of the pond			





**GRAMIN KRISHI MAUSAM SEWA**  
**ICAR RESEARCH COMPLEX FOR NEH REGION**  
**Nagaland Center, Medziphema-797106**  
**Dimapur, Nagaland**

Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in



Date: 26.03.19	District: Phek	Bulletin No: 25/2019
<b>Weather of preceding week</b>		<b>Weather forecast valid upto 31.03.2019</b>
<ul style="list-style-type: none"> <li>Light rain occurred the past week</li> <li>Max and min temperatures ranged 27<sup>0</sup>C to 31<sup>0</sup>C and 15<sup>0</sup>C to 18<sup>0</sup>C, respectively.</li> <li>Relative humidity varied from 24% to 80%.</li> <li>Wind speed ranged from 3 to 4 kmph</li> </ul>		<ul style="list-style-type: none"> <li>Probability of light rain</li> <li><b>Max temp</b> is likely to be 26<sup>0</sup>C - 28<sup>0</sup>C and <b>min temp</b> 11<sup>0</sup>C to 13<sup>0</sup>C</li> <li><b>Sky is likely to be mainly clear</b> the coming week</li> <li><b>Relative Humidity</b> is likely to range from 27% to 85%.</li> <li><b>Wind speed</b> may reach upto 2-4 kmph</li> <li><b>Wind may blow most southerly direction</b></li> <li><b>Weekly cumulative Rainfall:</b> 20 mm</li> </ul>

**WEATHER FORECAST BASED AGROMET ADVISORIES**

**General Advisories:** Short duration pulse crops and green manuring crops like dhaincha, sunnhemp etc should be planted as pre-kharif crop in fallow paddy lowland/ Terrace Rice field to enhance the soil fertility

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
TRC/WRC paddy	Storage		Place sticky traps in the drying and storage areas for rats
Potato	Vegetative stage	Early blight, leaf spot	Regular monitoring of disease should be done. Avoid cultivation of solanaceous crops like tomato, brinjal, chillies etc. nearby potato fields Fields should be kept neat and clean, free from weeds
Maize	Land preparation	White grubs	Deep ploughing should be done to expose the resting stages of pest. Collect and destroy the white grubs
Cucurbitaceous vegetable	Germination	Fruit fly	Apply neem cake @ 250kg/ha immediately after germination
Khasi mandarin			Prepare basin around the plants followed by mulching with local grasses.
Summer crops			Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. and certified seeds should be use to produce good yield.
Poultry	For day old chicks provide cool water and electrolytes on their arrival to farm before offering feed to avoid dehydration after transportation Supply of plenty of clean and cool water must be ensured during summer months. Provide Electral Energy (1-2 gm/liter) in water during hot hours.		
Piggery	As temperature and humidity rises, pay extra attention to the movement of animals. Provide adequate water Prevent overcrowding		
Fishery	As most of the pond is dry, take up renovation works like repairing and cleaning of bottom of the pond		



**GRAMIN KRISHI MAUSAM SEWA**  
**ICAR RESEARCH COMPLEX FOR NEH REGION**  
**Nagaland Center, Medziphema-797106**  
**Dimapur, Nagaland**

Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in



Date: 26.03.19		District: Tuensang		Bulletin No: 25/2019	
Weather of preceding week			Weather forecast valid upto 31.03.2019		
<ul style="list-style-type: none"><li>▪ Light rain occurred the past week</li><li>▪ Max and min temperatures ranged 24<sup>0</sup>C to 30<sup>0</sup>C and 14<sup>0</sup>C to 17<sup>0</sup>C, respectively.</li><li>▪ Relative humidity varied from 28% to 80%.</li><li>▪ Wind speed ranged from 3 to 4 kmph</li></ul>			<ul style="list-style-type: none"><li>▪ Probability of light rain</li><li>▪ <b>Max temp</b> is likely to be 25<sup>0</sup>C-27<sup>0</sup>C and <b>min temp</b> 10<sup>0</sup>C to 12<sup>0</sup>C</li><li>▪ <b>Sky is likely to be mainly clear</b> the coming week</li><li>▪ <b>Relative Humidity</b> is likely to range from 25% to 79%.</li><li>▪ <b>Wind speed</b> may reach upto 3-4 kmph</li><li>▪ <b>Wind may blow from mostly southerly direction</b></li></ul> <div>Weekly cumulative Rainfall: 9 mm</div>		
WEATHER FORECAST BASED AGROMET ADVISORIES					
General Recommendations: In the present weather, start sowing of summer crops					
Main crop		Growth Stage	Pest/Disease	Weather based Agro-Advisories	
Jhum paddy		Sowing	Blast	To manage blast, use high quality, disease free seeds for sowing, Destroy the infested residue, stubbles, grass and weeds present around the field. They act as a source for the infection	
TRC/WRC paddy		Storage		Storage rooms should be physically rodent and bird proof, if possible, place sticky traps in the drying and storage areas for rats.	
Potato		Vegetative stage	Early blight, leaf spot	Regular monitoring of disease should be done. Avoid cultivation of solanaceous crops like tomato, brinjal, chillies etc. nearby potato fields Fields should be kept neat and clean, free from weeds	
Rajmash		Vegetative stage		In the present weather, Gap filling should be carried out	
Cucurbitaceous vegetable		Germination	Fruit fly	Apply neem cake @ 250kg/ha immediately after germination	
Khasi Mandarin		Post harvesting stage		In the present weather, water sprout, diseases and dried/dead twigs of mandarin should be removed after harvesting and cut end pasted with Bordeaux paste	
Poultry		For day old chicks provide cool water and electrolytes on their arrival to farm before offering feed to avoid dehydration after transportation Supply of plenty of clean and cool water must be ensured during summer months. Provide Electral Energy (1-2 gm/liter) in water during hot hours.			
Piggery		As temperature and humidity rises, pay extra attention to the movement of animals. Provide adequate water Prevent overcrowding			
Fishery		As most of the pond is dry, take up renovation works like repairing and cleaning of bottom of the pond			



**GRAMIN KRISHI MAUSAM SEWA**  
**ICAR RESEARCH COMPLEX FOR NEH REGION**  
**Nagaland Center, Medziphema-797106**  
**Dimapur, Nagaland**

Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in



Date: 26.03.19	District: Wokha	Bulletin No: 25/2019
Weather of preceding week		Weather forecast valid upto 31.03.2019
<ul style="list-style-type: none"> <li>Light rain occurred the past week</li> <li>Max and min temperatures ranged 21<sup>0</sup>C to 23<sup>0</sup>C and 10<sup>0</sup>C to 11<sup>0</sup>C respectively.</li> <li>Relative humidity varied from 40% to 80%.</li> <li>Wind speed ranged from 5 to 6 kmph</li> </ul>		<ul style="list-style-type: none"> <li>Probability of light rain</li> <li><b>Max temp</b> is likely to be 28<sup>0</sup>C-30<sup>0</sup>C and <b>min temp</b> 13<sup>0</sup>C to 15<sup>0</sup>C</li> <li><b>Sky is likely to be mainly clear</b> the coming week</li> <li><b>Relative Humidity</b> is likely to range from 32% to 78%.</li> <li><b>Wind speed</b> may reach upto 3-4 kmph</li> <li><b>Wind may blow from easterly direction</b></li> </ul>
		<b>Weekly cumulative Rainfall:</b> 25mm

**WEATHER FORECAST BASED AGROMET ADVISORIES**

**General Recommendations:** Short duration pulse crops and green manuring crops like dhaincha, sunnhemp etc should be planted as pre-kharif crop in fallow paddy lowland/ Terrace Rice field to enhance the soil fertility

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
<i>Jhum paddy</i>	Sowing	Blast	<i>To manage blast, use high quality, disease free seeds for sowing, Destroy the infested residue, stubbles, grass and weeds present around the field. They act as a source for the infection</i>
<b>Potato</b>	Vegetative stage	Early blight, leaf spot	<i>Regular monitoring of disease should be done. Avoid cultivation of solanaceous crops like tomato, brinjal, chillies etc. nearby potato fields Fields should be kept neat and clean, free from weeds</i>
<i>Maize</i>	Land preparation	-	Use resistant varieties in consultation with your respective agri department.
<i>Brinjal</i>	Nursery		<i>Prepare the nursery bed, the soil should be fine and seeds should be sown thinly in rows 10cm apart and 0.5cm deep. Mulch with dry straw or grass whenever it is necessary.</i>
<i>Summer crops</i>			<i>Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. and certified seeds should be use to produce good yield.</i>
<i>Citrus</i>	Propagation	-	<i>Propagation through cuttings or seeds should be started before the onset of hot weather for citrus. For young trees to encourage good growth, a weed free area should be maintained atleast 4 feet diameter around the young trees.</i>
<b>Poultry</b>	Supply of plenty of clean and cool water must be ensured during summer months. Provide Electral Energy (1-2 gm/liter) in water during hot hours.		
<b>Piggery</b>	As temperature and humidity rises, pay extra attention to the movement of animals. Provide adequate water Prevent overcrowding		
<b>Fishery</b>	As most of the pond is dry, take up renovation works like repairing and cleaning of bottom of the pond		





**GRAMIN KRISHI MAUSAM SEWA**  
**ICAR RESEARCH COMPLEX FOR NEH REGION**  
**Nagaland Center, Medziphema-797106**  
**Dimapur, Nagaland**

Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in



Date: 26.03.19	District: Zunheboto	Bulletin No: 25/2019
Weather of preceding week		Weather forecast valid upto 31.03.2019
<ul style="list-style-type: none"> <li>Light rain occurred the past week</li> <li>Max and min temperatures ranged 21<sup>0</sup>C to 23<sup>0</sup>C and 9<sup>0</sup>C to 12<sup>0</sup>C, respectively.</li> <li>Relative humidity varied from 30% to 86%.</li> <li>Wind speed ranged from 3 to 4 kmph</li> </ul>		<ul style="list-style-type: none"> <li>Probability of light rain</li> <li><b>Max temp</b> is likely to be 28<sup>0</sup>C- 30<sup>0</sup>C and <b>min temp</b> 13<sup>0</sup>C to 14<sup>0</sup>C</li> <li><b>Sky is likely to be mainly clear</b> the coming week</li> <li><b>Relative Humidity</b> is likely to range from 27% to 83%.</li> <li><b>Wind speed</b> may reach upto 4-5 kmph</li> <li><b>Wind may blow from south easterly direction</b></li> </ul>
		<b>Weekly cumulative Rainfall:</b> 9 mm

**WEATHER FORECAST BASED AGROMET ADVISORIES**

**General Recommendations:** Short duration pulse crops and green manuring crops like dhaincha, sunnhemp etc should be planted as pre-kharif crop in fallow paddy lowland/ Terrace Rice field to enhance the soil fertility

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
<i>Jhum paddy</i>	Sowing	Blast	<i>To manage blast, use high quality, disease free seeds for sowing, Destroy the infested residue, stubbles, grass and weeds present around the field. They act as a source for the infection</i>
<b>Potato</b>	Vegetative stage	Early blight, leaf spot	<i>Regular monitoring of disease should be done. Avoid cultivation of solanaceous crops like tomato, brinjal, chillies etc. nearby potato fields Fields should be kept neat and clean, free from weeds</i>
<i>Maize</i>	Land preparation	-	<i>Early sowing of <b>maize</b> in March-April escapes the attack of most pests and crop rotation should be carried out with other crops as it reduces the pest infestation by interrupting the continuity of the food chain. Use resistant varieties in consultation with your respective agri department.</i>
<i>Brinjal</i>	Nursery		<i>Prepare the nursery bed, the soil should be fine and seeds should be sown thinly in rows 10cm apart and 0.5cm deep. Mulch with dry straw or grass whenever it is necessary.</i>
<i>Summer crops</i>			<i>Farmers are advised to go for sowing of summer vegetables like ladies finger, cucurbitaceous crops like gourd, pumpkin, ridge gourd, ash gourd, spine gourd etc. and certified seeds should be use to produce good yield.</i>
<i>Citrus</i>	Propagation	-	<i>Propagation through cuttings or seeds should be started before the onset of hot weather for citrus. For young trees to encourage good growth, a weed free area should be maintained atleast 4 feet diameter around the young trees.</i>
<i>Poultry</i>			<i>Supply of plenty of clean and cool water must be ensured during summer months. Provide Electral Energy (1-2 gm/liter) in water during hot hours.</i>
<i>Piggery</i>		Swine fever	<i>Vaccinate all newly born piglets against <b>swine fever</b> at the age of 3 months followed by annual booster dose. Keep floor dry to avoid skin diseases.</i>

<i><b>Fishery</b></i>			As most of the pond is dry, take up renovation works like repairing and cleaning of bottom of the pond
-----------------------	--	--	--