



## Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



# Agromet Advisory Bulletin

Date : 01-11-2019

Weather Forecast of TUENSANG(Nagaland) Issued On : 2019-11-01(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-11-02	0	32	18	86	36	3.3	112	1
2019-11-03	0	31	18	85	35	3.3	118	0
2019-11-04	0	30	18	83	33	3.3	123	0
2019-11-05	0	30	17	82	34	3.3	113	0
2019-11-06	0	29	16	83	33	3.5	117	1

### Weather Summary/Alert:

Probability of no rainfall the coming days
Pani etu hapta to nikiribolhe pareh. Din pal hobo

### General Advisory:

Contact nearby veterinarian Officer for measure against adverse conditions
Apni khan parishe koilhe Veterinarian Officer lakha number to rakhibi, enikha hoilhe gauri lagha kiba dhok khan hoilhe jolti matibo baribo

### SMS Advisory:

Avoid excessive drying of paddy to avoid breakage of the grains
Dhan bishi bi nusukabi nahoileh dhan goti tho mehin hobo

### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	HARVESTING STAGE: Do not dry the grains at each time more than 4 - 6 hours otherwise grains will shrivel and break.
	Hudai sukha somia tae 4-6 ghonta opor husukabi nanoileh dhan goti mehin hobo
GINGER	Harvesting stage. Grading should be done after harvest, this would generate more income in the market
	Athua han tho mati para olai kina bhal aro biya nahoiley chutu aro thangor majot de alak alak rhaki bo ley lage. Etu para bazaar de bikai homoi de lap babo.

### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
BANANA	Remove the affected pest and disease leaf, dried leaves from time to time.

Horticulture (Varieties)	Horticulture Specific Advisory
	Puka aru bimar lagijah pata khan hatabi aru sukha pata khan bi somoi somoi tae hatai thakibi
CABBAGE	VEGETATIVE STAGE: Earthing up should be done 30 days after transplanting
	Transplanting kurighenah 30 din pichetae mathi olop dhalibi

**Livestock Specific Advisory:**

Livestock(Varieties)	Livestock Specific Advisory
PIG	Routine vaccination programme is recommended and once any disease is diagnosed treat immediately with antibiotics
	Somoi tae vaccination kurithakibi aru bimar kunba lagileh joldi antibiotics dibi

**Fisheries Specific Advisory:**

Fisheries(Varieties)	Fisheries Specific Advisory
COMMON CARP	Check the movement of the fish every day. If abnormal behavior is observed consult an expert for suggestion
	Hudai mass khila tho sabo lageh. Thik nikhileh mass expert logotae alusnah kuribi

**Poultry Specific Advisory:**

Poultry(Varieties)	Poultry Specific Advisory
CHICKEN	Proper sanitary measures should be done to control the diseases. Slaughtering of all apparently ailing birds, segregating of in - contact in group of 10 to 15 each; removal of all infective materials such as droppings, residues of poultry cleanliness
	Bimar nalagiboh karoneh sapha parah murgi palibi



## Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



# Agromet Advisory Bulletin

Date : 01-11-2019

Weather Forecast of WOKHO(Nagaland) Issued On : 2019-11-01(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-11-02	1	30	18	93	60	4.9	83	2
2019-11-03	1	31	18	94	62	4.8	81	0
2019-11-04	0	30	18	93	60	4.7	77	0
2019-11-05	4	30	17	92	60	4.7	87	0
2019-11-06	1	29	16	92	61	4.8	86	2

### Weather Summary/Alert:

Probability of light rainfall the coming days
pani etia ahibolhe din to olop kiribolhe dikaiase

### General Advisory:

Contact nearby veterinarian Officer for measure against adverse conditions
Apni khan parishe koilhe Veterinarian Officer lakha number to rakhibi, enikha hoilhe gauri lagha kiba dhok khan hoilhe jolti matibo baribo

### SMS Advisory:

Avoid excessive drying of paddy to avoid breakage of the grains
Dhan bishi bi nusukabi nahoileh dhan goti tho mehin hobo

### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	RIPENING STAGE: For bird control erect scarecrows randomly in the field, scare the bird manually too. Tie old VHS tapes diagonally across and around the field. Use catapults
	For bird control erect scarecrows randomly in the field, scare the bird manually too. Tie old VHS tapes diagonally across and around the field. Use catapults
GINGER	MATURITY STAGE: For vegetable purpose , harvesting can be carried out
	Sobji te hali khena kabo kharoneh to etia para harvest khoribo pareh

### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
BRINJAL	HARVESTING STAGE: Timely harvesting should be done. Remove the infested fruit

Horticulture (Varieties)	Horticulture Specific Advisory
	Dhik somoi te bijon ulabi aru biya bijon to somoi te ulai khena bilabi
CHILLI	HARVESTING STAGE: Timely harvesting should be carried out to prevent loss Thik somoi te mircha to olabo lakhe
COLOCASIA/ TARO	HARVESTING STAGE: Leaves which have turned completely brown and have reduced or shirk can be harvested Patha khan brown, sukha hoilhe harvest khoribo pareh

#### Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	Protect the animals from cold draft using gunny bags and other locally available materials. Provide protein rich ration and ad libitum clean drinking water Protect the animals from cold draft using gunny bags and other locally available materials. Provide protein rich ration and ad libitum clean drinking water

#### Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
COMMON CARP	If the pond has minimum 1m depth water, apply lime @ 50kg/bigha along with raw cow dung @ 125 kg/bigha, Urea 3.5 kg/bigha, SSP 2.7 kg /bigha. If water depth is less than 1m harvest 50% of the stocked fish If the pond has minimum 1m depth water, apply lime @ 50kg/bigha along with raw cow dung @ 125 kg/bigha, Urea 3.5 kg/bigha, SSP 2.7 kg /bigha. If water depth is less than 1m harvest 50% of the stocked fish

#### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICKEN	Deworm the birds once every 2 months Deworm the birds once every 2 months



## Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



# Agromet Advisory Bulletin

Date : 01-11-2019

Weather Forecast of ZUNHEBOTO(Nagaland) Issued On : 2019-11-01(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-11-02	0	32	17	95	43	4.3	124	1
2019-11-03	0	32	17	96	44	4.4	125	0
2019-11-04	0	30	18	94	41	4.3	127	0
2019-11-05	0	30	17	94	44	4.3	124	0
2019-11-06	0	30	16	97	47	4.4	123	2

### Weather Summary/Alert:

Probability of no rainfall the coming days
pani etu hapta to nikiribo pareh

### General Advisory:

Contact nearby veterinarian Officer for measure against adverse conditions
Apni khan parishe koilhe Veterinarian Officer lakha number to rakhibi, enikha hoilhe gauri lagha kiba dhok khan hoilhe jolti matibo baribo

### SMS Advisory:

Avoid excessive drying of paddy to avoid breakage of the grains
Dhan bishi bi nusukabi nahoileh dhan goti tho mehin hobo

### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	RIPENING STAGE: For bird control erect scarecrows randomly in the field, scare the bird manually too. Tie old VHS tapes diagonally across and around the field. Use catapults
	For bird control erect scarecrows randomly in the field, scare the bird manually too. Tie old VHS tapes diagonally across and around the field. Use catapults
GINGER	MATURITY STAGE: For vegetable purpose , harvesting can be carried out
	Sobji te hali khena kabo kharoneh to etia para harvest khoribo pareh

### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
BRINJAL	HARVESTING STAGE: Timely harvesting should be done. Remove the infested fruit

Horticulture (Varieties)	Horticulture Specific Advisory
	Dhik somoi te bijon ulabi aru biya bijon to somoi te ulai khena bilabi
CHILLI	HARVESTING STAGE: Timely harvesting should be carried out to prevent loss Thik somoi te mircha to olabo lakhe
COLOCASIA/ TARO	HARVESTING STAGE: Leaves which have turned completely brown and have reduced or shirk can be harvested Patha khan brown, sukha hoilhe harvest khoribo pareh

#### Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	Protect the animals from cold draft using gunny bags and other locally available materials. Provide protein rich ration and ad libitum clean drinking water Protect the animals from cold draft using gunny bags and other locally available materials. Provide protein rich ration and ad libitum clean drinking water

#### Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
CATLA	If the pond has minimum 1m depth water, apply lime @ 50kg/bigha along with raw cow dung @ 125 kg/bigha, Urea 3.5 kg/bigha, SSP 2.7 kg /bigha. If water depth is less than 1m harvest 50% of the stocked fish If the pond has minimum 1m depth water, apply lime @ 50kg/bigha along with raw cow dung @ 125 kg/bigha, Urea 3.5 kg/bigha, SSP 2.7 kg /bigha. If water depth is less than 1m harvest 50% of the stocked fish

#### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICKEN	Deworm the birds once every 2 months Deworm the birds once every 2 months



## Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



# Agromet Advisory Bulletin

Date : 01-11-2019

Weather Forecast of DIMAPUR(Nagaland) Issued On : 2019-11-01(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-11-02	0	30	19	85	57	5.5	121	2
2019-11-03	0	30	19	88	55	5.7	125	0
2019-11-04	0	30	20	83	52	5.6	121	0
2019-11-05	0	29	19	85	54	5.4	125	0
2019-11-06	0	29	18	87	53	5.6	130	2

### Weather Summary/Alert:

Probability of no rain the coming days
Pani etu hapta to nikiribo

### General Advisory:

Contact nearby veterinarian Officer for measure against adverse conditions
Apni khan parishe koilhe Veterinarian Officer lakha number to rakhibi, enikha hoilhe gauri lagha kiba dhok khan hoilhe jolti matibo baribo

### SMS Advisory:

Avoid excessive drying of paddy to avoid breakage of the grains
Dhan bishi bi nusukabi nahoileh dhan goti tho mehin hobo

### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	RIPENING STAGE: Avoid excessive drying of paddy to avoid breakage of the grains
	RIPENING STAGE: Dhan bishi bi nusukabi nahoileh dhan goti tho mehin hobo
GROUNDNUT	POST HARVEST: When temperature is very high, avoid direct sun drying. Collect the pods in gunnies and store on the ground over a layer of sand to avoid any moisture coming in contact with dry pods
	Bishi gorom din tae pariatok thik dhup nichetae nusukabi. Sukha badam khan tae moisture nalagiboh karoneh, bosta tae dhalighenah, mati opor tae bhalu olop rakhighenah hi etu opor tae bosto tho rakhibi
MAIZE	SEEDLING STAGE: Provide mulching to conserve moisture
	SEEDLING STAGE: mati tae moisture thakibo karoneh, sukha patah khan rakhibi
SESAME (GINGELLY/TIL)	VEGETATIVE STAGE: When the plants are about 15 cm in height, thin the crop so as to give a spacing of 15-25 cm between plants.

Crop(Varieties)	Crop Specific Advisory
	Ghas 15 cm ocha hua pichetae, ghas majotae 15-25 cm khali thakiboleh ghas khan ulailobi
GINGER	HARVESTING STAGE: However, for making dry ginger, the matured rhizomes are harvested at full maturity i.e. when the leaves turn yellow and start drying.
	Sukha aduwa bonaboleh, pata tho yellow huighenah murijah somoi tae khundilobi aru sakahbi

#### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
FRENCH BEAN	VEGETATIVE STAGE: Weeding should be given 20 – 25 days and 40 – 45 days after sowing. The crop should be earthed up after each weeding.
	Ghas lagaigheanh 20 – 25 din and 40 – 45 din tae jonjol ghas sapha kuribi. Sapha kurah pichetae ghas khan tae mathi olop dhalibi
NAGA KING CHILLI	HARVESTING STAGE: Harvesting should be done when the sky is clear
	Din bal somoi te mircha do ulabi.
COLOCASIA/TARO	Harvesting stage. Clean the debris and sundry the harvested tubers
	Kochu kuti han tho bhal para sabha kuri kina, sun de sukhai dibo lage.

#### Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
COW	There is a change in the weather from this month onwards, hence, due arrangements should be made to protect the animals from the onset of the winter.
	Etu moinah para to hawa pani bodli hoi, apni khan bi ghoru lakha ghor dho dhanta hawa para bachabo karoneh bonailobi
PIG	Contact nearby veterinarian Officer for measure against adverse conditions
	Apni khan parishe koilhe Veterinarian Officer lakha number to rakhibi, enikha hoilhe gauri lagha kiba dhok khan hoilhe jolti matibo baribo

#### Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
COMMON CARP	As far as possible do not disturb the pond. Prune the tree branches along the pond embankment and allow enough sunlight to fall on the pond water surface
	Pukhuri pani tho parah thok disturb nukuribi. Pukhuri kinar tae ghas lagah branch khan katilobi. Pani opor tae dhup bhal parah pabo lageh

#### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICKEN	As the temperature is dropping, provide vitamin complex. Get ready to provide warmth by use of incandescent bulbs to protect them from cold winds by use of plastic sheets, jute sacks and plastic sacks
	Temperature tho thanda hui ase etu karoneh vitamin complex dibo lageh. Hawa thanda parah bachibo karoneh plastic nahoileh bosta parah ghor kinar bon kuribi aru bulb bi cholabi





## Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



# Agromet Advisory Bulletin

Date : 01-11-2019

Weather Forecast of KIPHIRE(Nagaland) Issued On : 2019-11-01(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-11-02	2	31	18	85	37	3.4	68	2
2019-11-03	0	31	18	83	33	3.4	76	0
2019-11-04	0	30	18	82	31	3.4	80	1
2019-11-05	3	29	17	81	36	3.6	62	0
2019-11-06	0	29	16	82	33	3.3	71	1

### Weather Summary/Alert:

probability of light rainfall the coming days
etu hapta bi pani olop kiribolhe diakaise

### General Advisory:

Contact nearby veterinarian Officer for measure against adverse conditions
Apni khan parishe koilhe Veterinarian Officer lakha number to rakhibi, enikha hoilhe gauri lagha kiba dhok khan hoilhe jolti matibo baribo

### SMS Advisory:

Avoid excessive drying of paddy to avoid breakage of the grains
Dhan bishi bi nusukabi nahoileh dhan goti tho mehin hobo

### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	HARVESTING STAGE: Do not dry the grains at each time more than 4 - 6 hours otherwise grains will shrivel and break.
	Hudai sukha somia tae 4-6 ghonta opor husukabi nanoileh dhan goti mehin hobo
FIELD PEA	VEGETATIVE STAGE: In the present weather, weeding should be done
	Etia somoi te habi khan pilabi
MUSTARD	VEGETATIVE STAGE: Weeding and thinning out should be done after third week of sowing. Another weeding should be done in the 6th week of sowing
	Weeding and thinning out should be done after third week of sowing. Another weeding should be done in the 6th week of sowing

### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
-----------------------------	--------------------------------

Horticulture (Varieties)	Horticulture Specific Advisory
CABBAGE	VEGETATIVE STAGE: Shallow hoeing should be done to remove weeds and loosen the soil for better aeration. Deep hoeing should be avoided, as the cabbage roots are surface feeders.
	Mati tho olup he khundi bo lage habi han bhelai homoi de. Etu para mati bithor de hawa dibov kuntu khas laga root karone bhal asae. Mati tho bishi bithor nakundibo lage kiley koiley kobi laga root tho mati opor de he tailaga khas karone root para loi thake.
ONION	Vegetative stage. Thrips is a common problem during this stage. A Use of reflective plastic silver colour and aluminum painted black mulches repel the thrips.
	Onion tey bi bishi ola problem doh chutu chutu puka(Thrips) para hui,etu han para ghas doh piya kori dea thake , etu puka han ke bhakai/ahiboley nadibo karone silver colour plastic aro aluminium painted black mulches doh use koribo lagey
COLOCASIA/TARO	Harvesting stage. Clean the debris and sundry the harvested tubers
	Kochu kuti han tho bhal para sabha kuri kina, sun de sukhai dibo lage.

#### Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	A separate house should be there to keep the animals suffering from contagious diseases.
	Bimar khaori han karone ekta alak khor bona bo lage.

#### Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
COMMON CARP	Partial harvest of fish can be done in order to reduce stocking density. While harvesting make sure to leave the small fishes in the pond itself as it may reduce in generating income.
	"Pond tey jaka komti huboley nadibo nimi tey maas han doh kitya ba olai thaki bo lagey. Maas han ola time tey juti chutu maas han pailey pond tey wapal thali debi."

#### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICKEN	Avoid overcrowding of birds. In every two months deworming should be done.
	Murki han tho bishi tight para narakibo lage. Aro hotai tui moina de kechu laga dawai kelabo lage



## Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



# Agromet Advisory Bulletin

Date : 01-11-2019

Weather Forecast of KOHIMA(Nagaland) Issued On : 2019-11-01(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-11-02	2	31	18	95	57	4.8	133	2
2019-11-03	1	31	18	96	54	5.0	135	0
2019-11-04	0	31	18	95	50	5.0	135	0
2019-11-05	4	30	18	95	58	4.8	134	0
2019-11-06	3	29	17	97	64	5.2	139	3

### Weather Summary/Alert:

Probability of light rainfall the coming days
Pani etu hapta bi olop kiribolhe dikaiase

### General Advisory:

Contact nearby veterinarian Officer for measure against adverse conditions
Apni khan parishe koilhe Veterinarian Officer lakha number to rakhibi, enikha hoilhe gauri lagha kiba dhok khan hoilhe jolti matibo baribo

### SMS Advisory:

Avoid excessive drying of paddy to avoid breakage of the grains
Dhan bishi bi nusukabi nahoileh dhan goti tho mehin hobo

### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	HARVESTING STAGE: Do not dry the grains at each time more than 4 - 6 hours otherwise grains will shrivel and break.
	Hudai sukha somia tae 4-6 ghonta opor husukabi nanoileh dhan goti mehin hobo
GINGER	Harvesting stage. Grading should be done after harvest, this would generate more income in the market.
	Athua han tho mati para olai kina bhal aro biya nahoiley chutu aro thangor majot de alak alak rhaki bo ley lage. Etu para bazzar de bikai homoi de lap babo.

### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
BANANA	Remove the affected pest and disease leaf, dried leaves from time to time.

Horticulture (Varieties)	Horticulture Specific Advisory
	Puka aru bimar lagijah pata khan hatabi aru sukha pata khan bi somoi somoi tae hatai thakibi
CABBAGE	VEGETATIVE STAGE: Earthing up should be done 30 days after transplanting. Transplanting kurighenah 30 din pichetae mathi olop dhalibi

**Livestock Specific Advisory:**

Livestock(Varieties)	Livestock Specific Advisory
PIG	Routine vaccination programme is recommended and once any disease is diagnosed treat immediately with antibiotics Somoi tae vaccination kurithakibi aru bimar kunba lagileh joldi antibiotics dibi

**Fisheries Specific Advisory:**

Fisheries(Varieties)	Fisheries Specific Advisory
COMMON CARP	Check the movement of the fish every day. If abnormal behavior is observed consult an expert for suggestion Hudai mass khila tho sabo lageh. Thik nikhileh mass expert logotae alusnah kuribi

**Poultry Specific Advisory:**

Poultry(Varieties)	Poultry Specific Advisory
CHICKEN	Proper sanitary measures should be done to control the diseases. Slaughtering of all apparently ailing birds, segregating of in - contact in group of 10 to 15 each; removal of all infective materials such as droppings, residues of poultry cleanliness Bimar nalagiboh karoneh sapha parah murgi palibi.



## Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



# Agromet Advisory Bulletin

Date : 01-11-2019

Weather Forecast of LONGLENG(Nagaland) Issued On : 2019-11-01(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-11-02	0	30	18	89	51	4.8	121	1
2019-11-03	0	31	18	89	54	4.8	121	0
2019-11-04	0	30	17	87	51	4.5	122	0
2019-11-05	0	30	17	85	50	4.6	128	0
2019-11-06	0	29	16	85	48	4.7	132	2

### Weather Summary/Alert:

Probability of no rainfall the coming days
Pani etu hapta to nikiribolhe pareh

### General Advisory:

Contact nearby veterinarian Officer for measure against adverse conditions
Apni khan parishe koilhe Veterinarian Officer lakha number to rakhibi, enikha hoilhe gauri lagha kiba dhok khan hoilhe jolti matibo baribo

### SMS Advisory:

Avoid excessive drying of paddy to avoid breakage of the grains
Dhan bishi bi nusukabi nahoileh dhan goti tho mehin hobo

### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	RIPENING STAGE: For bird control erect scarecrows randomly in the field, scare the bird manually too. Tie old VHS tapes diagonally across and around the field. Use catapults
	For bird control erect scarecrows randomly in the field, scare the bird manually too. Tie old VHS tapes diagonally across and around the field. Use catapults
GINGER	MATURITY STAGE: For vegetable purpose , harvesting can be carried out
	Sobji te hali khena kabo kharoneh to etia para harvest khoribo pareh

### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
BRINJAL	HARVESTING STAGE: Timely harvesting should be done. Remove the infested fruit

Horticulture (Varieties)	Horticulture Specific Advisory
	Dhik somoi te bijon ulabi aru biya bijon to somoi te ulai khena bilab
BRINJAL	HARVESTING STAGE: Timely harvesting should be carried out to prevent loss
	Thik somoi te mircha to olabo lakhe
COLOCASIA/TARO	HARVESTING STAGE: Leaves which have turned completely brown and have reduced or shirk can be harvested
	Patha khan brown, sukha hoilhe harvest khoribo pareh

#### Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	Protect the animals from cold draft using gunny bags and other locally available materials. Provide protein rich ration and ad libitum clean drinking water
	Protect the animals from cold draft using gunny bags and other locally available materials. Provide protein rich ration and ad libitum clean drinking water

#### Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
COMMON CARP	If the pond has minimum 1m depth water, apply lime @ 50kg/bigha along with raw cow dung @ 125 kg/bigha, Urea 3.5 kg/bigha, SSP 2.7 kg /bigha. If water depth is less than 1m harvest 50% of the stocked fish
	If the pond has minimum 1m depth water, apply lime @ 50kg/bigha along with raw cow dung @ 125 kg/bigha, Urea 3.5 kg/bigha, SSP 2.7 kg /bigha. If water depth is less than 1m harvest 50% of the stocked fish

#### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICKEN	Deworm the birds once every 2 months
	Deworm the birds once every 2 months



## Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



# Agromet Advisory Bulletin

Date : 01-11-2019

Weather Forecast of MOKOKCHUNG(Nagaland) Issued On : 2019-11-01(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-11-02	0	31	17	91	51	4.7	117	1
2019-11-03	0	31	18	92	53	4.7	117	0
2019-11-04	0	30	17	91	51	4.4	117	0
2019-11-05	0	30	17	88	50	4.4	123	0
2019-11-06	0	29	16	89	49	4.6	125	2

### Weather Summary/Alert:

Probability of no rainfall the coming days
pani etu hapta to nikiribo pareh.

### General Advisory:

Contact nearby veterinarian Officer for measure against adverse conditions
Apni khan parishe koilhe Veterinarian Officer lakha number to rakhibi, enikha hoilhe gauri lagha kiba dhok khan hoilhe jolti matibo baribo

### SMS Advisory:

Avoid excessive drying of paddy to avoid breakage of the grains
Dhan bishi bi nusukabi nahoileh dhan goti tho mehin hobo

### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	RIPENING STAGE: For bird control erect scarecrows randomly in the field, scare the bird manually too. Tie old VHS tapes diagonally across and around the field. Use catapults
	For bird control erect scarecrows randomly in the field, scare the bird manually too. Tie old VHS tapes diagonally across and around the field. Use catapults
GINGER	MATURITY STAGE: For vegetable purpose , harvesting can be carried out
	Sobji te hali khena kabo kharoneh to etia para harvest khoribo pareh

### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
BRINJAL	HARVESTING STAGE: Timely harvesting should be done. Remove the infested fruit

Horticulture (Varieties)	Horticulture Specific Advisory
	Dhik somoi te bijon ulabi aru biya bijon to somoi te ulai khena bilabi
CHILLI	HARVESTING STAGE: Timely harvesting should be carried out to prevent loss
	Thik somoi te mircha to olabo lakhe
COLOCASIA/TARO	HARVESTING STAGE: Leaves which have turned completely brown and have reduced or shirk can be harvested
	Patha khan brown, sukha hoilhe harvest khoribo pareh
TAPIOCA	VEGETATIVE STAGE: Install sticky cum light trap TO REDUCE WHITE FLY POPULATION
	Whitefly lagha population komti khoribo karoneh sticky cum light trap dho khete te jakha jakha te rakhibi

#### Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	Protect the animals from cold draft using gunny bags and other locally available materials. Provide protein rich ration and ad libitum clean drinking water
	Protect the animals from cold draft using gunny bags and other locally available materials. Provide protein rich ration and ad libitum clean drinking water

#### Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
COMMON CARP	If the pond has minimum 1m depth water, apply lime @ 50kg/bigha along with raw cow dung @ 125 kg/bigha, Urea 3.5 kg/bigha, SSP 2.7 kg /bigha. If water depth is less than 1m harvest 50% of the stocked fish
	If the pond has minimum 1m depth water, apply lime @ 50kg/bigha along with raw cow dung @ 125 kg/bigha, Urea 3.5 kg/bigha, SSP 2.7 kg /bigha. If water depth is less than 1m harvest 50% of the stocked fish

#### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICKEN	Deworm the birds once every 2 months
	Deworm the birds once every 2 months





## Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



# Agromet Advisory Bulletin

Date : 01-11-2019

Weather Forecast of MON(Nagaland) Issued On : 2019-11-01(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-11-02	2	31	18	91	57	4.9	125	1
2019-11-03	1	31	18	89	57	5.0	124	0
2019-11-04	1	30	17	87	54	4.8	123	0
2019-11-05	3	30	17	87	54	4.9	135	0
2019-11-06	0	29	16	86	51	4.9	136	1

### Weather Summary/Alert:

Probability of light rainfall the coming days
Kali purso pani olop kiribolhe dikaiase

### General Advisory:

Contact nearby veterinarian Officer for measure against adverse conditions
Apni khan parishe koilhe Veterinarian Officer lakha number to rakhibi, enikha hoilhe gauri lagha kiba dhok khan hoilhe jolti matibo baribo

### SMS Advisory:

Avoid excessive drying of paddy to avoid breakage of the grains
Dhan bishi bi nusukabi nahoileh dhan goti tho mehin hobo

### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	RIPENING STAGE: Avoid excessive drying of paddy to avoid breakage of the grains
	Dhan bishi bi nusukabi nahoileh dhan goti tho mehin hobo
GROUNDNUT	POST HARVEST: When temperature is very high, avoid direct sun drying. Collect the pods in gunnies and store on the ground over a layer of sand to avoid any moisture coming in contact with dry pods
	Bishi gorom din tae pariatok thik dhup nichetae nusukabi. Sukha badam khan tae moisture nalagiboh karoneh, bosta tae dhalighenah, mati opor tae bhalu olop rakhighenah hi etu opor tae bosto tho rakhibi
MAIZE	SEEDLING STAGE: Provide mulching to conserve moisture
	SEEDLING STAGE: mati tae moisture thakibo karoneh, sukha patah khan khan rakhibi
SESAME (GINGELLY/TIL)	VEGETATIVE STAGE: When the plants are about 15 cm in height, thin the crop so as to give a spacing of 15-25 cm between plants.

Crop(Varieties)	Crop Specific Advisory
	Ghas 15 cm ocha hua pichetae, ghas majotae 15-25 cm khali thakiboleh ghas khan ulailobi
GINGER	HARVESTING STAGE: However, for making dry ginger, the matured rhizomes are harvested at full maturity i.e. when the leaves turn yellow and start drying.
	Sukha aduwa bonaboleh, pata tho yellow huighenah murijah somoi tae khundilobi aru sakahbi

#### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
FRENCH BEAN	VEGETATIVE STAGE: Weeding should be given 20 – 25 days and 40 – 45 days after sowing. The crop should be earthed up after each weeding.
	Ghas lagaigheanh 20 – 25 din and 40 – 45 din tae jonjol ghas sapha kuribi. Sapha kurah pichetae ghas khan tae mathi olop dhalibi
NAGA KING CHILLI	HARVESTING STAGE: Harvesting should be done when the sky is clear
	Din bal somoi te mircha do ulabi.
COLOCASIA/TARO	Harvesting stage. Clean the debris and sundry the harvested tubers
	Kochu kuti han tho bhal para sabha kuri kina, sun de sukhai dibo lage.

#### Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
COW	There is a change in the weather from this month onwards, hence, due arrangements should be made to protect the animals from the onset of the winter.
	Etu moinah para to hawa pani bodli hoi, apni khan bi ghoru lakha ghor dho dhanta hawa para bachabo karoneh bonailobi
PIG	Contact nearby veterinarian Officer for measure against adverse conditions
	Apni khan parishe koilhe Veterinarian Officer lakha number to rakhibi, enikha hoilhe gauri lagha kiba dhok khan hoilhe jolti matibo baribo

#### Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
COMMON CARP	Remove pond bottom debris before the onset of winter preferably by October
	Apni lakha pukhuri do etu moinah dho sapa khoriboleh laghe

#### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICKEN	As the temperature is dropping, provide vitamin complex. Get ready to provide warmth by use of incandescent bulbs to protect them from cold winds by use of plastic sheets, jute sacks and plastic sacks
	Temperature tho thanda hui ase etu karoneh vitamin complex dibo lageh. Hawa thanda parah bachibo karoneh plastic nahoileh bosta parah ghor kinar bon kuribi aru bulb bi cholabi



## Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



# Agromet Advisory Bulletin

Date : 01-11-2019

Weather Forecast of PEREN(Nagaland) Issued On : 2019-11-01(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-11-02	1	32	20	96	52	4.0	99	2
2019-11-03	0	31	20	95	49	4.2	108	0
2019-11-04	0	31	21	93	47	4.1	112	1
2019-11-05	1	30	20	96	51	3.7	108	0
2019-11-06	0	30	18	96	48	4.2	105	3

### Weather Summary/Alert:

Probability of light rain the coming days
Pani olop kiribolhe dikaiase

### General Advisory:

Contact nearby veterinarian Officer for measure against adverse conditions
Apni khan parishe koilhe Veterinarian Officer lakha number to rakhibi, enikha hoilhe gauri lagha kiba dhok khan hoilhe jolti matibo baribo

### SMS Advisory:

Avoid excessive drying of paddy to avoid breakage of the grains
Dhan bishi bi nusukabi nahoileh dhan goti tho mehin hobo

### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	RIPENING STAGE: Avoid excessive drying of paddy to avoid breakage of the grains
	RIPENING STAGE: Dhan bishi bi nusukabi nahoileh dhan goti tho mehin hobo
GROUNDNUT	POST HARVEST: When temperature is very high, avoid direct sun drying. Collect the pods in gunnies and store on the ground over a layer of sand to avoid any moisture coming in contact with dry pods
	Bishi gorom din tae pariatok thik dhup nichetae nusukabi. Sukha badam khan tae moisture nalagiboh karoneh, bosta tae dhalighenah, mati opor tae bhalu olop rakhighenah hi etu opor tae bosto tho rakhibi
MAIZE	SEEDLING STAGE: Provide mulching to conserve moisture
	SEEDLING STAGE: mati tae moisture thakibo karoneh, sukha patah khan khan rakhibi
SESAME (GINGELLY/TIL)	VEGETATIVE STAGE: When the plants are about 15 cm in height, thin the crop so as to give a spacing of 15-25 cm between plants.

Crop(Varieties)	Crop Specific Advisory
	Ghas 15 cm ocha hua pichetae, ghas majotae 15-25 cm khali thakiboleh ghas khan ulailobi
GINGER	HARVESTING STAGE: However, for making dry ginger, the matured rhizomes are harvested at full maturity i.e. when the leaves turn yellow and start drying.
	Sukha aduwa bonaboleh, pata tho yellow huighenah murijah somoi tae khundilobi aru sakahbi

#### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
FRENCH BEAN	VEGETATIVE STAGE: Weeding should be given 20 – 25 days and 40 – 45 days after sowing. The crop should be earthed up after each weeding.
	Ghas lagaigheanh 20 – 25 din and 40 – 45 din tae jonjol ghas sapha kuribi. Sapha kurah pichetae ghas khan tae mathi olop dhalibi
NAGA KING CHILLI	HARVESTING STAGE: Harvesting should be done when the sky is clear
	Din bal somoi te mircha do ulabi
COLOCASIA/TARO	Harvesting stage. Clean the debris and sundry the harvested tubers
	Kochu kuti han tho bhal para sabha kuri kina, sun de sukhai dibo lageh.

#### Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	Contact nearby veterinarian Officer for measure against adverse conditions
	Apni khan parishe koilhe Veterinarian Officer lakha number to rakhibi, enikha hoilhe gauri lagha kiba dhok khan hoilhe jolti matibo baribo

#### Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
COMMON CARP	As far as possible do not disturb the pond. Prune the tree branches along the pond embankment and allow enough sunlight to fall on the pond water surface
	Pukhuri pani tho parah thok disturb nukuribi. Pukhuri kinar tae ghas lagah branch khan katilobi. Pani opor tae dhup bhal parah pabo lageh

#### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICKEN	As the temperature is dropping, provide vitamin complex. Get ready to provide warmth by use of incandescent bulbs to protect them from cold winds by use of plastic sheets, jute sacks and plastic sacks
	Temperature tho thanda hui ase etu karoneh vitamin complex dibo lageh. Hawa thanda parah bachibo karoneh plastic nahoileh bosta parah ghor kinar bon kuribi aru bulb bi cholabi



## Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



# Agromet Advisory Bulletin

Date : 01-11-2019

Weather Forecast of PHEK(Nagaland) Issued On : 2019-11-01(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-11-02	2	31	18	94	44	3.0	153	2
2019-11-03	1	31	18	92	39	3.1	154	0
2019-11-04	0	30	19	93	36	3.3	156	1
2019-11-05	3	29	18	93	45	3.4	160	1
2019-11-06	3	29	17	94	47	3.3	161	2

### Weather Summary/Alert:

Probability of light rainfall the coming days
Pani etu hapta bi olop kiribolhe dikaiase

### General Advisory:

Contact nearby veterinarian Officer for measure against adverse conditions
Apni khan parishe koilhe Veterinarian Officer lakha number to rakhibi, enikha hoilhe gauri lagha kiba dhok khan hoilhe jolti matibo baribo

### SMS Advisory:

Avoid excessive drying of paddy to avoid breakage of the grains
Dhan bishi bi nusukabi nahoileh dhan goti tho mehin hobo

### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	HARVESTING STAGE: Do not dry the grains at each time more than 4 - 6 hours otherwise grains will shrivel and break
	Hudai sukha somia tae 4-6 ghonta opor husukabi nanoileh dhan goti mehin hobo
GINGER	Harvesting stage. Grading should be done after harvest, this would generate more income in the market
	Athua han tho mati para olai kina bhal aro biya nahoiley chutu aro thangor majot de alak alak rhaki bo ley lage. Etu para bazaar de bikai homoi de lap babo.

### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
BANANA	Remove the affected pest and disease leaf, dried leaves from time to time.

Horticulture (Varieties)	Horticulture Specific Advisory
	Puka aru bimar lagijah pata khan hatabi aru sukha pata khan bi somoi somoi tae hatai thakibi
CABBAGE	VEGETATIVE STAGE: Earthing up should be done 30 days after transplanting
	Transplanting kurighenah 30 din pichetae mathi olop dhalibi

**Livestock Specific Advisory:**

Livestock(Varieties)	Livestock Specific Advisory
PIG	Routine vaccination programme is recommended and once any disease is diagnosed treat immediately with antibiotics
	Somoi tae vaccination kurithakibi aru bimar kunba lagileh joldi antibiotics dibi

**Fisheries Specific Advisory:**

Fisheries(Varieties)	Fisheries Specific Advisory
COMMON CARP	Check the movement of the fish every day. If abnormal behavior is observed consult an expert for suggestion
	Hudai mass khila tho sabo lageh. Thik nikhileh mass expert logotae alusnah kuribi

**Poultry Specific Advisory:**

Poultry(Varieties)	Poultry Specific Advisory
CHICKEN	Proper sanitary measures should be done to control the diseases. Slaughtering of all apparently ailing birds, segregating of in - contact in group of 10 to 15 each; removal of all infective materials such as droppings, residues of poultry cleanliness
	Bimar nalagiboh karoneh sapha parah murgi palibi.