



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 15-10-2019

Weather Forecast of ZUNHEBOTO(Nagaland) Issued On : 2019-10-15(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-10-16	1	34	20	96	46	4.2	127	3
2019-10-17	6	33	19	98	54	4.1	124	0
2019-10-18	0	34	19	98	48	4.2	124	3
2019-10-19	0	33	19	94	45	4.2	127	5
2019-10-20	6	34	19	96	48	4.5	125	3

Weather Summary/Alert:

Probability of light rainfall the coming days
Pani etia ahiboleh din bi pani kiriboleh dikaiase

General Advisory:

Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done
Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done

SMS Advisory:

Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region
Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
JHUM PADDY(LOCAL)	POST HARVEST: Paddy which have been harvested should be dried properly and keep in the storage area which have been cleaned properly Dhan khan katia khan to bhal para sukai khena rakhibi. Thailaka sukhai khena rakha jaka to pal bara sabha khoribi etu bichhe te hi rakhibi
TRC OR WRC(LOCAL)	MATURITY STAGE: For bird control erect scarecrows randomly in the field, scare the bird manually too. Tie old VHS tapes diagonally across and around the field. Use catapults For bird control erect scarecrows randomly in the field, scare the bird manually too. Tie old VHS tapes diagonally across and around the field. Use catapults
SOYBEAN(LOCAL)	PODDING STAGE: Monitor the plants against pest and diseases Etia khut bakia somoi te kheti te pukha bimar khan sabole lakhe

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
BRINJAL(LOCAL)	HARVESTING STAGE: Grow maize in the brinjal field as trap crop
	Makoi do begon lakha kheti te lagabi pukha kumtikoribo kharoneh
CHILLI(LOCAL)	HARVESTING STAGE: Harvesting should be carried out when the sky is clear
	Din bal somoi te mircha do ulabi.
COLOCASIA(LOCAL)	MATURITY STAGE: Remove any diseased leaves and daily monitoring should be done
	Pata biya, puchichakhan to olai khena bilabi
GINGER(LOCAL)	VEGETATIVE STAGE. Bacterial wilt. Drooping and wilting of leaves in the early morning when there is no sunshine. Base of the pseudostem turns grey with water soaked and soft . Provide proper drainage to avoid water stagnation. 4 to 5 years of crop rotation should be followed to avoid disease incidence
	VEGETATIVE STAGE. Bacterial wilt. Pani jabolhe nullah khan bal bara bonabi, pani jama hobo nedhibi. Khelhe koilhe etu bara bacterial wilt bimar hoi, pata khan sukhi khena girichai

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIGGERY(LOCAL)	Find out what diseases are prevalent in your area and timely arrange for vaccination and treatment if necessary
	Apni lakha jakha te ki bimar hoi etu lakha hisab te vaccination khan shoru khoribi

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
FISHERY(LOCAL)	After seven months of rearing or when the water level decreases, the fish should be harvested. Harvesting should be done either by cast net or drag net. Care has to be taken that more person does not enter the pond.
	Apni khan sabi ki pukhuri te pani kumti hoi nah nai. Kumti hoishe koilhe mass to durilobi. Sabo le lakhe ki mass duri somoi te pukhuri te manu bishi najabo lakhe

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
POULTRY(LOCAL)	Vitamin E and selenium supplement should be given to poultry for disease resistance. Mixed the supplement in clean drinking water. Change the water daily
	Etia thanda somoi ahiase, murghi khan bimar nohobo karoneh Vitamin E and supplement (ekta packet hea ase) dho pani te milai khena dibi. Pani to hudai bodli khuribo lageh



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 15-10-2019

Weather Forecast of DIMAPUR(Nagaland) Issued On : 2019-10-15(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-10-16	0	31	21	83	52	5.4	133	3
2019-10-17	0	31	21	84	48	4.8	126	2
2019-10-18	0	31	21	81	50	5.1	118	2
2019-10-19	0	32	21	82	52	5.1	124	3
2019-10-20	0	32	21	83	50	4.7	133	0

Weather Summary/Alert:

Probability of no rain the coming days
Etia ahiboleh din do pani nikiriboleh pareh

General Advisory:

Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done.
Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done.

SMS Advisory:

Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region
Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
LOWLAND PADDY(LOCAL)	PANICLE INITIATION STAGE.For bird control erect scarecrows randomly in the field, scare the bird manually too. Tie old VHS tapes diagonally across and around the field. Use catapults Kheti tae chiria khetaboleh kapra para manu nishinah bonaighenah jaga jaga tae rakhibi. (VHS tapes) color maika kheti bitor aru kinar tae cholabi. Gitis cholabi
GROUNDNUT(LOCAL)	HARVESTING STAGE: After harvest, dry the pods in the sun for 4 or 5 days. Repeat drying for 2 or 3 more days after an interval of 2 or 3 days to ensure complete drying kuti bakia bicche te, doob te baal para sukhaba char/pass din karoneh.
SOYABEAN(LOCAL)	PODDING STAGE: Yellowing of leaves and shedding , indicate the maturity of the crop. Cut the entire plant when most of the pods have turned yellow

Crop(Varieties)	Crop Specific Advisory
	Patha yellow hoikhenah kherah dho , ghoti bakhibohleh dhekaiaise.
SESAME(LOCAL)	VEGETATIVE STAGE: Keep the field weed free upto 40 DAS as the crop is very sensitive during this period
	lakha bicche te 40 DAS tak do habi khan bal para saba koriboleh lakhe

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
NAGA KING CHILLI(LOCAL)	HARVESTING STAGE: In the present weather harvesting should be carried out.
	Din bal somoi te mircha do ulabi
COLOCASIA(LOCAL)	MATURITY/ HARVESTING STAGE. While harvesting , uprooting should be done carefully and the mother corms and cormels are to be separated
	MATURITY/ HARVESTING STAGE. Etia bakia somoi hoishe, ulai somoi te hisab khoribi ki alu to ulai somoi te biya nohobo kharoneh
GINGER(LOCAL)	VEGETATIVE STAGE : Diseased plants should be identified while the crop is in field. Rhizomes from such plants should not be selected for the seed purpose
	Bimarghas khan dho kheti dhe dakha somoi dhe sailobhi. Aru aha sal karoneh bijon rakhibo karoneh etu bimar ghas bara dho narhikibi

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
CATTLE(LOCAL CROSS BREED)	A lot of green fodder is available at this time, hence, to protect animals from problems associated with excess grazing, restrict the time that they spend in pastures. Mix essential salts in the feed of the animals
	Etia to janwar khan kaboleh gass khan bishi ase, hoilebi hisab khoribo laghe ki bishi gass nagabo leh karoneh. Dana di somoi te nimok milaikhena dibi
PIGGERY(LOCAL)	Sick or injured animals should be attended to immediately, preferably moved to a Comfortable environment
	Gauri bimar khan dho alak bareh rakhibo lakhe

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
FISHERY(LOCAL)	Fertilization of pond should not be started as once you start it becomes a permanent part of your management programme. The disadvantages of fertilization usually outweighs the advantages.
	Bariadhak phokhori dhe dawai khan nadhalibi, khelhe khoilhe apni ekbar shoro khorilhe khori dhakiboleh olaboh

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
POULTRY(LOCAL)	Feed during early morning and late afternoon hours to prevent build up of heat. Feed balanced ration with adequate vitamins and minerals.
	Gao gorom nathakibo karoneh, morning aru sam tae hi dana khelabi. Vitamins aru minerals milaighenah thik hisap tae dana dibi



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 15-10-2019

Weather Forecast of KIPHIRE(Nagaland) Issued On : 2019-10-15(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-10-16	4	34	20	91	45	3.1	65	3
2019-10-17	2	33	19	88	41	3.2	56	1
2019-10-18	2	34	19	87	40	3.0	66	2
2019-10-19	3	34	19	84	38	3.1	63	4
2019-10-20	3	34	19	84	37	3.4	60	1

Weather Summary/Alert:

Probability of light rainfall the coming days
Pani etu ahibleh din bi olop kiriboleh dikaiase

General Advisory:

Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done.
Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done.

SMS Advisory:

Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region
Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
JHUM PADDY(LOCAL)	HARVESTING STAGE: Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
	Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
TRC PADDY(LOCAL)	MILK DEVELOPMENT STAGE: For management of leaf folder, keep the bunds clean, spray NSKE 5%, but should be done when the sky is clear
	Leaf folder bishi nohobo karoneh, kheti dhe habi khan sabah rakhibi, aru NSKE 5% bi spray khoribo pareh

Horticulture Specific Advisory:

Horticulture (Varities)	Horticulture Specific Advisory
KING CHILLI(LOCAL)	MATURITY/ HARVESTING STAGE: Weight loss and damage of fresh ripen chillies can be minimized by packing in perforated planstic bags of 1 kg capacity and further storing it in bamboo baskets or carton
	Mircha pakia khan laka ujon aru pojicha para bacha bo karoneh , mircha khan to maika/ bag chenta thakia aru etu ke bamboo laka basket de nahoiley carton te rakely bhal hobo
COLOCASIA(LOCAL)	VEGETATIVE STAGE: For leaf blight, remove the diseased plants, taking them far from the planting area and destroy them by burying, burning or composting
	leaf blight karoneh bata bimar khan ulai khena cholai debi naholai mati dhe rhobai dhebi
LARGE CARDAMOM(LOCAL)	HARVESTING STAGE: Matured fruits should be harvested without delay
	khuti pakia khan to dhik somoi te olai dibo lage

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIGGERY(LOCAL)	Regular deworming should be carried out to check the parasitic infestation. This practice must be started at the age of two weeks followed after 21 days and repeated 3-4 times a year at regular interval
	Gahori baja khan to kichu laga dawai khelai thakibo lageh. Etu dho gahori tui hapta hoi somoi te shuru khuri dibo lageh, aru 21 din piche te 3-4 times ek sal te dibo lage

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
FISHERY (LOCAL)	Do not let large animals such as cows, buffalo graze on the banks of your pond, they are too heavy and may break the banks down
	Janwar dangor khan pukhuri lakha osor te, kinar phaley ghas khan kabo nedhibo laghe. Etu janwar lagha ojon para pukhuri lagha kinar tho bagibo pareh

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
POULTRY(LOCAL)	Vitamin E and Selenium supplement should be given for disease resistance. Mix the supplements in clean drinking water and change the water daily.
	Etia thanda somoi ahiase, murghi khan bimar nohobo karoneh Vitamin E and selenium supplement (exta packet hea ase) dho pani te milai khena dibi. Pani to hudai bodli khiribo lageh



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 15-10-2019

Weather Forecast of KOHIMA(Nagaland) Issued On : 2019-10-15(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-10-16	7	33	20	92	55	5.1	139	3
2019-10-17	6	32	19	94	56	4.7	136	1
2019-10-18	4	33	19	94	56	4.8	132	3
2019-10-19	6	33	20	94	57	4.9	134	5
2019-10-20	13	33	19	94	61	4.8	142	1

Weather Summary/Alert:

Probability of light rainfall the coming days
Etia ahiboleh din bi pani kiribolhe dikaiase

General Advisory:

Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done.
Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done.

SMS Advisory:

Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region
Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
JHUM PADDY(LOCAL)	POST HARVEST: Paddy which have been harvested should be dried properly and keep in the storage area which have been cleaned properly Dhan khan katia khan to bhal para sukai khena rakhibi. Thailaka sukhai khena rakha jaka to pal bara sabha khoribi etu bichhe te hi rakhibi
TRC OR WRC(LOCAL)	MILK DEVELOPMENT STAGE: Keep the bunds clean to prevent infestation of pest and disease incidence Kheti lagha khenar khan bariadhok saba rakhibi, pukha bimar para dhik dhar nedibo karoneh
FOXTAIL MILLET(LOCAL)	FLOWERING STAGE: Proper monitoring should be done for pest and disease. Management practices should be carried out during the initial days

Crop(Varieties)	Crop Specific Advisory
	Pukha bimar khan ase nah nai somoi dhe sai dakhibo lakhe. Takhele cholti management practices khoribo lakhe
SOYBEAN(LOCAL)	MATURITY/HARVESTING STAGE: Seeds should be harvested at field maturity. Delayed harvesting should be avoided. As rain has occurred the preceeding week, after harvesting proper drying in the sun should be done. keep the harvested product protected from rain khoti khan dhik somoi dhe ula bo lakhe. Dheri ulai lhe khoti loss hobo. Apni para paba nishena napabo

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
BANANA(LOCAL)	FRUITING STAGE: As rain has occurred the preceding week, carry out earthing up and and propping should be done to provide support Chaha hapta bi pani bishi kirishe,khul gass lakha niche te mati dabi aru Khol, ghas dhe ulaighena dakhelhe bamboo para support dhebi
GINGER(LOCAL)	VEGETATIVE STAGE. Bacterial wilt. Drooping and wilting of leaves in the early morning when there is no sunshine.Base of the pseudostem turns grey with water soaked and soft . Provide proper drainage to avoid water stagnation. 4 to 5 years of crop rotation should be followed to avoid disease incidence VEGETATIVE STAGE. Bacterial wilt. Pani jabolhe nullah khan bal bara bonabi, pani jama hobo nedhibi. Khelhe koilhe etu bara bacterial wilt bimar hoi, pata khan sukhi khena girichai

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIGGERY(LOCAL)	Find out what diseases are prevalent in your area and timely arrange for vaccination and treatment if necessary Apni lakha jakha te ki bimar hoi etu lakha hisab te vaccination khan shoru khoribi

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
FISHERY(LOCAL)	Liming during winter months is essential even if the pH is within the desired range. This is because fish are susceptible to diseases and lime act as a prophylactic agent. Etia thanda moinah ahiase, to pukhuri dhe pH dho dhik ase koilebi liming to khoriboleh lakhe kalhekoileh mass khan to bimar hobo bishe easy ase aru apni liming khoria kharoneh bimar para bachhai

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
POULTRY(LOCAL)	Vitamin E and Selenium supplement should be given for disease resistance. Mix the supplements in clean drinking water and change the water daily. Etia thanda somoi ahiase, murghi khan bimar nohobo karoneh Vitamin E and selenium supplement (exta packet hea ase) dho pani te milai khena dibi. Pani to hudai bodli khiribo lageh



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 15-10-2019

Weather Forecast of LONGLENG(Nagaland) Issued On : 2019-10-15(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-10-16	0	32	20	87	50	4.6	136	2
2019-10-17	0	32	20	91	53	4.5	133	0
2019-10-18	0	32	19	89	52	4.5	121	0
2019-10-19	0	32	20	85	49	4.4	132	4
2019-10-20	1	33	20	87	50	4.4	140	5

Weather Summary/Alert:

Probability of light rain the coming days
Probability of light rain the coming days

General Advisory:

Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done
Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done

SMS Advisory:

Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region
Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
JHUM PADDY(LOCAL)	POST HARVEST: Paddy which have been harvested should be dried properly and keep in the storage area which have been cleaned properly Dhan khan katia khan to bhal para sukai khena rakhibi. Thailaka sukhai khena rakha jaka to pal bara sabha khoribi etu bichhe te hi rakhibi
TRC OR WRC(LOCAL)	MATURITY STAGE: For bird control erect scarecrows randomly in the field, scare the bird manually too. Tie old VHS tapes diagonally across and around the field. Use catapults For bird control erect scarecrows randomly in the field, scare the bird manually too. Tie old VHS tapes diagonally across and around the field. Use catapults
SOYBEAN(LOCAL)	PODDING STAGE: Monitor the plants against pest and diseases Etia khut bakia somoi te kheti te pukha bimar khan sabole lakhe

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
BRINJAL(LOCAL)	HARVESTING STAGE: Grow maize in the brinjal field as trap crop
	Makoi do begon lakha kheti te lagabi pukha kumtikoribo kharoneh
CHILLI(LOCAL)	HARVESTING STAGE: Harvesting should be carried out when the sky is clear
	Din bal somoi te mircha do ulabi.
CHILLI(LOCAL)	MATURITY STAGE: Remove any diseased leaves and daily monitoring should be done
	Pata biya, puchichakhan to olai khena bilabi
GINGER(LOCAL)	VEGETATIVE STAGE. Bacterial wilt. Drooping and wilting of leaves in the early morning when there is no sunshine. Base of the pseudostem turns grey with water soaked and soft . Provide proper drainage to avoid water stagnation. 4 to 5 years of crop rotation should be followed to avoid disease incidence
	VEGETATIVE STAGE. Bacterial wilt. Pani jabolhe nullah khan bal bara bonabi, pani jama hobo nedhibi. Khelhe koilhe etu bara bacterial wilt bimar hoi, pata khan sukhi khena girichai

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIGGERY(LOCAL)	Find out what diseases are prevalent in your area and timely arrange for vaccination and treatment if necessary
	Apni lakha jakha te ki bimar hoi etu lakha hisab te vaccination khan shoru khoribi

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
FISHERY(LOCAL)	After seven months of rearing or when the water level decreases, the fish should be harvested. Harvesting should be done either by cast net or drag net. Care has to be taken that more person does not enter the pond.
	Apni khan sabi ki pukhuri te pani kumti hoi nah nai. Kumti hoishe koilhe mass to durilobi. Sabo le lakhe ki mass duri somoi te pukhuri te manu bishi najabo lakhe

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
POULTRY(LOCAL)	Vitamin E and selenium supplement should be given to poultry for disease resistance. Mixed the supplement in clean drinking water. Change the water daily
	Etia thanda somoi ahiase, murghi khan bimar nohobo karoneh Vitamin E and supplement (ekta packet hea ase) dho pani te milai khena dibi. Pani to hudai bodli khuribo lageh



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 15-10-2019

Weather Forecast of MOKOKCHUNG(Nagaland) Issued On : 2019-10-15(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-10-16	1	33	20	91	52	4.5	126	2
2019-10-17	6	32	20	94	56	4.3	128	0
2019-10-18	0	33	19	93	53	4.4	116	1
2019-10-19	0	33	20	89	50	4.3	125	4
2019-10-20	6	33	20	91	52	4.3	137	5

Weather Summary/Alert:

Probability of light rainfall the coming days
Etia ahiboleh din bi pani olop hi kiriboleh dikaiase

General Advisory:

Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done
Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done

SMS Advisory:

Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region
Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
JHUM PADDY(LOCAL)	POST HARVEST: Paddy which have been harvested should be dried properly and keep in the storage area which have been cleaned properly
	POST HARVEST: Paddy which have been harvested should be dried properly and keep in the storage area which have been cleaned properly
TRC OR WRC(LOCAL)	MATURITY STAGE: For bird control erect scarecrows randomly in the field, scare the bird manually too. Tie old VHS tapes diagonally across and around the field. Use catapults
	For bird control erect scarecrows randomly in the field, scare the bird manually too. Tie old VHS tapes diagonally across and around the field. Use catapults
SOYBEAN(LOCAL)	PODDING STAGE: Monitor the plants against pest and diseases
	Etia khut bakia somoi te kheti te pukha bimar khan sabole lakhe

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
BRINJAL(LOCAL)	HARVESTING STAGE: Grow maize in the brinjal field as trap crop
	Makoi do begon lakha kheti te lagabi pukha kumtikoribo kharoneh
CHILLI(LOCAL)	HARVESTING STAGE: Harvesting should be carried out when the sky is clear
	Din bal somoi te mircha do ulabi
GINGER(LOCAL)	VEGETATIVE STAGE: Bacterial wilt. Drooping and wilting of leaves in the early morning when there is no sunshine. Base of the pseudostem turns grey with water soaked and soft . Provide proper drainage to avoid water stagnation. 4 to 5 years of crop rotation should be followed to avoid disease incidence
	VEGETATIVE STAGE: Bacterial wilt. Pani jabolhe nullah khan bal bara bonabi, pani jama hobo nedhibi. Khelhe koilhe etu bara bacterial wilt bimar hoi, pata khan sukhi khena girichai
COLOCASIA(LOCAL)	MATURITY STAGE: Remove any diseased leaves and daily monitoring should be done
	Pata biya, puchichakhan to olai khena bilabi

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIGGERY(LOCAL)	Find out what diseases are prevalent in your area and timely arrange for vaccination and treatment if necessary
	Apni lakha jakha te ki bimar hoi etu lakha hisab te vaccination khan shoru khoribi

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
FISHERY(LOCAL)	After seven months of rearing or when the water level decreases, the fish should be harvested. Harvesting should be done either by cast net or drag net. Care has to be taken that more person does not enter the pond.
	Apni khan sabi ki pukhuri te pani kumti hoi nah nai. Kumti hoishe koilhe mass to durilobi. Sabo le lakhe ki mass duri somoi te pukhuri te manu bishi najabo lakhe

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
POULTRY(LOCAL)	Vitamin E and selenium supplement should be given to poultry for disease resistance. Mixed the supplement in clean drinking water. Change the water daily
	Etia thanda somoi ahiase, murghi khan bimar nohobo karoneh Vitamin E and supplement (ekta packet hea ase) dho pani te milai khena dibi. Pani to hudai bodli khuribo lageh



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 15-10-2019

Weather Forecast of MON(Nagaland) Issued On : 2019-10-15(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-10-16	8	33	20	88	54	4.8	135	2
2019-10-17	5	32	20	91	58	4.5	133	1
2019-10-18	2	32	19	88	55	4.7	126	0
2019-10-19	3	33	20	86	54	4.7	136	3
2019-10-20	5	33	20	87	55	4.4	140	6

Weather Summary/Alert:

Probability of light rainfall the coming days

Pani etu hapta bi olop hi kiribolhe dikaiase

General Advisory:

Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done

Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done

SMS Advisory:

Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region

Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
JHUM PADDY(LOCAL)	POST HARVEST: Follow proper sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds
	Apni khan dhan sukhai somoi te aru khuntia somoi te saboleh lakhe ki dhan khan to biya nohobo laghe.
TRC PADDY(LOCAL)	MILK DEVELOPMENT STAGE:Fungus can attack the leaves, collars, nodes and panicles of rice plants. Scout fields for leaf blast symptoms. If leaf blast is present, a preventative fungicide application should be considered. For rice blast management, destroy the infested residue, stubbles, grass and weeds present in the bunds of the field as they act as a source for the infection

Crop(Varieties)	Crop Specific Advisory
	Etia lakha somoi te blast bimar olabo pare, kheti te phal bare sabi. Bishi dakishe koileh ekta fungicide cholabo lakhe . Kheti lakha kinar khan te habi, jongol khan sabha khoridakibo lageh
GROUNDNUT(LOCAL)	MATURITY STAGE: Observe the crop, drying and falling of older leaves and yellowing of the top leaves indicate maturity Ghas tho saithakibi. Pakiboleh somoi tae pata sukibo, bura pata khan giribo aru pata opor khan yellow hobo
SOYBEAN(LOCAL)	MATURITY/HARVESTING STAGE:Seeds should be dried to safe moisture levels soon after harvest Khoti bijon khan do ulabishi dhe bhal bara sukhaba

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
CARDAMON(LOCAL)	MATURITY/HARVESTING STAGE: Matured fruits should be harvested without delay khuti pakia khan to dhik somoi te olai dibo lage
COLOCASIA(LOCAL)	MATURITY STAGE: Remove any diseased leaves and daily monitoring should be done Pata biya, puchichakhan to olai khena bilabi

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIGGERY(LOCAL)	Contact nearby veterinarian officer for measures against adverse conditions Gauri te kiba bimarkhan hoishe kuilhe cholti veterinarian officer logotae alusnah khorilobi

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
FISHERY(LOCAL)	Remove pond bottom debris before the onset of winter preferably this month Pukhuri lakha niche te litera khan dakato parishe koile etu moinah korilobi

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
POULTRY(LOCAL)	Vitamin E and selenium supplement should be given to poultry for disease resistance. Mixed the supplement in clean drinking water. Change the water daily Etia thanda somoi ahiase, murghi khan bimar nohobo karoneh Vitamin E and supplement (ekta packet hea ase) dho pani te milai khena dibi.Pani to hudai bodli khuribo lageh



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 15-10-2019

Weather Forecast of PEREN(Nagaland) Issued On : 2019-10-15(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-10-16	17	33	22	96	58	3.6	103	3
2019-10-17	7	34	21	96	56	3.4	90	2
2019-10-18	6	34	21	93	58	3.8	97	2
2019-10-19	6	34	21	94	57	3.5	104	2
2019-10-20	2	35	21	94	50	3.8	107	1

Weather Summary/Alert:

Probability of light rainfall the coming days
Pani etu hapta bi olop hi kiribolhe dikaiase

General Advisory:

Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done.
Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done

SMS Advisory:

Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region
Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
JHUM PADDY(LOCAL)	HARVESTING STAGE: As there is probability of rain the coming days too, do not keep the harvested product in the open, keep it in a place where it is protected from rain
	HARVESTING STAGE: As there is probability of rain the coming days too, do not keep the harvested product in the open, keep it in a place where it is protected from rain
TRC OR WRC(LOCAL)	PANICLE INITIATION STAGE.For bird control erect scarecrows randomly in the field, scare the bird manually too. Tie old VHS tapes diagonally across and around the field. Use catapults
	Kheti tae chiria khetaboleh kapra para manu nishinah bonaighenah jaga jaga tae rakhibi. (VHS tapes) color maika kheti bitor aru kinar tae cholabi. Gitis cholabi

Crop(Varieties)	Crop Specific Advisory
GROUNDNUT(LOCAL)	HARVESTING STAGE: After harvest, dry the pods in the sun for 4 or 5 days. Repeat drying for 2 or 3 more days after an interval of 2 or 3 days to ensure complete drying
	kuti bakia bicche te, doob te baal para sukhabhi char/pass din karoneh.
SOYBEAN(LOCAL)	PODDING STAGE: Yellowing of leaves and shedding , indicate the maturity of the crop. Cut the entire plant when most of the pods have turned yellow
	Patha yellow hoikhenah kherah dho , ghoti bakhbohleh dhekaiaase

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
NAGA CHILLI(LOCAL)	HARVESTING STAGE: Harvesting should be done when the sky is clear
	Din bal somoi te mircha do ulabi.
COLOCASIA(LOCAL)	MATURITY/ HARVESTING STAGE. While harvesting , uprooting should be done carefully and the mother corms and cormels are to be separated
	MATURITY/ HARVESTING STAGE. Etia bakia somoi hoishe, ulai somoi te hisab khoribi ki alu to ulai somoi te biya nohobo kharoneh

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIGGERY(LOCAL)	Sick or injured animals should be attended to immediately, preferably moved to a Comfortable environment
	Bimarghas khan dho kheti dhe dakha somoi dhe sailobhi. Aru aha sal karoneh bijon rakhibo karoneh etu bimar ghas bara dho narhikibi

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
FISHERY(LOCAL)	Fertilization of pond should not be started as once you start it becomes a permanent part of your management programme. The disadvantages of fertilization usually outweighs the advantages.
	Bariadhak phokhori dhe dawai khan nadhalibi, khelhe khoilhe apni ekbar shoro khorilhe khori dhakiboleh olaboh

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
POULTRY(LOCAL)	Feed during early morning and late afternoon hours to prevent build up of heat.Feed balanced ration with adequate vitamins and minerals
	Gao gorom nathakibo karoneh, morning aru sam tae hi dana khelabi. Vitamins aru minerals milaighenah thik hisap tae dana dibi.



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 15-10-2019

Weather Forecast of PHEK(Nagaland) Issued On : 2019-10-15(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-10-16	10	33	20	96	46	3.2	156	3
2019-10-17	6	33	20	94	44	2.8	145	1
2019-10-18	5	33	19	93	44	2.9	153	3
2019-10-19	6	34	20	93	45	3.2	158	4
2019-10-20	9	34	19	90	47	2.9	171	0

Weather Summary/Alert:

Probability of light rain the coming days
Etia ahiboleh din bi pani olop kiriboleh dikaiase

General Advisory:

Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done
Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done

SMS Advisory:

Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region
Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
JHUM PADDY(LOCAL)	POST HARVEST: Paddy which have been harvested should be dried properly and keep in the storage area which have been cleaned properly Dhan khan katia khan to bhal para sukai khena rakhibi. Thailaka sukhai khena rakha jaka to pal bara sabha khoribi etu bichhe te hi rakhibi
TRC OR WRC(LOCAL)	MILK DEVELOPMENT STAGE: Keep the bunds clean to prevent infestation of pest and disease incidence Kheti lagha khenar khan bariadhok saba rakhibi, pukha bimar para dhik dhar nedibo karoneh
FOXTAIL MILLET(LOCAL)	FLOWERING STAGE: Proper monitoring should be done for pest and disease. Management practices should be carried out during the initial days

Crop(Varieties)	Crop Specific Advisory
	Pukha bimar khan ase nah nai somoi dhe sai dakhibo lakhe. Takhele cholti management practices khoribo lakhe

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
GINGER(LOCAL)	VEGETATIVE STAGE. Bacterial wilt. Drooping and wilting of leaves in the early morning when there is no sunshine. Base of the pseudostem turns grey with water soaked and soft. Provide proper drainage to avoid water stagnation. 4 to 5 years of crop rotation should be followed to avoid disease incidence
	VEGETATIVE STAGE. Bacterial wilt. Pani jabolhe nullah khan bal bara bonabi, pani jama hobo nedhibi. Khelhe koilhe etu bara bacterial wilt bimar hoi, pata khan sukhi khena girichai
SOYBEAN(LOCAL)	MATURITY/HARVESTING STAGE: Seeds should be harvested at field maturity. Delayed harvesting should be avoided. As rain has occurred the preceeding week, after harvesting proper drying in the sun should be done. keep the harvested product protected from rain"
	khoti khan dhik somoi dhe ula bo lakhe. Dheri ulai lhe khoti loss hobo. Apni para paba nishena napabo

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIGGERY(LOCAL)	Find out what diseases are prevalent in your area and timely arrange for vaccination and treatment if necessary
	Apni lakha jakha te ki bimar hoi etu lakha hisab te vaccination khan shoru khoribi

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
FISHERY(LOCAL)	Liming during winter months is essential even if the pH is within the desired range. This is because fish are susceptible to diseases and lime act as a prophylactic agent
	Etia thanda moinah ahiase, to pukhuri dhe pH dho dhik ase koilebi liming to khoriboleh lakhe kalhekoileh mass khan to bimar hobo bishe easy ase aru apni liming khoriboleh lakhe para bachhai

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
POULTRY(LOCAL)	Vitamin E and Selenium supplement should be given for disease resistance. Mix the supplements in clean drinking water and change the water daily.
	Etia thanda somoi ahiase, murghi khan bimar nohobo karoneh Vitamin E and selenium supplement (exta packet hea ase) dho pani te milai khena dibi. Pani to hudai bodli khiribo lageh



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 15-10-2019

Weather Forecast of TUENSANG(Nagaland) Issued On : 2019-10-15(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-10-16	8	34	20	93	48	3.0	106	2
2019-10-17	2	33	19	93	49	2.9	108	1
2019-10-18	1	34	19	90	40	3.1	114	0
2019-10-19	0	34	19	85	36	3.2	115	2
2019-10-20	0	34	19	88	38	3.1	111	2

Weather Summary/Alert:

Probability of light rainfall the coming day
Pani olop hi kiribolhe dikaiase

General Advisory:

Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done.
Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done.

SMS Advisory:

Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region
Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
JHUM PADDY(LOCAL)	POST HARVEST: Paddy which have been harvested should be dried properly and keep in the storage area which have been cleaned properly Dhan khan katia khan to bhal para sukai khena rakhibi. Thailaka sukhai khena rakha jaka to pal bara sabha khoribi etu bichhe te hi rakhibi
TRC OR WRC(LOCAL)	MILK DEVELOPMENT STAGE: For management of leaf folder, keep the bunds clean, spray NSKE 5% but should be done when the sky is clean Leaf folder bishi nohobo karoneh, kheti dhe habi khan sabah rakhibi, aru NSKE 5% bi spray khoribo pareh

Horticulture Specific Advisory:

Horticulture (Varities)	Horticulture Specific Advisory
KING CHILLI(LOCAL)	MATURITY/ HARVESTING STAGE: Weight loss and damage of fresh ripen chillies can be minimized by packing in perforated plastic bags of 1 kg capacity and further storing it in bamboo baskets or cartons
	Mircha pakia khan laka ujon aru pojicha para bacha bo karoneh , mircha khan to maika/ bag chenta thakia aru etu ke bamboo laka basket de nahoiley carton te rakely bhal hobo
COLOCASIA(LOCAL)	VEGETATIVE STAGE: For leaf blight, remove the diseased plants, taking them far from the planting area and destroy them by burying, burning or compost
	leaf blight karoneh bata bimar khan ulai khena cholai debi naholai mati dhe rhobai dhebi

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIGGERY(LOCAL)	Regular deworming should be carried out to check the parasitic infestation. This practice must be started at the age of two weeks followed after 21 days and repeated 3-4 times a year at regular interval
	Gahori baja khan to kichu laga dawai khelai thakibo lageh. Etu dho gahori tui hapta hoi somoi te shuru khuri dibo lageh, aru 21 din piche te 3-4 times ek sal te dibo lage

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
FISHERY(LOCAL)	Do not let large animals such as cows, buffalo graze on the banks of your pond, they are too heavy and may break the banks down
	Janwar dangor khan pukhuri lakha osor te, kinar phaley ghas khan kabo nedhibo laghe. Etu janwar lagha ojon para pukhuri lagha kinar tho bagibo pareh

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
POULTRY(LOCAL)	Vitamin E and Selenium supplement should be given for disease resistance. Mix the supplements in clean drinking water and change the water daily.
	Etia thanda somoi ahiase, murghi khan bimar nohobo karoneh Vitamin E and selenium supplement (extra packet hea ase) dho pani te milai khena dibi. Pani to hudai bodli khiribo lageh



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 15-10-2019

Weather Forecast of WOKHO(Nagaland) Issued On : 2019-10-15(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-10-16	4	32	20	91	60	4.7	94	4
2019-10-17	6	32	20	93	63	4.6	97	1
2019-10-18	4	32	19	94	61	4.7	77	2
2019-10-19	5	32	20	90	60	4.7	87	4
2019-10-20	13	33	19	91	61	4.4	99	2

Weather Summary/Alert:

Probability of light rain the coming days
Etia ahilbolhe din bi pani olop kiribolhe pareh

General Advisory:

Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done
Nursery preparation should be started for vegetables like cabbage, cauliflower, knolkhol, broccoli, radish, tomato and arrangement for quality seeds should be done

SMS Advisory:

Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region
Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
JHUM PADDY(LOCAL)	POST HARVEST: Paddy which have been harvested should be dried properly and keep in the storage area which have been cleaned properly
	Dhan khan katia khan to bhal para sukai khena rakhibi. Thailaka sukhai khena rakha jaka to pal bara sabha khoribi etu bichhe te hi rakhibi
TRC OR WRC(LOCAL)	MATURITY STAGE: For bird control erect scarecrows randomly in the field, scare the bird manually too. Tie old VHS tapes diagonally across and around the field. Use catapults
	For bird control erect scarecrows randomly in the field, scare the bird manually too. Tie old VHS tapes diagonally across and around the field. Use catapults
SOYBEAN(LOCAL)	PODDING STAGE: Monitor the plants against pest and diseases
	Etia khut bakia somoi te kheti te pukha bimar khan sabole lakhe

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
BRINJAL(LOCAL)	HARVESTING STAGE: Grow maize in the brinjal field as trap crop
	Makoi do begon lakha kheti te lagabi pukha kumtikoribo kharoneh
CHILLI(LOCAL)	HARVESTING STAGE: Harvesting should be carried out when the sky is clear
	Din bal somoi te mircha do ulabi
COLOCASIA(LOCAL)	MATURITY STAGE: Remove any diseased leaves and daily monitoring should be done
	Pata biya, puchichakhan to olai khena bilabi
GINGER(LOCAL)	VEGETATIVE STAGE. Bacterial wilt. Drooping and wilting of leaves in the early morning when there is no sunshine. Base of the pseudostem turns grey with water soaked and soft . Provide proper drainage to avoid water stagnation. 4 to 5 years of crop rotation should be followed to avoid disease incidence
	VEGETATIVE STAGE. Bacterial wilt. Pani jabolhe nullah khan bal bara bonabi, pani jama hobo nedhibi. Khelhe koilhe etu bara bacterial wilt bimar hoi, pata khan sukhi khena girichai

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIGGERY(LOCAL)	Find out what diseases are prevalent in your area and timely arrange for vaccination and treatment if necessary
	Apni lakha jakha te ki bimar hoi etu lakha hisab te vaccination khan shoru khoribi

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
FISHERY(LOCAL)	After seven months of rearing or when the water level decreases, the fish should be harvested. Harvesting should be done either by cast net or drag net. Care has to be taken that more person does not enter the pond.
	Apni khan sabi ki pukhuri te pani kumti hoi nah nai. Kumti hoishe koilhe mass to durilobi. Sabo le lakhe ki mass duri somoi te pukhuri te manu bishi najabo lakhe

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
POULTRY(LOCAL)	Vitamin E and selenium supplement should be given to poultry for disease resistance. Mixed the supplement in clean drinking water. Change the water daily
	Etia thanda somoi ahiase, murghi khan bimar nohobo karoneh Vitamin E and supplement (ekta packet hea ase) dho pani te milai khena dibi. Pani to hudai bodli khuribo lageh