



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 22-10-2019

Weather Forecast of ZUNHEBOTO(Nagaland) Issued On : 2019-10-22(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-10-23	0	31	18	95	52	4.5	119	1
2019-10-24	2	31	19	99	50	4.1	118	6
2019-10-25	3	31	18	99	51	3.0	70	8
2019-10-26	4	27	17	99	66	3.8	71	8
2019-10-27	20	25	19	100	99	2.1	106	8

Weather Summary/Alert:

Probability of light rainfall the coming days
Pani etu hapta bi olop kiribolhe dikaiase

General Advisory:

Harvesting of ginger and turmeric should be done. Weeding and earthing up should be carried out.
Adua, haldi khan harvest khoribo pareh. Pakan te habi khan pila biche te mati dho utabi

SMS Advisory:

To control the incidence of sucking pest in the vegetable crops farmer are advised to spray neem seed kernal extract 5 percent
Sobji te sucking puka komti koribo karoneh neem seed kernal extract 5 percent alibi

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
JHUM PADDY(LOCAL)	POST HARVEST: Paddy which have been harvested should be dried properly and keep in the storage area which have been cleaned properly
	Dhan khan katia khan to bhal para sukai khena rakhibi. Thailaka sukhai khena rakha jaka to pal bara sabha khoribi etu bichhe te hi rakhibi
TRC OR WRC(LOCAL)	MATURITY STAGE: As by next month paddy will be harvested, store house should be kept ready by cleaning up of old grain and grain debris and proper sanitation should be maintained
	Aha moinah para to dhan katibo shoruh hobo, to dhan rakhiboleh ghor to pal para saba khorilobi
SOYBEAN(LOCAL)	HARVESTING STAGE: In the present weather , harvesting should be done before the pods are shattered
	Etia lakha somoi te akhoni lakha khoti dho nabagia agotae harvest khorilobi

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
BRINJAL(LOCAL)	HARVESTING STAGE: Timely harvesting should be done
	Dhik somoi te bijon ulabi
CHILLI(LOCAL)	HARVESTING STAGE: Timely harvesting should be carried out to prevent loss
	Thik somoi te mircha to olabo lakhe
GINGER(LOCAL)	MATURITY STAGE: For vegetable purpose , harvesting can be carried out
	Sobji te hali khena kabo kharoneh to etia para harvest khoribo pareh
COLOCASIA(LOCAL)	HARVESTING STAGE: Leaves which have turned completely brown and have reduced or shirk can be harvested
	Patha khan brown, sukha hoilhe harvest khoribo pareh

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIGGERY(LOCAL)	Vaccination against Foot and Mouth disease in pig should be done. Prior to vaccination, deworm all your animals with appropriate anti-helmentric drugs one week before vaccinating your animals
	Mukh aro dhang laka bimari para paja boley karone vaccine doh mari debi. Vaccine namara ake tey ek hapta ake tey janwar han ke kichu laga dawai di debi anti-helmentric drugs para.

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
FISHERY(LOCAL)	Remove pond bottom debris before the onset of winter preferably by October
	Apni lakha pukhuri do etu moinah dho sapa khoriboleh laghe

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
POULTRY(LOCAL)	Vaccination should be started for prevalent disease like Ranikhet disease and deworm after every two months for backyard and intensive system both.
	Ranikhet bimar karoneh Vaccination dho shuru khorianai khoileh, ajikali shurukhoribi aru parishekoileh dui moinah bichhedeh kichu dawai to dibo lageh



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 22-10-2019

Weather Forecast of DIMAPUR(Nagaland) Issued On : 2019-10-22(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-10-23	0	32	20	86	48	3.9	123	0
2019-10-24	0	31	21	88	54	5.0	108	4
2019-10-25	0	30	18	94	55	4.2	64	6
2019-10-26	1	24	17	93	72	6.9	32	8
2019-10-27	4	26	17	98	90	5.4	64	8

Weather Summary/Alert:

Probability of light rainfall on 27th Oct

Pani 27th te olop kiriboleh pareh

General Advisory:

Harvesting of ginger and turmeric should be done. Weeding and earthing up should be carried out.

Adua, haldi khan harvest khoribo pareh. Pakan te habi khan pila biche te mati dho utabi

SMS Advisory:

To control the incidence of sucking pest in the vegetable crops farmer are advised to spray neem seed kernal extract 5 percent

Sobji te sucking puka komti koribo karoneh neem seed kernal extract 5 percent alibi

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
LOWLAND PADDY(LOCAL)	MILK DEVELOPMENT STAGE: From heading till harvesting, inspect field, remove and destroy off type plants and weeds. Dhan to nagata dhak, somoi te sai dhakibolageh ki ghas te bimar pukha khan ase nah nai, jongol ghas aru habi khan bi somoi te ulai dibo lageh
GROUNDNUT(LOCAL)	POST HARVEST: After harvest, dry the pods in the sun for 4 or 5 days. Repeat drying for 2 or 3 more days after an interval of 2 or 3 days to ensure complete drying Kuti bakia biche te, doob te baal para sukhabi char/pass din karoneh
SOYABEAN(LOCAL)	HARVESTING STAGE: In the present weather harvesting should be carried out. Threshing should be done carefully to prevent damage of seed coat Etia lakha somoi te harvesting to shoru khorilobo laghe.

Crop(Varieties)	Crop Specific Advisory
RABI MAIZE(RCM 76)	LAND PREPARATION: Land preparation should be started for rabi maize. Soils should have adequate drainage. Field should be ploughed properly so as to expose the pupae of red hairy caterpillar. Sow certified seeds from a reliable source to prevent seed rot and seedling blight. If one wants to use his own seed, seed treatment is a must. Some of the varieties recommended for our region are HQPM-1, Allrounder and RCM-76.
	Makoi lagha bo lthe mati bonailobi. Mati to pal bara kuntibi red hairy caterpillar lagha pupae to expose khoribo karoneh. Seed rot aru seedling blight nahabo karoneh bijon to pal dokan para hi khenibolhe laghe. Apni khan nijor lagha bijon hi jobo monjailhe seed treatment to khoribo laghe
SESAME(LOCAL)	VEGETATIVE STAGE: In the present weather timely weeding should be done
	Etia lakha somoi te habi pilabo laghe
TORIA(TS 38)	Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region
	Toria lakha Bijon laghabo lthe shoru khoribi. M-27, TS-36 and TS-38 bijon to moikan lagha jakha karoneh pal ase

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
FRENCH BEAN(LOCAL)	SOWING STAGE: Treat the seeds with Thiram or carbendazim @2g/kg of seeds 24 hours before sowing to control fungal diseases
	Bijon ghoti nalagha agotae bijon ghoti to Thiram nahoileh carbendazi 2g/kg of seeds logo te milai khena lagabi
NAGA KING CHILLI(LOCAL)	HARVESTING STAGE: Timely harvesting should be carried out to prevent loss
	Thik somoi te mircha to olabo lakhe
COLOCASIA(LOCAL)	HARVESTING STAGE: Leaves which have turned completely brown and have reduced or shirk can be harvested
	Patha khan brown, sukha hoilhe harvest khoribo pareh
GINGER(LOCAL)	MATURITY STAGE: When the leaves turn yellow and start gradually drying up, the clumps should be lifted carefully with a spade or digging fork and the adhering soil removed
	Patha sukha aru yellow hoilhe, baorah bara asti para konti bi

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
CATTLE(LOCAL CROSS BREED)	There is a change in the weather from this month onwards, hence, due arrangements should be made to protect the animals from the onset of the winter.
	Etu moinah para to hawa pani bodli hoi, apni khan bi ghoru lakha ghor dho dhanta hawa para bachabo karoneh bonailobi
PIGGERY(LOCAL)	Contact nearby veterinarian Officer for measure against adverse conditions
	Apni khan parishe koilhe Veterinarian Officer lakha number to rakhibi, enikha hoilhe gauri lagha kiba dhok khan hoilhe jolti matibo baribo

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
FISHERY(LOCAL)	Remove pond bottom debris before the onset of winter preferably by October
	Apni lakha pukhuri do etu moinah dho sapa khoriboleh laghe

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
--------------------	---------------------------

Poultry(Varieties)	Poultry Specific Advisory
POULTRY(LOCAL)	Vaccination should be started for prevalent disease like Ranikhet disease and deworm after every two months for backyard and intensive system both.
	Ranikhet bimar karoneh Vaccination dho shuru khorianai khoileh, ajikali shurukhoribi aru parishekoileh dui moinah bichhedeh kichu dawai to dibo lageh



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 22-10-2019

Weather Forecast of KIPHIRE(Nagaland) Issued On : 2019-10-22(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-10-23	1	31	18	78	41	3.9	59	1
2019-10-24	3	32	20	83	34	3.1	127	4
2019-10-25	3	29	19	89	45	3.0	35	6
2019-10-26	2	30	20	92	48	2.9	93	8
2019-10-27	17	27	20	96	80	2.6	80	8

Weather Summary/Alert:

Probability of light rainfall the coming days
Pani etia ahiboleh din bi olop kiribolhe dikaiase

General Advisory:

Harvesting of ginger and turmeric should be done. Weeding and earthing up should be carried out.
Adua, haldi khan harvest khoribo pareh. Pakan te habi khan pila biche te mati dho utabi

SMS Advisory:

To control the incidence of sucking pest in the vegetable crops farmer are advised to spray neem seed kernal extract 5 percent
Sobji te sucking puka komti koribo karoneh neem seed kernal extract 5 percent alibi

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
JHUM PADDY(LOCAL)	STORAGE: Care should be taken to store different varieties of paddy separately Sabolhe lakhe ki dhan variety hisam te alak alak rakhibi
TRC PADDY(LOCAL)	DOUGH DEVELOPMENT STAGE: As by next month paddy will be harvested, store house should be kept ready by cleaning up of old grain and grain debris and proper sanitation should be maintained Aha moinah para to dhan katibo shoruh hobo, to dhan rakhiboleh ghor to pal para saba khorilobi

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
PEA(LOCAL)	SOWING STAGE: Before sowing treat the seeds with rhizobium culture Bijon nalagha agotae, rhizobium culture logotae milai khena he lagabi

Horticulture (Varieties)	Horticulture Specific Advisory
CABBAGE(LOCAL)	TRANSPLANTING STAGE: In the present weather, transplant 40-45 days old seedlings, it should be done in the morning or late evening Etia lakha somoi te poli 40-45 days howa to transplant khorilobi
ONION(LOCAL)	SOWING: Before sowing, seeds should be treated with thiram @ 2 g/kg of seed to avoid damage from damping off disease. Bijon to thiram@2g/kg of seed te treat khorikhena hi lagabi, enika hoile damping off bimar para bachibo
KING CHILLI(LOCAL)	MATURITY/ HARVESTING STAGE: Weight loss and damage of fresh ripen chillies can be minimized by packing in perforated plastic bags of 1 kg capacity and further storing it in bamboo baskets or cartons Mircha pakia khan laka ujon aru pojicha para bacha bo karoneh , mircha khan to maika/ bag chenta thakia aru etu ke bamboo laka basket de nahoiley carton te rakely bhal hobo
COLOCASIA(LOCAL)	HARVESTING STAGE: Leaves which have turned completely brown and have reduced or shirk can be harvested Patha khan brown, sukha hoilhe harvest khoribo pareh
LARGE CARDAMOM(LOCAL)	POST HARVEST: Fruits should be properly dried in the sun Bijon khoti khan dho harvest khoria biche te dhob te pal bara sukhabi.

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIGGERY(LOCAL)	Contact nearby veterinarian Officer for measure against adverse conditions Apni khan parishe koilhe Veterinarian Officer lakha number to rakhibi, enikha hoilhe gauri lagha kiba dhok khan hoilhe jolti matibo baribo

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
FISHERY (LOCAL)	Liming during winter months is essential even if the pH is within the desired range. This is because fish are susceptible to diseases and lime act as a prophylactic agent Etia thanda moinah ahiase, to pukhuri dhe pH dho dhik ase koilebi liming to khoriboleh lakhe kalhekoileh mass khan to bimar hobo bishe easy ase aru apni liming khoria kharoneh bimar para bachhai

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
POULTRY(LOCAL)	Vaccination should be started for prevalent disease like Ranikhet disease and deworm after every two months for backyard and intensive system both. Ranikhet bimar karoneh Vaccination dho shuru khorianai khoileh, ajikali shurukhoribi aru parishekoileh dui moinah bichhedeh kichu dawai to dibo lageh



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 22-10-2019

Weather Forecast of KOHIMA(Nagaland) Issued On : 2019-10-22(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-10-23	0	32	19	94	55	4.3	132	1
2019-10-24	5	31	20	97	63	4.4	120	5
2019-10-25	5	30	18	98	62	3.7	87	7
2019-10-26	5	26	17	98	80	4.9	62	8
2019-10-27	16	26	18	99	97	3.5	120	8

Weather Summary/Alert:

Probability of light rain the coming days too

Pani etia ahiboleh din bi kiriboleh dikaiase

General Advisory:

Harvesting of ginger and turmeric should be done. Weeding and earthing up should be carried out

Adua, haldi khan harvest khoribo pareh. Pakan te habi khan pila biche te mati dho utabi

SMS Advisory:

To control the incidence of sucking pest in the vegetable crops farmer are advised to spray neem seed kernal extract 5 percent

Sobji te sucking puka komti koribo karoneh neem seed kernal extract 5 percent alibi

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
JHUM PADDY(LOCAL)	STORAGE: Storage of paddy should be such that the rice is kept safe against humidity, moisture, insects, rats, birds, and other contaminants Dhan rakhiya jakha to sukha hobo lakhe, juha aru jiriya para diktar nidiya jakha te he rakhibi
TRC OR WRC(LOCAL)	DOUGH DEVELOPMENT STAGE: For bird control erect scarecrows randomly in the field, scare the bird manually too. Tie old VHS tapes diagonally across and around the field. Use catapults Kheti tae chiria khetaboleh kapra para manu nishinah bonaighenah jaga jaga tae rakhibi. (VHS tapes) color maika kheti bitor aru kinar tae cholabi. Gitis cholabi
FOXTAIL MILLET(LOCAL)	HARVESTING STAGE : In the present weather, start harvesting those which have been ripened Etia lagha somoi te bijon bakia khan to khati bo leh shoru khoribi
SOYBEAN(LOCAL)	HARVESTING STAGE: In the present weather , harvesting should be continued before the pods are shattered

Crop(Varieties)	Crop Specific Advisory
	Etia lakha somoi te akhoni lakha khoti dho nabagia agotae harvest khorilobi

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
CABBAGE(LOCAL)	TRANSPLANTING STAGE: In the present weather, transplant 40-45 days old seedlings, it should be done in the morning or late evening. etia lakha somoi te poli 40-45 days howa to transplant khorilobi
BANANA(LOCAL)	MATURITY STAGE: To reduce the population of fruit flies, collect and destroy the fallen, damaged and overripe fruits Fruit fly lakha population kumti koriboh karoneh, ghol baki khenakhheria khan dho utai khenah jholai dhebi nahoileh mati ghonti khenah alibi.
GINGER(LOCAL)	MATURITY STAGE: For vegetable purpose, harvesting can be carried out Sobji te hali khenah kabo kharoneh to etia para harvest khoribo pareh

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIGGERY(LOCAL)	Vaccination against Foot and Mouth disease in pig should be done. Prior to vaccination, deworm all your animals with appropriate anti-helmentric drugs one week before vaccinating your animals Mukh aro dhang laka bimari para paja boley karone vaccine doh mari debi. Vaccine namara ake tey ek hapta ake tey janwar han ke kichu laga dawai di debi anti-helmentric drugs para

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
FISHERY(LOCAL)	Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water. Take the advice of experts and information from mass media Mass khan ke daana khila time tey pond doh bhal para care koribo lagey, pond tey pani doh bhorta thaki bo lagey. Advice doh kuriboley jana manu han lo lui thakibi aro media han tey para khobor bi lui thakibi

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
POULTRY(LOCAL)	Prevailing high humidity may cause ammonia fumes which irritate the respiratory tract and eyes of the poultry. Good ventilation provides a comfortable environment without draft Kitya murki laga ghor tey humidity doh bishi hubo janwar khan doh sas lobeley dukh pai aro suku bi piya hui jai etu karone hawa jabley karone ventilation bhal aro ghor laga environment doh theek hubo lagey.



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 22-10-2019

Weather Forecast of LONGLENG(Nagaland) Issued On : 2019-10-22(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-10-23	2	31	18	93	56	4.6	138	1
2019-10-24	1	31	19	97	58	4.4	119	6
2019-10-25	1	29	18	94	70	3.2	66	8
2019-10-26	8	25	16	96	74	5.4	63	8
2019-10-27	10	26	17	99	96	4.2	85	8

Weather Summary/Alert:

Probability of light rainfall the coming days

Pani etia ahiboleh din bi kiriboleh dikaiase

General Advisory:

Harvesting of ginger and turmeric should be done. Weeding and earthing up should be carried out.

Adua, haldi khan harvest khoribo pareh. Pakan te habi khan pila biche te mati dho utabi

SMS Advisory:

To control the incidence of sucking pest in the vegetable crops farmer are advised to spray neem seed kernal extract 5 percent

Sobji te sucking puka komti koribo karoneh neem seed kernal extract 5 percent alibi

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
JHUM PADDY(LOCAL)	POST HARVEST: Paddy which have been harvested should be dried properly and keep in the storage area which have been cleaned properly
	Dhan khan katia khan to bhal para sukai khena rakhibi. Thailaka sukhai khena rakha jaka to pal bara sabha khoribi etu bichhe te hi rakhibi
TRC OR WRC(LOCAL)	MATURITY STAGE: As by next month paddy will be harvested, store house should be kept ready by cleaning up of old grain and grain debris and proper sanitation should be maintained
	Aha moinah para to dhan katibo shoruh hobo, to dhan rakhiboleh ghor to pal para saba khorilobi
SOYBEAN(LOCAL)	HARVESTING STAGE: In the present weather , harvesting should be done before the pods are shattered
	Etia lakha somoi te akhoni lakha khoti dho nabagia agotae harvest khorilobi

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
BRINJAL(LOCAL)	HARVESTING STAGE: Timely harvesting should be done
	Dhik somoi te bijon ulabi
CHILLI(LOCAL)	HARVESTING STAGE: Timely harvesting should be carried out to prevent loss
	Thik somoi te mircha to olabo lakhe
COLOCASIA(LOCAL)	HARVESTING STAGE: Leaves which have turned completely brown and have reduced or shirk can be harvested
	Patha khan brown, sukha hoilhe harvest khoribo pareh
GINGER(LOCAL)	MATURITY STAGE: For vegetable purpose , harvesting can be carried out
	Sobji te hali khena kabo kharoneh to etia para harvest khoribo pareh

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIGGERY(LOCAL)	Vaccination against Foot and Mouth disease in pig should be done. Prior to vaccination, deworm all your animals with appropriate anti-helmentric drugs one week before vaccinating your animals
	Mukh aro dhang laka bimari para paja boley karone vaccine doh mari debi. Vaccine namara ake tey ek hapta ake tey janwar han ke kichu laga dawai di debi anti-helmentric drugs para.

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
FISHERY(LOCAL)	Remove pond bottom debris before the onset of winter preferably by this month
	Apni lakha pukhuri do etu moinah dho sapa khoriboleh laghe

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
POULTRY(LOCAL)	Vaccination should be started for prevalent disease like Ranikhet disease and deworm after every two months for backyard and intensive system both.
	Ranikhet bimar karoneh Vaccination dho shuru khorianai khoileh, ajikali shurukhoribi aru parishekoileh dui moinah bichhedeh kichu dawai to dibo lageh



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 22-10-2019

Weather Forecast of MOKOKCHUNG(Nagaland) Issued On : 2019-10-22(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-10-23	4	31	18	95	58	4.6	134	1
2019-10-24	1	31	19	97	58	4.4	123	6
2019-10-25	3	29	18	97	68	3.2	62	8
2019-10-26	14	26	17	97	74	5.4	61	8
2019-10-27	17	26	17	99	97	4.2	84	8

Weather Summary/Alert:

Probability of light rainfall the coming days
Pani etia ahiboleh hapta bi olop kiribolhe dikaiase

General Advisory:

Harvesting of ginger and turmeric should be done. Weeding and earthing up should be carried out
Adua, haldi khan harvest khoribo pareh. Pakan te habi khan pila biche te mati dho utabi

SMS Advisory:

To control the incidence of sucking pest in the vegetable crops farmer are advised to spray neem seed kernal extract 5 percent
Sobji te sucking puka komti koribo karoneh neem seed kernal extract 5 percent alibi

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
JHUM PADDY(LOCAL)	POST HARVEST: Paddy which have been harvested should be dried properly and keep in the storage area which have been cleaned properly
	Dhan khan katia khan to bhal para sukai khena rakhibi. Thailaka sukhai khena rakha jaka to pal bara sabha khoribi etu bichhe te hi rakhibi
TRC OR WRC(LOCAL)	MATURITY STAGE: As by next month paddy will be harvested, store house should be kept ready by cleaning up of old grain and grain debris and proper sanitation should be maintained
	Aha moinah para to dhan katibo shoruh hobo, to dhan rakhiboleh ghor to pal para saba khorilobi
SOYBEAN(LOCAL)	HARVESTING STAGE: In the present weather , harvesting should be done before the pods are shattered
	Etia lakha somoi te akhoni lakha khoti dho nabagia agotae harvest khorilobi

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
BRINJAL(LOCAL)	HARVESTING STAGE: Timely harvesting should be done
	Dhik somoi te bijon ulabi
CHILLI(LOCAL)	HARVESTING STAGE: Timely harvesting should be carried out to prevent loss
	Thik somoi te mircha to olabo lakhe
GINGER(LOCAL)	MATURITY STAGE: For vegetable purpose , harvesting can be carried out
	Sobji te hali khena kabo kharoneh to etia para harvest khoribo pareh
COLOCASIA(LOCAL)	HARVESTING STAGE: Leaves which have turned completely brown and have reduced or shirk can be harvested
	Patha khan brown, sukha hoilhe harvest khoribo pareh
TAPIOCA(LOCAL)	VEGETATIVE STAGE: Install sticky cum light trap TO REDUCE WHITE FLY POPULATION
	Whitefly lagha population komti khoribo karoneh sticky cum light trap dho kheti te jakha jakha te rakhibi

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIGGERY(LOCAL)	Vaccination against Foot and Mouth disease in pig should be done. Prior to vaccination, deworm all your animals with appropriate anti-helmentric drugs one week before vaccinating your animals
	Mukh aro dhang laka bimari para paja boley karone vaccine doh mari debi. Vaccine namara ake tey ek hapta ake tey janwar han ke kichu laga dawai di debi anti-helmentric drugs para.

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
FISHERY(LOCAL)	Remove pond bottom debris before the onset of winter preferably by October
	Apni lakha pukhuri do etu moinah dho sapa khoriboleh laghe

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
POULTRY(LOCAL)	Vaccination should be started for prevalent disease like Ranikhet disease and deworm after every two months for backyard and intensive system both
	Ranikhet bimar karoneh Vaccination dho shuru khorianai khoileh, ajikali shurukhoribi aru parishekoileh dui moinah bichhedeh kichu dawai to dibo lageh



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 22-10-2019

Weather Forecast of MON(Nagaland) Issued On : 2019-10-22(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-10-23	2	31	18	93	61	4.9	145	2
2019-10-24	4	31	20	95	60	4.3	111	6
2019-10-25	6	29	18	92	73	3.6	88	8
2019-10-26	8	26	16	97	74	4.8	85	8
2019-10-27	12	27	17	99	96	3.8	109	8

Weather Summary/Alert:

Probability of light rainfall the coming days

Etia ahibleh din bi pani kiriboleh dikaiase

General Advisory:

Harvesting of ginger and turmeric should be done. Weeding and earthing up should be carried out

Adua, haldi khan harvest khoribo pareh. Pakan te habi khan pila biche te mati dho utabi

SMS Advisory:

To control the incidence of sucking pest in the vegetable crops farmer are advised to spray neem seed kernal extract 5 percent

Sobji te sucking puka komti koribo karoneh neem seed kernal extract 5 percent alibi

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
JHUM PADDY(LOCAL)	POST HARVEST: Paddy which have been harvested should be dried properly and keep in the storage area which have been cleaned properly
	Dhan khan katia khan to bhal para sukai khena rakhibi. Thailaka sukhai khena rakha jaka to pal bara sabha khoribi etu bichhe te hi rakhibi
TRC PADDY(LOCAL)	MATURITY STAGE: As the paddy is to be harvested the coming days, storage structure should be cleaned properly. Take precautions to avoid infestation by the the stored grain insects and the fungal infections
	Etia ahibleh din te dhan katiboleh ase, dho etia para dai lakha rakha jaka to pal para sabha khoribi
GROUNDNUT(LOCAL)	HARVESTING STAGE: In the present weather, harvesting should be carried out
	etia lakha somoi te , khoti to harvest khoribi

Crop(Varieties)	Crop Specific Advisory
TORIA(LOCAL)	Field preparation should be started for toria and arrangement for quality seeds should be done. M-27, TS-36 and TS-38 are some of the varieties recommended for our region
	Toria lakha Bijon laghabo lhe shoru khoribi. M-27, TS-36 and TS-38 bijon to moikan lagha jakha karoneh pal ase
SOYBEAN(LOCAL)	HARVESTING STAGE: Completely cover plant residue by clean ploughing the field soon after harvest.
	Harvest khorabiche te, ghas to mati te dahlibi

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
POTATO(LOCAL)	SOWING STAGE: The seed to be sown should be free a disease free area to ensure that the pathogen is not carried through seed tuber
	Etia bijo lakha bo lhe somoi ase. Bijon dho bimar nathakha jakha para rakha khan to heh lagabi
PEA(LOCAL)	SOWING STAGE: Before sowing treat the seeds with rhizobium culture
	Bijon nalagha agotae, rhizobium culture logotae milai khena he lagabi
GARLIC(LOCAL)	SOWING STAGE: Sowing should be done in a well drained soil. IT should be done at 15 x 10cm . Before planting treat the cloves with Carbendazim 2 g/kg to prevent clove rot
	Mati dho pal bara pani ja lakha te he lagabi. Spacing to 15 x 10cm maintain khoribi
ONION(LOCAL)	SOWING: Before sowing, seeds should be treated with thiram @ 2 g/kg of seed to avoid damage from damping off disease.
	Bijon to thiram@2g/kg of seed te treat khorikhena hi lagabi, enika hoile damping off bimar para bachibo
CARDAMON(LOCAL)	MATURITY/HARVESTING STAGE: Matured fruits should be harvested without delay
	khuti pakia khan to dhik somoi te olai dibo laghe
COLOCASIA(LOCAL)	HARVESTING STAGE: Leaves which have turned completely brown and have reduced or shirk can be harvested
	Patha khan brown, sukha hoilhe harvest khoribo pareh

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIGGERY(LOCAL)	Sick or injured animals should be attended to immediately, preferably moved to a Comfortable environment
	Gauri bimar khan dho alak bareh rakhibo lakhe

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
FISHERY(LOCAL)	During extreme weather, frequent inspections and monitoring should be done as it helps identify consequences
	Bishi hawa pani biya din khan dhe, mass khan kinikha ase saidhakibolhe lakhe

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
POULTRY(LOCAL)	Vaccination should be started for prevalent disease like Ranikhet disease and deworm after every two months for backyard and intensive system both
	Ranikhet bimar karoneh Vaccination dho shuru khorianai khoileh, ajikali shurukhoribi aru parishekoileh dui moinah bichhedeh kichu dawai to dibo lageh



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 22-10-2019

Weather Forecast of PEREN(Nagaland) Issued On : 2019-10-22(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-10-23	1	32	21	96	50	3.8	104	2
2019-10-24	1	32	21	97	50	3.6	60	4
2019-10-25	3	31	21	99	50	3.6	20	7
2019-10-26	5	26	20	99	80	5.3	192	8
2019-10-27	21	26	20	100	96	2.6	94	8

Weather Summary/Alert:

Probability of light rainfall the coming days
Etia ahiboleh din bi pani olop kiriboleh dikaiase

General Advisory:

Harvesting of ginger and turmeric should be done. Weeding and earthing up should be carried out
Adua, haldi khan harvest khoribo pareh. Pakan te habi khan pila biche te mati dho utabi

SMS Advisory:

To control the incidence of sucking pest in the vegetable crops farmer are advised to spray neem seed kernal extract 5 percent
Sobji te sucking puka komti koribo karoneh neem seed kernal extract 5 percent alibi

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
JHUM PADDY(LOCAL)	POST HARVEST: Paddy which have been harvested should be dried properly and keep in the storage area which have been cleaned properly
	Dhan khan katia khan to bhal para sukai khena rakhibi. Thailaka sukhai khena rakha jaka to pal bara sabha khoribi etu bichhe te hi rakhibi
TRC OR WRC(LOCAL)	MILK DEVELOPMENT STAGE: From heading till harvesting, inspect field, remove and destroy off type plants and weeds.
	Dhan to nagata dhak, somoi te sai dhakibolageh ki ghas te bimar pukha khan ase nah nai, jongol ghas aru habi khan bi somoi te ulai dibo lageh
GROUNDNUT(LOCAL)	HARVESTING STAGE: After harvest, dry the pods in the sun for 4 or 5 days. Repeat drying for 2 or 3 more days after an interval of 2 or 3 days to ensure complete drying
	kuti bakia bicche te, doob te baal para sukhabi char/pass din karoneh.
SOYBEAN(LOCAL)	HARVESTING STAGE: In the present weather harvesting should be carried out. Threshing should be done carefully to prevent damage of seed coat

Crop(Varieties)	Crop Specific Advisory
	Etia lakha somoi te harvesting to shoru khorilobo laghe.

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
COLOCASIA(LOCAL)	HARVESTING STAGE: Leaves which have turned completely brown and have reduced or shirk can be harvested Patha khan brown, sukha hoilhe harvest khoribo pareh
NAGA CHILLI(LOCAL)	HARVESTING STAGE: Timely harvesting should be carried out to prevent loss Thik somoi te mircha to olabo lakhe

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIGGERY(LOCAL)	Contact nearby veterinarian Officer for measure against adverse conditions Apni khan parishe koilhe Veterinarian Officer lakha number to rakhibi, enikha hoilhe gauri lagha kiba dhok khan hoilhe jolti matibo baribo

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
FISHERY(LOCAL)	Remove pond bottom debris before the onset of winter preferably by October Apni lakha pukhuri do etu moinah dho sapa khoriboleh laghe

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
POULTRY(LOCAL)	Vaccination should be started for prevalent disease like Ranikhet disease and deworm after every two months for backyard and intensive system both. Ranikhet bimar karoneh Vaccination dho shuru khorianai khoileh, ajikali shurukhoribi aru parishekoileh dui moinah bichhedeh kichu dawai to dibo lageh



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 22-10-2019

Weather Forecast of TUENSANG(Nagaland) Issued On : 2019-10-22(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-10-23	0	31	18	86	43	3.9	114	1
2019-10-24	1	32	20	90	35	2.9	90	5
2019-10-25	2	30	19	93	44	2.7	61	7
2019-10-26	2	29	19	95	54	3.3	89	8
2019-10-27	12	27	19	98	88	2.8	121	8

Weather Summary/Alert:

Probability of light rainfall the coming days

Etia ahiboleh din bi pani kiriboleh dikaiase

General Advisory:

Harvesting of ginger and turmeric should be done. Weeding and earthing up should be carried out.

Adua, haldi khan harvest khoribo pareh. Pakan te habi khan pila biche te mati dho utabi

SMS Advisory:

To control the incidence of sucking pest in the vegetable crops farmer are advised to spray neem seed kernal extract 5 percent

Sobji te sucking puka komti koribo karoneh neem seed kernal extract 5 percent alibi

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
JHUM PADDY(LOCAL)	STORAGE: Care should be taken to store different varieties of paddy separately Sabolhe lakhe ki dhan variety hisam te alak alak rakhibi
TRC OR WRC(LOCAL)	DOUGH DEVELOPMENT STAGE: As by next month paddy will be harvested, store house should be kept ready by cleaning up of old grain and grain debris and proper sanitation should be maintained Aha moinah para to dhan katibo shoruh hobo, to dhan rakhiboleh ghor to pal para saba khorilobi
MUSTARD(LOCAL)	SOWING: Sowing should be done in fine field land preparation for good germination Khoti dho mati dhik para bona jaka te hi dhalibi

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
COLOCASIA(LOCAL)	HARVESTING STAGE: Leaves which have turned completely brown and have reduced or shirk can be harvested Patha khan brown, sukha hoilhe harvest khoribo pareh
KING CHILLI(LOCAL)	MATURITY/ HARVESTING STAGE: Weight loss and damage of fresh ripen chillies can be minimized by packing in perforated plastic bags of 1 kg capacity and further storing it in bamboo baskets or cartons. Mircha pakia khan laka ujon aru pojicha para bacha bo karoneh , mircha khan to maika/ bag chenta thakia aru etu ke bamboo laka basket de nahoiley carton te rakely bhal hobo

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIGGERY(LOCAL)	Contact nearby veterinarian Officer for measure against adverse conditions Apni khan parishe koilhe Veterinarian Officer lakha number to rakhibi, enikha hoilhe gauri lagha kiba dhok khan hoilhe jolti matibo baribo

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
FISHERY(LOCAL)	Liming during winter months is essential even if the pH is within the desired range. This is because fish are susceptible to diseases and lime act as a prophylactic agent. Etia thanda moinah ahiase, to pukhuri dhe pH dho dhik ase koilebi liming to khoriboleh lakhe kalhekoileh mass khan to bimar hobo bishe easy ase aru apni liming khoria kharoneh bimar para bachhai

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
POULTRY(LOCAL)	Vaccination should be started for prevalent disease like Ranikhet disease and deworm after every two months for backyard and intensive system both. Ranikhet bimar karoneh Vaccination dho shuru khorianai khoileh, ajikali shurukhoribi aru parishekoileh dui moinah bichhedeh kichu dawai to dibo lageh



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Complex Branch

Umiam, Meghalaya

(Project Sponsored by IMD, Ministry of Earth Sciences, Govt. of India)



Agromet Advisory Bulletin

Date : 22-10-2019

Weather Forecast of WOKHO(Nagaland) Issued On : 2019-10-22(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2019-10-23	4	32	19	93	58	4.5	107	1
2019-10-24	4	31	20	95	64	5.2	112	5
2019-10-25	5	29	19	97	76	3.8	102	8
2019-10-26	13	26	17	97	81	6.7	31	8
2019-10-27	18	25	17	99	97	5.6	51	8

Weather Summary/Alert:

Probability of light rainfall the coming days
Etia ahiboleh din bi pani kiriboleh dikaiase

General Advisory:

Harvesting of ginger and turmeric should be done. Weeding and earthing up should be carried out
Adua, haldi khan harvest khoribo pareh. Pakan te habi khan pila biche te mati dho utabi

SMS Advisory:

To control the incidence of sucking pest in the vegetable crops farmer are advised to spray neem seed kernal extract 5 percent
Sobji te sucking puka komti koribo karoneh neem seed kernal extract 5 percent alibi

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
JHUM PADDY(LOCAL)	POST HARVEST: Paddy which have been harvested should be dried properly and keep in the storage area which have been cleaned properly
	Dhan khan katia khan to bhal para sukai khena rakhibi. Thailaka sukhai khena rakha jaka to pal bara sabha khoribi etu bichhe te hi rakhibi
TRC OR WRC(LOCAL)	MATURITY STAGE: As by next month paddy will be harvested, store house should be kept ready by cleaning up of old grain and grain debris and proper sanitation should be maintained
	Aha moinah para to dhan katibo shoruh hobo, to dhan rakhiboleh ghor to pal para saba khorilobi
SOYBEAN(LOCAL)	HARVESTING STAGE: In the present weather , harvesting should be done before the pods are shattered
	Etia lakha somoi te akhoni lakha khoti dho nabagia agotae harvest khorilobi

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
BRINJAL(LOCAL)	HARVESTING STAGE: Timely harvesting should be done
	Dhik somoi te bijon ulabi
CHILLI(LOCAL)	HARVESTING STAGE: Timely harvesting should be carried out to prevent loss
	Thik somoi te mircha to olabo lakhe
GINGER(LOCAL)	MATURITY STAGE: For vegetable purpose , harvesting can be carried out
	Sobji te hali khena kabo kharoneh to etia para harvest khoribo pareh
COLOCASIA(LOCAL)	HARVESTING STAGE: Leaves which have turned completely brown and have reduced or shirk can be harvested
	Patha khan brown, sukha hoilhe harvest khoribo pareh

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIGGERY(LOCAL)	Vaccination against Foot and Mouth disease in pig should be done. Prior to vaccination, deworm all your animals with appropriate anti-helmentric drugs one week before vaccinating your animals
	Mukh aro dhang laka bimari para paja boley karone vaccine doh mari debi. Vaccine namara ake tey ek hapta ake tey janwar han ke kichu laga dawai di debi anti-helmentric drugs para.

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
FISHERY(LOCAL)	Remove pond bottom debris before the onset of winter preferably by October
	Apni lakha pukhuri do etu moinah dho sapa khoriboleh laghe

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
POULTRY(LOCAL)	Vaccination should be started for prevalent disease like Ranikhet disease and deworm after every two months for backyard and intensive system both.
	Ranikhet bimar karoneh Vaccination dho shuru khorianai khoileh, ajikali shurukhoribi aru parishekoileh dui moinah bichhedeh kichu dawai to dibo lageh