




# STATE COMPOSITE AGRO-MET ADVISORY BULLETIN FOR THE STATE OF SIKKIM PERIOD: 12<sup>th</sup> AUGUST – 14<sup>th</sup> AUGUST 2014



*IMD, GANGTOK IN COLLABORATION WITH ICAR, GANGTOK, FSAD AND HCCD, GANGTOK*



**ISSUED ON TUESDAY 12<sup>th</sup> AUGUST 2014 BY**

## **Meteorological Sub-division of Sikkim**

As per India Meteorological Department's classification, Sikkim is a part of Met Sub-Division of Sub Himalayan West Bengal & Sikkim (SHWB & SKM) and is divided into four (4) Districts:

- 1) North District with HQ at Mangan
- 2) East District with HQ at Gangtok
- 3) West District with HQ at Gyalsing
- 4) South District with HQ at Namchi

# Part I: Weather

**Significant Past Weather in different districts during the period 08/8/14 TO 11/8/14**

**Weather Forecast for different districts for the period 12/8/14 to 16/8/14**

## North Sikkim

Mangan (AWS)				North Sikkim
Date	Rainfall (mm)	Max. Temp (°C)	Min. Temp (°C)	<ul style="list-style-type: none"> <li>Generally Cloudy sky during the period 12<sup>th</sup> August to 16<sup>th</sup> August is expected.</li> <li>Moderate to rather heavy rain is expected during the period 12<sup>th</sup> August to 16<sup>th</sup> August.</li> <li>Wind from southeasterly to south-south-westerly with an average speed of 1-2 km/hr is expected.</li> <li>Maximum and minimum temperatures are expected to be around 27 – 29 °C and 20 °C respectively.</li> <li>Maximum and minimum relative humidity is expected to be 92 - 98 % and 77 - 87 % respectively.</li> </ul>
08-08-2014	002.0	30.7	20.2	
09-08-2014	012.0	30.4	20.7	
10-08-2014	013.0	29.7	20.5	
11-08-2014	005.0	28.8	20.2	

## East Sikkim

Gangtok				East Sikkim
Date	Rainfall (mm)	Max. Temp (°C)	Min. Temp (°C)	<ul style="list-style-type: none"> <li>Generally Cloudy sky during the period 12<sup>th</sup> August to 16<sup>th</sup> August is expected.</li> <li>Moderate to rather heavy rain is expected during the period 12<sup>th</sup> August to 16<sup>th</sup> August.</li> <li>Wind from east-south-easterly to south southwesterly with an average speed of 1-2 km/hr is expected.</li> <li>Maximum and minimum temperatures are expected to be around 21 – 23 °C and 18 °C respectively.</li> <li>Maximum and minimum relative humidity is expected to be 95 - 98 % and 82 – 88 % respectively.</li> </ul>
08-08-2014	023.0	023.4	018.2	
09-08-2014	011.8	023.1	018.7	
10-08-2014	012.2	021.7	018.7	
11-08-2014	013.3	023.0	018.5	




Significant Past Weather in different districts during the period 08/8/14 TO 11/8/14				Weather Forecast for different districts for the period 12/8/14 to 16/8/14	
South Sikkim					
Namthang				South Sikkim	
Date	Rainfall (mm)	Max. Temp (°C)	Min. Temp (°C)	<div><div></div> Generally Cloudy sky during the period 12<sup>th</sup> August to 16<sup>th</sup> August is expected.</div> <div><div></div> Moderate to rather heavy rain is expected during the period 12<sup>th</sup> August to 16<sup>th</sup> August.</div> <div><div></div> Wind from easterly to southwesterly with an average speed of 2-4 km/hr is expected.</div> <div><div></div> Maximum and minimum temperatures are expected to be around 24 – 25 °C and18 - 19 °C respectively.</div> <div><div></div> Maximum and minimum relative humidity is expected to be 92 - 98 % and 80 – 87 % respectively.</div>	
08-08-2014	000.0	25	19.6		
09-08-2014	003.0	24.2	20.2		
10-08-2014	006.0	23.1	19.8		
11-08-2014	013.5	22.9	18		
West Sikkim					
Gyalshing				West Sikkim	
Date	Rainfall (mm)	Max. Temp (°C)	Min. Temp (°C)	<div><div></div> Generally Cloudy sky during the period 12<sup>th</sup> August to 16<sup>th</sup> August is expected.</div> <div><div></div> Moderate to rather heavy rain is expected during the period 12<sup>th</sup> August to 16<sup>th</sup> August.</div> <div><div></div> Wind from easterly to southwesterly with an average speed of 1-3 km/hr is expected.</div> <div><div></div> Maximum and minimum temperatures are expected to be around 23 - 25 °C and 18 °C respectively.</div> <div><div></div> Maximum and minimum relative humidity is expected to be 95 - 98 % and 78 – 87 % respectively.</div>	
08-08-2014	004.0	017.2	027.4		
09-08-2014	003.0	018.0	025.5		
10-08-2014	017.0	018.1	024.8		
11-08-2014	018.0	018.2	025.3		





## **Part II**

### **AGRO-METEREOLOGICAL ADVISORIES**





#### **General Advisories for all the crops**



- ◆ Proper drainage to be maintained in pulse/beans/ginger/turmeric/Dalley chlli.
- ◆ Avoid water stagnation in guava/ mandarin crop.
- ◆ Avoid top dressing of organic manure.
- ◆ Preventive spray and soil drenching with copperoxychloride@2.5 g/l of water immediate after rain subside.
- ◆ Spray sulphur powder@ 2g/l in case of powdery mildew disease.
- ◆ Avoid spraying of plant protection biopesticides during rain and morning Hours.

Name of the crop/Animal	Stage	Pest/ Disease	Agro-Met Advisories
<b>Crops</b>			
Rice	Seedling	Leaf folder	<ul style="list-style-type: none"> <li>✿ Weeding and thinning should be done.</li> <li>✿ Maintain 2-3 cm water in the field.</li> <li>✿ Spray neem based formulation (1500ppm) @ 3ml per litre of water for controlling leaf folder.</li> <li>✿ Monitor regularly and unfold the leaves folded by the leaf folder; and manually collect and destroy.</li> <li>✿ Spray copper oxychloride @2.5 g/l for rice blast.</li> </ul>
		 Blast of rice 	
Soybean	Flowering	Blister beetle 	<ul style="list-style-type: none"> <li>✿ Proper drainage should be maintained.</li> <li>✿ Spray neem based formulation (1500ppm) @ 3ml per litre of water for controlling leaf roller.</li> <li>✿ Monitor regularly and unfold the leaves folded by the leaf roller. Manually collect and destroy both the</li> </ul>

		<p>Leaf roller damage</p> 	<p>insects.</p>
<p>Pahenlo dal /Kalo Dal (<i>urd</i> bean)</p>	<p>vegetative</p>		<ul style="list-style-type: none"> <li>✳ Proper drainage should be maintained</li> <li>✳ Weeding and thinning should be done.</li> <li>✳ Spray neem based formulation (1500 ppm) @ 3ml per litre of water for controlling leaf feeding beetles.</li> </ul>
<p>Finger millet</p>	<p>Vegetative</p>		<ul style="list-style-type: none"> <li>✳ Transplanting and gap filling should be completed as earliest. Maintain optimum plant population for good yield.</li> </ul>
<p>Maize</p>	<p>Harvesting</p>		<ul style="list-style-type: none"> <li>✳ Harvest mature cobs and sun dry. After proper drying, store grains/seeds in well ventilated godown.</li> </ul>
<p>Beans</p>	<p>Vegetative/ Flowering</p>	<ul style="list-style-type: none"> <li>• Blister beetle may occur and destroy flowers.</li> </ul>  <ul style="list-style-type: none"> <li>• Powdery mildew</li> </ul>  <ul style="list-style-type: none"> <li>• Damping off</li> </ul> 	<ul style="list-style-type: none"> <li>✳ Handpicking is effective to control this pest.</li> <li>✳ Collection and destruction of initial infected plant parts.</li> <li>✳ Spray out sulphur based fungicide @ 2 g/l to control powdery mildew.</li> <li>✳ Proper drainage should be maintained</li> <li>✳ Seed treatment with <i>Trichoderma viride</i> culture @ 5g/kg of seed prior to re-sowing for gap filling.</li> <li>✳ Soil drenching with copper oxychloride @2.5 g/l for damping off in infected area.</li> </ul>



Large cardamom	Capsule formation	Infestation of shoot flies and stem borer.	<ul style="list-style-type: none"> <li>Collection and destruction of infested tillers. Spraying of neem formulation (1500 ppm) @ 3 ml/l.</li> <li>Keep cardamom fields weed free.</li> <li>Collection and destruction of disease infected plants/ tillers.</li> </ul>
Cole crops	--		<ul style="list-style-type: none"> <li>Time for nursery raising of cabbage, cauliflower and broccoli under low cost structure.</li> <li>Soil drenching of nursery bed with copper oxychloride @5 g/ l to prevent damping off.</li> <li>Prepare 15cm raised bed nursery of 1m width.</li> <li>Well-decomposed FYM @ 2.5 kg/sqm to be applied during nursery bed preparation.</li> </ul>
Dalley Chilli	Flowering and Fruiting	Fruit fly damage 	<ul style="list-style-type: none"> <li>Field should be cleaned and weed free.</li> <li>Proper drainage should be maintained.</li> <li>Fallen fruits should be collected and destroyed to minimize fruit fly.</li> </ul>
Sikkim Mandarin	Fruit growth	Aphid  Leaf miner  Mealy bug 	<ul style="list-style-type: none"> <li>Proper drainage should be maintained.</li> <li>Basin of plants should be cleaned.</li> <li>Spray petroleum oil based agro-spray @ 10 ml/l for managing aphid, leaf miner, mealy bug etc.</li> <li>Regular monitoring and cleaning of infested trunk. Plug the holes with cotton soaked in kerosene/petrol and plaster with mixture of cow dung and soil.</li> <li>Spray copper oxychloride @ 2.5 g/ l to manage citrus scab.</li> </ul>

		<p>Trunk borer</p>  <p>Scab</p> 	
--	--	--	--

### ***Animal Sciences***

Tibetan Sheep	Full grown wool	Chances of soiling and picking infection.	✿ Go for shearing of you Sheep followed by spraying or Dipping in ectoparasitic solution.
Yak, Tibetan Sheep, Goat Cattle	High humidity and Moderate temperature	Favorable condition for ectoparasitic infection.	✿ Go for deworming with fenbedazole @10mg/kg.
Cattle, Sheep and Goat	Comfortable temperature	Plenty of green posture/ fodder available in the field.	✿ Chances of coming into estrus so farmers are advised to look for heat sign especially during early morning or late evening and ensure breeding of the animal after 12 hr of 1 <sup>st</sup> heat observe.