




# AGRO-MET ADVISORY BULLETIN FOR THE DISTRICT OF NORTH SIKKIM

## PERIOD:

**09<sup>th</sup> NOVEMBER to 13<sup>th</sup> NOVEMBER' 2019**

*IMD, GANGTOK IN COLLABORATION WITH ICAR, GANGTOK, FSAD  
AND HCCD, GANGTOK*



**ISSUED ON 8<sup>th</sup> November 2019**

### Meteorological Sub-division of Sikkim

As per India Meteorological Department's classification, Sikkim is a part of Met Sub-Division of Sub Himalayan West Bengal & Sikkim (SHWB & SKM) and is divided into four (4) Districts:

- 1) North District with HQ at Mangan
- 2) East District with HQ at Gangtok
- 3) West District with HQ at Gyalsing
- 4) South District with HQ at Namchi

Issued by: Santanu Das, SMS (Agril. Meteorology), ICAR-KVK, East Sikkim, Ranipool

### Value Added forecast for NORTH-DISTRICT for next five days

Parameter/Date	09/11/2019	10/11/2019	11/11/2019	12/11/2019	13/11/2019
Rainfall (mm)	No rainfall	Light rainfall	Light rainfall	No rainfall	No rainfall
Max Temp (°C)	25	23	22	23	24
Min Temp (°C)	14	13	12	12	13
Sky condition	Generally cloudy	Generally cloudy	Generally cloudy	Generally cloudy	Generally cloudy
Max RH (%)	87	91	93	89	85
Min RH (%)	70	75	80	72	70
Wind Speed (Kmph)	003	003	004	003	003
Wind Direction (deg)	SE	SE	SE	SE	S

Northerly- **N**, North-Easterly- **N-E**, Easterly- **E**, South-Easterly- **S-E**, Southerly- **S**, South-Westerly- **S-W**, Westerly- **W**, North-westerly- **N-W**.

Weather for preceding week			
Date	Rainfall	Max. Temp	Min. Temp
	(mm)	(°C)	(°C)
05/11/2019	002	N/A	N/A
06/11/2019	000	23.9	12.0
07/11/2019	000	25.5	13.8

### Weather forecast valid up to 13<sup>th</sup> November 2019


- Probability of no rain.
- Max temp may be varying from 22-25°C, whereas, Min temp may vary from 12-14°C
- Sky will remain generally cloudy.
- Max RH is likely to be 85-93 %, whereas, Min RH may vary from 70-80 %
- Wind speed may likely vary from 3-4 km/hr.
- Wind direction may blow from southeasterly direction.
- Weekly cumulative rainfall= 006.0 mm

## AGRO-METEOREOLOGICAL ADVISORIES

### General Advisories for all the crops/ livestock

- ➡ Complete the sowing of rabi crops as early as possible before residual soil moisture lost.
- ➡ Harvest mature crops before unexpected rain.
- ➡ While selecting site for construction of animal shed a site that allows sun light exposure as well as good wind control is preferable

Name of the crop/Animal	Stage	Cultural practices/ Pest/ Diseases	Agro-Met Advisories
<b><i>Agricultural and Horticultural crops:</i></b>			
Rice	Grain maturity to harvesting and postharvest		<ul style="list-style-type: none"><li>◆ Drain out water completely 10 days before harvesting.</li><li>◆ Harvesting should start when the crops are 80-90% mature</li><li>◆ Maintain seed moisture about 8-10 % for storage.</li></ul>
Rjmash	Pre flowering/flowering		<ul style="list-style-type: none"><li>◆ Irrigate crop at flowering stage.</li></ul>
Ginger/ Turmeric	Tillering stage		<ul style="list-style-type: none"><li>◆ Allow minimum entry to crop field to reduce disease infestation.</li><li>◆ Allow proper drainage to avoid stagnation of water in the field.</li><li>◆ Timely monitor the field to check the attack of bacterial wilt due to high humidity and temperature.</li></ul>
Red cherry pepper/Brinjal/ Toamto	Fruiting	Blight	<ul style="list-style-type: none"><li>◆ Allow proper drainage to remove excess rain water from the field.</li><li>◆ Timely monitor the field to check the attack of bacterial will due to high humidity and temperature.</li><li>◆ Due to high temperature and high rainfall may cause late blight. Apply COC @ 0.25% at weekly interval for 4-5 sprays will reduce the disease.</li></ul>
Cole crops	Nursery preparation		<ul style="list-style-type: none"><li>◆ The nursery site should be dry and free from water stagnation as well as high moisture in the soil.</li><li>◆ To protect from heavy rain the nursery should be in semi protected area.</li></ul>

Sikkim Mandarin	Planting/fruiting		<ul style="list-style-type: none"> <li>◆ As there is a possibility of light to moderate rainfall so apply soil nutrients @ 25-35 kg/ha as maximum nutrient losses may occur due to rainfall.</li> </ul>
<b><i>Animal Sciences and livestock management</i></b>			
Poultry		Coccidiosis, Ranikhet disease	<ul style="list-style-type: none"> <li>◆ Keep the Vanaraja chicks in warm condition (32-33°C) during the first four weeks.</li> <li>◆ Low cost brooder should be made to prevent chick mortality due to cold stress.</li> <li>◆ Maintain brooding temperature through continuous electricity supply and for emergency Sigiri/Bukhari can be used.</li> </ul>
Sheep/goat			<ul style="list-style-type: none"> <li>◆ During winter months special care should be taken for sheep, goat for external parasites.</li> </ul>

*Agro-Met Advisory member of ICAR-NOFRI and KVK- East Sikkim*

<b>Name</b>	<b>Designation</b>	<b>Department</b>
<b><i>Dr. R. K. Avasthe</i></b>	<b>Joint Director</b>	<b>Soil Science</b>
<i>Dr. Raghavendra Singh</i>	Senior scientist and Head (I/C)	Agronomy
<i>Dr. P. K. Pathak</i>	SMS	Animal science
<i>Dr. J.K. Singh</i>	SMS	Plant Breeding
<i>Dr. Manoj Kumar</i>	SMS	Agronomy
<i>Sh. Boniface Lepcha</i>	SMS	Horticulture
<i>Sh. Santanu Das</i>	SMS	Agril. Meteorology
<i>Miss Tirtha Kumari Chettri</i>	Met. Observer	Biochemistry




# AGRO-MET ADVISORY BULLETIN FOR THE DISTRICT OF EAST SIKKIM

**PERIOD:**

**09<sup>th</sup> NOVEMBER to 13<sup>th</sup> NOVEMBER 2019**

*IMD, GANGTOK IN COLLABORATION WITH ICAR, GANGTOK, FSAD  
AND HCCD, GANGTOK*



**ISSUED ON 08<sup>th</sup> November 2019**

## Meteorological Sub-division of Sikkim

As per India Meteorological Department's classification, Sikkim is a part of Met Sub-Division of Sub Himalayan West Bengal & Sikkim (SHWB & SKM) and is divided into four (4) Districts:

- 1) North District with HQ at Mangan
- 2) East District with HQ at Gangtok
- 3) West District with HQ at Gyalsing
- 4) South District with HQ at Namchi

Issued by: Santanu Das, SMS (Agril. Meteorology), ICAR-KVK, East Sikkim, Ranipool

### Value Added forecast for EAST-DISTRICT for next five days

Parameter/Date	09/11/2019	10/11/2019	11/11/2019	12/11/2019	13/11/2019
Rainfall (mm)					
Max Temp ( $^{\circ}$ C)					
Min Temp ( $^{\circ}$ C)					
Sky condition					
Max RH (%)					
Min RH (%)					
Wind Speed (Kmph)					
Wind Direction (deg)					

Northerly- **N**, North-Easterly- **N-E**, Easterly- **E**, South-Easterly- **S-E**, Southerly- **S**, South-Westerly- **S-W**, Westerly- **W**, North-westerly- **N-W**.

Weather for preceding week			
Date	Rainfall	Max. Temp	Min. Temp
05/11/2019			
06/11/2019			
07/11/2019			

### Weather forecast valid up to 13<sup>th</sup> November 2019

- There is no probability of rainfall.
- Max temp may be varying from 17-20 $^{\circ}$ C, whereas, Min temp may vary from 12-13  $^{\circ}$ C
- Sky will remain partially clear
- Max RH is likely to be 85-91 %, whereas, Min RH may vary from 65-70 %
- Wind speed may likely vary from 2-3 km/hr.
- Wind direction may blow from northeasterly
- Weekly cumulative rainfall= 001.0 mm


## AGRO-METEOREOLOGICAL ADVISORIES

### General Advisories for all the crops/ livestock

- Complete the sowing of rabi crops as early as possible before residual soil moisture lost.
- Harvest mature crops before unexpected rain.
- While selecting site for construction of animal shed a site that allows sun light exposure as well as good wind control is preferable

Name of the crop/Animal	Stage	Cultural practices/ Pest/ Diseases	Agro-Met Advisories
<b><i>Agricultural and Horticultural crops:</i></b>			
Rice	Grain maturity to harvesting and postharvest		<ul style="list-style-type: none"> <li>◆ Drain out water completely 10 days before harvesting.</li> <li>◆ Harvesting should start when the crops are 80-90% mature</li> <li>◆ Maintain seed moisture about 8-10 % for storage.</li> </ul>
Rjmash	Pre flowering/flowering		<ul style="list-style-type: none"> <li>◆ Irrigate crop at flowering stage.</li> </ul>
Ginger/ Turmeric	Tillering stage		<ul style="list-style-type: none"> <li>◆ Allow minimum entry to crop field to reduce disease infestation.</li> <li>◆ Allow proper drainage to avoid stagnation of water in the field.</li> <li>◆ Timely monitor the field to check the attack of bacterial wilt due to high humidity and temperature.</li> </ul>
Red cherry pepper/Brinjal/Toamto	Fruiting	Blight	<ul style="list-style-type: none"> <li>◆ Allow proper drainage to remove excess rain water from the field.</li> <li>◆ Timely monitor the field to check the attack of bacterial will due to high humidity and temperature.</li> <li>◆ Due to high temperature and high rainfall may cause late blight. Apply COC @ 0.25% at weekly interval for 4-5 sprays will reduce the disease.</li> </ul>
Cole crops	Nursery preparation		<ul style="list-style-type: none"> <li>◆ The nursery site should be dry and free from water stagnation as well as high moisture in the soil.</li> <li>◆ To protect from heavy rain the nursery should be in semi protected area.</li> </ul>



Sikkim Mandarin	Planting/fruiting		<ul style="list-style-type: none"> <li>◆ As there is a possibility of light to moderate rainfall so apply soil nutrients @ 25-35 kg/ha as maximum nutrient losses may occur due to rainfall.</li> </ul>
<i>Animal Sciences and livestock management</i>			
Poultry		Coccidiosis, Ranikhet disease	<ul style="list-style-type: none"> <li>◆ Keep the Vanaraja chicks in warm condition (32-33°C) during the first four weeks.</li> <li>◆ Low cost brooder should be made to prevent chick mortality due to cold stress.</li> <li>◆ Maintain brooding temperature through continuous electricity supply and for emergency Sigiri/Bukhari can be used.</li> </ul>
Sheep/goat			<ul style="list-style-type: none"> <li>◆ During winter months special care should be taken for sheep, goat for external parasites.</li> </ul>

*Agro-Met Advisory member of ICAR-NOFRI and KVK- East Sikkim*

<b>Name</b>	<b>Designation</b>	<b>Department</b>
<b><i>Dr. R. K. Avasthe</i></b>	<b>Joint Director</b>	<b>Soil Science</b>
<i>Dr. Raghavendra Singh</i>	Senior scientist and Head (I/C)	Agronomy
<i>Dr. P. K. Pathak</i>	SMS	Animal science
<i>Dr. J.K. Singh</i>	SMS	Plant Breeding
<i>Dr. Manoj Kumar</i>	SMS	Agronomy
<i>Sh. Boniface Lepcha</i>	SMS	Horticulture
<i>Sh. Santanu Das</i>	SMS	Agril. Meteorology
<i>Miss Tirtha Kumari Chettri</i>	Met. Observer	Biochemistry




# AGRO-MET ADVISORY BULLETIN FOR THE DISTRICT OF SOUTH SIKKIM

**PERIOD:**

**09<sup>th</sup> NOVEMBER to 13<sup>th</sup> NOVEMBER 2019**

*IMD, GANGTOK IN COLLABORATION WITH ICAR, GANGTOK, FSAD  
AND HCCD, GANGTOK*



**ISSUED ON 08<sup>th</sup> November 2019**

## **Meteorological Sub-division of Sikkim**

As per India Meteorological Department's classification, Sikkim is a part of Met Sub-Division of Sub Himalayan West Bengal & Sikkim (SHWB & SKM) and is divided into four (4) Districts:

- 1) North District with HQ at Mangan
- 2) East District with HQ at Gangtok
- 3) West District with HQ at Gyalsing
- 4) South District with HQ at Namchi

Issued by: Santanu Das, SMS (Agril. Meteorology), ICAR-KVK, East Sikkim, Ranipool

## **Value Added forecast for SOUTH-DISTRICT for next five days**

Parameter/Date	09/11/2019	10/11/2019	11/11/2019	12/11/2019	13/11/2019
Rainfall (mm)	No rainfall	Light rainfall	Light rainfall	No rainfall	No rainfall
Max Temp ( $^{\circ}\text{C}$ )	25	23	22	23	24
Min Temp ( $^{\circ}\text{C}$ )	14	13	12	12	13
Sky condition	Generally cloudy	Generally cloudy	Generally cloudy	Generally cloudy	Generally cloudy
Max RH (%)	85	90	91	85	82
Min RH (%)	70	75	80	75	72
Wind Speed (Kmph)	003	004	004	003	003
Wind Direction (deg)	NE	N	N	NE	NE

Northerly- **N**, North-Easterly- **N-E**, Easterly- **E**, South-Easterly- **S-E**, Southerly- **S**, South-Westerly- **S-W**, Westerly- **W**, North-westerly- **N-W**.

Weather for preceding week			
Date	Rainfall	Max. Temp	Min. Temp
	(mm)	( $^{\circ}\text{C}$ )	( $^{\circ}\text{C}$ )
05/11/2019	N/A	23.8	N/A
06/11/2019	N/A	23.0	12.0
07/11/2019	N/A	25.2	12.6

### **Weather forecast valid up to 13<sup>th</sup> November 2019**


- There will be no rainfall
- Max temp may be varying from 22-25 $^{\circ}\text{C}$ , whereas, Min temp may vary from 12-14 $^{\circ}\text{C}$
- Sky will remain generally cloudy.
- Max RH is likely to be 82-91 %, whereas, Min RH may vary from 70-80 %
- Wind speed may likely vary from 3-4 km/hr.
- Wind direction may blow from northeasterly direction
- Weekly cumulative rainfall= 007.0 mm

## AGRO-METEOREOLOGICAL ADVISORIES

### General Advisories for all the crops/ livestock

- Complete the sowing of rabi crops as early as possible before residual soil moisture lost.
- Harvest mature crops before unexpected rain.
- While selecting site for construction of animal shed a site that allows sun light exposure as well as good wind control is preferable

Name of the crop/Animal	Stage	Cultural practices/ Pest/ Diseases	Agro-Met Advisories
<b><i>Agricultural and Horticultural crops:</i></b>			
Rice	Grain maturity to harvesting and postharvest		<ul style="list-style-type: none"><li>◆ Drain out water completely 10 days before harvesting.</li><li>◆ Harvesting should start when the crops are 80-90% mature</li><li>◆ Maintain seed moisture about 8-10 % for storage.</li></ul>
Rjmash	Pre flowering/flowering		<ul style="list-style-type: none"><li>◆ Irrigate crop at flowering stage.</li></ul>
Ginger/ Turmeric	Tillering stage		<ul style="list-style-type: none"><li>◆ Allow minimum entry to crop field to reduce disease infestation.</li><li>◆ Allow proper drainage to avoid stagnation of water in the field.</li><li>◆ Timely monitor the field to check the attack of bacterial wilt due to high humidity and temperature.</li></ul>
Red cherry pepper/Brinjal/ Toamto	Fruiting	Blight	<ul style="list-style-type: none"><li>◆ Allow proper drainage to remove excess rain water from the field.</li><li>◆ Timely monitor the field to check the attack of bacterial will due to high humidity and temperature.</li><li>◆ Due to high temperature and high rainfall may cause late blight. Apply COC @ 0.25% at weekly interval for 4-5 sprays will reduce the disease.</li></ul>
Cole crops	Nursery preparation		<ul style="list-style-type: none"><li>◆ The nursery site should be dry and free from water stagnation as well as high moisture in the soil.</li><li>◆ To protect from heavy rain the nursery should be in semi protected area.</li></ul>

Sikkim Mandarin	Planting/fruiting		<ul style="list-style-type: none"> <li>As there is a possibility of light to moderate rainfall so apply soil nutrients @ 25-35 kg/ha as maximum nutrient losses may occur due to rainfall.</li> </ul>
<i>Animal Sciences and livestock management</i>			
Poultry		Coccidiosis, Ranikhet disease	<ul style="list-style-type: none"> <li>Keep the Vanaraja chicks in warm condition (32-33°C) during the first four weeks.</li> <li>Low cost brooder should be made to prevent chick mortality due to cold stress.</li> <li>Maintain brooding temperature through continuous electricity supply and for emergency Sigiri/Bukhari can be used.</li> </ul>
Sheep/goat			<ul style="list-style-type: none"> <li>During winter months special care should be taken for sheep, goat for external parasites.</li> </ul>

*Agro-Met Advisory member of ICAR-NOFRI and KVK- East Sikkim*

<b>Name</b>	<b>Designation</b>	<b>Department</b>
<b><i>Dr. R. K. Avasthe</i></b>	<b>Joint Director</b>	<b>Soil Science</b>
<i>Dr. Raghavendra Singh</i>	Senior scientist and Head (I/C)	Agronomy
<i>Dr. P. K. Pathak</i>	SMS	Animal science
<i>Dr. J.K. Singh</i>	SMS	Plant Breeding
<i>Dr. Manoj Kumar</i>	SMS	Agronomy
<i>Sh. Boniface Lepcha</i>	SMS	Horticulture
<i>Sh. Santanu Das</i>	SMS	Agril. Meteorology
<i>Miss Tirtha Kumari Chettri</i>	Met. Observer	Biochemistry




# AGRO-MET ADVISORY BULLETIN FOR THE DISTRICT OF WEST SIKKIM

## PERIOD:

**09<sup>th</sup> NOVEMBER to 13<sup>th</sup> NOVEMBER' 2019**

*IMD, GANGTOK IN COLLABORATION WITH ICAR, GANGTOK, FSAD  
AND HCCD, GANGTOK*



**ISSUED ON 08<sup>th</sup> November 2019**

### Meteorological Sub-division of Sikkim

As per India Meteorological Department's classification, Sikkim is a part of Met Sub-Division of Sub Himalayan West Bengal & Sikkim (SHWB & SKM) and is divided into four (4) Districts:

- 1) North District with HQ at Mangan
- 2) East District with HQ at Gangtok
- 3) West District with HQ at Gyalsing
- 4) South District with HQ at Namchi

Issued by: Santanu Das, SMS (Agril. Meteorology), ICAR-KVK, East Sikkim, Ranipool



### Value Added forecast for WEST-DISTRICT for next five days

Parameter/Date	09/11/2019	10/11/2019	11/11/2019	12/11/2019	13/11/2019
Rainfall (mm)	No rainfall	Light rainfall	Very light rainfall	No rainfall	No rainfall
Max Temp ( $^{\circ}\text{C}$ )	22	21	19	20	21
Min Temp ( $^{\circ}\text{C}$ )	13	12	11	11	12
Sky condition	Generally cloudy	Generally cloudy	Generally cloudy	Generally cloudy	Generally cloudy
Max RH (%)	87	90	91	86	84
Min RH (%)	72	76	80	74	70
Wind Speed (Kmph)	002	003	004	003	003
Wind Direction (deg)	NW	NW	NW	NW	NW

Northerly- **N**, North-Easterly- **N-E**, Easterly- **E**, South-Easterly- **S-E**, Southerly- **S**, South-Westerly- **S-W**, Westerly- **W**, North-westerly- **N-W**.

Weather for preceding week			
Date	Rainfall	Max. Temp	Min. Temp
	(mm)	( $^{\circ}\text{C}$ )	( $^{\circ}\text{C}$ )
05/11/2019	005	21.4	N/A
06/11/2019	000	21.9	10.4
07/11/2019	000	N/A	11.5

### Weather forecast valid up to 13<sup>th</sup> November 2019


- Probability of no rainfall
- Max temp may be varying from 20-22 $^{\circ}\text{C}$ , whereas, Min temp may vary from 11-13 $^{\circ}\text{C}$
- Sky will remain generally cloudy.
- Max RH is likely to be 84-91 %, whereas, Min RH may vary from 70-80 %
- Wind speed may likely vary from 2-4 km/hr.
- Wind direction may blow from northwesterly.
- Weekly cumulative rainfall= 002.0 mm

## AGRO-METEOREOLOGICAL ADVISORIES

### General Advisories for all the crops/ livestock (English)

- ➡ Complete the sowing of rabi crops as early as possible before residual soil moisture lost.
- ➡ Harvest mature crops before unexpected rain.
- ➡ While selecting site for construction of animal shed a site that allows sun light exposure as well as good wind control is preferable

Name of the crop/Animal	Stage	Cultural practices/ Pest/ Diseases	Agro-Met Advisories
<b><i>Agricultural and Horticultural crops:</i></b>			
Rice	Grain maturity to harvesting and postharvest		<ul style="list-style-type: none"><li>◆ Drain out water completely 10 days before harvesting.</li><li>◆ Harvesting should start when the crops are 80-90% mature</li><li>◆ Maintain seed moisture about 8-10 % for storage.</li></ul>
Rjmash	Pre flowering/flowering		<ul style="list-style-type: none"><li>◆ Irrigate crop at flowering stage.</li></ul>
Ginger/ Turmeric	Tillering stage		<ul style="list-style-type: none"><li>◆ Allow minimum entry to crop field to reduce disease infestation.</li><li>◆ Allow proper drainage to avoid stagnation of water in the field.</li><li>◆ Timely monitor the field to check the attack of bacterial wilt due to high humidity and temperature.</li></ul>
Red cherry pepper/Brinjal/ Toamto	Fruiting	Blight	<ul style="list-style-type: none"><li>◆ Allow proper drainage to remove excess rain water from the field.</li><li>◆ Timely monitor the field to check the attack of bacterial will due to high humidity and temperature.</li><li>◆ Due to high temperature and high rainfall may cause late blight. Apply COC @ 0.25% at weekly interval for 4-5 sprays will reduce the disease.</li></ul>
Cole crops	Nursery preparation		<ul style="list-style-type: none"><li>◆ The nursery site should be dry and free from water stagnation as well as high moisture in the soil.</li><li>◆ To protect from heavy rain the nursery should be in semi protected area.</li></ul>

Sikkim Mandarin	Planting/fruiting		<ul style="list-style-type: none"> <li>◆ As there is a possibility of light to moderate rainfall so apply soil nutrients @ 25-35 kg/ha as maximum nutrient losses may occur due to rainfall.</li> </ul>
<b><i>Animal Sciences and livestock management</i></b>			
Poultry		Coccidiosis, Ranikhet disease	<ul style="list-style-type: none"> <li>◆ Keep the Vanaraja chicks in warm condition (32-33°C) during the first four weeks.</li> <li>◆ Low cost brooder should be made to prevent chick mortality due to cold stress.</li> <li>◆ Maintain brooding temperature through continuous electricity supply and for emergency Sigiri/Bukhari can be used.</li> </ul>
Sheep/goat			<ul style="list-style-type: none"> <li>◆ During winter months special care should be taken for sheep, goat for external parasites.</li> </ul>

*Agro-Met Advisory member of ICAR-NOFRI and KVK- East Sikkim*

<b>Name</b>	<b>Designation</b>	<b>Department</b>
<b><i>Dr. R. K. Avasthe</i></b>	<b>Joint Director</b>	<b>Soil Science</b>
<i>Dr. Raghavendra Singh</i>	Senior scientist and Head (I/C)	Agronomy
<i>Dr. P. K. Pathak</i>	SMS	Animal science
<i>Dr. J.K. Singh</i>	SMS	Plant Breeding
<i>Dr. Manoj Kumar</i>	SMS	Agronomy
<i>Sh. Boniface Lepcha</i>	SMS	Horticulture
<i>Sh. Santanu Das</i>	SMS	Agril. Meteorology
<i>Miss Tirtha Kumari Chettri</i>	Met. Observer	Biochemistry