




AGRO-MET ADVISORY BULLETIN FOR THE DISTRICT OF NORTH SIKKIM

PERIOD:

02nd OCTOBER to 06th OCTOBER' 2019

*IMD, GANGTOK IN COLLABORATION WITH ICAR, GANGTOK, FSAD
AND HCCD, GANGTOK*



ISSUED ON 01st October 2019

Meteorological Sub-division of Sikkim

As per India Meteorological Department's classification, Sikkim is a part of Met Sub-Division of Sub Himalayan West Bengal & Sikkim (SHWB & SKM) and is divided into four (4) Districts:

- 1) North District with HQ at Mangan
- 2) East District with HQ at Gangtok
- 3) West District with HQ at Gyalsing
- 4) South District with HQ at Namchi

Issued by: Santanu Das, SMS (Agril. Meteorology), ICAR-KVK, East Sikkim, Ranipool

Value Added forecast for NORTH-DISTRICT for next five days

Parameter/Date	02/10/2019	03/10/2019	04/10/2019	05/10/2019	06/10/2019
Rainfall (mm)	Light rainfall	Light rainfall	Light rainfall	Light rainfall	Light rainfall
Max Temp ($^{\circ}\text{C}$)	28	27	26	26	27
Min Temp ($^{\circ}\text{C}$)	17	17	16	16	17
Sky condition	Partially clear	Generally cloudy	Generally cloudy	Cloudy	Cloudy
Max RH (%)	92	93	88	93	92
Min RH (%)	72	75	77	74	75
Wind Speed (Kmph)	6	6	4	5	4
Wind Direction (deg)	S	S	S	S	S

Northerly- **N**, North-Easterly- **N-E**, Easterly- **E**, South-Easterly- **S-E**, Southerly- **S**, South-Westerly- **S-W**, Westerly- **W**, North-westerly- **N-W**.

Weather for preceding week			
Date	Rainfall	Max. Temp	Min. Temp
	(mm)	($^{\circ}\text{C}$)	($^{\circ}\text{C}$)
28/9/2019	007	21.2	17.5
29/9/2019	005	20.9	16.9
30/9/2019	008	24.7	17.2

Weather forecast valid up to 6th October 2019


- Probability of light rainfall.
- Max temp may be varying from 26-28 $^{\circ}\text{C}$, whereas, Min temp may vary from 16-17 $^{\circ}\text{C}$
- Sky will remain generally cloudy.
- Max RH is likely to be 88-93 %, whereas, Min RH may vary from 72-77 %
- Wind speed may likely vary from 4-6 km/hr.
- Wind direction may blow from southerly direction.
- Weekly cumulative rainfall= 34 mm

AGRO-METEOREOLOGICAL ADVISORIES

General Advisories for all the crops/ livestock

- ➡ If there is a probability of light rainfall, farmers are advised to keep drainage facilities on their field and harvest all the matured crops.
- ➡ During high humidity and cloudy weather there may be infestation of insect pest, so the farmers are suggested to apply neem oil.
- ➡ Bunds to be provide to reduce nutrient lost due to high rainfall.
- ➡ Rain water harvesting through *jalkund* should be done to conserve excess rain water during high rainfall.
- ➡ Excess water stagnation may cause fungal and bacterial diseases to the crops.
- ➡ While selecting site for construction of shed a site that allows sun light exposure as well as good wind control is preferable.

Name of the crop/Animal	Stage	Cultural practices/ Pest/ Diseases	Agro-Met Advisories
<i>Agricultural and Horticultural crops:</i>			
Rice	Tillering		<ul style="list-style-type: none">◆ Maintain 5 cm of water in the paddy field.◆ First weeding should be completed at 30-35 days after transplanting.
Black gram (Pahelo dal) /Rjmash/Cowpe a	Vegetative		<ul style="list-style-type: none">◆ Farmers are advised to provide proper drainage in the field to avoid wilting in high rainfall areas.◆ First weeding should be completed after 25-30 days of sowing to avoid weed competition.
Ginger/ Turmeric	Tillering stage		<ul style="list-style-type: none">◆ Allow minimum entry to crop field to reduce disease infestation.◆ Allow proper drainage to avoid stagnation of water in the field.◆ Timely monitor the field to check the attack of bacterial wilt due to high humidity and temperature.
Red cherry pepper/Brinjal/ Toamto	Fruiting	Blight	<ul style="list-style-type: none">◆ Allow proper drainage to remove excess rain water from the field.◆ Timely monitor the field to check the attack of bacterial will due to high humidity and temperature.◆ Due to high temperature and high

			rainfall may cause late blight. Apply COC @ 0.25% at weekly interval for 4-5 sprays will reduce the disease.
Cole crops	Nursery preparation		<ul style="list-style-type: none"> ◆ The nursery site should be dry and free from water stagnation as well as high moisture in the soil. ◆ To protect from heavy rain the nursery should be in semi protected area.
Sikkim Mandarin	Planting/fruiting		<ul style="list-style-type: none"> ◆ As there is a possibility of light to moderate rainfall so apply soil nutrients @ 25-35 kg/ha as maximum nutrient losses may occur due to rainfall.
Animal Sciences and livestock management			
Cattle			<ul style="list-style-type: none"> ◆ Provide plenty of cold clean drinking water to your livestock at the time between 01 to 3 pm during the peak summer months. ◆ Maintain hygienic condition of animal shed and disinfect the animal shed with 10% formalin solution.
Poultry		Coccidiosis, Ranikhet disease	<ul style="list-style-type: none"> ◆ Provide proper ventilation to poultry birds to prevent from heat stress during summer months ◆ Periodic supplementation of vitamins and minerals to your livestock is very much essential during hot weather. ◆ Special care should be taken for mite and Gastro Intestinal tape worm in case of poultry during the hot and humid summer months. ◆ Take preventive measures against Coccidiosis of poultry during summer months.
Pig	All stages	Swine fever	<ul style="list-style-type: none"> ◆ Provide creep area (separate enclosure, where the piglets have access but the sow does not) with heat source (200 watt bulb) to reduce piglet mortality due to cold weather.

Agro-Met Advisory member of ICAR-NOFRI and KVK- East Sikkim

Name	Designation	Department
<i>Dr. R. K. Avasthe</i>	Joint Director	Soil Science
<i>Dr. Raghavendra Singh</i>	Senior scientist and Head (I/C)	Agronomy
<i>Dr. P. K. Pathak</i>	SMS	Animal science
<i>Dr. J.K. Singh</i>	SMS	Plant Breeding
<i>Dr. Manoj Kumar</i>	SMS	Agronomy
<i>Sh. Boniface Lepcha</i>	SMS	Horticulture
<i>Sh. Santanu Das</i>	SMS	Agril. Meteorology
<i>Miss Tirtha Kumari Chettri</i>	Met. Observer	Biochemistry




AGRO-MET ADVISORY BULLETIN FOR THE DISTRICT OF EAST SIKKIM

PERIOD:

02nd OCTOBER to 06th OCTOBER' 2019

*IMD, GANGTOK IN COLLABORATION WITH ICAR, GANGTOK, FSAD
AND HCCD, GANGTOK*



ISSUED ON 1st October 2019

Meteorological Sub-division of Sikkim

As per India Meteorological Department's classification, Sikkim is a part of Met Sub-Division of Sub Himalayan West Bengal & Sikkim (SHWB & SKM) and is divided into four (4) Districts:

- 1) North District with HQ at Mangan
- 2) East District with HQ at Gangtok
- 3) West District with HQ at Gyalsing
- 4) South District with HQ at Namchi

Issued by: Santanu Das, SMS (Agril. Meteorology), ICAR-KVK, East Sikkim, Ranipool

Value Added forecast for EAST-DISTRICT for next five days

Parameter/Date	02/10/2019	03/10/2019	04/10/2019	05/10/2019	06/10/2019
Rainfall (mm)	Light rainfall	Light rainfall	Light rainfall	Light rainfall	Light rainfall
Max Temp (°C)	24	25	24	24	27
Min Temp (°C)	16	17	16	16	17
Sky condition	Partially clear	Generally cloudy	Generally cloudy	Generally cloudy	Cloudy
Max RH (%)	94	93	90	93	92
Min RH (%)	59	61	59	54	57
Wind Speed (Kmph)	3	3	2	3	3
Wind Direction (deg)	NE	NE	NE	NE	NE

Northerly- **N**, North-Easterly- **N-E**, Easterly- **E**, South-Easterly- **S-E**, Southerly- **S**, South-Westerly- **S-W**, Westerly-**W**, North-westerly- **N-W**.

Weather for preceding week			
Date	Rainfall	Max. Temp	Min. Temp
28/9/2019	010	17.7	15.3
29/9/2019	015	16.8	14.7
30/9/2019	015	16.9	14.7

Weather forecast valid up to 06th October 2019

- Probability of light rainfall
- Max temp may be varying from 24-27 °C, whereas, Min temp may vary from 16-17 °C
- Sky will remain generally cloudy
- Max RH is likely to be 90-94 %, whereas, Min RH may vary from 54-61 %
- Wind speed may likely vary from 2-3 km/hr.
- Wind direction may blow from northeasterly
- Weekly cumulative rainfall= 29mm

AGRO-METEREOLOGICAL ADVISORIES


General Advisories for all the crops/ livestock

- If there is a probability of light rainfall, farmers are advised to keep drainage facilities on their field and harvest all the matured crops.
- During high humidity and cloudy weather there may be infestation of insect pest, so the farmers are suggested to apply neem oil.
- Bunds to be provide to reduce nutrient lost due to high rainfall.
- Rain water harvesting through *jalkund* should be done to conserve excess rain water during high rainfall.
- Excess water stagnation may cause fungal and bacterial diseases to the crops.
- While selecting site for construction of shed a site that allows sun light exposure as well as good wind control is preferable.

Name of the crop/Animal	Stage	Cultural practices/ Pest/ Diseases	Agro-Met Advisories
<i>Agricultural and Horticultural crops:</i>			
Rice	Tillering		<ul style="list-style-type: none">◆ Maintain 5 cm of water in the paddy field.◆ First weeding should be completed at 30-35 days after transplanting.
Black gram (Pahelo dal) /Rjmash/Cowpe a	Vegetative		<ul style="list-style-type: none">◆ Farmers are advised to provide proper drainage in the field to avoid wilting in high rainfall areas.◆ First weeding should be completed after 25-30 days of sowing to avoid weed competition.
Ginger/ Turmeric	Tillering stage		<ul style="list-style-type: none">◆ Allow minimum entry to crop field to reduce disease infestation.◆ Allow proper drainage to avoid stagnation of water in the field.◆ Timely monitor the field to check the attack of bacterial wilt due to high humidity and temperature.
Red cherry pepper/Brinjal/ Toamto	Fruiting	Blight	<ul style="list-style-type: none">◆ Allow proper drainage to remove excess rain water from the field.◆ Timely monitor the field to check the attack of bacterial will due to high humidity and temperature.◆ Due to high temperature and high

			rainfall may cause late blight. Apply COC @ 0.25% at weekly interval for 4-5 sprays will reduce the disease.
Cole crops	Nursery preparation		<ul style="list-style-type: none"> ◆ The nursery site should be dry and free from water stagnation as well as high moisture in the soil. ◆ To protect from heavy rain the nursery should be in semi protected area.
Sikkim Mandarin	Planting/fruiting		<ul style="list-style-type: none"> ◆ As there is a possibility of light to moderate rainfall so apply soil nutrients @ 25-35 kg/ha as maximum nutrient losses may occur due to rainfall.

Animal Sciences and livestock management

Cattle			<ul style="list-style-type: none"> ◆ Provide plenty of cold clean drinking water to your livestock at the time between 01 to 3 pm during the peak summer months. ◆ Maintain hygienic condition of animal shed and disinfect the animal shed with 10% formalin solution.
Poultry		Coccidiosis, Ranikhet disease	<ul style="list-style-type: none"> ◆ Provide proper ventilation to poultry birds to prevent from heat stress during summer months ◆ Periodic supplementation of vitamins and minerals to your livestock is very much essential during hot weather. ◆ Special care should be taken for mite and Gastro Intestinal tape worm in case of poultry during the hot and humid summer months. ◆ Take preventive measures against Coccidiosis of poultry during summer months.
Pig	All stages	Swine fever	<ul style="list-style-type: none"> ◆ Provide creep area (separate enclosure, where the piglets have access but the sow does not) with heat source (200 watt bulb) to reduce piglet mortality due to cold weather.

Agro-Met Advisory member of ICAR-NOFRI and KVK- East Sikkim

Name	Designation	Department
<i>Dr. R. K. Avasthe</i>	Joint Director	Soil Science
<i>Dr. Raghavendra Singh</i>	Senior scientist and Head (I/C)	Agronomy
<i>Dr. P. K. Pathak</i>	SMS	Animal science
<i>Dr. J.K. Singh</i>	SMS	Plant Breeding
<i>Dr. Manoj Kumar</i>	SMS	Agronomy
<i>Sh. Boniface Lepcha</i>	SMS	Horticulture
<i>Sh. Santanu Das</i>	SMS	Agril. Meteorology
<i>Miss Tirtha Kumari Chettri</i>	Met. Observer	Biochemistry




AGRO-MET ADVISORY BULLETIN FOR THE DISTRICT OF SOUTH SIKKIM

PERIOD:

02nd OCTOBER to 06th OCTOBER' 2019



*IMD, GANGTOK IN COLLABORATION WITH ICAR, GANGTOK, FSAD AND HCCD,
GANGTOK*



ISSUED ON 01st October 2019

Meteorological Sub-division of Sikkim

As per India Meteorological Department's classification, Sikkim is a part of Met Sub-Division of Sub Himalayan West Bengal & Sikkim (SHWB & SKM) and is divided into four (4) Districts:

- 1) North District with HQ at Mangan
- 2) East District with HQ at Gangtok
- 3) West District with HQ at Gyalsing
- 4) South District with HQ at Namchi

Issued by: Santanu Das, SMS (Agril. Meteorology), ICAR-KVK, East Sikkim, Ranipool

Value Added forecast for SOUTH-DISTRICT for next five days

Parameter/Date	02/10/2019	03/10/2019	04/10/2019	05/10/2019	06/10/2019
Rainfall (mm)	Light rainfall	Very light rainfall	Light rainfall	Light rainfall	Light rainfall
Max Temp (°C)	25	25	24	24	25
Min Temp (°C)	16	17	16	16	17
Sky condition	Clear sky	Generally cloudy	Generally cloudy	Cloudy	Cloudy
Max RH (%)	88	84	78	84	86
Min RH (%)	72	74	74	68	70
Wind Speed (Kmph)	003	003	002	002	002
Wind Direction (deg)	N	NE	NE	NE	N

Northerly- **N**, North-Easterly- **N-E**, Easterly- **E**, South-Easterly- **S-E**, Southerly- **S**, South-Westerly- **S-W**, Westerly-**W**, North-westerly- **N-W**.

Weather for preceding week			
Date	Rainfall	Max. Temp	Min. Temp
	(mm)	(°C)	(°C)
28/9/2019	N/A	19.9	17.0
29/9/2019	N/A	17.6	15.9
30/9/2019	N/A	21.5	15.1

Weather forecast valid up to 6th October 2019

- There will be chances of light rainfall
- Max temp may be varying from 24-25°C, whereas, Min temp may vary from 16-17°C
- Sky will remain generally cloudy
- Max RH is likely to be 78-86 %, whereas, Min RH may vary from 68-74 %
- Wind speed may likely vary from 2-3 km/hr.
- Wind direction may blow from north easterly direction
- Weekly cumulative rainfall= 21 mm

AGRO-METEREOLOGICAL ADVISORIES


General Advisories for all the crops/ livestock

- If there is a probability of light rainfall, farmers are advised to keep drainage facilities on their field and harvest all the matured crops.
- During high humidity and cloudy weather there may be infestation of insect pest, so the farmers are suggested to apply neem oil.
- Bunds to be provide to reduce nutrient lost due to high rainfall.
- Rain water harvesting through *jalkund* should be done to conserve excess rain water during high rainfall.
- Excess water stagnation may cause fungal and bacterial diseases to the crops.
- While selecting site for construction of shed a site that allows sun light exposure as well as good wind control is preferable.

Name of the crop/Animal	Stage	Cultural practices/ Pest/ Diseases	Agro-Met Advisories
<i>Agricultural and Horticultural crops:</i>			
Rice	Tillering		<ul style="list-style-type: none">◆ Maintain 5 cm of water in the paddy field.◆ First weeding should be completed at 30-35 days after transplanting.
Black gram (Pahelo dal) /Rjmash/Cowpe a	Vegetative		<ul style="list-style-type: none">◆ Farmers are advised to provide proper drainage in the field to avoid wilting in high rainfall areas.◆ First weeding should be completed after 25-30 days of sowing to avoid weed competition.
Ginger/ Turmeric	Tillering stage		<ul style="list-style-type: none">◆ Allow minimum entry to crop field to reduce disease infestation.◆ Allow proper drainage to avoid stagnation of water in the field.◆ Timely monitor the field to check the attack of bacterial wilt due to high humidity and temperature.
Red cherry pepper/Brinjal/ Toamto	Fruiting	Blight	<ul style="list-style-type: none">◆ Allow proper drainage to remove excess rain water from the field.◆ Timely monitor the field to check the attack of bacterial will due to high humidity and temperature.◆ Due to high temperature and high

			rainfall may cause late blight. Apply COC @ 0.25% at weekly interval for 4-5 sprays will reduce the disease.
Cole crops	Nursery preparation		<ul style="list-style-type: none"> ◆ The nursery site should be dry and free from water stagnation as well as high moisture in the soil. ◆ To protect from heavy rain the nursery should be in semi protected area.
Sikkim Mandarin	Planting/fruiting		<ul style="list-style-type: none"> ◆ As there is a possibility of light to moderate rainfall so apply soil nutrients @ 25-35 kg/ha as maximum nutrient losses may occur due to rainfall.

Animal Sciences and livestock management

Cattle			<ul style="list-style-type: none"> ◆ Provide plenty of cold clean drinking water to your livestock at the time between 01 to 3 pm during the peak summer months. ◆ Maintain hygienic condition of animal shed and disinfect the animal shed with 10% formalin solution.
Poultry		Coccidiosis, Ranikhet disease	<ul style="list-style-type: none"> ◆ Provide proper ventilation to poultry birds to prevent from heat stress during summer months ◆ Periodic supplementation of vitamins and minerals to your livestock is very much essential during hot weather. ◆ Special care should be taken for mite and Gastro Intestinal tape worm in case of poultry during the hot and humid summer months. ◆ Take preventive measures against Coccidiosis of poultry during summer months.
Pig	All stages	Swine fever	<ul style="list-style-type: none"> ◆ Provide creep area (separate enclosure, where the piglets have access but the sow does not) with heat source (200 watt bulb) to reduce piglet mortality due to cold weather.

Agro-Met Advisory member of ICAR-NOFRI and KVK- East Sikkim

Name	Designation	Department
<i>Dr. R. K. Avasthe</i>	Joint Director	Soil Science
<i>Dr. Raghavendra Singh</i>	Senior scientist and Head (I/C)	Agronomy
<i>Dr. P. K. Pathak</i>	SMS	Animal science
<i>Dr. J.K. Singh</i>	SMS	Plant Breeding
<i>Dr. Manoj Kumar</i>	SMS	Agronomy
<i>Sh. Boniface Lepcha</i>	SMS	Horticulture
<i>Sh. Santanu Das</i>	SMS	Agril. Meteorology
<i>Miss Tirtha Kumari Chettri</i>	Met. Observer	Biochemistry




AGRO-MET ADVISORY BULLETIN FOR THE DISTRICT OF WEST SIKKIM

PERIOD:

02nd OCTOBER to 06th OCTOBER' 2019

*IMD, GANGTOK IN COLLABORATION WITH ICAR, GANGTOK, FSAD
AND HCCD, GANGTOK*



ISSUED ON 01st October 2019

Meteorological Sub-division of Sikkim

As per India Meteorological Department's classification, Sikkim is a part of Met Sub-Division of Sub Himalayan West Bengal & Sikkim (SHWB & SKM) and is divided into four (4) Districts:

- 1) North District with HQ at Mangan
- 2) East District with HQ at Gangtok
- 3) West District with HQ at Gyalsing
- 4) South District with HQ at Namchi

Issued by: Santanu Das, SMS (Agril. Meteorology), ICAR-KVK, East Sikkim, Ranipool

Value Added forecast for WEST-DISTRICT for next five days

Parameter/Date	02/10/2019	03/10/2019	04/10/2019	05/10/2019	06/10/2019
Rainfall (mm)	Light rainfall	Light rainfall	Light rainfall	Light rainfall	Light rainfall
Max Temp (°C)	25	25	24	24	25
Min Temp (°C)	15	16	16	15	15
Sky condition	Partially cloudy	Generally cloudy	Generally cloudy	Cloudy	Cloudy
Max RH (%)	94	93	92	93	91
Min RH (%)	75	76	78	72	73
Wind Speed (Kmph)	007	006	006	006	006
Wind Direction (deg)	NW	NW	NW	N	N

Northerly- **N**, North-Easterly- **N-E**, Easterly- **E**, South-Easterly- **S-E**, Southerly- **S**, South-Westerly- **S-W**, Westerly-**W**, North-westerly- **N-W**.

Weather for preceding week			
Date	Rainfall	Max. Temp	Min. Temp
	(mm)	(°C)	(°C)
28/9/2019	N/A	17.8	16.0
29/9/2019	N/A	17.1	14.3
30/9/2019	N/A	21.2	14.0

Weather forecast valid up to 06th October 2019


- Probability of light rainfall
- Max temp may be varying from 24-25°C, whereas, Min temp may vary from 15-16°C
- Sky will remain generally cloudy
- Max RH is likely to be 91-94 %, whereas, Min RH may vary from 72-78 %
- Wind speed may likely vary from 6-7 km/hr.
- Wind direction may blow from northwesterly
- Weekly cumulative rainfall= 23 mm

AGRO-METEOROLOGICAL ADVISORIES

General Advisories for all the crops/ livestock (English)

- ➡ If there is a probability of light rainfall, farmers are advised to keep drainage facilities on their field and harvest all the matured crops.
- ➡ During high humidity and cloudy weather there may be infestation of insect pest, so the farmers are suggested to apply neem oil.
- ➡ Bunds to be provide to reduce nutrient lost due to high rainfall.
- ➡ Rain water harvesting through *jalkund* should be done to conserve excess rain water during high rainfall.
- ➡ Excess water stagnation may cause fungal and bacterial diseases to the crops.
- ➡ While selecting site for construction of shed a site that allows sun light exposure as well as good wind control is preferable.

Name of the crop/Animal	Stage	Cultural practices/ Pest/ Diseases	Agro-Met Advisories
<i>Agricultural and Horticultural crops:</i>			
Rice	Tillering		<ul style="list-style-type: none">◆ Maintain 5 cm of water in the paddy field.◆ First weeding should be completed at 30-35 days after transplanting.
Black gram (Pahelo dal) /Rjmash/Cowpe a	Vegetative		<ul style="list-style-type: none">◆ Farmers are advised to provide proper drainage in the field to avoid wilting in high rainfall areas.◆ First weeding should be completed after 25-30 days of sowing to avoid weed competition.
Ginger/ Turmeric	Tillering stage		<ul style="list-style-type: none">◆ Allow minimum entry to crop field to reduce disease infestation.◆ Allow proper drainage to avoid stagnation of water in the field.◆ Timely monitor the field to check the attack of bacterial wilt due to high humidity and temperature.
Red cherry pepper/Brinjal/ Toamto	Fruiting	Blight	<ul style="list-style-type: none">◆ Allow proper drainage to remove excess rain water from the field.◆ Timely monitor the field to check the attack of bacterial will due to high humidity and temperature.◆ Due to high temperature and high

			rainfall may cause late blight. Apply COC @ 0.25% at weekly interval for 4-5 sprays will reduce the disease.
Cole crops	Nursery preparation		<ul style="list-style-type: none"> ◆ The nursery site should be dry and free from water stagnation as well as high moisture in the soil. ◆ To protect from heavy rain the nursery should be in semi protected area.
Sikkim Mandarin	Planting/fruiting		<ul style="list-style-type: none"> ◆ As there is a possibility of light to moderate rainfall so apply soil nutrients @ 25-35 kg/ha as maximum nutrient losses may occur due to rainfall.
Animal Sciences and livestock management			
Cattle			<ul style="list-style-type: none"> ◆ Provide plenty of cold clean drinking water to your livestock at the time between 01 to 3 pm during the peak summer months. ◆ Maintain hygienic condition of animal shed and disinfect the animal shed with 10% formalin solution.
Poultry		Coccidiosis, Ranikhet disease	<ul style="list-style-type: none"> ◆ Provide proper ventilation to poultry birds to prevent from heat stress during summer months ◆ Periodic supplementation of vitamins and minerals to your livestock is very much essential during hot weather. ◆ Special care should be taken for mite and Gastro Intestinal tape worm in case of poultry during the hot and humid summer months. ◆ Take preventive measures against Coccidiosis of poultry during summer months.
Pig	All stages	Swine fever	<ul style="list-style-type: none"> ◆ Provide creep area (separate enclosure, where the piglets have access but the sow does not) with heat source (200 watt bulb) to reduce piglet mortality due to cold weather.

Agro-Met Advisory member of ICAR-NOFRI and KVK- East Sikkim

Name	Designation	Department
<i>Dr. R. K. Avasthe</i>	Joint Director	Soil Science
<i>Dr. Raghavendra Singh</i>	Senior scientist and Head (I/C)	Agronomy
<i>Dr. P. K. Pathak</i>	SMS	Animal science
<i>Dr. J.K. Singh</i>	SMS	Plant Breeding
<i>Dr. Manoj Kumar</i>	SMS	Agronomy
<i>Sh. Boniface Lepcha</i>	SMS	Horticulture
<i>Sh. Santanu Das</i>	SMS	Agril. Meteorology
<i>Miss Tirtha Kumari Chettri</i>	Met. Observer	Biochemistry