




AGRO-MET ADVISORY BULLETIN FOR THE DISTRICT OF NORTH SIKKIM

PERIOD:

05th OCTOBER to 09th OCTOBER' 2019

*IMD, GANGTOK IN COLLABORATION WITH ICAR, GANGTOK, FSAD
AND HCCD, GANGTOK*



ISSUED ON 04th October 2019

Meteorological Sub-division of Sikkim

As per India Meteorological Department's classification, Sikkim is a part of Met Sub-Division of Sub Himalayan West Bengal & Sikkim (SHWB & SKM) and is divided into four (4) Districts:

- 1) North District with HQ at Mangan
- 2) East District with HQ at Gangtok
- 3) West District with HQ at Gyalsing
- 4) South District with HQ at Namchi

Issued by: Santanu Das, SMS (Agril. Meteorology), ICAR-KVK, East Sikkim, Ranipool

Value Added forecast for NORTH-DISTRICT for next five days

Parameter/Date	05/10/2019	06/10/2019	07/10/2019	08/10/2019	09/10/2019
Rainfall (mm)	Light rainfall	Light rainfall	Light rainfall	Light rainfall	Light rainfall
Max Temp ($^{\circ}\text{C}$)	25	26	26	25	23
Min Temp ($^{\circ}\text{C}$)	17	18	18	17	16
Sky condition	Generally cloudy	Generally cloudy	Generally cloudy	Generally cloudy	Generally cloudy
Max RH (%)	93	91	91	92	94
Min RH (%)	83	80	80	82	85
Wind Speed (Kmph)	006	004	004	007	006
Wind Direction (deg)	S	S	SE	S	S

Northerly- **N**, North-Easterly- **N-E**, Easterly- **E**, South-Easterly- **S-E**, Southerly- **S**, South-Westerly- **S-W**, Westerly- **W**, North-westerly- **N-W**.

Weather for preceding week			
Date	Rainfall	Max. Temp	Min. Temp
	(mm)	($^{\circ}\text{C}$)	($^{\circ}\text{C}$)
01/10/2019	000	N/A	17.3
02/10/2019	005	23.3	18.0
03/10/2019	008	25.7	17.7

Weather forecast valid up to 9th October 2019


- Probability of light rainfall.
- Max temp may be varying from 23-26 $^{\circ}\text{C}$, whereas, Min temp may vary from 16-18 $^{\circ}\text{C}$
- Sky will remain generally cloudy.
- Max RH is likely to be 91-94 %, whereas, Min RH may vary from 80-85 %
- Wind speed may likely vary from 4-6 km/hr.
- Wind direction may blow from southerly direction.
- Weekly cumulative rainfall= 34 mm

AGRO-METEOREOLOGICAL ADVISORIES

General Advisories for all the crops/ livestock

- ➡ If there is a probability of light rainfall, farmers are advised to keep drainage facilities on their field and harvest all the matured crops.
- ➡ During high humidity and cloudy weather there may be infestation of insect pest, so the farmers are suggested to apply neem oil.
- ➡ Bunds to be provide to reduce nutrient lost due to high rainfall.
- ➡ Rain water harvesting through *jalkund* should be done to conserve excess rain water during high rainfall.
- ➡ Excess water stagnation may cause fungal and bacterial diseases to the crops.
- ➡ While selecting site for construction of shed a site that allows sun light exposure as well as good wind control is preferable.

Name of the crop/Animal	Stage	Cultural practices/ Pest/ Diseases	Agro-Met Advisories
<i>Agricultural and Horticultural crops:</i>			
Rice	Tillering		<ul style="list-style-type: none">◆ Maintain 5 cm of water in the paddy field.◆ First weeding should be completed at 30-35 days after transplanting.
Black gram (Pahelo dal) /Rjmash/Cowpe a	Vegetative		<ul style="list-style-type: none">◆ Farmers are advised to provide proper drainage in the field to avoid wilting in high rainfall areas.◆ First weeding should be completed after 25-30 days of sowing to avoid weed competition.
Ginger/ Turmeric	Tillering stage		<ul style="list-style-type: none">◆ Allow minimum entry to crop field to reduce disease infestation.◆ Allow proper drainage to avoid stagnation of water in the field.◆ Timely monitor the field to check the attack of bacterial wilt due to high humidity and temperature.
Red cherry pepper/Brinjal/ Toamto	Fruiting	Blight	<ul style="list-style-type: none">◆ Allow proper drainage to remove excess rain water from the field.◆ Timely monitor the field to check the attack of bacterial will due to high humidity and temperature.◆ Due to high temperature and high

			rainfall may cause late blight. Apply COC @ 0.25% at weekly interval for 4-5 sprays will reduce the disease.
Cole crops	Nursery preparation		<ul style="list-style-type: none"> ◆ The nursery site should be dry and free from water stagnation as well as high moisture in the soil. ◆ To protect from heavy rain the nursery should be in semi protected area.
Sikkim Mandarin	Planting/fruiting		<ul style="list-style-type: none"> ◆ As there is a possibility of light to moderate rainfall so apply soil nutrients @ 25-35 kg/ha as maximum nutrient losses may occur due to rainfall.
Animal Sciences and livestock management			
Cattle			<ul style="list-style-type: none"> ◆ Provide plenty of cold clean drinking water to your livestock at the time between 01 to 3 pm during the peak summer months. ◆ Maintain hygienic condition of animal shed and disinfect the animal shed with 10% formalin solution.
Poultry		Coccidiosis, Ranikhet disease	<ul style="list-style-type: none"> ◆ Provide proper ventilation to poultry birds to prevent from heat stress during summer months ◆ Periodic supplementation of vitamins and minerals to your livestock is very much essential during hot weather. ◆ Special care should be taken for mite and Gastro Intestinal tape worm in case of poultry during the hot and humid summer months. ◆ Take preventive measures against Coccidiosis of poultry during summer months.
Pig	All stages	Swine fever	<ul style="list-style-type: none"> ◆ Provide creep area (separate enclosure, where the piglets have access but the sow does not) with heat source (200 watt bulb) to reduce piglet mortality due to cold weather.

Agro-Met Advisory member of ICAR-NOFRI and KVK- East Sikkim

Name	Designation	Department
<i>Dr. R. K. Avasthe</i>	Joint Director	Soil Science
<i>Dr. Raghavendra Singh</i>	Senior scientist and Head (I/C)	Agronomy
<i>Dr. P. K. Pathak</i>	SMS	Animal science
<i>Dr. J.K. Singh</i>	SMS	Plant Breeding
<i>Dr. Manoj Kumar</i>	SMS	Agronomy
<i>Sh. Boniface Lepcha</i>	SMS	Horticulture
<i>Sh. Santanu Das</i>	SMS	Agril. Meteorology
<i>Miss Tirtha Kumari Chettri</i>	Met. Observer	Biochemistry




AGRO-MET ADVISORY BULLETIN FOR THE DISTRICT OF EAST SIKKIM

PERIOD:

05th OCTOBER to 09th OCTOBER' 2019

*IMD, GANGTOK IN COLLABORATION WITH ICAR, GANGTOK, FSAD
AND HCCD, GANGTOK*



ISSUED ON 4th October 2019

Meteorological Sub-division of Sikkim

As per India Meteorological Department's classification, Sikkim is a part of Met Sub-Division of Sub Himalayan West Bengal & Sikkim (SHWB & SKM) and is divided into four (4) Districts:

- 1) North District with HQ at Mangan
- 2) East District with HQ at Gangtok
- 3) West District with HQ at Gyalsing
- 4) South District with HQ at Namchi

Issued by: Santanu Das, SMS (Agril. Meteorology), ICAR-KVK, East Sikkim, Ranipool

Value Added forecast for EAST-DISTRICT for next five days

Parameter/Date	05/10/2019	06/10/2019	07/10/2019	08/10/2019	09/10/2019
Rainfall (mm)	Very light rainfall	Very light rainfall	Light rainfall	Light rainfall	Light rainfall
Max Temp (°C)	20	21	21	20	19
Min Temp (°C)	16	16	16	15	15
Sky condition	Generally cloudy	Generally cloudy	Generally cloudy	Generally cloudy	Overcast
Max RH (%)	93	92	92	94	95
Min RH (%)	83	82	82	85	86
Wind Speed (Kmph)	004	005	006	006	005
Wind Direction (deg)	NE	NE	NE	NE	NE

Northerly- **N**, North-Easterly- **N-E**, Easterly- **E**, South-Easterly- **S-E**, Southerly- **S**, South-Westerly- **S-W**, Westerly-**W**, North-westerly- **N-W**.

Weather for preceding week			
Date	Rainfall	Max. Temp	Min. Temp
01/10/2019	001	20.8	14.5
02/10/2019	031	17.6	15.6
03/10/2019	006	18.6	15.6

Weather forecast valid up to 09th October 2019

- Probability of light rainfall
- Max temp may be varying from 19-21°C, whereas, Min temp may vary from 15-16 °C
- Sky will remain generally cloudy
- Max RH is likely to be 92-95 %, whereas, Min RH may vary from 82-86 %
- Wind speed may likely vary from 4-6 km/hr.
- Wind direction may blow from northeasterly
- Weekly cumulative rainfall= 21 mm

AGRO-METEREOLOGICAL ADVISORIES


General Advisories for all the crops/ livestock

- ➡ If there is a probability of light rainfall, farmers are advised to keep drainage facilities on their field and harvest all the matured crops.
- ➡ During high humidity and cloudy weather there may be infestation of insect pest, so the farmers are suggested to apply neem oil.
- ➡ Bunds to be provide to reduce nutrient lost due to high rainfall.
- ➡ Rain water harvesting through *jalkund* should be done to conserve excess rain water during high rainfall.
- ➡ Excess water stagnation may cause fungal and bacterial diseases to the crops.
- ➡ While selecting site for construction of shed a site that allows sun light exposure as well as good wind control is preferable.

Name of the crop/Animal	Stage	Cultural practices/ Pest/ Diseases	Agro-Met Advisories
<i>Agricultural and Horticultural crops:</i>			
Rice	Tillering		<ul style="list-style-type: none">◆ Maintain 5 cm of water in the paddy field.◆ First weeding should be completed at 30-35 days after transplanting.
Black gram (Pahelo dal) /Rjmash/Cowpe a	Vegetative		<ul style="list-style-type: none">◆ Farmers are advised to provide proper drainage in the field to avoid wilting in high rainfall areas.◆ First weeding should be completed after 25-30 days of sowing to avoid weed competition.
Ginger/ Turmeric	Tillering stage		<ul style="list-style-type: none">◆ Allow minimum entry to crop field to reduce disease infestation.◆ Allow proper drainage to avoid stagnation of water in the field.◆ Timely monitor the field to check the attack of bacterial wilt due to high humidity and temperature.
Red cherry pepper/Brinjal/ Toamto	Fruiting	Blight	<ul style="list-style-type: none">◆ Allow proper drainage to remove excess rain water from the field.◆ Timely monitor the field to check the attack of bacterial will due to high humidity and temperature.◆ Due to high temperature and high

			rainfall may cause late blight. Apply COC @ 0.25% at weekly interval for 4-5 sprays will reduce the disease.
Cole crops	Nursery preparation		<ul style="list-style-type: none"> ◆ The nursery site should be dry and free from water stagnation as well as high moisture in the soil. ◆ To protect from heavy rain the nursery should be in semi protected area.
Sikkim Mandarin	Planting/fruiting		<ul style="list-style-type: none"> ◆ As there is a possibility of light to moderate rainfall so apply soil nutrients @ 25-35 kg/ha as maximum nutrient losses may occur due to rainfall.

Animal Sciences and livestock management

Cattle			<ul style="list-style-type: none"> ◆ Provide plenty of cold clean drinking water to your livestock at the time between 01 to 3 pm during the peak summer months. ◆ Maintain hygienic condition of animal shed and disinfect the animal shed with 10% formalin solution.
Poultry		Coccidiosis, Ranikhet disease	<ul style="list-style-type: none"> ◆ Provide proper ventilation to poultry birds to prevent from heat stress during summer months ◆ Periodic supplementation of vitamins and minerals to your livestock is very much essential during hot weather. ◆ Special care should be taken for mite and Gastro Intestinal tape worm in case of poultry during the hot and humid summer months. ◆ Take preventive measures against Coccidiosis of poultry during summer months.
Pig	All stages	Swine fever	<ul style="list-style-type: none"> ◆ Provide creep area (separate enclosure, where the piglets have access but the sow does not) with heat source (200 watt bulb) to reduce piglet mortality due to cold weather.

Agro-Met Advisory member of ICAR-NOFRI and KVK- East Sikkim

Name	Designation	Department
<i>Dr. R. K. Avasthe</i>	Joint Director	Soil Science
<i>Dr. Raghavendra Singh</i>	Senior scientist and Head (I/C)	Agronomy
<i>Dr. P. K. Pathak</i>	SMS	Animal science
<i>Dr. J.K. Singh</i>	SMS	Plant Breeding
<i>Dr. Manoj Kumar</i>	SMS	Agronomy
<i>Sh. Boniface Lepcha</i>	SMS	Horticulture
<i>Sh. Santanu Das</i>	SMS	Agril. Meteorology
<i>Miss Tirtha Kumari Chettri</i>	Met. Observer	Biochemistry




AGRO-MET ADVISORY BULLETIN FOR THE DISTRICT OF SOUTH SIKKIM

PERIOD:

05th OCTOBER to 09th OCTOBER' 2019



*IMD, GANGTOK IN COLLABORATION WITH ICAR, GANGTOK, FSAD AND HCCD,
GANGTOK*



ISSUED ON 09th October 2019

Meteorological Sub-division of Sikkim

As per India Meteorological Department's classification, Sikkim is a part of Met Sub-Division of Sub Himalayan West Bengal & Sikkim (SHWB & SKM) and is divided into four (4) Districts:

- 1) North District with HQ at Mangan
- 2) East District with HQ at Gangtok
- 3) West District with HQ at Gyalsing
- 4) South District with HQ at Namchi

Issued by: Santanu Das, SMS (Agril. Meteorology), ICAR-KVK, East Sikkim, Ranipool

Value Added forecast for SOUTH-DISTRICT for next five days

Parameter/Date	05/10/2019	06/10/2019	07/10/2019	08/10/2019	09/10/2019
Rainfall (mm)	Light rainfall	Light rainfall	Light rainfall	Light rainfall	Light rainfall
Max Temp (°C)	24	25	24	23	22
Min Temp (°C)	16	17	17	16	16
Sky condition	Generally cloudy	Generally cloudy	Generally cloudy	Generally cloudy	Generally cloudy
Max RH (%)	92	90	91	92	93
Min RH (%)	80	78	78	80	83
Wind Speed (Kmph)	003	004	005	006	005
Wind Direction (deg)	NE	N	N	N	N

Northerly- **N**, North-Easterly- **N-E**, Easterly- **E**, South-Easterly- **S-E**, Southerly- **S**, South-Westerly- **S-W**, Westerly-**W**, North-westerly- **N-W**.

Weather for preceding week			
Date	Rainfall	Max. Temp	Min. Temp
	(mm)	(°C)	(°C)
01/10/2019	N/A	25.8	14.6
02/10/2019	N/A	23.2	16.9
03/10/2019	N/A	23.8	16.1

Weather forecast valid up to 09th October 2019

- There will be chances of light rainfall
- Max temp may be varying from 22-25°C, whereas, Min temp may vary from 16-17°C
- Sky will remain generally cloudy
- Max RH is likely to be 90-92 %, whereas, Min RH may vary from 78-83 %
- Wind speed may likely vary from 3-6 km/hr.
- Wind direction may blow from northerly direction
- Weekly cumulative rainfall= 26 mm

AGRO-METEREOLOGICAL ADVISORIES


General Advisories for all the crops/ livestock

- ➡ If there is a probability of light rainfall, farmers are advised to keep drainage facilities on their field and harvest all the matured crops.
- ➡ During high humidity and cloudy weather there may be infestation of insect pest, so the farmers are suggested to apply neem oil.
- ➡ Bunds to be provide to reduce nutrient lost due to high rainfall.
- ➡ Rain water harvesting through *jalkund* should be done to conserve excess rain water during high rainfall.
- ➡ Excess water stagnation may cause fungal and bacterial diseases to the crops.
- ➡ While selecting site for construction of shed a site that allows sun light exposure as well as good wind control is preferable.

Name of the crop/Animal	Stage	Cultural practices/ Pest/ Diseases	Agro-Met Advisories
<i>Agricultural and Horticultural crops:</i>			
Rice	Tillering		<ul style="list-style-type: none">◆ Maintain 5 cm of water in the paddy field.◆ First weeding should be completed at 30-35 days after transplanting.
Black gram (Pahelo dal) /Rjmash/Cowpe a	Vegetative		<ul style="list-style-type: none">◆ Farmers are advised to provide proper drainage in the field to avoid wilting in high rainfall areas.◆ First weeding should be completed after 25-30 days of sowing to avoid weed competition.
Ginger/ Turmeric	Tillering stage		<ul style="list-style-type: none">◆ Allow minimum entry to crop field to reduce disease infestation.◆ Allow proper drainage to avoid stagnation of water in the field.◆ Timely monitor the field to check the attack of bacterial wilt due to high humidity and temperature.
Red cherry pepper/Brinjal/ Toamto	Fruiting	Blight	<ul style="list-style-type: none">◆ Allow proper drainage to remove excess rain water from the field.◆ Timely monitor the field to check the attack of bacterial will due to high humidity and temperature.◆ Due to high temperature and high

			rainfall may cause late blight. Apply COC @ 0.25% at weekly interval for 4-5 sprays will reduce the disease.
Cole crops	Nursery preparation		<ul style="list-style-type: none"> ◆ The nursery site should be dry and free from water stagnation as well as high moisture in the soil. ◆ To protect from heavy rain the nursery should be in semi protected area.
Sikkim Mandarin	Planting/fruiting		<ul style="list-style-type: none"> ◆ As there is a possibility of light to moderate rainfall so apply soil nutrients @ 25-35 kg/ha as maximum nutrient losses may occur due to rainfall.

Animal Sciences and livestock management

Cattle			<ul style="list-style-type: none"> ◆ Provide plenty of cold clean drinking water to your livestock at the time between 01 to 3 pm during the peak summer months. ◆ Maintain hygienic condition of animal shed and disinfect the animal shed with 10% formalin solution.
Poultry		Coccidiosis, Ranikhet disease	<ul style="list-style-type: none"> ◆ Provide proper ventilation to poultry birds to prevent from heat stress during summer months ◆ Periodic supplementation of vitamins and minerals to your livestock is very much essential during hot weather. ◆ Special care should be taken for mite and Gastro Intestinal tape worm in case of poultry during the hot and humid summer months. ◆ Take preventive measures against Coccidiosis of poultry during summer months.
Pig	All stages	Swine fever	<ul style="list-style-type: none"> ◆ Provide creep area (separate enclosure, where the piglets have access but the sow does not) with heat source (200 watt bulb) to reduce piglet mortality due to cold weather.

Agro-Met Advisory member of ICAR-NOFRI and KVK- East Sikkim

Name	Designation	Department
<i>Dr. R. K. Avasthe</i>	Joint Director	Soil Science
<i>Dr. Raghavendra Singh</i>	Senior scientist and Head (I/C)	Agronomy
<i>Dr. P. K. Pathak</i>	SMS	Animal science
<i>Dr. J.K. Singh</i>	SMS	Plant Breeding
<i>Dr. Manoj Kumar</i>	SMS	Agronomy
<i>Sh. Boniface Lepcha</i>	SMS	Horticulture
<i>Sh. Santanu Das</i>	SMS	Agril. Meteorology
<i>Miss Tirtha Kumari Chettri</i>	Met. Observer	Biochemistry




AGRO-MET ADVISORY BULLETIN FOR THE DISTRICT OF WEST SIKKIM

PERIOD:

05th OCTOBER to 09th OCTOBER' 2019

*IMD, GANGTOK IN COLLABORATION WITH ICAR, GANGTOK, FSAD
AND HCCD, GANGTOK*



ISSUED ON 09th October 2019

Meteorological Sub-division of Sikkim

As per India Meteorological Department's classification, Sikkim is a part of Met Sub-Division of Sub Himalayan West Bengal & Sikkim (SHWB & SKM) and is divided into four (4) Districts:

- 1) North District with HQ at Mangan
- 2) East District with HQ at Gangtok
- 3) West District with HQ at Gyalsing
- 4) South District with HQ at Namchi

Issued by: Santanu Das, SMS (Agril. Meteorology), ICAR-KVK, East Sikkim, Ranipool

Value Added forecast for WEST-DISTRICT for next five days

Parameter/Date	05/10/2019	06/10/2019	07/10/2019	08/10/2019	09/10/2019
Rainfall (mm)	Very light rainfall	Very light rainfall	Light rainfall	Light rainfall	Light rainfall
Max Temp (°C)	23	24	24	23	22
Min Temp (°C)	16	17	17	16	15
Sky condition	Generally cloudy	Generally cloudy	Generally cloudy	Generally cloudy	Generally cloudy
Max RH (%)	92	91	92	93	93
Min RH (%)	82	81	82	85	85
Wind Speed (Kmph)	006	006	006	007	007
Wind Direction (deg)	N	NW	N	N	NW

Northerly- **N**, North-Easterly- **N-E**, Easterly- **E**, South-Easterly- **S-E**, Southerly- **S**, South-Westerly- **S-W**, Westerly-**W**, North-westerly- **N-W**.

Weather for preceding week			
Date	Rainfall	Max. Temp	Min. Temp
	(mm)	(°C)	(°C)
01/10/2019	N/A	24.7	14.1
02/10/2019	000	19.9	16.1
03/10/2019	009	23.4	15.9

Weather forecast valid up to 09th October 2019


- Probability of light rainfall
- Max temp may be varying from 22-24°C, whereas, Min temp may vary from 15-17°C
- Sky will remain generally cloudy
- Max RH is likely to be 91-93 %, whereas, Min RH may vary from 81-85 %
- Wind speed may likely vary from 6-7 km/hr.
- Wind direction may blow from northerly.
- Weekly cumulative rainfall= 20 mm

AGRO-METEOREOLOGICAL ADVISORIES

General Advisories for all the crops/ livestock (English)

- ➡ If there is a probability of light rainfall, farmers are advised to keep drainage facilities on their field and harvest all the matured crops.
- ➡ During high humidity and cloudy weather there may be infestation of insect pest, so the farmers are suggested to apply neem oil.
- ➡ Bunds to be provide to reduce nutrient lost due to high rainfall.
- ➡ Rain water harvesting through *jalkund* should be done to conserve excess rain water during high rainfall.
- ➡ Excess water stagnation may cause fungal and bacterial diseases to the crops.
- ➡ While selecting site for construction of shed a site that allows sun light exposure as well as good wind control is preferable.

Name of the crop/Animal	Stage	Cultural practices/ Pest/ Diseases	Agro-Met Advisories
<i>Agricultural and Horticultural crops:</i>			
Rice	Tillering		<ul style="list-style-type: none">◆ Maintain 5 cm of water in the paddy field.◆ First weeding should be completed at 30-35 days after transplanting.
Black gram (Pahelo dal) /Rjmash/Cowpe a	Vegetative		<ul style="list-style-type: none">◆ Farmers are advised to provide proper drainage in the field to avoid wilting in high rainfall areas.◆ First weeding should be completed after 25-30 days of sowing to avoid weed competition.
Ginger/ Turmeric	Tillering stage		<ul style="list-style-type: none">◆ Allow minimum entry to crop field to reduce disease infestation.◆ Allow proper drainage to avoid stagnation of water in the field.◆ Timely monitor the field to check the attack of bacterial wilt due to high humidity and temperature.
Red cherry pepper/Brinjal/ Toamto	Fruiting	Blight	<ul style="list-style-type: none">◆ Allow proper drainage to remove excess rain water from the field.◆ Timely monitor the field to check the attack of bacterial will due to high humidity and temperature.◆ Due to high temperature and high

			rainfall may cause late blight. Apply COC @ 0.25% at weekly interval for 4-5 sprays will reduce the disease.
Cole crops	Nursery preparation		<ul style="list-style-type: none"> ◆ The nursery site should be dry and free from water stagnation as well as high moisture in the soil. ◆ To protect from heavy rain the nursery should be in semi protected area.
Sikkim Mandarin	Planting/fruiting		<ul style="list-style-type: none"> ◆ As there is a possibility of light to moderate rainfall so apply soil nutrients @ 25-35 kg/ha as maximum nutrient losses may occur due to rainfall.
Animal Sciences and livestock management			
Cattle			<ul style="list-style-type: none"> ◆ Provide plenty of cold clean drinking water to your livestock at the time between 01 to 3 pm during the peak summer months. ◆ Maintain hygienic condition of animal shed and disinfect the animal shed with 10% formalin solution.
Poultry		Coccidiosis, Ranikhet disease	<ul style="list-style-type: none"> ◆ Provide proper ventilation to poultry birds to prevent from heat stress during summer months ◆ Periodic supplementation of vitamins and minerals to your livestock is very much essential during hot weather. ◆ Special care should be taken for mite and Gastro Intestinal tape worm in case of poultry during the hot and humid summer months. ◆ Take preventive measures against Coccidiosis of poultry during summer months.
Pig	All stages	Swine fever	<ul style="list-style-type: none"> ◆ Provide creep area (separate enclosure, where the piglets have access but the sow does not) with heat source (200 watt bulb) to reduce piglet mortality due to cold weather.

Agro-Met Advisory member of ICAR-NOFRI and KVK- East Sikkim

Name	Designation	Department
<i>Dr. R. K. Avasthe</i>	Joint Director	Soil Science
<i>Dr. Raghavendra Singh</i>	Senior scientist and Head (I/C)	Agronomy
<i>Dr. P. K. Pathak</i>	SMS	Animal science
<i>Dr. J.K. Singh</i>	SMS	Plant Breeding
<i>Dr. Manoj Kumar</i>	SMS	Agronomy
<i>Sh. Boniface Lepcha</i>	SMS	Horticulture
<i>Sh. Santanu Das</i>	SMS	Agril. Meteorology
<i>Miss Tirtha Kumari Chettri</i>	Met. Observer	Biochemistry